

This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

### Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + Refrain from automated querying Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + Keep it legal Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

#### **About Google Book Search**

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <a href="http://books.google.com/">http://books.google.com/</a>

J. Hill 1991

THE

# ALPHEUS FELCH HISTORICAL LIBRARY

**BEQUEATHED** 

TO THE

UNIVERSITY OF MICHIGAN

BY THE

HON. ALPHEUS FELCH.

1896.

151.3 .L68 1778 32 Indoneny. gal 8

# New Dispensatory:

CONTAINING,

I.

The ELEMENTS OF PHARMACY.

II. 60/33

The MATERIA MEDICA:

O R.

An Account of the Substances employed in MEDICINE; with the Virtues and Uses of each Article, so far as they are warranted by Experience and Observation.

III.

The PREPARATIONS and Compositions of the

3

New London and Edinburgh Pharmacopæias;

#### WITH

Such of the old ones as are kept in the Shops; the most celebrated Foreign MEDICINES; the most useful of those directed in the Hospitals; sundry elegant extemporaneous Forms, &c. digested in such a Method as to compose a regular System of PHARMACY; with Remarks on their Preparation and Uses; the Means of distinguishing Adulterations; of performing the more difficult and dangerous Processes with Ease and Sasety, &c.

The Whole interspersed with

Practical Cautions and Observations.

By WILLIAM LEWIS, M. B. F. R. S.

The FOURTH EDITION, corrected, with large Additions.

D U B L I N:

Printed by JAMES POTTS, at No. 74, in DAME-STREET.

M DCC LXXVIII.

• •

# PREFACE.

HE New Dispensatory was intended as a regular book of practical and scientific pharmacy; composed on principles agreeable to those, on which the colleges of London and Edinburgh have proceeded, in the late reformation of their officinal pharmacoposias; containing full and clear directions, drawn from actual experience, for the preparation of the several medicines, particularly where accompanied with any difficulty or danger; and affigning every-where, as far as possible, their real virtues and uses; intentions, which though of primary importance in a work of this kind, do not seem to have been at all segarded in the other dispensatories that have hitherto appeared.

The author has had the satisfaction of finding that his endeavours have not been in vain; that though the work fell very short of the perfection which he wished for, it was distinguished with approbations even beyond his hopes; with approbations, which have induced the compilers of the other dispensatories to borrow very considerable parts of it in their last editions, in one of which, besides many paragraphs and entire pages here and there, the greatest part of two hundred pages together is illiberally copied from this work.

In this fourth edition, I have made many material corrections and additions; and retrenched fundry exceptionable particulars, which, in compliance with common prejudices, had been admitted in the first attempt.

THE first part contains the Elements of Pharmacy, or what is commonly called Pharmaceutical Chemistry. general neglect of this interesting and useful study, as applied to medicinal subjects, has engaged me to greatly enlarge this part, and to labour it with more care and precision. I have endeavoured to give a concile and systematic view of the general properties and relations of vegetable, animal, and mineral bodies; the different medicinal principles they contain; the means of extracting and fe--parating their native component parts, without making any alteration in their qualities; and the different forms and powers which they assume, from different natural or artificial operations, or from the mixture and coalition of one with another; avoiding every-where all hypothetical reasonings, and delivering only the direct result of expe-To this history is added a pracriment and observation. tical account of the instruments and operations of the art, which, it is hoped, will give the reader a full idea of them. without the tediousness of minute details.

The next part contains the Materia Medica, or medicinal simples, which, for reasons assigned in the introduction to this part, are all ranged in alphabetic order. Rationales of the operations of medicines, which are at best but conjectural and unsatisfactory, have no place in this practical work: but some general observations, of the sensible effects of certain classes of medicines, in Cartheuser's manner, it has been thought expedient to retain, with some amendments from the former editions.

In treating of the several simples themselves, I have given, where necessary, a description of the simple, with the marks of its genuineness and goodness; and pointed out the

the diffinguishing characters of such, as, from a resemblance in external appearance, are liable to be confounded with others of different qualities. With regard to their virtues, particular care has been taken to reject the fabulous ones, which are still preserved in other books of this kind; and to give only those, which have either been confirmed by repeated experience, or may be rationally inferred from the sensible qualities of the subject, or from its agreement in smell, taste, &c. with others of known virtue. Under each simple are mentioned all the preparations made from it, and all the compositions in which it is an ingredient, in the London and Edinburgh pharmacopæias. Many of the capital articles I have examined pharmaceutically. and shewn in what separable part of the mixed its virtue resides, by what means the active principle is best extracted or preserved, and in what form the substance itself or its preparations are most commodiously and advantageously exhibited. At the end of this part, the directions for the collection and preservation of medicinal substances are reconfidered.

THE third and fourth parts contain the preparations and compositions of the new London and Edinburgh pharmacopæias, with a few of the old ones, which I am informed are still kept in some shops, and occasionally called for; several of the more celebrated medicines which have come into esteem in France and Germany; many from our hospitals; and some elegant extemporaneous prescriptions, such as are directed in practice.

In the distribution of these materials, it has been found necessary to depart from the order hitherto received. In other dispensatories, and in the former edition of this, medicines are divided into two general heads, officinal and extemporaneous. This division is apparently faulty; for many of those called officinal are strictly extemporaneous, being made only as they are wanted; and many of those, which are called extemporaneous, are very well sitted for keeping: if we should appropriate the term officinal to those

THE Author is sufficiently sensible, that there are still many impersections in this performance; but hopes it will appear, that he has every where consulted the dignity of the art, the ease and advantage of the operator, and the health of the patient.

CONTENTS.

# PART I.

# ELEMENTS of PHARMACY.

CHAP. II. General view of the properties and relations of	age t
medicinal substances	4
Sect. i. Vegetables	ib.
1. Productions from vegetables by fer mentation	<b>*-</b>
	5
2. Productions from vegetables by fire 3. Substances naturally contained in vege	, <b>y</b>
ables, and separable by art without	ut
alteration of their native qualities	
1. Gross oils	ib.
2. Gross sebaceous matter	11
3. Essential oils	ib.
4. Concrete essential oils	12
5. Camphor	ib.
6. Resin	13
8. Gum-resin	14
9. Saline matter	ib.
General observations on the foregoing	
principles	15
Sect. ii. Animals	18
Sect. iii. Minerals	21
Oils and bitumens	ib.
Earths	ib.
Metals	
Acids	23
	24
Of the affinities of bodies	25
Tables of affinities	28
CHAP. II. Of the pharmaceutical apparatus	.30
Furnaces	ib.
Baths	33
Coating of glasses, and lutes	34
Veffels	35
Weights and measures	36
Table of the weight of different fluids	_ 37
	CHAP.

CHAP. III.	Of the pharmaceutical operations		p. 39
	Sect. i.	Solution -	ib.
		Tables of the solubility of salts	39-41
,	Sect. ii.	Extraction	43
	Sect. iii.	Depuration	45
	Sect. iv.	Crystallization	46
	Sect. v.	Precipitation	47
	Sect. vi.	Evaporation .	48
	Sect. vii.	Distillation	49
•	Se&. viii.		51
	Sect. ix.		52
	Sect. x.	Exficcation	ib.
	Sect. xi.	Comminution	53
	Sect. xii.	Fusion	55
	Sect. xiii.	Calcination, &c.	55 56

# PART II.

# The MATERIA MEDICA.

General observations on the materia	medica	9
Acids		ě
Absorbent earths	i	6
Tables of the powers of dif	ferent absorbents	. 6
Indissoluble earths	Gibbones	ě
Glutinou and mucilaginous	Juojianies	
Un uous and oily substance	S	6
Astringents		il
Sweets		6
Acrids		. 6
Aromatics		i
Bitters .	•	• 7
Emetics and cathartics		i
Account of the particular simples, r	anged in alphabetic	order ;
General titles including several simp	les	24
Seneral littles including feed at finish	F Gambles	
Of the collection and preservation of	Journies	25

# PART III.

A fystem of pharmaceutical preparations, officinal and extemporaneous.

CHAP. I. The more simple preparations

25**5** 

CHAP.	II. Substances	extracted from vegetables by expression	p. 263
	Sect. i.	Juices	ib.
	Se&. ii.	Y 4 44	265
		Essential oils by expression	26 <b>6</b>
CHAP.	III. Infusions.	in different menstrua	268
	Sect. i.	Infusions and decostions in water	ib.
		1. Infusions in cold water	269
		2. Infusions in boiling water	273
		3. Decoctions	278
	Se&. ii.	Wheys	288
	Se&. iii.	Vinegars	289
	Se&. iv.	Medicated wines	29 t
	Sea. v.		301
	Se&. vi.	Spirituous tinctures, elixirs, esfenc	es,
		balfams	303
_	Sect. vii.	Oils by infusion and decostion	. 330
CHAP.	IV. Conservat.	ion of recent vegetables, &c. by sugar	and
	boney		333
	Sect. i.	Conserves	ib.
	Sect. ii.	Preferves	336
	Sect. iii.	Gellies	33 <b>7</b>
	Sea. iv.	Syrups	338
C	Sect. v.	Honeys and oxymels	349
CHAP.	v . separation	and collection of those parts of veget	avie
	ana an	nimal substances which are volatile in f boiling water	
	Sect. i.		<b>3</b> 5 <b>3</b>
	Sect. ii.		ib.
	Se&t. iii.	Distilled spirits	370
	Sect. iv.		38 t 38 <b>9</b>
Снар:		tion of the medicinal parts of juices	20 <b>9</b>
~		ns by evaporation	
	Sect. i.		39 <b>9</b> ib.
		Extracts with water	403
	Sect. iii.		ntial 403
		extracts, resins	409
	Se&. iv.	Extracts with water and spirit	412
	Sect. v.	Extracts by long digestion	415
CHAP.	VII. Empyreus		418
		ion of empyreumatic oils	421
CHAP.	VIII. Salts and	Saline preparations	423
	Sect. i.	Fixed alkaline salts	ib.
	Sect. ii.	Volatile alkaline salts and spirits	420
	Sect. iii.	Combination of alkalies with oils and	d in-
		flammable spirits	436
	Sect. iv.	Acid spirits	443
			Cuan

CHAP. VIII	Sect. v.	Combination of acid with vinous spir	its p. 452
	Sect. vi.	Neutral and compound salts	458
		Anomalous salts	473
CHAP. IX.		ons of sulpbur	484
		reparations	489
		Preparations of gold	ib,
		Preparations of silver	490
		Preparations of iron	492
		Preparations of copper	498
		Preparations of lead	500
•		Preparations of tin	503
		Preparations of mercury	507
		Preparations of antimony	525
		Preparations of bifmuth	541
		Prepa ations of zinc	542
		Compound metallic preparations	54 <b>3</b>

# PART IV.

# MEDICINAL COMPOSITIONS, officinal and extemporaneous.

	_	٠,
CHAP. I.	Powders	547
	Troches and lozenges	560
CHAP. III.		<u>5</u> 67.
CHAP. IV.		579
CHAP. V.		585
CHAP. VI.		603
CHAP. VII.	Emulsions	605
CHAP. VIII.	Juleps, mixtures, draughts	609
CHAP. IX.	Lotions, gargarisms, injections, &c.	619
CHAP. X.	Plasters	624
CHAP. XI.	Ointments, liniments, and cerates	634
CHAP. XII.	Epithems .	648

The useful compositions of the Pharmacopæia pauperum, which made a distinct part in the former edition, are here distributed in the two foregoing parts; all the medicines of similar forms being now, for the convenience of the reader, placed together.

## THE

# NEW DISPENSATORY.

# PART L

ELEMENTS OF PHARMACY.

# CHAPTER L

Definition and Division of PHARMACY.

PHARMACY is the art of preparing and compounding natural and artificial substances for medicinal purposes, in a manner suitable to their respective properties, and the intentions of cure:

THIS art has been commonly divided into two branches, GALENICAL and CHEMICAL; but no rational principle of diffinction between them has as yet been fixed on. If it be a chemical process to evaporate juice of plantane over a gentle fire till it becomes thick, it is surely not less chemical to evaporate the juice of sloes in the same manner; and yet the former only is ranked among the chemical, and the latter among the galenical preparations. Prequently, also, one and the same preparation is, in different Pharmacopæias, referred to the different branches: thus, distilled waters and distilled spirits, which make the first of the galenical articles in one Pharmacopæia, make the first of the chemical in another.

It is agreed on both fides, that effential oils, extracts, refins, volatile and fixed falts, the artificial neutral falts, metallic preparations, and other like productions, belong to the chemical pharmacy; and pills, boluses, troches, electaries, draughts, ointments, plasters, poultices, &c. to the galenical; as if the distinction was founded, neither neither on the nature of the operation, nor of the materials, nor of the effect produced, but merely on the form, in which the medicine is intended to be taken or applied. Thus, the diffolution of mercury in aquafortis is ranked among the chemical preparations; while the very same process, with the additional circumstance of uniting an unctuous material, which renders it, if any thing, still more chemical, is nevertheless reckoned a galenical one, because the product is used as an ointment. It cannot surely be supposed, that this is a just division; or that the same process or preparation can become chemical or not chemical, according to the intention to which they are applied, or the form in which the product is used.

If vitriol of iron (that is, iron united with a certain acid) and any volatile alkaline salt, as that of hartshorn or sal ammoniac, be put together into water, in due proportions; the pungent smell of the volatile falt will be immediately suppressed, this falt uniting with the acid of the vitriol into a new compound, while the iron This is undoubtedly a chemical is separated and thrown out. effect; and this effect will happen, wherever these two ingredients meet together in a moist state, whatever the form of the medicine be. It is obvious, therefore, that the Galenical forms are by no means independent of chemistry; and that this science extends to mixtures of the most simple kind.

The London college has very judiciously rejected this division; a division, apparently derived from prejudice and superficial knowledge, and which has been continued only in compliance with cuf-Pharmacy, in its full extent, is no other than a branch of chemistry; and the most simple pharmaceutical preparations are so far chemical, as they have any dependence upon the properties or re-

lations of the materials.

PHARMACY, according to our definition, may be divided into THEORETICAL and PRACTICAL. The former teaches the knowledge of the medicinal substances themselves, their various properties, qualities, and relations to one another, and their general effects on the human body: the latter, the skilful performance of the several processes, or operations, by which they are adapted to particular uses.

What is here called theory is not to be understood as consisting of speculative truths, or philosophical investigations, calculated for explaining the phenomena, or teaching the rationale of the effects pro-The theory of pharmacy is the direct result of experiment and observation, or rather a general and comprehensive view of experiments and facts themselves. It may be termed SCIENTIFIC PHAR-MACY, in distinction from mere manual labour.

Scientific pharmacy includes all those facts which relate to-the reduction of medicinal substances into different forms, and the forms in which particular substances are most commodiously or advantageoufly advantageously used;—their relations to one another in regard to missibility, and the means by which those, that of themselves are not missible, may be made to unite;—the separation of the medicinal from the inactive matter, and of different kinds of medicinal matter from one another when combined together in the same subject, on the principle of one being dissoluble in liquors which will not dissolve the other; of one being exhalable by heat while the other remains fixed, &c.—the alterations which the medicinal parts themselves undergo, in different circumstances, and by different themselves undergo, in the production of new properties and medicinal powers from the coalition of diffimilar things;—with many other particulars analagous to these.

It is obvious, that a perfect acquaintance with pharmacy, confidered in this light, is effentially necessary to the due exercise of the art of physick. Without it, the prescriber must often err, in the choice of materials for the different forms of preparation or composition, or in adapting a manner of preparation to given materials; and often be deceived also in the medicinal effects, which the known

powers of the ingredients, separately, gave room to expect.

It would be inconsistent with the nature of a Dispensatory, to wholly detach the scientistic part of Pharmacy from that which is more directly practical; for the science gradually results in the course of the practical details. In this first part of the work it has been thought expedient to premise a summary view of the general elements of the art, both practical and scientists, that the reader may be the better prepared for the particular subjects and processes which sollow in the second and third parts.

CHAP.

## CHAPTER IL

A general view of the properties and relations of medicinal substances.

## SECT. I.

## Vegetables.

VEGETABLES are organized bodies, containing, in certain vessels, different kinds of substances, in which their medicinal virtues consist, and which are found so differ greatly, not only in their quantity, but likewise in their quality, according to the age of the plant, the season of the year, and the soil in which it is

produced.

Thus some herbs in their infancy abound most with odoriferous matter; of which others yield little or none till they have attained to a more advanced age. Many fruits, in their immature state. contain an austere acid juice, which by maturation is changed into a fweet: others, as the orange, are first warm and aromatic, and afterwards, by degrees, become filled with a strong acid. common grain, and fundry other feeds, when beginning to vegetate, are in taste remarkably sweet; yet the kernels of certain fruits prove, at the same period, extremely acid. The roots of some of our indigenous plants, whose juice is, during the summer, thin and watery, if wounded early in the spring, yield rich balsamic juices, which, exposed to a gentle warmth, soon concrete into solid gummy-refins, superior to many of those brought from abroad. In open exposures, dry soils, and fair warm seasons, aromatic plants prove stronger and more fragrant, and setid ones weaker in smell, than in the opposite circumstances. To these particulars, therefore, due regard ought to be had in the collecting of plants for medicinal ules.

It may be proper to observe also, that the different parts of one plant are often very different in quality from one another. Thus the bitter herb wormwood rises from an aromatic root; and the narcotic poppy-head includes seeds which have no narcotic power. These differences, though very obvious in the common culinary plants, do not seem to have been sufficiently observed, or attended to in the medicinal ones.

The medicinal juices of vegetables, and the active parts with which they are impregnated, may, generally, be extracted and fepa-

rated, by simple operations, without any alteration being made in their native qualities. They may, likewife, be variously altered and transformed, by operations not less simple. By sermentation and the power of fire vegetables, and all the substances that exist in them, (the pure watery part excepted) totally change their acture, and are converted or resolved into products of another order. It will be proper to take a view of these productions first, some of them being substantiant to the separation of the native principles, and to the better-understanding of their properties.

### 1. Productions from vegetables by fermentation.

THE sweet and ascescent jaices of fruits, infesions of malted grain, and almost all vegetable juices or insusions that are either simply sweet or of a sweetness mixed with acidity, on being kept in a place of temperate warmth, in a vessel not closely stopt, ferment, grow turbid, throw off a large quantity of gross matter, and are converted by degrees into a vinous liquous; from which may be separated, by processes hereaster described, a pure implementable spirit.

It is needless to observe how different these productions are, in their medicinal as well as their more obvious properties, from the liquors that afforded them. The native juices of fruit attenuate the animal fluids, and relax the solids, so as to prove in some cases useful aperient medicines, and to occasion, when imprudently taken, dangerous fluxes; whereas the vinous and spirituous liquors, produced from them by fermentation, have the opposite effects, constringing the solids, and thickening or coagulating the sluids.

In vinous liquors there are great diversities, independently of their being more or less watery; for some of the native qualities of vegetable juices and infusions, as colour, flavour, viscidity, &c. often remain in the wine, not being totally subduable by that degree of fermentation by which the liquor is rendered vinous: but of these diversities the spirit is never found to paraske: this, separated from the wine and properly purified, is always one and the same thing, whatever kind of vegetable liquor it was produced from.

Besides the gross matter thrown off during the fermentation, there separates from sundry wines, after the fermentation is completed, another kind of substance. The sides and bottom of the cask become gradually incrustated with a saline concrete, called TARTAR, of an acid taste, and of a reddish or white colour, according to that of the wine. The colour is adventitious to the salt; for the tartar may be purished from it by solution in water: when thus purished, the tartar

of all wines is found to be the fame.

There is separated also, in fermentation, a substance of a much more active nature than any of the preceding. When the fermentation is at its height, a subtile, pungent, elastic, incoercible va-POUR is discharged; which, when copiously accumulated in close tooms, extinguishes fire, and instantaneously suffocates animals, without producing any apparent disease, or any injury that can be perceived upon diffection. Boerhaave says, he does not remember that so immediate, mortal, and subtle a poison has been hitherto discovered: that if a large vessel, full of the juice of grapes in high fermentation, should discharge its accumulated vapour through a small orifice, and a strong healthy man should draw in the vapour at his nostrils, he would instantly fall down dead; or, if he rereceived but little thereof, become apoplectic; or, if still less, would remain an ideot during life, or become paralytic: and that there accidents befall those, who imprudently remain long in close vaults where large quantities are fermenting. It may be observed, that this vapour, when not collected in such quantity as to extinguish a small flame, at that of a candle, is generally not dangerous, or at leaft not mortal, to animals. ...

There are several substances, of themselves not susceptible of fermentation, which, nevertheless, may be brought into it by the admixture of those that are; as, by adding to them, along with a proper quantity of water, a portion of the yeast or head thrown up to the surface of fermencing liquors. This expedient is sometimes had recourse to; for unlocking the texture of certain compact ve table matters, in order to enable them to give out more readily forme of their medicinal principles. In these cases, the fermentation must be continued but for a little time; lest the resolution of the subject should proceed beyond the intended limits, and the principles, ex-

pected from it. be converted into other products.

The fermentable juices of fruits; boiled till they become thick, are found to be indisposed to ferment, and this not only in their thick state, but when distuted again with water; though there appears to be scarcely any other alteration produced in them by the boiling. Hence liquids, prone to fermentation, may thus be preserved. How far this diminution of their fermentability may affect their medical virtues, is not as yet clear.

THE degree or the species of fermentation, by which vines and inflammable spirits are produced, is called winous fermentation. the process is further protracted, more gross matter is thrown off, and new changes succeed, but in a slower and less tumultuary manner than before. The heating inebriating wine becomes by degrees a cool acid vinggar, which seems to counteract the effects of the other: the more the wine abounded with inflammable spirit, the more does the vinegar abound with uninflammable acid.

Chap. 2

There are, however, certain qualities of vegetables, which are not completely subdued even by this second stage of sermentation; some vinegars being apparently more coloured, and containing more of an oily and viscid matter than others. By adding to the sermentable liquor subjects of other kinds, the qualities both of wines and vinegars may be still further diversified, so as to adapt them to particular medicinal uses.

It is observable, that though the acctous fermentation will always fucceed the vinous, unless industriously prevented, yet it is not always preceded thereby; for many, perhaps all, fermentable liquors may be made to pass to the acctous state, without any intermediate period of true vinosity.

Ir the process is still further continued, further changes take place. The matter putrefies: and at length, what little liquor remains unevaporated, is found to be mere water, and the folid substance at the bottom appears to be the same with common mould,

This is reckoned by the chemists one of the stages of fermentation, and distinguished by the name of the purefactive stage: it is far more general in its object than the other two; every vegetable matter being susceptible of putrefaction, but some particular kinds

only being adapted to vinous or acetous fermentation.

Putrefaction discovers one difference in vegetables, which, though not taken notice of, so far as I know, by any writer, seems worthy of being remarked. The generality of vegetables rot and turn to mould, without yielding any very offensive smell from the beginning to the end of the resolution: but there are which emit, throughout the whole process, a strong setor, very nearly of the same kind with that which accompanies the putrefaction of animal substances.

## 2. Productions from vegetables by fire.

Fire, the other grand agent in the refolution of bodies, produces in vegetables decompositions of a different kind. Its general effects are the following:

VEGETABLE substances, burned in the open air, are reduced partly into Asses, and partly into FLAME and SMOKE; which last, condensed in long canals or otherwise, forms a nauseous bitter black appr. In the burning of most vegetables; an acid vapour accompanies the smoke; but the soot is never found to partake of it.

Vegetables urged with a red heat in close vessels (the vessel containing the subject being made to communicate with another placed, beyond the action of the fire for receiving the matters forced out.

by the heat) give over a WATERT LIQUOR called phiegm; an ACID Liquon called spirit; an elastic incoercible vapour, which appears to be AIR, and to which an exit much be occasionally allowed, left it burft the veffels or blow off the receiver; a thin oil, and at length a very dark thick coloured oil, both which are of an acrimonious taste, and a burnt fetid fmell, whence they are called empyreumatic oils." There remains behind a black COAL, not dissoluble in any kind of liquors, not susceptible of putrefaction, not alterable by the most vehement degree of fire so long as the air is excluded. but which, on admitting air to it, burns without flaming, and with little or no smoke, and leaves a very small quantity of white

The white ashes of vegetables, infused or boiled in water, impart to it a pungent faine substance, called FIXED ACKALINE SALT, which may be separated in a solid form by evaporating the water. The remaining part of the ashes, which is by far the largest in quantity, is a pure BARTH, differing from that which is the result of puttefaction, in being readily dissoluble by every acid liquor, while the other is not afted upon by any acid...

Such is the general analysis of regetables by sire. But there are some vegetables, which, as they seem to shew, during putrefaction, some analogy in their matter with that which constitutes animal bedies, discover also a like analogy in the present resolution, yielding listle or no acid, and, instead of a fixed alkaline fait which remains In the after, affording a volatile alkaline exert which arises along with the aqueous and oily principles.

ALKABINE falts, and acid or four fubftances, are looked upon as being opposite in their nature to one another. Most of the bodies. which are dissoluble in alkaline liquors, are precipitated or thrown out from the foliation on the addition of an acid; and most of those, which are diffoluble in acids, are in like manner precipitated by alkalies. If an acid and an alkali be directly mixed together, there generally enfues an efferveffence or tumultuary discharge of air-bub-bles; though alkalies, both fixed and volatile, may be so prepared as to make no effervescence with acids; and in this case they are far more pungent than in their common state.

In all cases, the alkali and acid, uniting together, compose a new body, called a NEUTRAL SALT, which has neither the fourness of the one ingredient, nor the peculiar pungency of the other, and which will not diffolve those substances which either the acid or the alkali separately would diffolve. separately would dissolve.

To these characters it may be added, that alkaline salts change the colour of blue flowers or their infusions, as of violets to a green, and acids to a red, while the neutral compound, formed by the coalition of the two, makes no alteration in the colour.

It must be observed, however, that to change blue slowers to a green, is not universally a mark of alkalies, for some solutions of earthy bodies in acids have the same effect: these last may be distinguished from alkalies, by adding to them a known alkali, which will immediately precipitate the earth, and form a neutral compound with the acid.

Fixt alkaline falts, perfectly purified, appear to be one and the fame, whatever kind of vegetable they were produced from; those of fome marine plants excepted, of which hereafter. In volatile alkalies, and in the pure earthy part of the askes, there appears to be, respectively, the like identity.

Empyreumatic oils differ somewhat in the degree of acrimony and fetidacis, and the acid spirits differ in degree of strength, or in the quantity of water they are diluted with; how far they may differ in any other respects, is little known, these preparations having been

rarely used or examined.

It may be observed, that the alkaline salts, both of the fixt and of the volatile kind, are entirely creatures of the fire; being never found to exist naturally in any vegetable; the oil, doubtless, præ-existed in the subject, but owes its acrimony and setidness to the fire; for the most mild and insipid oils receive the same qualities on being urged with the same degree of heat; the acid, which is likewise naturally contained in vegetable subjects, proves always tainted, in the present process, with the ill smell and taste of the oil that accompanies it; but whether the acid itself suffers any change in its nature, is unknown.

When chemistry began first to be formed into a rational science, and to examine the compotent parts and internal confliction of bodies. it was imagined, that this resolution of vegetables by are, discovering to us all their active principles, unclorged and unmixed with one another, would afford the furest means of judging of their medicinal powers. But on profecuting these experiments, it was soon found that they were insufficient for that end: that the analyses of poisonous and esculent plants agreed often as nearly with one another as two analyses of one plant: that by the action of a burning heat, the principles of vegetables are not barely separated, but altered, transposed, and combined into new forms; infomuch, that it was impossible to know, what form they existed in, and what qualities they were endowed with, before these changes and transpositions happened. If, for example, thirty-two ounces of a certain vegetable substance are found to yield ten ounces and a half of acid liquor, above one ounce and five drams of oil, and three drams and a half of fixt alkaline falt; what idea can this analysis give of the medicinal qualities of gum Arabic ? : \_:: ... :

3. Substances naturally contained in vegetables, and separable by art without alteration of their native qualities.

#### 1. Gross oils.

GROSS oils abound chiefly in the kernels of fruits and in certain feeds; from which they are commonly extracted by expression, and hence are distinguished by the name of expressed oils. They are contained also in all the parts of all vegetables that have been examined, and may be forced out by vehemence of fire; but here their qualities are greatly altered in the process of which they are extracted or dis-

covered, as we have feen under the foregoing head.

These oils, in their common sase, are not dissoluble either in vinous spirits or in water, though, by means of certain intermedia, they may be united both with one and the other. Thus a skilful interposition of sugar renders them miscible with water into what are called lohochs and oily draughts: by the intervention of sum or mucilage they unite with water into a milky. Suid: by alkaline salts they are changed into a scap, which is miscible both with watery and spirituous liquors, and is perfectly dissolved by the latter into an uniform transparent sluid. The addition of any acid to the scapy solution absorbe the alkaline salt; and the oil, which of course separates, is found to have undergone this remarkable change, that it now dissolves, without any intermedium, in pure spirit of wine.

Expressed oils, exposed to the cold, lose greatly of their studity: some of them, in a small degree of cold, congeal into a consistent mass. Kept for some time in a warm sir, they become thin and highly rancid: their soft, lubricating, and relaxing quality is changed into a sharp acrimonious one: and in this state, instead of allaying, they occasion irritation; instead of obtunding corrosive humours, they corrode and instance. These oils are liable to the same noxious alteration, while contained in the original subject: hence the rancidity which the oily seeds and kernels, as almonds and those called the cold seeds, are so liable to contract in keeping. Nevertheless on triturating these seeds or kernels with water, the oil, by the intervention of the other matter of the subject, unites with the water into an emultion or milky liquor, which, instead of growing rancid, turns sour on standing.

In the beat of boiling water, and even in a degree of heat as much exceeding this as the heat of boiling water does that of the human body, these oils suffer little dissipation of their parts. In a greater heat, they emit a pungent vapour, seemingly of the acid kind; and when suffered to grow cold again, they are found to have acquired a greater degree of consistence than they had before, together with an acrid tasse. In a heat approaching to ignition, in close vessels, greatest

greatest part of the oil arises in an empyreumatic state, a black coal remaining behind.

#### 2. Gross sebaceous matter.

From the kernels of some fruits, as that of the chocolate nut, we obtain, instead of a sluid oil, a substance of a butyraceous confishence; and from others, as the nutmeg, a solid matter as firm as tallow. These concretes are most commodiously extracted by boiling the subject in water; the sebaceous matter, liquested by the heat, separates and arises to the surface, and resumes its proper consistence as the liquor cools.

The substances of this class have the same general properties with expressed oils, but are less disposed to become rancid in keeping than most of the common shuid oils. It is supposed by the chemists, that their thick consistence is owing to a larger admixture of an acid principle: for, in their resolution by sire, they yield a vapour more sensibly acid than the shuid oils; and shuid oils, by the admixture of concentrated acids, are reduced to a thick or solid mass.

#### 3. Essential oils.

ESSENTIAL oils are obtained only from those vegetables, or parts of vegetables, that are considerably odorous. They are the direct principle, in which the odour, and oftentimes the warmth, pungency, and other active powers of the subject, reside; whence their name of essences or essential oils.

Effential oils unite with rectified spirit of wine, and compose with it one homogene transparent fluid; though some of them require for this purpose a much larger proportion of the spirit than others. Water also, though it does not dissolve their whole substance, may be made to imbibe some portion of their more subtile matter, so as to become considerably impregnated with their flavour: by the admixture of sugar, gum, the yolk of an egg, or alkaline salts, they are made totally dissoluble in water. Digested with volatile alkalies, they undergo various changes of colour, and some of the less odorous acquire considerable degrees of fragrance; whilst sixt alkalies universally impair their odour.

In the heat of boiling water, these oils totally exhale; and on this principle they are commonly extracted from subjects that contain them: for no other sluid; that naturally exists in vegetables, is exhalable by that degree of heat, except the aqueous moisture, from which greatest part of the oil is easily separated. Some of these oils arise with a much less heat, a heat little greater than that in which water begins visibly to evaporate. In their resolution by a burning

heat, they differ little from expressed oils.

Effential oils, exposed for some time to a warm air, suffer an alteration very different from that which the expressed undergo.

Infead

Instead of growing thin, rancid, and acrimonious, they gradually become thick, and at length harden into a solid brittle concrete; with a remarkable diminution of their volatility, fragrancy, pungency, and warm stimulating quality. In this state they are found to consist of two kinds of matter; a stuid oil, volatile in the heat of boiling water, and nearly of the same quality with the original oil; and of a grosser substance which remains behind, not exhalable without a burning heat, or such a one as changes its nature, and resolves it into an acid, an empyreumatic oil, and a black coal.

The admixture of a concentrated acid infantly produces, in effectial oils, a change nearly fimilar to that which time effects. In making shele kinds of commixtures, the operator ought to be on his guard: for when a strong acid, particularly that of nitre (of which hereaster) is poured hastily into an essential oil, a great heat and ebullition ensue, and often an explosion happens, or the mixture bursts into stame. The union of expressed oils with acids is accompanied

with much less confict.

#### 4. Concrete essential oil.

Some vegetables, as roses and elecampane roots, instead of a stuid essential oil, yield a substance possessing the same general properties, but of a thick or sebaceous consistence. This substance appears to be of as great volatility, and subtility of parts, as the suid oils: it equally exhales in the heat of boiling water, and concretes upon the surface of the collected vapour. The total exhalation of this matter, and its concreting again into its original consistent state, without any separation of it into a sluid and a solid part, dissinguishes it from essential oils that have been thickened or indurated by age or by acids.

### 5. Camphor.

CAMPHOR is a folid concrete, obtained chiefly from the woody parts of certain Indian trees. It is volatile like effential oils, and foluble both in oils and inflammable spirits: it unites freely with water by the intervention of gum, but very sparingly and imperfectly by the other intermedia that renders oils miscible with watery liquors. It differs from the sebaceous as well as fluid effential oils, in suffering no sensible alteration from long keeping; in being totally exhalable, not only by the heat of boiling water, but in a warm air, without any change or separation of its parts, the last particle that remains unexhaled appearing to be of the same nature with the original camphor: in its receiving no empyreumatic impression, and suffering no refolution, from any degree of fire to which it can be exposed in close reffels, though readily combustible in the open air; it being dissolved by concentrated acids into a liquid form; and in several other properties which it is needless to specify in this place. 6. Refin

#### 6. Refin.

Effectial oils, indurated by age or acids, are called refins. When the indurated mass has been exposed to the heat of boiling water. till its more subtile part, or the pure essential oil that remained in it, has exhaled; the gross.matter, lest behind, is likewise called resin. We find, in many vegetables, refins analogous both to one and the other of these concretes; some containing a subtile oil, separable by a heat of boiling water; others containing nothing that is capable of

exhaling in that heat.

Refins in general dissolve in reclified spirit of wine, though some of them much more difficultly than others: it is chiefly by means of this dissolvent, that they are extracted from the subjects in which they are contained. They dissolve also in oils both expressed and esfential; and may be united with watery liquors by means of the same intermedia which render the fluid oils miscible with water. In a heat less than that of boiling water, they melt into an oily fluid, and in this state they may be incorporated one with another. In their refolution by fire, in close vessels, they yield a manifest acid, and a large quantity of empyreumatic oil.

#### 7. Gum.

Gum differs from the foregoing substances, in being uninflammable: for though it may be burnt to a coal, and thence to ashes, it never yields any flame. It differs remarkably also in the proportion of the principles into which it is resolved by fire; the quantity of empyreumatic oil being far less, and that of acid far greater. In the heat of boiling water, it suffers so distipation: nor does it liquely like refins; but continues unchanged, till the heat is so far increased as to fcorch or turn it to a coal.

By a little quantity of water, it is softened into a viscous adhesive mais, called mucilage: by a larger quantity it is dissolved into a stuid, which proves more or less glutinous, according to the proportion of gum. It does not dissolve in vinous spirits, or in any kind of oil: nevertheless, when softened with water into a mucilage, it is easily miscible both with the fluid oils and with refins, which, by this means, become foluble in watery liquors along with the gum, and are thus excellently fitted out for medicinal purposes.

This elegant method of uniting oils with aqueous liquors, which has been kept a secret in few hands, appears to have been known to Dr. Grew. "I took (says he) oil of aniseeds, and pouring it upon 41 amother body, I so ordered it, that it was thereby turned into a per-44 feet milk-white balfam or butter; by which means the oil became 46 mingleable with any vinous or watery liquor; easily and instan-. taneously dissolving therein, in the form of a milk. And note,

"this is done without the least alteration of the smell, taste, nature or operation of the said oil. By somewhat the like means any other stillatitious oil may be transformed into a milk-white butter, and in like manner be mingled with water or any other liquor; which is of various use in medicine, and what I find oftentimes very converient and advantageous to be done." (Grew of mixture, chap. v. inst. i. § 7.) This enquiry has lately been further prosecuted, in the first volume of the Medical Observations published by a society of physicians in London; where a variety of experiments is related, of rendering oils both essential and expressed, and different unctuous and resinous bodies, soluble in water by the mediation of gum.

As oily and refinous substances are thus united to water by the means of gum, so gums may in like manner be united to spirit of wine by the intervention of resins and essential oils; though the spirit does not take up near so much of the gum, as water does of the oil

of refin.

Acid liquors, though they thicken pure oils or render them confishent, do not impede the dissolution of gum, or of oils blended with gum. Alkaline salts, on the contrary, both fixt and volatile, though they render pure oils dissoluble in water, prevent the solution of gum, and of mixtures of gum and oil. If any pure gum be dissolved in water, the addition of any alkali will occasion the gum to separate, and fall to the bottom in a consistent form: if any oily or resinous body was previously blended with the gum, this also separates, and either finks to the bottom, or rises to the top, according to its gravity.

#### 8. Gum-refin.

By gum-resin is understood a mixture of gum and resin. Many vegetables contain mixtures of this kind, in which the component parts are so intimately united, with the interposition perhaps of some other matter, that the compound, in a pharmaceutical view, may be considered as a distinct kind of principle; the whole mass dissolving almost equally in aqueous and in spirituous liquors; and the solutions being not turbid or milky, like those of the grosser mixtures of gum and resin, but perfectly transparent. Such is the astringent matter of bistort root, and the bitter matter of gentian. It were to be wished that we had some particular name for this kind of matter; as the term gum-resin is appropriated to the grosser mixtures, in which the gummy and resinous part are but loosely joined, and easily separable from one another.

#### 9. Saline matter.

Or the faline juices of vegetables there are different kinds, which have hitherto been but little examined: the sweet and the acid ones are the most plentiful, and those which are the most known.

These

These juices, exposed to a heat equal to that of boiling water, suffer generally no other change, than the evaporation of their watery moisture; the saline matter remaining behind, along with such of the other not volatile parts as were blended with it in the juice. From many, after the exhalation of great part of the water, the saline matter gradually separates in keeping, and concretes into little solid masses, leaving the other substances dissolved or in a moist state: from others, no means have yet been found of obtaining a pure concrete salt.

These salts dissolve not only in water like other saline bodies, but many of them, particularly the sweet, in rectified spirit also. The gross oily and gummy matter, with which they are almost always accompanied in the subject, dissolves freely along with them in water, but is by spirit in great measure lest behind. Such heterogeneous matters, as the spirit takes up, are almost completely retained by it, while the salt concretes; but of those, which water takes up, a considerable part always adheres to the salt. Hence essential salts, as they are called, prepared in the common manner from the watery juices of vegetables, are always sound to partake largely of the other soluble principles of the subject; whilst those extracted by spirit of wine prove far more pure. By means of rectified spirit, some productions of this kind may be excellently freed from their impurities; and perfect saccharine concretions obtained from many of our indigenous sweets.

There is another kind of saline matter, obtained from some refinous bodies, particularly from benzoine. of a different nature from the foregoing, and supposed by some of the chemists to be a part of the effential oil of the resin, coagulated by an acid, with the acid more predominant, or more disengaged, than in the other kinds of coagulated or indurated oils. These concretes dissolve both in water and in vinous spirits, though difficultly and sparingly in both: they shew some marks of acidity, have a considerable share of smell like that of the resin they are obtained from, exhale in a heat equal to that of boiling water or a little greater, and prove instammable in

the fire.

### General observations on the foregoing principles.

1. ESSENTIAL oils, as already observed, are obtainable only from a few vegetables, and camphor from a much smaller number: but gross oil, resin, gum, and saline matter, appear to be common in greater or less proportion, to all; some abounding more with one, and others with another.

2. The several principles are in many cases intimately combined; so as to be extracted together from the subject, by those dissolvents, in which some of them, separately, could not be dissolved. Hence watery infusions, and spirituous tinctures of a plant contain, respectively, more than water or spirit is the proper dissolvent of.

3. After

3. After a plant has been sufficiently infosed in water, all that spirit extracts from the residuum may be looked upon as consisting wholly of such matter as directly belongs to the action of spirit. And contrariwise, when spirit is applied first, all that water extracts afterwards may be looked upon as consisting only of that matter of which water is the direct dissolvent.

4. If a vegetable substance, containing all the principles we have been speaking of, be boiled in water; the effential oil, whether stuid or concrete, and the camphor, and volatile essential salt, will gradually exhale with the steam of the water, and may be collected by receiving the steam in proper vessels placed beyond the action of the heat. The other principles not being volatile in this degree of heat, remain behind: the gross oil and sebaceous matter stoat on the top: the gummy and saline substance, and a part of the resin, are dissolved by the water, and may be obtained in a folid form by straining the liquor, and exposing it to a gentle heat till the water has exhaled. The rest of the resin, still retained by the subject, may be extracted by spirit of wine, and separated in its proper form, by exhaling the spirit. On these soundations, most of the substances contained in vegetables may be extracted, and obtained in a pure state, however they may be compounded together in the subject.

Sometimes one or more of the principles is found naturally difengaged from the others, lying in dillinct receptacles within the fubject, or extravasated and accumulated on the surface. Thus, in the dried roots of angelica, cut longitudinally, the microscope discovers veins of refin. In the flower cups of hypericum, and the leaves of the orange tree, transparent points are distinguished by the naked eye, which on the first view seem to be holes, but on a closer examination are found to be little vehicles filled with effential oil. In the bark of the fir, pine, larch, and some other trees, the oily receptacles are extremely numerous, and so copiously supplied with the oily and refinous fluid, that they frequently burft, especially in the warm climates, and discharge their contents in great quantities. acacia tree in Egypt, and the plum and cherry among ourselves, yield almost pure gummy exudations. From a species of ash is secreted the faline sweet substance manna; and the only kind of sugar which the ancients were acquainted with, appears to have been a natural ex-

adation from the cane.

6. The foregoing principles are, so far as is known, all that naturally exist in vegetables; and all that art can extract from them, without such operations as change their nature, and destroy their original qualities. In one or more of these principles, the colour, smell, taste, and medicinal virtues of the subject, are almost always found consequenced.

found concentrated.

7. In fome vegetables, the whole medicinal activity refides in one principle. Thus, in sweet almonds, the only medicinal principle

ciple is a gross oil; in horse-raddish root, an essential oil; in jalap root, a resin; in marshmallow root, a gum; in the leaves of sorrel, a saline acid sustance.

8. Others have one kind of virtue residing in one principle, and another in another. Thus Peruvian bark has an astringent resin, and a bitter gum; wormwood, a strong slavoured essential oil, and a bitter gum resin.

9. The gross insipid oils and sebaceous matters, the simple insipid gums, and the sweet and acid saline substances, appear to nearly agree, respectively, among themselves, in their medicinal qualities, as well

as in their pharmaceutic properties.

10. But effential oils, refins, and gum-refins, differ greatly in different subjects. As effential oils are universally the principle of odour in vegetables, it is obvious that they must differ in this respect as much as the subjects from which they are obtained. Refins frequently partake of the oil, and consequently of the differences depending thereon; with this further diversity, that the gross resinous part often contains other powers than those which reside in oils. Thus from a wormwood a resin may be prepared, containing not only the strong smell and slavour, but likewise the whole bitterness of the herb; which last quality the oil is entirely free from. The bitter, aftringent, purgative and emetic virtues of vegetables reside generally in different fort of resinous matter, either pure, or blended with gummy and saline parts; of which kind of combinations there are many so intimate, that the component parts can scarcely be separated from one another, the whole compound dissolving almost equally in aqueous and spirituous menstrua.

11. There are some substances also, which from their being totally dissoluble in water, and not at all in spirit, may be judged to be mere gums, but which, nevertheless, pesses virtues never to be sound in the simple gums. Such are the aftringent gum called acacia, and the

purgative gum extracted from aloes.

12. It is supposed that vegetables contain certain subtile principles or presiding spirits, different in different plants, of too great tenuity to be collected in their pure state, and of which oils, gums, and refins are only the matrices or vehicles. This enquiry is foreign to the purposes of pharmacy, which is concerned only about grosser and more sensible objects. When we obtain from an odoriserous plant an essential oil, containing in a small compass the whole fragrance of a large quantity of the subject, our intentions are equally answered, whether the substance of the oil be the direct odorous matter, or whether it has diffused through it a fragrant principle more subtile than itself. And when this oil, in long keeping, loses its odour, and becomes a resin, it is equal, in regard to the present considerations, whether the effect happens from the avolation of a subtile

fubtile principle, or from a change produced in the fubflance of the oil itself.

### SECT. II.

#### Animals.

IN animal bodies we find certain substances, which have a great refemblance, in their general properties, to those of the vegetable

kingdom.

Animal oils and fats, like the gross oils of vegetables, are not, of themselves, dissoluble either in water or vinous spirits; but they may be united with water by the intervention of gum or mucilage; and most of them may be changed into soap, and thus rendered miscible with spirit, as well as water, by fixed alkaline salts.

The odorous matter of some odoriferous animal substances, as musk, civet, castor, is, like essential oil, soluble in spirit of wine, and volatile in the heat of boiling water. Cartheuser relates, that from castor an actual essential oil has been obtained, in a very small quantity,

but of an exceedingly strong diffusive smell.

The vesicating matter of cantharides, and those parts of sundry animal substances, in which their peculiar tastes reside, are dissolved by rectified spirit, and seem to have some analogy with resins and gummy resins.

The gelatinous principle of animals, like the gum of vegetables, dissolves in water, but not in spirit or in oils; like gums also, it ren-

ders oils and fats miscible with water into a milky liquor.

Some infects, particularly the ant, are found to contain an acid

juice, which approaches nearly to the nature of vegetable acids.

There are however fundry animal juices, which differ greatly, even in these general kinds of properties, from the corresponding ones of vegetables. Thus animal serum, which appears analogous to vegetable gummy juices, has this remarkable difference, that though it mingles uniformly with cold or warm water, yet on considerably heating the mixture, the animal matter separates from the watery sluid, and concretes into a solid mass. Some have been apprehensive, that the heat of the body, in some distempers, might rise to such a degree, as to produce this dangerous or mortal concretion of the serous humours: but the heat requisite for this effect is greater than the human body appears capable of sustaining; being nearly about the middle point between the greatest human heat commonly observed and that of boiling water.

THE foft and fluid parts of animals are strongly disposed to run into putrefaction: they putrefy much sooner than vegetable matters, and, when corrupted, prove more offensive.

This

This process takes place, in some degree, in the bodies of living animals; as often as the juices stagnate long, or are prevented, by an obstruction of the natural emunctories, from throwing off their more

volatile and corruptible parts.

The doctrine of putrefaction, both in living and in dead animals, has lately received great light from the curious and interesting experiments and observations of Dr. Pringle. He observes, that if the corruption is great and sudden, a fever or a flux ensue; but that, if the accumulation of corrupted matter is so slow, that the body becomes habituated to the putrefaction, a scurvy prevails. Hence the frequency of this last distemper, in long voyages, on board unventilated ships, from corrupted air and provisions; in marshy countries, from similar causes; and in a lesser degree, in all northern climates, in moist situations, from a want of due perspiration.

During putrefaction, a quantity of air is generated; all the humours become gradually thinner, and the fibrous parts more lax and tender. Hence the tympany, which succeeds the corruption of any of the viscera, or the imprudent suppression of dysenteries by aftringents; and the weakness and laxity of the vessels observable in

scurvies, &c.

The crassamentum of human blood changes by putrefaction in to a dark livid-coloured liquor; a few drops of which tinge the ferum of a tawny hue; like that of the ichor of fores and dysenteric fluxes, and of the white of the eye, the saliva, the serum of blood drawn from a vein, and that which oozes from a blister, in deep scurvies, and the advanced state of malignant severs.

The putrid crassamentum changes a large quantity of recent urine to a slame-coloured water, so common in severs and in the scurvy. This mixture, after standing an hour or two, gathers a cloud, resembling what is seen in the crude water of acute distempers, with some oily matter on the surface, like the scum which

floats on scorbutic urine.

The ferum of blood deposits, in putrefaction, a sediment resembling well digested pus, and changes to a faint olive green. A serum, so far putressed as to become green is perhaps never to be seen in the vessels of living animals: but in dead bodies this serum is to be distinguished by the green colour which the stesh acquires in corrupting. In salted meats, this is commonly ascribed to the brine, but erroneously; for that has no power of giving this colour, but only of qualifying the taste, and in some degree the ill effects of corrupted aliments. In soul ulcers, and other fores, where the serum is lest to stagnate long, the matter is likewise found of this colour, and is then always acrimonious.

The putrefaction of animal substances is prevented or retarded by all saline matters, even by the fixt and volatile alkaline salts, which C 2

have generally been supposed to produce a contrary effect. Of all the salts that have been made trial of, sea salt seems to resist putrefaction the least: in small quantities, it even accelerates the process. The vegetable bitters, as chamomile flowers, are much stronger antiseptics, not only preserving sless long uncorrupted, but likewise somewhat correcting it when putrid: the mineral acids have this last effect in a more remarkable degree. Vinous spirits, aromatic and warm substances, most of the diaphoretic drugs, and the acrid plant salfely called alkalesent, as scurvygrass and horseradish, are also found to reast putrefaction; and some of the absorbent earths, as chalk, to promote it.

It is observable, that notwithstanding the strong tendency of animal matters to putrefaction, yet broths made from them with the admixture of vegetables, instead of putrefying, turn sour. Dr. Pringle sinds, that when animal sless in substance is beaten up with bread, or other farinaceous vegetables, and a proper quantity of water, into the consistence of a pap, this mixture likewise, kept in a heat equal to that of the human body, grows in a little time sour; whilst the vegetable matters, with the sless, suffer no such change. See the appendix to

his observations on the diseases of the army.

Animal substances, burnt in the open air, are resolved, like vegetables, into soot and ashes, but with this difference, that no fixt alkaline salt can be obtained from the ashes, and that no acid vapour accompanies the smoke. They emit, during the burning, a setid smell, of a peculiar kind, by which animal substances may be distinguished at once from all those of the vegetable kingdom. In close vessels, they give over, after the watery moisture, a volatile alkaline salt, which either concretes into a folid form, or dissolves in the water, and thus composes what is called spirit; together with an empyreumatic oil, of a more setid kind than the oils of vegetables; without the least souther of acid throughout the whole process. A black coal remains, which, in the open air, burns into white ashes void of saline matter.

It was observed in the preceding section, that some sew vegetables, in this resolution of them by fire, discover some agreement, in their matter, with bodies of the animal kingdom: yielding a volatile alkaline salt in considerable quantity, with little or nothing of the acid or fixed alkali, which the generality of vegetables afford. In animal substances also there are some exceptions to the general alalysis; from animal sats, instead of a volatile alkali, an acid liquor is obtained, and their empyreumatic oil wants the peculiar offensiveness of the

other animal oils.

## SECT. III.

#### Minerals.

#### Oils and Bitumens.

IN the mineral kingdom is found a fluid oil, called naphtha or petroleum, floating on the furface of waters, or issuing from clefts of rocks, particularly in the eastern countries, of a strong smell very different from that of vegetable or animal oils, limpid almost as water, highly inflammable, not soluble in spirit of wine, and more averse to upion with water than any other oils.

There are different forts of these mineral oils, more or less tinged, of a more or less agreeable, and a stronger or weaker smell. By the admixture of concentrated acids, which raise no great heat or conflict with them, they may become thick, and at length consistent; and

in these states are called bitumens.

These thickened or concreted oils, like the corresponding products of the vegetable kingdom, are generally soluble in spirit of wine, but much more disticultly, more sparingly, and for the most part only partially: they liquefy by heat, but require the heat to be considerably stronger. In a proper degree of heat, they give out a sluid oil, greatly resembling the native patrolea; a small quantity of a black coaly matter remaining behind. Their smells are various; but all of them, either in their natural state, or when melted, or set on sire, yield a peculiar kind of strong scent, called, from them, bituminous.

Earths.

In treating of vegetables and animals, we forbore to speak of their earthy matters, that the distinguishing characters of the several classes of earthy bodies might be the easter apprehended, by having them placed here in one synoptical view: the little impropriety, of joining the vegetable and animal earths to the mineral must be overlooked for the sake of that advantage. Under the mineral earths are included stones, these being no other than earths in an indurated state.

The different kinds of these bodies hitherto taken notice of, are the following:

I. Earths foluble in the nitrous, marine, and vegetable acids, but not at all or exceeding sparingly in the vitriolic acid. When previously disfolved in other acids, they are precipitated by the addition of this last, which thus unites with them into inspid, or nearly insipid concretes, not dissolve in any liquor. Of this kind are,

1. The mineral calcareous earth: distinguished by its being convertible, in a strong fire without addition, into an acrimonious calx, called quicklime. The earth occurs in a variety of forms in the mi-

neral kingdom. The fine foft chalk, the coarfer limestones, the hard marbles, the transparent spars, the earthy matter contained in waters, and which, separating from them, incrustates the sides of caverns or hangs in icicles from the top, receiving from its different appearances different appellations; how strongly soever some of these bodies have been recommended for particular medicinal purposes, are at bottom no other than different forms of this calcareous earth, fimple pulverization depriving them of the superficial characters by which they were distinguished in the mass. Most of them contain generally a greater or less admixture of some of the indissoluble kinds of earth; which, however, affects their medicinal qualities no otherwise, than by the addition which it makes to their bulk. Chalk appears to be one of the purest, and is therefore in general preferred. They all burn into a strong quicklime: in this state a part of them dissolves in water, which thus becomes impregnated with the aftringent and lithontriptic powers that have been erroneously ascribed to some of the earths in their natural state.

2. The animal calcareous earth: burning into quicklime, like the mineral. Of this kind are oyster shells, and all the marine shells that have been examined; though with some variation in the strength of

the quicklime produced from them.

3. The earth of bones and horns: not at all burning into quicklime. This kind of earth is more difficult of folution in acids than either of the preceding. It is accompanied in the subject with a quantity of gelatinous matter, which may be separated by long boiling in water, and more perfectly by burning in the open air: the earth may be extracted also from the bone or horn, though difficultly, by means of acids; whereas vegetables, and the soft parts of animals, yield their pure easth by burning only.

II. Earths soluble with ease in the vitriolic as well as wher acids; and yielding, in all their combinations therewith, saline concretes

soluble in water:

1. Magnefia alba: composing with the vitriolic acid a bitter purgative liquor. This earth has not yet been found naturally in a pure state. It is obtained from the purging mineral waters and their salts, from the bitter liquor which remains after the crystallization of sea salt from sea water, and from the sluid which remains uncrystallized in the putrefaction of some forts of rough nitre. The ashes of vegetables appear to be nearly the same kind of earth.

2. Alumenous earth: composing with the vitriolic acid a very aftringent liquor. This earth also has not been found naturally pure. It is obtained from alum, which is no other than a combination of it with the vitriolic acid: it may likewise be extracted, by strong boil-

ing in that acid, from clays and boles.

III. Earths,

III. Earths, which by digestion in acids, either in the cold or in a moderate warmth, are not at all dissolved.

1. Argillaceous earth; becoming bard, or acquiring an additional bardness in the fire. Of this kind of earth there are several varieties, differing in some particular properties: as the purer clays, which when moistened with water form a very viscous mass, difficultly diffusible through a larger quantity of the fluid, and slowly subsiding from it: boles, less viscous, more readily miscible with water, and more readily subsiding: and ochers, which have little or nothing of the viscosity of the two foregoing, and are commonly impregnated with a yellow or red ferugineous calx.

2. Crystalline earth: naturally bard, so as to strike sparks with steel: becoming friable in a strong fire. Of this kind are slints, crystals, &c. which appear to consist of one and the same earth, dif-

fering in purity, hardness, and transparency of the mass.

- 3. Gypseous earth: reducible by a gentle beat into a soft powder, which unites with water into a mass, somewhat viscous and tenacious while moist, but quickly drying and becoming bard. A greater heat deprives the powder of this property, without occasioning any other alteration. Such are the transparent selenitæ; the sibrous stony masses, improperly called English tale; and the granulated gypsa or plaster of Paris stones. Though these bodies, however, have been commonly looked upon as mere earths, of a distinct kind from the rest, they appear, both from analytical and synthetical experiments, to be no other than combinations of the mineral calcareous earth with vitrolic acid. See the characters of the earths of the first class.
- 4. Talky earth: fcarcely alterable by a webenent fire. The masses of this earth are generally of a sibrous or leafy texture; more or less pellucid, bright or glittering; smooth and unctuous to the touch; too slexible and elastic to be easily pulverized; soft, so as to be cut with a knife. In these respects some of the gypseous earths greatly resemble them, but the difference is readily discovered by fire; a weak heat reducing the gypseous to powder, while the strongest makes no other alteration in the talky, than somewhat diminishing their flexibility, brightness, and unctuosity.

#### Metals.

Or metals, the next division of mineral bodies, the most obvious characters are, their peculiar bright aspect, perfect opacity, and great weight; the lightest of them is six, and the heaviest upwards of nineteen times heavier than an equal bulk of water.

They all melt in the fire, except platina, a metallic body, which has not been applied to any medical use, and which is therefore ex-

cluded from this general view of medical subjects.

Gold.

w =

Gold and filver, how long soever they are continued in sustaining remain unchanged and undiminished. The others, if air is admitted to them, are gradually converted, with different degrees of facility, into a powdery or friable substance, called calx, destitute of the metallic aspect, and much lighter, in proportion to its bulk, than the metal itself. This change in their obvious properties is generally accompanied with a notable alteration in their medical virtues: thus quickfilver, which taken into the body in its crude state and undivided, seems inactive; when calcined by fire proves, even in small doses, a strong emeric and cathartic, and in smaller ones, a powerful alterative in chronical disorders; while regulus of antimony, on the contrary, is changed, by the same treatment, from a high degree of virulence to a state of inactivity.

Calces of mercury and arienic exhale in a heat below ignition: those of lead and bismuth, in a red or low white heat, run into a transparent glass: the others are not at all vitrescible, or not without extreme vehemence of fire. Both the calces and glasses recover their metallic form and qualities again, by the skilful addition of any kind of inflammable substance that does not contain a mineral

acid,

All metallic bodies dissolve in acids; some only in particular acids, as silver and lead in the nitrous; some only in compositions of acids, as gold in a mixture of the nitrous and marine; and others, as iron and zinc, in all acids. Some likewise dissolve in alkaline liquors, as copper; and others, as lead, in expressed oils. Fused with a composition of sulphur and fixt alkaline salt, they are all, ex-

cept zinc, made soluble in water.

All metallic substances, dissolved in saline liquors, have powerful effects in the human body, though many of them appear in their pure state to be inactive. Their activity is generally in proportion to the quantity of acid combined with them: Thus lead, which in its crude form has no sensible effect, when united with a small portion of vegetable acid into cerus, discovers a low degree of the styptic and malignant quality, which it so strongly exerts when blended, with a larger quantity of the same acid, into what is called saccharum salurni: and thus mercury, with a certain quantity of the marine acid, forms the violent corrosive sublimate, which by diminishing the proportion of acid, becomes the mild medicine called mercurius dulcis.

Acids.

The mineral acids are distinguished by the names of the concretes from which they have been principally extracted; the vitriolic from vitriol, the nitrous from nitre or saltpetre, and the marine from common sea salt. They are all highly corrosive, infomuch as not to be safely touched, unless largely diluted with

water, or united with such substances as obtund or suppress their acidity. Mixed hastily with vinous spirits, they raise a violent ebullition and heat, accompanied with a copious discharge of noxious sumes: a part of the acid unties intimately with the vinous spirit into a new compound, void of acidity, called dulcised spirit. It is observable, that the marine acid is much less disposed to this union with spirit of wine, than either of the other two: nevertheless, many of the compound salts resulting from the combination of earthy and metallic bodies with this acid, are soluble in that spirit, while those with the other acids are not. All these acids effervesce strongly with alkaline salts, both sixt and volatile, and form with them neutral salts, that is, such as discover no marks either of an acid or alkaline quality.

The nitrous and marine acids are obtained in the form of a thin liquor, the acid part being blended with a large proportion of water, without which it would be diffused into an incoercible vapour: the vitriolic stands in need of so much less water for its condensation as to assume commonly an oily consistence (whence it is called oil of vitriol) and in some circumstances even a solid one. Alkaline salts, and the soluble earths and metals, absorb from the acid liquors only the pure acid part; so that the water may now be evaporated by heat.

and the compound falt left in a dry form.

We have already taken notice of two forts of alkaline falt, the volatile alkali of animals, and the fixt alkali of vegetables. In the mineral kingdom, another species of fixt alkali, different in several respects from the vegetable, is found sometimes in a detached state, but more plentifully in combination with the marine acid, with which it composes sea salt. From the coalition of the different acids with these three alkalies, and with the several soluble earths and metallic bodies, result a variety of saline compounds, the principal of which will be particularized in the sequel of this work.

The vitriolic acid, in its concentrated liquid state, is much more ponderous than the other two, emits no visible vapours in the heat of the atmosphere, but imbibes moisture therefrom, and in creases in its weight: the nitrous and marine emit copious corrosive sume, the nitrous yellowish red, and the marine white ones. If bottles containing the three acids are stopt with cork, the cork is found in a little time tinged black with the vitriolic, corroded into a yellow substance by the nitrous, and into a whitish one by the

marine.

It is above laid down as a character of one of the classes of earths, that the vitriolic acid precipitates them when they are previously dissolved in any other acid: it is obvious, that on the same principle this particular acid may be distinguished from all others. This character serves not only for the acid in its pure state, but like-

wise for all its combinations that are soluble in water: if a solution of any compound salt, whose acid is the vitriolic, be added to a solution of chalk in any other acid, the vitriolic acid will part from the substance it was before combined with, and join itself to the chalk, forming therewith a compound, which being no longer dissoluble in the liquor, renders the whole milky for a time, and then

gradually subsides.

This acid may be diffinguished also, in compound salts, by another criterion not less strongly marked: if any salt containing it be mixed with powdered charcoal, and the mixture exposed, in a close vessel, to a moderately strong sire, the acid will unite with the directly inflammable part of the charcoal, and compose therewith a genuine sulphur. Common brimstone is no other than a combination of the vitriolic acid with a small proportion of inflammable matter. With any kind of inflammable matter that is not volatile in close vessels, as the coal of vegetables, of animals, or of bitumens, this acid composes always the same identical sulphur.

The nitrous acid also, whatever kind of body it be combined with, is both distinguished, and extricated therefrom, by means of any inflammable substance brought to a state of ignition: if the subject be mixed with a little powdered charcoal, and made red-hot, a deslagratio nor submination ensues, that is, a bright slame with a hissing noise, and the inflammable matter and the acid being thus consumed or dissipated together, there remains only the substance that was before combined with the acid, and the small quantity of assessments.

by the coal.

This property of the nitrous acid, deflagrating with inflammable fubfiances, and that of the vitriolic of forming sulphur with them, serve not only as criteria of the respective acids in their various forms and disguises, but likewise for discovering inflammable matter in bodies, when its quantity is too small to be sensible on other trials.

If a fixt alkaline salt be united with a vegetable acid, as that of vinegar, into a neutral salt; on adding to this compound some marine acid, the acetous acid will be disengaged, so as to exhale totally in a moderate heat, leaving the marine in possession of the alkali: the addition of the nitrous will in like manner disposses the marine, which now arises in its proper white sames, though without such an addition it could not be extricated from the alkali by any degree of heat: on the addition of the vitriolic acid, the nitrous gives way in its turn, exhaling in red sumes, and leaving only the vitriolic acid and the alkali united together.

Again, if any metallic body be dissolved in an acid, the addition of any earthy body that is dissolved in that acid will precipitate the metal: a volatile alkaline falt will in like manner precipitate the earth; and a fixt alkali will dislodge the volatile; which

which last being readily exhalable by heat, the remaining salt will be the same as if the acid and fixt alkali had been joined together at first, without the intervention of any of the other bodies.

THE power in bodies, on which these various transpositions and combinations depend, is called by the chemists APPINITY; a term, like the Newtonian attraction, designed to express, not the cause, but the effect. When an acid spontaneously quits a metal to unite with an alkali, they say it has a greater affinity to the alkali than to the metal: and when, conversively, they say it has a greater affinity to faxt alkalies than to those of the volatile kind, they mean only that it will unite with the fixt in preference to the volatile, and that if previously united with a volatile alkali, it will forsake this for a fixt one.

The doctrine of the affinities of bodies is of very extensive use in the chemical pharmacy: many of the officinal processes, as we shall see hereaster, are sounded on it: several of the preparations turn out very different from what would be expected by a person unacquainted with these properties of bodies; and several of them, if, from an error in the process, or other causes, they prove unsit for the use intended, may be rendered applicable to other purposes, by such transpositions of their component parts as are pointed out by the knowledge of their affinities.

I shall here therefore subjoin a table of the principal affinities obferved in pharmaceutical operations, formed chiefly on that of Mr. Geoffroy (which was published in the Memoirs of the French academy for the year 1718) with such corrections and additions as later expe-

riments have furnished.

The table is thus to be understood. The substance printed in capitals, on the top of each series, has the greatest affinity with that immediately under it, a less affinity with the next, and so on to the end of the series; that is, if any of the remote bodies has been combined with the top one, the addition of any of the intermediate bodies will disunite them; the intermediate body uniting with the uppermost body of the series, and throwing out the remote one. Thus in the first series of the affinities of water, a fixt alkali being placed between the water and inflammable spirit, it is to be concluded, that wherever water and spirit are mixt together, the addition of any fixed alkaline salt will absorb the water, and occasion the pure spirit to be separated. Where several substances are expressed in one series, it is to be understood, that any one of those bodies, which are nearest to the uppermost, will in like manner disengage from it any one of those which are more remote.

I. WATER:

Fixt alkaline salt: Inflammable spirit.

2. WATER: Inflummable spirit: Volatile alkaline falt.

3. WATER: Inflammable spirit: Sandry compound salts.

4. Inflammable Spirit: Water:
Oils and Refins.

5. VITRIOLIC ACID:
Inflammable principle:
Fixt alkaline falts:
Calcareous earths calcined:
Volatile alkaline falts:
Calcareous earths uncalcined:
Zinc and Iron:
Copper:
Sifter.

6. NITROUS ACID:
Inflammable principle:
Pixt alkaline falts:
Calcareous earths calcined:
Volatile alkaline falts:
Calcareous earths uncalcined:
Zinc:
Iton:
Copper:
Lead:
Mercury:
Silver:
Camphor.

7. MARINE ACID:
Fixt alkaline falts:
Calcareous earths calcined:
Volatile alkaline falts:
Calcareous earths uncalcined:

Zinc:
Iron:
Tin:
Regulus of antimony:
Copper:
Lead:
Silver:
Mercury.

8. ACETOUS ACID: Iron: Copper.

9. ALKALINE SALTS:
Vitriolic acid:
Nitrous acid:
Marine acid:
Vinegar:
Tartar:
Oils and Sulphur.

10. Soluble Earths: Vitriolic acid: Nitrous acid: Marine acid.

11. INPLAMMABLE PRINCIPLE: Nitrous acid: Vitriolic acid: Metallic fubstances: Fixt alkaline falts.

I2. SULPHUR:
Fixt alkali, and Quicklime:
Iron:
Copper:
Lead:
Silver:
Regalus of antimony:
Mercury:
Arfenic.

13. Gold: Ethereal spirit: Acids.

14. MER-

14. MERCURY:

Marine acid: Vitriolic acid: Nitrous acid.

Vitriolic acid: Marine acid: Nitrons acid: Vinegar: Oils.

16. SILVER :

Marine acid: Vitriolic acid: Nitrous acid.

17. COPPER: Vitriolic acid:

Marine acid: Nitrous acid.

18. IRON &

Vitriolic acid: Marine acid: Nitrous acid.

19 Recutus of Antimony: Vitriolic acid:

Nitrous acid: Marine acid.

## CHAPTER III.

# Of the Pharmaceutical Apparatus.

NE of the principal parts of the pharmaceutic apparatus confifts in contrivances for containing and applying fire, and for directing and regulating its power. Of these contrivances, called furnaces, there are different kinds, according to the conveniency of the place, and the particular purposes they are intended to answer. I shall here endeavour to give a general idea of the structure of those which are employed in pharmaceutical operations, and of the principles on which they are built.

urnaces

THE most simple furnace is the common stove, otherwise called the furnace for OPEN FIRE. This is usually made of an iron hoop, sive or six inches deep; with a grate or some iron bars across the bottom, for supporting the suel. It either stands upon seet, so as to be moveable from place to place, or is fixt in brickwork. In this last case, a cavity is lest under the grate, for receiving the assess that drop through it; and an aperture or door, in the forepart of this assembly serves both for allowing the assess to be occasionally raked out, and for admitting air to pass up through the fuel. This surnace is designed for such operations as require only a moderate heat; as insusion, decostion, and the evaporation of liquids. The vessel, containing the subject matter, is supported over the fire by a trevet.

A deeper hoop or body, cylindrical, parallelopipedal, widening upwards, elliptical, or of other figures; formed of, or lined with, such materials as are capable of sustaining a strong sire; with a grate and ash-pit beneath, as in the preceding; and communicating at the top with a perpendicular pipe, or chimney; makes a WIND

FURNACE.

The greater the perpendicular height of the chimney, the greater will be the draught of air through the furnace, and the more intensely will the fire burn; provided the width of the chimney is sufficient to allow a free passage to all the air that the furnace can receive through the grate: for which purpose, the area of the aperture of the chimney should be nearly equal to the area of the interstices of the grate.

Hence, where the chimney confifts of moveable pipes, made to fit upon one another at the ends, fo that the length can be occasionally increased or diminished, the vehemence of the fire will be increased

or diminished in the same proportion.

In furnaces whose chimney is fixed, the same advantage may be procured on another principle. As the intensity of the fire depends wholly upon the quantity of air successively passing through and animating the burning suel, it is obvious, that the most vehement fire may be suppressed or restrained at pleasure, by more or less closing either the ash-pit door by which the air is admitted, or the chimney by which it passes off; and that the fire may be more or less raised again, by more or less opening those passages. A moveable plate or REGISTER, in any convenient part of the chimney, affords commodious means of varying the width of the passage, and consequently of regulating the heat.

THERE are two general kinds of these wind surnaces; one with the chimney on the top, over the middle of the surnace; the other

with the chimney on one fide, and the mouth clear.

In the first, either the upper part of the furnace is contracted to fuch an aperture, that the chimney may sit upon it; or it is covered with an arched dome, or with a flat plate, having a like aperture in the middle. As in this disposition of the chimney, the inside of the furnace cannot be come at from above, a door is made in the side, a little above the grate, for supplying suel, inspecting the matter in the sire, &c.

For performing pusions in this furnace, the crucible, or melting vessel, is placed immediately among the sue; with a slip of brick, or some other like support, between it and the grate, to keep the cold air, which enters underneath, from striking on its bottom.

When defigned as a REVERBERATORY, that is, for distillation in long necks or coated glass retorts, two iron bars are placed across, above the fire, for supporting the vessel, whose neck comes out at an aperture made for that purpose in the side. This aperture should be made in the side opposite to that in which is the door above-mentioned, or at least so remote from it, that the receiver, sitted on the neck of the distilling vessel without the furnace, may not lie in the operator's way when he wants to stir the fire, or throw in fresh suel.

The other kind of wind furnace communicates, by an aperture in its back part near the top, either with an upright pipe of its own, or with the chimney of the room; in which last case, all other passages into the chimney must be closed up. Here the mouth of the furnace serves for a door, which may be occasionally covered with a plate or tile. Of this kind is the furnace most commonly used for suspense for a crucible.

This last construction, by leaving the mouth of the furnace clear, affords the conveniency of letting into it a boiling or evaporating pan, a copper still, an iron pot for distilling hartshorn, an

iron fand pot, or other like vessels, of such a size, that they may be supported on the surnace by their rims. The mouth being thus occupied by the vessels, a door must be made in the side for supplying and

firring the fuel.

When a furnace of this kind is designed only for a sand bath, it is most commodious to have the fand placed on a long iron plate furnished with a ledge of freestone or brickwork at each side. The mouth of the furnace is to be closely covered by one end of this plate; and the canal, by which the furnace communicates with its chimney, is to be lengthened and carried along under the plate; the plate forming the upper side of the canal. In this kind of sandbath, digestions, &c. requiring different degrees of heat, may be carried on at once; for the heat decreases gradually from the end over the surnace to the other.

When large vessels, as STILLS, and iron pots for distilling hartshorn and aqua fortis, are fixed in furnaces, a considerable part of the bottom of the vessel is commonly made to rest upon solid brick work.

The large still, whose bottom is narrow in proportion to its height, and whose weight when charged with liquor requires great part of it to be thus supported, exposes but a small surface to the action of the sire underneath. To make up for this disadvantage, the heat, which rises at the surther end of a long narrow grate, is conveyed all round the sides of the vessel, by a spiral canal, which communicates at top with a common chimney.

The pots for distilling hartshorn and aqua-fortis in the large way have part of their great weight borne up by three strong pins or trunnions, at equal distances round the pot towards the middle, reaching into a brickwork; so that less support being necessary underneath, a greater surface of the wide bottom lies exposed to the

immediate action of the fire.

If a surface, communicating with its chimney by a lateral canal, as in the sand surface above-mentioned, be carried to a considerable height above the part where this canal enters it; and if it be filled with such to the top, and closely covered; the such will burn no higher than up to the upper side of the canal through which the air passes off; and in proportion as this lower part of the such consumes, it will be supplied by that above, which falls down in its place. Hence in this surface, called an ATHANOR, a constant heat may be kept up for a considerable length of time, without attendance.

The tower of the athanor, or that part which receives the fuel, is commonly made to widen a little downwards, that the coals may fall the more freely; but not so much as that the part on fire at bottom may be too strongly pressed. A small aperture is made opposite

opposite to the canal or slew, or a number of openings according to the size of the furnace and the degree of heat required, for supplying air, which is more conveniently admitted in this manner than through the grate, as the interestices of the grate are in time choaked up by the ashes.

This furnace is defigned only for heating bodies exterior to it. Its canal or flew, as in the fand furnace already described, passes under a sand-bath or water-bath; at the further end of which, it rises perpendicularly to such a height, as may occasion a sufficient draught of

air through the fire.

The flew may be so wide, as to correspond to the whole height of the fire-place. A register or sliding plate, placed between the flew and the surnace, enables us to increase or diminish this height, and consequently the quantity of fire, at pleasure. If the space beneath the flew be inclosed to the ground, the heat in this cavity will be considerable enough to be applicable to some useful purposes.

WITH regard to the materials of furnaces, the fixt ones are built of bricks, cemented together by some good loam or clay. Any kind of loam or clayey composition that is of a proper degree of tenacity, which, when made into a passe with water and well worked, does not stick to the fingers, and which, when thoroughly dried, neither cracks nor melts in a vehement fire, is sit for this use: the purer and more tenacious clays require to have their tenacity lessened by an admixture of sand, or rather of the same kind of clay burnt and grossly powdered.

Smaller portable furnaces are made of strong iron or copper places, lined to the thickness of an inch or more with the same kind of clayey composition; which, for this use, may be beaten with some horse-

dung, chopt fraw, or cut hair or tow.

Very commodious portable furnaces, for a business of moderate extent, may be formed also of the larger kind of the common black-lead melting pots; by cutting a door at the bottom of the pot for the ashpit, another above this for the sire-place, and introducing a circular iron grate, of such a size, that it may rest between the two doors. A particular account of the method of preparing these surnaces for different uses may be seen in the sirst part of the Commercium philosophico-technicum, lately published.

#### Baths.

WHERE a strong degree of heat is requisite, as in the sustion of metals, &c. the vessel containing the subject matter is placed among the burning suel, or immediately over it: this is called operating in a naked sire. Where a smaller heat is sufficient, and the vessel employed is either of glass, or of the more tender kinds of earthen ware, the sand-bath or water-bath is used, to defend the vessel from

the immediate action of the fire, and to render the heat less fluc-

tuating.

Both these baths have their particular advantages and inconveniencies. In water the heat is equal through every part of the fluid; whereas in fand, it varies in different parts of one perpendicular line, decreasing from the bottom to the top. Water cannot be made to receive, or to transmit to vessels immersed in it, above a certain degree of heat, viz. that which is sufficient to make it boil, and hence it secures effectually against any danger of an excess of heat, in those operations wherein the product would be injured by a heat greater than that of boiling water; but this advantage renders it useless for processes which require a greater heat, and for which sand or other like folid intermedia are necessarily employed. There is this convenience also in the fand bath, that the heat may be readily diminished or increased about any particular vessel, by raising it higher out of the land, or finking it deeper; that different subjects may be exposed to different degrees of heat from one fire; and that it keeps the vessels steady. The sand made choice of should be a large coarsegrained kind, separated from the finer parts by washing, and from little stones by the sieve.

#### Coating of glasses, Lutes.

Some processes require to be performed with glass vessels in a naked fire. For these purposes, vessels made of the thinnest glass should be chosen; for these bear the fire, without cracking, much better than

those which are thicker and in appearance stronger.

All glasses, or other vessels that are apt to crack in the fire, must be cautiously nealed, that is, heated by flow degrees: and when the process is finished, they should be as slowly cooled, unless where the vessel is to be broken to get out the preparation, as in some sublimations: in this case it is more adviseable to expose the hot glass suddenly to the cold air, which will soon occasion it to crack, than to endanger throwing down the sublimed matter among the seces by a blow.

As a defence from the violence of the fire, and to prevent the contact of cold air on supplying fresh suel, &c. the glass is to be coated over, to the thickness of about half a crown, with Windsor loam sostened with water into a proper consistence and beaten up with some horsedung, or with the other clayey compositions abovementioned.

These compositions serve also as a lute, for securing the junctures of the vessels in the distillation of the volatile salts and spirits of animals: for the distillation of acid spirits, the matter may be moistened with a solution of fixed alkaline salt instead of water. For most other purposes, a piece of wet bladder, or a paste of flour and water, or of linseed meal (that is, the cake left after the expression

of oil of linseed) are sufficient lutes. The sew simple lutes, here described, will be found to answer all the purposes of the more operose compositions recommended for these intentions by the chemical writers.

#### Vessels.

It would be needless to enter here into a particular detail of the pharmaceutical inftruments; as we shall have occasion to mention the principal of them in the following chapter, in speaking of the several operations to which they are respectively subservient. In this place, I shall only give the operator a few general cautions with regard to the matter of the vessels designed for containing the

fubject.

Metalline vessels, except those made of gold or filver, are corroded by acids, even by the milder ones of the vegetable kingdom. Copper ones are corroded also by alkaline liquors, and by some neutral ones, as solutions of sal ammoniac: it is observable, that vegetable acids do not act upon this metal by boiling, so much as by standing in the cold; for even lemon-juice may be boiled in a clean copper vessel, without receiving from it any taste or ill quality, whereas, in the cold, it soon dissolves so much as to contract a pernicious taint. The ctin, with which copper vessels are usually lined, gives likewise a sensible impregnation to acid juices; and this impregnation also is probably not innocent, more especially as a quantity of lead is commonly mixed with the tin.

The common EARTHEN vessels are of a loose porous texture, and hence are apt to imbibe a considerable quantity of certain liquids, particularly of those of the saline kind; which soon discover their penetrating the vessel, by shooting into saline esslorescences on the outside. Those which are GLAZED have their glazing corroded by acids, by vinegar, and the acid juices of fruits, as well as by the stronger acids of the mineral kingdom. And as this glazing consists chiefly of vitresied lead, the impregnation, which it communicates to these liquors, is of a very dangerous kind: if vinegar be boiled for some time in a glazed earthen vessel, it will yield, on being inspissated, a true saccharum saturni, that is, a salt composed of lead and the acetous acid, of which hereaster.

The vessels called, from their hardness and compactness, STONE WARE, are in good measure free from the inconveniencies of the coarser earthen ones. Their glazing, being a part of the clay itself superficially vitrested by means of the sumes of common salt, appears to be proof against acids.

GLASS vessels suffer no corrosion, and give no taint, in any of the pharmaceutic operations. These therefore, in such processes as will

admit their use, ought always to be preferred.

#### Weights.

Two different kinds of weights are made use of in this country; one in the merchandize of gold and filver; the other for almost all goods besides. The sirst we call Troy, the latter Averdupois weight.

The goldsmiths divide the Troy pound into twelve ounces; the ounce into twenty pennyweights, and the pennyweight into twenty-four grains. The Averdupois pound is divided into sixteen ounces;

and the ounce into fixteen parts, called drams.

The pound of the London and Edinburgh dispensatories (which is the only one made use of in this work) is that of the goldsmiths, divided in the following manner;

The Pound
The Ounce
The Dram
The Scruple
The grain is equal to the goldsmiths grain.

The medical or Troy pound is less than the Averdupois, but the ounce and the dram greater. The Troy pound contains \$760 grains; the Averdupois 7000 grains The Troy ounce contains 480 grains; the Averdupois only 437½. The Troy dram 60; the Averdupois dram somewhat more than 27. Eleven drams Averdupois are equal to five drams Troy; twelve ounces Averdupois to nearly eleven ounces Troy; and nineteen pounds Averdupois to somewhat more than twenty-three pounds Troy.

These differences in our weights have occasioned great consusion in the practice of pharmacy. As the druggists and grocers sell by the Averdupois weight, the apothecaries have not in general kept any weights adjusted to the Troy pound greater than two drams, using for all above Averdupois. By this means it is apparent, that in all compositions, where the ingredients are prescribed some by pounds and others by ounces, they are taken in a wrong proportion to each other; and the same happens when any are directed in lesser denominations than the ounce, as these subdivisions, used by the apothecaries, are made to a different ounce. The mercurial plaster of the late Pharmacopæia, and the mercurial cerate of the present, if compounded by the Averdupois weight, contain about one fixth less quicksilver than if made, as they ought to be, by the Troy. This error prevailed so far as to be received in some former editions of the London pharmacopæia itself; but is now happily removed.

#### Measures.

The measures employed with us in pharmacy are sie common wine measures.

A Gallon
The Pint
The Ounce
Contains
The Ounce

By a spoonful is understood in the London Dispensatory the measure of half an ounce; in the Edinburgh, half an ounce weight in syrups,

and three drams in distilled waters.

Though the pint is called by Latin writers libra or pound, there is not any known liquor of which a pint measure answers to that weight. A pint of the highest rectified spirit of wine exceeds a pound by above half an ounce; a pint of water exceeds it by upwards of three ounces; and a pint of oil of vitriol weighs more than two pounds and a

quarter.

A table of the weights of certain measures of different fluids may on many occasions be useful, both for affisting the operator in regulating their proportions in certain cases, and for shewing the comparative gravities of the fluids themselves. I have therefore drawn up such a table for a pint, an ounce, and a dram measure of those liquids, whose gravity has been determined by experiments that can be relied on. The wine gallon contains 231 cubic inches; whence the pint contains 287; the ounce  $1\frac{1}{128}$ ; and the dram  $1\frac{2}{14}$  of a cubic inch.

		Ounce Dram
•	Pint weighs	
		weighs weighs
•	ounces drams grains	grains grains
T C C	ounce: drams grains	grains grains
INPLAMMABLE SPIRITS.		
Æthereal Spirit of Wine	11 1 36	336 42
Highly-rectified Spirit of Wine	12 5 20	
Common-rectified Spirit of Wine	13 2 40	
Proof Spirit	14 1 36	
Dulcified Spirit of Salt		
Dulcified Spirit of Nitre	14 4 48	
Dutchied Spirit of Petits	15 2 40	460 57 <del>1</del>
***	i :	
Wines.		
Burgundy	14 1 36	426 53
Red port	15 1 36	
Canary	15 6 40	
	1, 0 4	475 59\$
Expressed Oils.		
Oil Olive		
	1400	420 523
Linked Oil	1428	428 531
,	1	
Essential Oils.	i	
Oil of Turpentine	12 1 4	364 45 <del>1</del>
of Orange Peel	·- • ¶	
of Juniper Berries		1 - 1
	, ,	419   52
<b>D</b> 3		Oil

	Pint	weig	gh:	Ounce meaf.	Dram mesf.
•				weighs	weighs
Oil of Rosemary	onucee	drams	grains	430 432 436 443 443 448 450 457 458 476	weighs suitable 4 544 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
of Sassafras	}			503	63
Al'KALINE LIQUORS.  Lixivium faponarium, Pharm. Lond.  Spirit of Sal ammoniae	16 17 17 24	6	( 10 <b>2</b> 4 (	534	60 64 <del>1</del> 67 90
ACID LIQUORS.  Wine Vinegar	15 15 17 20 28	2	44 56 C 40 20	476 525 610	58 59½ 65½ 76 107¼
Animal Fluids.  Urine	15 15 16 16	o	20 40 0	475	59 59½ 60 60½
WATERS.  Distilled water	15 15 15 15 214	3 5	50 40 12 20 20	460 462	57 57½ 58 59 805

# [ 39 ]

#### CHAPTER IV.

## Of the Pharmaceutical Operations.

#### SECT. I.

#### Solution.

SOLUTION is an intimate commixture of folid bodies with fluids into one feemingly homogene liquor. The dissolving fluid is called a MENSTRUUM OF SOLVENT.

The principal menstrua made use of in pharmacy, are, water, vi-

nous spirits, oils, acid and alkaline liquors.

WATER is the menstruum of all salts, of vegetable gums, and of animal gellies. Of salts, it dissolves only a determinate quantity, though of one kind of salt more than another; and being thus saturated, leaves any additional quantity of the same salt untouched.

Experiments have been made for determining the quantities of water which different falts require for their diffolution: Mr. Eller has given a large set in the Memoirs of the royal academy of sciences of Berlin for the year 1750, from which the following table is extracted.

Eight ounces by weight of distilled water dissolved.

	OZ.	dr.	gr.
Of Refined Sugar	24	0	o ·
Green Vitriol	9	4	0
Blue Vitriol	Í	ò	0
White Vitriol	4	4	0
Epfom Salt	1 4	ö	0
Purified Nitre	1 4	0	0
Soluble Tartar	1 4	0	0
Common Sait	3	4	0
Sal gemmæ	3	4	0
Sal catharticus Glauberi	3	4	0
Seignette's Salt	3	ŏ	ō
Alum	2	4	o
Sal ammoniac	1	Ā	0
Vitriolated Tartar	1 -	4	٥
Salt of Hartshorn	1	4	0
Sugar of Lead	1 :	2	0
Cream of Tartar	1 :	õ	0
Borax	۱ .	4	
DVIRA	. 0	4	20

Though

Though great care appears to have been taken in making these experiments, it is not to be expected, that the proportions of the several falts, soluble in a certain quantity of water, will always be sound exactly the same with those above set down. Salts differ in their solubility, according to the degree of their purity, perfection, and dryness: the vitriols, and the artificial compound salts in general, differ remarkably in this respect, according as they are more or less impregnated with the acid ingredient. Thus vitriolated tartar, perfectly neutralized, is extremely difficult of folution: the matter which remains in making Glauber's spirit of Nitre (See Part III. Chap. viii. Sect. 6.) is no other than a vitriolated tartar, and it dissolves. fo difficultly, that the operator is obliged to break the retort in order to get it out; but on adding more of the vitriolic acid, it dissolves with ease. Hence many have been tempted to use an over-proportion of acid in this preparation; and we frequently find in the shops, under the name of vitriolated tartar, this acid soluble salt. gree of heat occasions also a notable difference in the quantity of falt taken up: in very cold weather, eight ounces of water will dissolve only about one ounce of nitre; whereas, in warm weather, the fame quantity will take up three ounces or more. To these circumstances are probably owing, in great part, the remarkable differences in the proportional folubilities of falts, as determined by different authors: it is observable, that common salt is less affected in its solubility, by a variation of heat, than any other, water in a temperate state dissolving nearly as much of it as very hot water; and accordingly this is the falt in which the different experiments agree the best: in the experiments of Hoffmann, Neumann, and Petit, the proportion of this falt, on a reduction of the numbers, comes out exactly the same, viz. three ounces of the salt to eight of water; Dr. Brownrigg makes the quantity of falt a little more; Dr. Grew, a dram and a scruple more; and Eller, as appears in the above table, four drams more; so that in the trials of fix different persons, made probably in different circumstances, the greatest difference is only one-fixth of the whole quantity of falt; whereas in some other salts there are differences of twice or thrice the quantity of the falt. In the experiments from which the table is drawn, the water was of the temperature of between 40 and 42 degrees of Fahrenheit's thermometer; or above freezing by about one-seventh of the interval between freezing and the human heat.

Some falts omitted by Eller are here subjoined: the first is taken

from Dr. Grew, and the other four from Neumann.

#### Eight ounces of water dissolved.

Of fixed alkaline Salt	•	above	02. 8	dr. O	gr. O	ĺ
Sal diureticus	•		8	0	0	ĺ
Sugar-candy, both brown and white	-	•	9	0	0	ı
Sugar of Milk	-	-	0	2	40	ı
Essential Salt of Sorrel	•	-	0	I	20	ł

Though water takes up only a certain quantity of one kind of falt, yet when faturated with one, it will fill dissolve some portion of another; and when it can bear no more of either of these, it will still take up a third, without letting go any of the former. The principal experiments of this kind, that have been made, relative to pharmaceutic subjects, are exhibited in the following table; of which the two sirst articles are from Grew, and the others from Eller.

#### Water, 32 parts by weight

fully faturated with	Diffolved afterwards.	
Nitre	Sal ammoniac - 10	
Common Salt	Nitre 10 Sal ammoniae - 2	
Nitre	Fixed Alkali - 7   Common Salt - 2	
Common Salt -	Nitre, near - 2 Fixed Alkali - 23	
	Nitre 4 Sugar 2	
Sal ammoniac -	Common Salt - $2\frac{1}{2}$	
Soluble Tartar -	Nitre 2	
	Fixed Alkali - 2	
Glauber's Salt -	Nitre 1 Sugar 1	
Epfom Salt	Sugar 6	
Borax	Fixed Alkali - 2	

In regard to the other class of bodies which water is a menstruum for, viz. those of the gummy and gelatinous kind, there is no determinate point of saturation: the water unites readily with any proportions of them, forming, with different quantities, liquors of different consistencies. This sluid takes up likewise, when assisted by triruration, the vegetable gummy resins, as ammoniacum and myrrh; the solutions of which, though imperfect, that is, not transparent, but turbid, and of a milky hue, are nevertheless applicable to valuable purposes in medicine. It mingles with vinous spirits, with acid and alkaline liquors, not with oils, but imbibes some of the more subtle parts of essential oils, so as to become impregnated with their smell and taste.

Rectified SPIRIT OF WINE is the menstruum of the essential oils, refins, and camphor of vegetables; of the pure distilled oils, and several

feveral of the colouring and medicinal parts of animals; of some mineral bituminous substances, as of ambergris; and of soaps, though it does not act upon the expressed oil and fixed alkaline salt, of which soap is composed; whence, if soap contains any superstuous quantity of either the oil or salt, it may, by means of this menstruum, be excellently purished therefrom. It dissolves, by the affistance of heat, volatile alkaline salts, and, more readily, the neutral ones, composed either of sixed alkali and the actous acid, as the sal diureticus, or of volatile alkali and the nitrous acid, as also the salt of amber, &c. It mingles with water and with acids, not with alkaline lixivia.

Oils dissolve vegetable refins and balfams, wax, animal fats, mineral bitumens, sulphur, and certain metallic substances, particularly lead. The expressed oils are, for most of these bodies, more powerful menstrua than those obtained by distillation; as the former are more capable of sustaining, without injury, a strong heat, which is, in most cases, necessary to enable them to act. It is said, that one ounce of sulphur will dissolve in three ounces of expressed oil, particularly that of linseed, but requires six ounces of essential oil, as that of turpentine.

All Agids diffolve alkaline falts, alkaline earths, and metallic fubstances. The different acids differ greatly in their action upon these last; one diffolving only some particular metals, and another others.

The vegetable acids dissolve a considerable quantity of zinc, iron, copper, lead, and tin, and extract so much from the metallic part of antimony, as to become powerfully emetic: they dissolve lead more readily, if the metal is previously calcined by fire, than in its metallic state.

The marine acid dissolves zinc, iron, and copper; and though it scarce acts on any other metallic substance in the common way of making solutions, it may nevertheless be artfully combined with them all except gold: the corrosive sublimate, and antimonial caustic of the shops, are combinations of it with mercury and the metallic part of antimony, effected by applying the acid, in the form of sume, to the subjects, at the same time also strongly heated.

The nitrous acid is the common mentruum of all metallic subflances, except gold and the metallic part of antimony; of which two the proper solvent is a mixture of the nitrous and marine acids,

called aqua regia.

The vitriolic acid, diluted with water, easily dissolves zinc and iron: in its concentrated state, and assisted by a boiling heat, it may be made to corrode, or impersectly dissolve, most of the other metals.

ALKALINE

ALKALINE lixivia dissolve oils, resinous substances, and sulphur. Their power is greatly promoted by the addition of quick lime; instances of which occur in the preparation of soap, and in the common caustic. Thus actuated, they reduce the slesh, bones, and other solid parts of animals, into a gelatinous matter.

Solutions made in water, and in spirit of wine, possess the virtues of the body dissolved; whilst oils generally sheathe its activity, and acids and alkalies vary its quality. Hence watery and spirituous liquors are the proper mensions of the native virtues of vegetable and

animal matter.

Most of the foregoing folutions are easily effected, by pouring the menstruum on the body to be dissolved, and suffering them to stand together, for some time, exposed to a suitable warmth. A strong heat is generally requisite to enable oils and alkaline liquors to perform their office: nor will acids act on some metallic bodies without its assistance. The action of watery and spirituous menstrua is likewise expedited by a moderate heat; though the quantity, which they afterwards keep dissolved, is not, as some suppose, by this means increased: all that heat occasions these to take up, more than they would do in a longer time in the cold, will, when the heat ceases, subside again: this at least is most commonly the case, though there may be some instances of the contrary.

The action of acids on the bodies which they dissolve, is generally accompanied with heat, effervescence, and a copious discharge of sumes. The sumes which arise during the dissolution of some metals in the vitriolic acid, prove inflammable: hence, in the preparation of the artificial vitriols of iron and zinc, the operator ought to be careful, especially where the solution is made in a narrow-mouthed vessel, lest, by the imprudent approach of a candle, the exhaling va-

pour be set on fire.

There is another species of solution, in which the moisture of the air is the menstruum. Fixed alkaline salts, and those of the neutral kind, composed of alkaline salts and the vegetable acids, or of soluble earths and any acid, except the vitriolic, and some metallic salts, on being exposed, for some time, to a moist air, gradually attract its humidity, and, at length, become siquid. Some substances, not disoluble by the application of water in its grosser form, as the butting antimony, are easily siquesied by this slow action of the aerial moisture. This process is termed Delignation.

## SECT. II.

#### Extraction.

THE liquors which dissolve certain substances in their pure state, serve likewise to extract them from admixtures of other matter. Thus rectified spirit of wine, the menstruum of effential

oils and refins, takes up the virtues of the refinous and oily vegetables; as water does those of the mucilaginous and saline; the inactive earthy parts remaining untouched by both. Water extracts likewise, from many plants, substances which by themselves it has little effect upon; even essential oils being, as we have formerly observed, rendered soluble in that sluid, by the admixture of gummy and saline matter, of which all vegetables participate in a greater or less degree. Thus many of the aromatic plants, and most of the bitters and astringents, yield their virtues to this menstruum.

Extraction is performed, by MACERATING OF STEEPING the fubject in its appropriated menstruum, in the cold; or DIGESTING or CIRCULATING them, in a moderate warmth; or INFUSING the plant in the boiling liquor, and suffering them to stand in a covered vessel till grown cold; or actually BOILING them together for some

time.

The term digestion is sometimes used for maceration; and in this case the process is directed to be performed without heat: where this circumstance is not expressed, digestion always implies the use of heat. Circulation differs from digestion only in this, that the steam. into which a part of the liquor is resolved by the heat, is, by means of a proper disposition of the vessels, condensed and conveyed back again upon the subject. Digestion is usually performed in a matrals (or bolthead) Florence flask, or the like; either of which may be converted into a circulatory weffel, by inverting another into the mouth, and fecuring the juncture with a piece of wet bladder. A fingle matrals, if its neck is very long and narrow, will answer the purpole as effectually; the vapour cooling and condensing before it can rise to the top. In a vessel of this kind, even spirit of wine, one of the most volatile liquors we know of, may be boiled without any confiderable loss. The use of this instrument is likewise free from an inconvenience, which may, in some cases, attend the other, of the uppermost vessel being burst or thrown off. As the longnecked matrasses, here recommended, are difficultly filled or emptied. and likewise very dear, a long glass pipe may be occasionally luted to the shorter ones.

Heat greatly expedites extraction; but by this means proves as injurious to some substances, by occasioning the monstruum to take up their grosser and more ungrateful parts, as it is necessary for enabling it to extract the virtues of others. Thus guiacum or logwood impart little to aqueous liquors, without a boiling heat, whilst even a small degree of warmth proves greatly prejudicial to the fine bitter of carduus benedictus. This plant, which, infused in boiling, or digested in sensibly hot water, gives out a nauseous taste, so offensive to the stomach as to promote vomiting, yields to the cold element a grateful balsamic bitter, the most elegant stomachic of the shops.

As heat promotes the diffolving power of liquids; fo cold, on the other hand, diminishes it. Hence tinctures, or extractions made by a considerable heat, deposite in cold weather a part of their contents, and thus become proportionably weaker; a circumstance which deferves particular regard.

#### SECT. TTT.

## Depuration.

HERE are different methods of depurating or purifying liquors from their feculencies, according as the liquor itself is more or less tenacious, or the feculent matter of greater or less gravity.

Thin fluids readily deposite their more ponderous impurities, upon standing at rest, for some time, in a cool place; and may then be pr-

CANTED, or poured off clear, by inclining the veffel.

Glutinous, unctuous, or thick substances, are to be liquested by a fuitable heat; when the groffer feculencies will fall to the bottom, the lighter arising to the furface, to be DESPUMATED, or scummed off.

Where the impurities are neither fo ponderous as to subfide freely to the bottom, nor so light as to arise readily to the surface, they may be separated in great measure by COLATURE through strainers of linen, woollen, or other cloth; and more perfectly by FILTRATION through

a fost bibulous kind of paper made for this use.

The grey paper which covers pill-boxes as they come from abroad, is one of the best for this purpose: it does not easily break when wetted, or tinge the liquor which passes through it, which the reddish fort, called bloffom paper, frequently does. The paper is supported by a sunnel, or piece of canvas fixed in a frame. When the funnel is used, it is convenient to put some straws or small sticks between the paper and its sides, to prevent the weight of the liquor from pressing the paper so close to it, as not to allow room for this fluid to transude. In some cases a funnel made of wire is put betwixt the paper and the glass sunnel. There is also a kind of glass funnel, with ridges down its sides, made on purpose for

Glutinous and unctuous liquors, which do not easily pass through the pores of a filter or strainer, are CLARIFIED, by beating them up with whites of eggs, which concreting or growing hard when heated, and entangling the impure matter, arise with it to the surface : the mixture is to be gently boiled, till the scum begins to break, when the vessel is to be removed from the fire, the crust taken off, and the liquor passed through a flannel bag.

Decantation,

Decantation, colature, and filtration, are applicable to most of the medicated liquors that stand in need of purisscation. Despumation and clarification very rarely have place; fince these, along with the impurities of the liquor, frequently separate its medicinal parts. Thus, if the decocion of poppy heads, for making diacodium, be solicitously scummed or clarified (as some have been accustomed to do) the medicine will lose almost all that the poppies communicated, and, instead of a mild opiate, turn out little other than a plain syrup of sugar.

It may be proper to observe, that the common sorts of filtering paper are apt to communicate a disagreeable flavour: and hence in filtering fine bitters, or other liquors, whose gratefulness is of primary consequence, the part which passes through first ought to be

kept apart for inferior purpofes.

#### SECT. IV.

## Crystallization.

ATER, affished by heat, dissolves a larger proportion of faline substances, than it can retain when grown cold: hence, on the abatement of the heat, a part of the salt separates from the menstruum, and concretes at the sides and bottom of the vessel. These concretions, unless too hastily formed by the sudden cooling of the liquor, or dissurbed in their coalescence by agitation, or other like causes, prove transparent, and of regular sigures, resembling in

appearance the natural sprig CRYSTALS.

Salts, dissolved in a large quantity of water, may, in like manner, be recovered from it in their crystalline form, by boiling down the solution, till so much of the sluid has exhaled, as that the remainder will be too little to keep the salt dissolved when grown perfectly cold. It is customary to continue the evaporation, till the salt shews a disposition to concrete even from the hot water, by forming a pellicle on that which is least hot, viz. on the surface. If large, beautiful, and perfectly sigured crystals are required, this point is somewhat too late: for if the salt thus begins to coalesce whilst considerably hot, on being removed into a cold place, its particles will run too hallily and irregularly together; the pellicle at the same time salling down through the liquor, and thus proving a farther disturbance to the regularity of the crystallization.

In order to perform this process in perfection, the evaporation must be gentle, and continued no longer than till some drops of the liquor, let fall on a cold glass plate, discover crystalline filaments. When this mark of sufficient exhalation appears, the vefsel is to be immed ately removed from the fire into a less warm, but not cold place, and covered with a cloth, to prevent the access of cold

air, and confequently the formation of a pellicle.

All the alkaline falts are excluded from this operation; fixed alkalies never affuming a crystalline form, and the volatile ones escaping before the mentruum exhales. Some even of the neutral kind, particularly those, of which certain metallic bodies are the bass, are so strongly retained by the aqueous sluid, as not to exhibit any appearance of crystallization, unless some other substance be added, with which the water has a greater affinity. The table of affinity shews, that such a substance is spirit of wine; by the prudent addition of which, these kinds of salts separate freely from the menstruum, and form large and beautiful crystals, scarcely attainable by any other means.

The operator must be careful not to add too much of the spirit; lest, instead of a gradual and regular crystallization, the basis of the salt be hastily precipitated in a powdery form. One twentieth part of the weight of the liquor will, in most cases, be a sufficient, and

in some too large a quantity.

Different salts require different quantities of water to keep them dissolved: and hence, if a mixture of two or more be dissolved in this sluid, they will begin to separate and crystallize at different periods of the evaporation. Upon this soundation, salts are freed, not only from such impurities, as water is not capable of dissolving and carrying through the pores of a filter, but likewise from admixtures of one another; that which requires most water to dissolve in, shooting first into crystals.

## SECT. V.

## Precipitation.

BY this operation, bodies are recovered from their solutions, by means of the addition of some other substance, with which either the menstruum, or the body dissolved, have a greater affinity than they

have with one another.

Precipitation, therefore, is of two kinds; one, where the substance superadded unites with the menstruum, and occasions that before dissolved to be thrown down: the other, in which it unites with the dissolved body, and falls along with it to the bottom. Of the first we have an example in the precipitation of sulphur from alkaline lixivia by the means of acids; of the second, in the precipitation of mercury from aqua-fortis by sea-falt, or its acid.

The subjects of this operation, as well those which are capable of being precipitated as those which precipitate them, will readily appear from inspection of the table of affinity. The manner of performing it is so simple, as not to stand in need of any particular di-

rections;

rections; no more being required, than to add the precipitant by degrees, so long as it continues to occasion any precipitation. When the whole of the powder has fallen, it is to be well EDULCORATED, that is, washed in several fresh parcels of water, and asterwards dried for use.

Where metals are employed as precipitants, as in the purification of martial vitriol from copper by the addition of fresh iron, they ought to be persectly clean and free from any rusty or greasy matter; otherwise they will not readily, if at all, dissolve, and consequently the precipitation will not succeed: for the substance to be precipitated separates only by the additional one dissolving and taking its place. The separated powder, oftentimes, instead of falling to the bottom, lodges upon the precipitant; from which it must be occasi-

onally shaken off, for reasons sufficiently obvious.

Though, in this operation, the precipitated powder is generally the part required for use, yet some advantage may frequently be made of the liquor remaining after the precipitation. Thus when fixed alkaline salt is dissolved in water, and sulphur dissolved in this lixivium, the addition of acids separates and throws down the sulphur, only in virtue of the acid uniting with, and neutralizing the alkali by which the sulphur was held dissolved: confequently, if the precipitation is made with the vitriolic acid, and the acid gradually dropped in till the alkali is completely satiated, that is, so long as it continues to occasion any precipitation or turbidness, the liquor will yield, by proper evaporation and crystallization, a neutral salt, composed of the vitriolic acid and fixed alkali, that is, vitriolated tartar. In like manner, if the precipitation is made with the nitrous acid, a true nitre may be recovered from the liquor; if with the marine, the salt called spiritus salis marini congulatus; and if with the acid of vinegar, the sal diureticus.

## SECT. VI.

## Evaporation.

Hit S is a third method of recovering folid bodies from their folutions, effected by the means of heat; which evaporating the fluid part, that is, forcing it off in steam, the matter which was

dissolved therein is left behind in its solid form

This process is applicable to the solutions of all those substances which are less volatile than the menstruum, or which will not exhale by the heat requisite for the evaporation of the sluid; as the solutions of fixed alkaline salts; of the gummy, gelatinous, and other inodorous parts of vegetables and animals in water; and of many resinous and odorous substances in spirit of wine.

Water

t

Water extracts the virtues of fundry fragrant aromatic herbs, almost as perfectly as rectified spirit of wine; but the aqueous insusions are far from being equally suited to this process, with those made in spirit; water carrying off the whole odour and slavour of the subject, which that lighter liquor leaves entire behind it. Thus a watery insusion of mint loses in evaporation the smell, taste, and virtues of the herb; whilst a tincture drawn with pure spirit, yields, on the same treatment, a thick balsamic liquid, or solid gummy resin, extremely

rich in the peculiar qualities of the mint.

In evaporating these kinds of liquors, particular care must be had, towards the end of the process, that the heat be very gentle; otherwise the matter as it grows thick will burn to the vessel, and contract a disagreeable smell and taste: this burned slavour is called an empyreuma. The liquor ought to be kept stirring during the evaporation; otherwise a part of the matter concretes on the surface exposed to the air, and forms a pellicle which impedes the farther evaporation. More particular directions for performing this operation to the greatest advantage, will be given hereafter in the second part.

## SECT. VII.

# Distillation.

IN the foregoing operation fluids are rarefied by heat into steam or vapour, which is suffered to exhale in the air, but which the business of this is to collect and preserve. For this purpose the steam is received in proper vessels, luted to that in which the subject is contained, and being there cooled, condenses into a sluid form again.

There are two kinds of distillation: by the one, the more subtile and volatile parts of liquors are elevated from the grosser; by the other, liquids, incorporated with solid bodies, are forced out from

them by vehemence of fire.

To the first belong, the distillation of the pure instammable spirit from vinous liquors; and of such of the active parts of vegetables as are capable of being extracted by boiling water or spirit, and at the

fame time of arising along with their steam.

As boiling water extracts or dissolves the effential oils of vegetables, whilst blended with the other principles of the subject, without saturation, but imbibes only a determinate, and that a small proportion of them in their pure state; as these oils are the only substances, contained in common vegetables, which prove totally volatile in that degree of heat, and as it is in them, that the virtues of aromatics, and the peculiar odour and slavour of all plants reside: it is evident, that water may be impregnated, by distillation, with the more valuable E

parts of many vegetables: that this impregnation is limited, the oil arifing in this process pure from those parts of the plant which before rendered it soluble in water without limitation; hence greatest part of the oil separates from the distilled aqueous liquor, and according to its greater or less gravity, either sinks to the bottom or swims on the surface: that consequently insusions and distilled waters are greatly different from one another: that the first may be rendered stronger and stronger by pouring the liquor on fresh parcels of the subject; but that the latter cannot be in like manner improved by cobabiting, or redistilling them from fresh ingredients. See Part II. Chap. v. Sect. 2.

As the oils of many vegetables do not freely distil with a less hear, than that in which water boils; as rectified spirit of wine is not susceptible of this degree of heat; and as this menstruum totally dissolves these oils in their pure state: it follows, that spirit elevates far less from most vegetables than water; but that nevertheless the distilled spirit, by keeping all that it does elevate perfectly dissolved, may, in in some cases, prove as strong of the subject as the distilled water.

The apparatus made use of for distilling spirits, waters, and oils, consists of a still, or copper vessel, for containing the subject, on which is luted a large bead with a swan neck. The vapour arising into the head, is hence conveyed through a worm, or long spiral pipe, placed in a vessel of cold water called a refrigeratory; and being there condensed, runs down into a receiver. In the second part of this work, we shall give some improvements in this apparatus for particular purposes; with directions for performing the several processes to the greatest advantage.

It may be observed, that as the parts which are preserved in evaporation cannot arise in distillation, the liquor remaining after the distillation, properly depurated and inspissated, will yield the same extracts as those prepared from the tincture or decoction of the subject made on purpose for that use; the one of these operations collecting only the volatile parts, and the other the more fixed; so that where one subject contains medicinal parts of both kinds, they may thus be obtained distinct, without one being injured by the process which col-

lects the other.

THE subjects of the second kind of distillation are, the gross oils of vegetables and animals, the mineral acid spirits, and the metallic suid quicksilver; which as they require a much stronger degree of heat to elevate them than the foregoing liquors can sustain, so they likewise condense without arising so far from the action of the sire. The distillation of these is performed in low glass vessels, called from their neck being bent to one side, resorts: to the farther end of the neck a receiver is luted, which standing without the surnace the vapours soon condense in it, without the use of a refrigeratory: nevertheless, to promote this effect, some are accustomed, especially in warm

warm weather, to cool the receiver, by occasionally applying wet cloths

to it, or keeping it partly immerfed in a veffel of cold water.

The vapours of some substances are so sluggish, or strongly retained by a mixed matter, as scarce to arise even over the low neck of the retort. These are most commodiously distilled in streight-necked earthen vessels, called longnecks, laid on their sides, so that the vapour passes off laterally with little or no ascent: a receiver is luted to the end of the neck without the furnace: in this manner, the acid spirit of vitriol is distilled. The matter which remains in the retort or longneck, after the distillation, is vulgarly called caput mortuum.

In these distillations, a quantity of elastic air is frequently generated; which, unless an exit is allowed it, blows off, or bursts the receiver. The danger of this may in good measure be prevented, by slowly raising the sire; but more effectually, by leaving a small hole in the luting, to be occasionally opened or stopped with a wooden plug; or inserting at the juncture an upright pipe of such a height, that the steam of the distilling liquor may not be able to rise to the

top.

# SECT. VIII.

## Sublimation.

As all fluids are volatile by heat, and confequently capable of being separated, in most cases, from fixed matters, by the foregoing process; so various solid bodies as subjected to a similar treatment. Fluids are said to distil, and solids to sublime; though sometimes both are obtained in one and the same operation. If the subliming matter concretes into a mass, it is commonly called a sublimate; if into a powdery form, slowers.

The principal subjects of this operation are, volatile alkaline salts; neutral salts composed of volatile alkalies and acids, as sal ammoniac; the salt of amber, and slowers of benzoine; mercurial preparations; and sulphur. Bodies of themselves not volatile are frequently made to sublime by the mixture of volatile ones: thus iron is carried up by sal

ammoniac in the preparation of the flores martiales.

The fumes of folid bodies, in close vessels, rise but little way, and adhere to that part of the vessel where they concrete. Hence a receiver or condenser is less necessary here than in the preceding operation; a single vessel, as a matrass, or tall vial, or the like, being frequently sufficient. The most commodious apparatus for the sublimation of particular substances, and the most advantageous method of conducting the several processes, will be delivered in the second part.

## SECT. IX.

# Expression.

HE press is chiefly made use of forcing out the juices of succulent herbs and fruits; and the insipid oils of the uncluous seeds and kernels.

The harder fruits, as quinces, require to be previously well beat or ground; but herbs are to be only moderately bruised. The subject is then included in a hair bag, and pressed betwixt wooden plates, in the common screw-press, as long as any juice runs from it. See Part III. Chap. ii.

THE expression of oils is performed nearly in the same manuer as that of juices; only here, iron plates are substituted to the wooden ones there made use of. The subject is well pounded, and included in a strong canvass bag, betwixt which, and the plates of the press a hair cloth is interposed.

The infipid oils of all the unctuous feeds are obtained, uninjured, by this operation, if performed without the use of heat; which though it greatly promotes the extraction of the oil, at the same time impresses an ungrateful flavour, and increases its disposition to grow

rancid.

The oils expressed from aromatic substances generally carry with them a portion of their essential oil: hence the smell and slavour of the expressed oils of nutmegs and mace. They are very rarely found impregnated with any of the other qualities of the subject: oil of nustrard seed, for instance, is as soft and void of acrimony, as that of almonds, the pungency of the mustard remaining entire in the cake left after the expression.

## SECT. X.

# Exficcation.

HERE are two general methods of exficcating or drying moint bodies: in the one, their humid parts are exhaled by heat; is the other, they are imbibed or absorbed by substances, whose soft and spongy texture adapts them to that use. Bodies intimately combined with, or dissolved in a sluid, as recent vegetables and their juices, require the first: such as are only superficially mixed, as when earthy or indissoluble powders are ground with water, are commodiously separated from it by the second.

Vegetables

Vegetables and their parts are usually exficcated by the natural warmth of the air: the affistance of a gentle artificial heat, may, nevertheless, in general, be not only safely, but advantageously had recourse to. By a moderate fire, even the more tender flowers may be dried, in a little time, without any considerable loss, either of their odour or lively colour: which would, both, be greatly injured or destroyed, by a more slow exsiccation in the air. Some plants indeed, particularly those of the acrid kind, as horse-radish, scurvy-grass, and arum, loose their virtues by this process, however carefully performed: but far the greater number retain them unimpaired, and oftentimes improved

improved.

The thicker vegetable juices may be exsiccated by the heat of the sun; or, where this is not sufficient, by that of a water-bath, or an oven moderately warm. The thinner juices may be gently boiled till they begin to thicken, and then treated as the foregoing; this process, termed INSPISSATION OF EVAPORATION has been spoken of already. The juices of some plants, as arum root, briony root, orris root, wild cucumbers, &c. separate, upon standing for some time, into a thick part, which falls to the bottom; and a thin aqueous one, which swims above it: this last is to be poured off, and the sirst exsiccated by a gentle warmth: preparations of this kind have been usually called PECULE; that of the wild cucumber, to be spoken of in its place, is the only one which practice now retains.

Indistoluble bodies mixed with water into a thick consistence, may be easily freed from the greatest of it, by dropping them on a chalk stone, or some powdered chalk pressed into a smooth mass, which readily imbibes their humidity. Where the quantity of sluid is large, as in the edulcoration of precipitates, it may be separated by decan-

tation or filtration.

## SECT. XI.

#### Comminution.

OMMINUTION is the bare reduction of folid coherent bodies into small particles or powder. The methods of effecting this are various, according to the texture of the subject.

Dry friable bodies, are such as are brittle and not very hard, and mixtures of these with somewhat moist ones, are easily PULVERIZED

in a mortar.

For very light, dry substances, resins, and the roots of tenacious sexture, the mortar may in some cases be previously rubbed with a little sweet oil, or a sew drops of oil to be occasionally added: this prevents the sacr powder of the sirst from slying off, and the others from cohering under the pessle. Camphor is most commodiously powdered, by rubbing it with a little rectified spirit of wine.

e 3

Though

Though substances, as woods, the peels of oranges and lemons, &c. are most conveniently rasped; and soft oily bodies, as nutmegs,

passed through a grater.

The comminution of the harder minerals, as calamine, crystal, slint, &c. is greatly facilitated by EXTINCTION; that is, by heating them red hot, and quenching them in water: by repeating this process a few times, most of the hard stones become easily pulverable. This process, however, is not to be applied to any of the alkaline or calcareous stones; lest, instead of an insipid powder, we produce an acrimonious calx or lime.

Some metals, as tin, though strongly cohering in their natural state, prove extremely brittle when heated, insomuch as to be easily divided into small particles by dextrous agitation. Hence the officinal method of pulverizing tin, by melting it, and, at the instant of its beginning to return into a state of solidity, briskly shaking it in a wooden box. The comminution of metals, in this manner, is termed by the metallurgists GRANULATION.

On a similar principle, certain salts, as nitre, may be reduced into powder in a large quantity, by dissolving them in boiling water, setting the solution over a moderate fire, and keeping the salt constantly stirring during its exsecution, so as to prevent its particles, disjoined by

the fluid, from re-uniting together into larger masses.

Powders are reduced to a great degree of fineness by TRITURATING, or rubbing them, for a length of time, in a mortar. Such as are not dissoluble in water, or injured by the admixture of that sluid, are moistened with it into the consistence of a paste, and Levigated, or ground, on a stat smooth marble or iron plate; or where a large quan-

tity is to be prepared at a time, in mills made for that use.

Comminution, though one of the most simple operations of pharmacy, has, in many cases, very considerable effects. The resumble purgatives, when sinely triturated, are more easily soluble in the animal sluids, and consequently prove more cathartic, and less irritating, than in their grosser state. Crude antimony, which, when reduced to a tolerable sine powder, discovers little medicinal virtue, if levigated, to a great degree of subtility, proves a powerful alterative in many chronical disorders.

By comminution, the heaviest bodies may be made to float in the lightest suids, for a longer or shorter time, according to their greater

Some attribute this effect to a diminution of the specific gravity of the body; and, at the same time, suppose the peculiar virtues of certain medicines, particularly mercury, to be in great measure owing to their gravity. If these hypotheses were just, it should follow, that the mercurial preparations, by being sinely comminuted, would lose proportionably of their efficacy; and so indeed mercurius dulcis, for instance,

greater or less degree of tenuity. Hence we are furnished with an excellent criterion of the fineness of certain powders, and a method of separating the more subtile parts from the grosser, distinguished by the name of ELUTRIATION, or washing over. See Part III. Chap. i.

#### SECT. XIL

## Fusion.

PUSION is the reduction of folid bodies into a state of sluidity by fire. Almost all natural substances, the pure earths, and the solid parts of animals and vegetables excepted, melt in proper degrees of fire; some in a very gentle heat, whilst others require its utmost violence.

Turpentine, and other fost resinous substances, LIQUETY in a gentle warmth; wax, pitch, sulphur, and the mineral bitumens, require a heat too great for the hand to support, fixed alkaline salts, common salt, nitre, require a red, or almost white heat to MELT them; and

glass, a full white heat.

Among metallic substances, tin, bismuth, and lead, slow long before ignition: antimony likewise melts before it is visibly red-hot, but not before the vessel is considerably so: the regulus of antimony demands a much stronger sire. Zinc begins to melt in a red heat; gold and silver require a low white heat; copper a bright white heat; and

iron, an extreme white heat.

One body, rendered fluid by heat, becomes sometimes a menstruum for another, not susible of itself in the same degree of sire. Thus red-hot silver melts, on being thrown into melted lead less hot than itself: and thus is steel, heated to whiteness, be taken out of the surnace, and applied to a roll of sulphur; the sulphur instantly lique-sping, occasions the steel to melt with it; hence the chalybs cum sulphure of the shops. This concrete, nevertheless, remarkably impedes the susion of some other metals, as lead, which when united with a certain quantity of sulphur, is scarce to be perfectly melted by a very strong sire: hence the method, described in its place, of purifying zinc, a metal which sulphur has no effect upon, from the lead so frequuently mixed with it.

instance, has been supposed to do. But experience shews, that this is far from being the case; and that comminution by no means lessens, but rather increases its power: when reduced to a great degree of subtility, it passes readily into the habit, and operates, according to its quantity, as an alterative or stalogogue; whilst in a grosser form, it is apt to irritate the stomach and bowels, and run off by the intestines, without being conveyed into the blood.

Sulphur is the only unmetallic substance which mingles in suston with metals. Earthy, saline, and other like matters, even the calces and glasses prepared from metals themselves, float distinct upon the surface, and form what is called scoria or dross. Where the quantity of this is large in proportion to the metal, it is most commodiously separated by pouring the whole into a conical mould: the pure metal or regulus, though small in quantity, occupies a considerable height in the lower narrow part of the cone, and when congealed, may be easily freed from the scorize by a hammer. The mould should be previously greased, or rather smoked, to make the metal come freely out; and thoroughly dried and heated, to prevent the explosion which sometimes happens from the sudden contact of melted metals with moist bodies.

#### SECT. XIII.

#### Calcination.

PY calcination is understood, the reduction of solid bodies, by the means of sire, from a coherent to a powdery state, accompanied with a change of their quality; in which last respect, this process differs from comminution.

To this head belong, the burning of vegetable and animal matters, otherwise called ustrow, incineration, or concremation; and the change of metals into a powder, which in the fire either does not

melt, or VITRIPIES, that is, runs into glass.

The metals which melt before ignition, are calcined by keeping them in fusion for some time. The free admission of air is essentially necessary to the success of this operation; and hence, when the surface of the metal appears covered with calx, this must be taken off, or raked to one side; otherwise, the remainder, excluded from the air, will not undergo the change intended. If any coal, or other instammable matter that does not contain a mineral acid, be suffered to fall into the vessel, the effect expected from this operation will not be produced, and part of what is already calcined, will be REVIVED OF REDUCED; that is, it will return into its metallic form again.

Those metals which require a strong fire to melt in, calcine with a much less heat than is sufficient to make them slow. Hence the burning or SCORIFICATION of such iron or copper vessels, as are long exposed to a considerable sire without defence from the air. Gold and

filver are not calcinable by any degree of fire.

In calcination, the metals visibly emit sumes; nevertheless the weight of the calx proves greater than that of the metal employed. The antimonial regulus gains about one-eleventh part of its weight; zinc, sometimes one-tenth; tin, above one-sixth; and lead, in its conversion into minium, oftentimes one-fourth.

The

The calcination of metallic bodies (gold, filver, and mercury, excepted) is greatly promoted by nitre. This falt exposed to the fire in conjunction with any instammable substances, extricates their instammable matter, and bursts with it into slame, accompanied with a hissing noise: this process is usually termed DEFLAGRATION OF DETONATION.

All the metallic calces and scoriz are revived into their metallic flate, by fusion with any vegetable or animal inflammable matter. They are all more difficult of fusion than the respective metals them. selves; and scarcely any of them, those of lead and bismuth excepted, can be made to melt at all, without some addition, in the strongest fire that can be produced in the common furnaces. The additions. called fluxes, employed for promoting the fusion, confist chiefly of fixed alkaline falts: a mixture of alkaline falt with inflammable matter, as powdered charcoal, is called a reducing flux, as contributing at the same time to bring the calx into fusion and to revive it into metal. Such a mixture is commonly prepared from one part of nitre, and two parts of tartar, by grinding them well together, fetting the powders on fire with a bit of coal, or a red-hot iron, then covering the vessel, and suffering them to deslagrate or burn, till they are changed into a black alkaline coaly mass. This is the common reducing flux of the chemists, and is called from its colour, the black flux. Metallic calces, or scoriz, mingled with twice their weight of this compound. and exposed to a proper fire, in a close covered crucible, melt and resume their metallic form; but though they received an increase of weight in the calcination, the revived metal is always found to weigh confiderably less than the quantity which the calx was made from.

-. • • - ·

## PART II.

THE

# MATERIA MEDICA.

RITERS on the Materia Medica have taken great pains in arranging the warious articles, of which it is composed, into different divisions and subdivisions, according to their real

or reputed medicinal powers.

It has been imagined, that, "the whole Materia Medica is reducible under the three diffinctions of alteratives, evacuants, and reftoratives: the first comprehending all that has any power to alter the
constitution, without sensibly increasing or diminishing any of the
natural evacuations; the second, whatever visiby promotes those
discharges; and the third, all that contributes to lessen them, and
make the increase greater than the waste." These divisions being
too general, they are broken into subdivisions; and these again are
further divided into different classes, under more restrained denominations, as cardiac, carminative, hysteric, stomachic, &c.

Specious as this plan may appear to be, I am afraid that the execution of it, to any useful purpose, would require a far more extensive knowledge of the nature and operation of medicines, than has yet been attained to. A just and useful method of simples is scarcely to be expected, while those properties, on which the method is founded,

are imperfectly known, and in many articles only conjectural.

In all the arguments that have been hitherto contrived upon this plan, there appears a striking incongruity among the several articles of which even the ultimate subdivisions are composed; substances extremely dissimilar being classed together, as cantharides and tea, to-bacco and bran, hemlock and cowslips, scurvigrass and rains, arum root and liquorice, wormwood and parsneps, cinnamon and nettles, rasherries and chalk, artichokes and alum, cloves and coffee, mustard seed and black cherries, &c. Nor are these incongruities to be laid always to the charge of the authors; the nature of the system itself rendering

rendering them often unavoidable: for the particular effect, which entitles a medicine to a particular class, may be produced by substances very dissimilar, and even opposite in their general powers: thus the alvine excretions are restrained by starch, wax, tormentil root, opium: among the capital diuretics are cantharides, nitre, fixt alkaline salts, squills. It should seem, that the method of arrangement cannot be a just one, which requires substances so discordant to be ranked together; and which further requires each of these substances to be ranked over again, in other classes, along with other substances to which they are equally discordant.

There is also a material impersection in this scheme, even in the primary divisions. Steel and its preparations act, in different circumstances, both as evacuants and restoratives. Mercury and antimony afford, in their different preparations, both evacuants and alteratives; and there are many other drugs which are sometimes used as alteratives, and sometimes as evacuants: indeed all evacuants, in diminished doses, seem to act merely as alteratives. It should seem therefore that "the division of the whole Materia Medica into alteratives, "evacuants, and restoratives," is a division not sounded in nature, even if there was no objection to the vague meaning of the appellati-

ons themselves.

Carthenfer has divided the Maseria Medica on a plan which appears more rational. Instead of the operations of medicines in the human body, which are precarious, complicated, and greatly diversified according to the desie, the preparation, and the ciscumstances of the patient, he takes for the basis of his arrangement, their more simple, obvious, and constant properties, as bitterness, sweetness, astringency, acidity, &c. Having considered the nature of bitterness, for instance, in general, he examines what effects medicines possessed of this property are capable of producing in the body, and in what ciscumstances they may be expected to be serviceable, and then proceeds as

an account of the particular bitters.

This method is of real use; but its use is limitted to a small part of the Materia Medica. There are many of the medicinal simples, in which we can distinguish no prevailing qualities of this kind, there are many, in which different qualities are blended together; and many, which though similar in these kinds of qualities, are very different are in their operations in the human body: thus though gentian and aloes agree in having a bitter taste, and sagar and manna in being sweet, their medicinal virtues are respectively very different. Accordingly the author is obliged in some cases to depart from his general plan, and sound the division on the medicinal effects: he makes one class of puspatives and emetics, and another of vaporose inchriants and narcotics: this last class consists of tobacco, elder-slowers, saffran, opium, and poppy-seeds, substances certainly very discordant in all their qualities that relate to medicinal intentions.

In this work, instead of attempting a medicinal distribution of the simples, which I apprehend not to be practicable to any good purpose, and which, as hitherto executed, seems more likely to missead the reader than to promote true knowledge, I shall take them in the order of the alphabet; and even in this order we shall seldom perhaps find substances more dissimilar come together, than those which have been joined into one class by some of the systematic writers. It may be proper, however, to premise some general observations on certain classes of medicines, in Cartheuser's manner, and thus to preserve the less exceptionable parts of his plan, with some amendments.

#### Acips.

Class 1. Vegetable

[native; as forrel, wood forrel, juice of lemons, oranges, barberries, and other fruits.

produced by fermentation; as vinegar, and tartar.

Class 2. Mineral: the acids of vitrial, nitre, and common falt.

THE medical effects of acids. duly diluted and given in proper doses, are, to cool, quench thirft, correct a tendency to putrefaction, and allay inordinate motions of the blood. By these qualities, in hot bilious temperaments and inflammatory diforders, they frequently reftrain immoderate hæmorrhages, and promote the natural secretions; in some kinds of fevers, they excite a copious diaphoresis, where the warm medicines, called alexipharmac, tend rather to prevent this salutary difcharge.

Vegetable acids, particularly the native juices of certain plants and fruits, have fome degree of a faponaceous quality; by means of which they attenuate or diffolve viscid phlegm, and deterge the vessels; and thus prove serviceable in sundry chronical disorders. Inveterate scurvies have sometimes yielded to their continued use, especially when given in conjunction

with medicines of the acrid or pungent kind: experience has shewn, that the acrid antiscorbutics have much better effects when thus managed, than when exhibited by themselves; hence in the fucci scorbutici of our dispensatory, Seville orange juice is usefully joined to that of the cochlearia and nasturtium.

The mineral acids inftantly coagulate blood: the vegetable dilute it, even when inspissated or thickened by heat; in which state, watery liquors will not mingle with it. Hence in some fevers, where water runs off by the kidneys almost as pale and insipid as it was drank, vegetable acids render the urine of the due colour and quality. The mineral acids (the spirit of nitre in particular) combined with vinous spirits, have a like effect.

Acids are prejudicial in cold, pale, phlegmatic habits, where the vessels are lax, the circulation languid languid, bile deficient, and the power of digestion weak. In these cases, an acid is often generated in the stomach, from milk and most vegetable soods, which, whilst it

continues in the first passages, occasions uneasiness about the stomach, statulencies, sometimes griping pains of the bowels, and vomitings.

## INSIPID EARTHS capable of ABSORBING ACIDS.

Oyster shells, Crabs claws, and eyes so called, Coral, red and white, Pearls, Bezoar, Chalk, Some marles, Lime stones, Marbles, Spars.

THE virtues of these substances are, to absorb or destroy acidities in the first passages, and consequently to remove such diforders as proceed from that cause. The cordial, alexipharmac, antifebrile, and other like virtues attributed to these medicines, appear to have little foundation; or at best, are only secondary ones. When united with the acid, they form a neutral faline compound, possessing some degree of an aperient and detergent quality, though too inconsiderable to be in general regarded.

The absorbent earths were all strangers to medicine in the earlier times; and their use does not seem to have been established before the last century; when some practitioners, from an opinion that most kinds of disease proceeded from a preternatural acid, introduced a great variety of antacid bodies, both of the earthy and saline kind; and very liberally exhibited them on almost every occasion.

It is certain that in children, and adults of a weak conflictation, and whose food is chiefly of the vegetable acescent kind, sundry disorders are occasioned by accidities; these readily discover themselves by sour erustations, the pale colour of the face, and in children by the sour smell and green colour of the

alvine feces, which are fometimes fo manifefuly acid as to raife a strong effervescence with alkaline falts. In these cases, and these only, the use of absorbent earths is indicated.

If there are really no acid juices in the ventricle, these earths are apt to concrete with the mucous matter usually lodged there, into hard indissoluble masses; which have fometimes been thrown up by vomit, or found in the stomach upon diffection. Hence indigeftion, loss of appetite, nausez, vomiting, obstructions of the bowels, and other diforders. Sometimes the stomach and intestines have been found lined with a crust, as it were, of these earthy bodies. which must not only have prevented the separation of the gastric liquor, but likewise closed the orifices of the lacteal vessels, so as to obstruct the passage of the chyle into the mass of blood.

Some suppose the earthy powders capable (without the concurrence of any acid) of passing the lacteals along with the chyle; and alledge, in support of this opinion, that when friturated with water, they are in part taken up, and carried with it through a filter of paper; the filtrated liquor leaving, upon evaporation, a portion of whitish earthy matter. This ex-

periment

periment (allowing the consequence to be justly drawn from it) is itself erroneous; the residuum proceeds from the earth naturally contained in the water, not from that employed in the experiment; for if pure distilled water be made use of, it will leave no residuum, though long triturated, or digested with the earth.

All these bodies, particularly those of the animal kind, contain, besides their purely alkaline earth, a portion of glutinous matter. instance of this we have in crabs eyes, which if macerated in the weaker acids, or the stronger sufficiently diluted with water; the earthy part will be dissolved, and the animal glue remain in form of a soft transparent mucilage. glutinous substance increases their tendency to concrete in the stomach; and hence those which contain least thereof should be preferred to the others. The mineral earths contain the least of this kind of matter, and some of them are very easy of solution; chalk for instance, which may therefore be given with greater safety than the animal absorbents. These substances, divested of their conglutinating matter by means of site, are reduced into acrimonious calces or limes, and thus become medicines of a different class.

The teeth, bones, hoo's, and horns of animals, confist of the fame principles with the animal absorbents above mentioned, but combined in different proportions: the quantity of gelatinous matter is so large, as to defend the earthy part from the action of weak acids; whilt the earth, in its turn, protects the gluten from being easily diffolved by watery liquors. Hence these bodies in their crude state, though recommended as possessing singular virtues, are not found to have any virtue at all.

Experiments have been made for determining the degree of folubility, or comparative itrength of these earths; the principal of which are arranged in the two following tables, one taken from Langius, and the other from Homberg.

Table of the quantity of acid destroyed by different absorbents.

Ten grains of	Some kinds of Limestones Oyster shells Chalk Shells of Garden Snails Calcined Cray Fish Pearl Tooth of the Sea Horse Volatile salts Fixt salts Coral, red and white Crabs eyes Eggshells Mother of Pearl Crabs claws Jawbone of the Pike fish	deflioyed the acidity of	160 120 100 100 80 80 80 60 60 50 50 40	Drops of Spirit of Salt.
---------------	---	--------------------------	--	-----------------------------

Table

Table of the quantity of absorbent earths soluble in acids.

576 grains of Spirit of Salt- diffolved of	Crabs eyes Mother of Pearl Pearls Oyfter shells Hartshorn Coral Oriental Bezoar Occidental Bezoar Quick Lime Slacked Lime	216 144 128 156 165 186 118 123 199
576 grains of Spirit of Nitred diffolved of	Crabs eyes Mother of Pearl Pearls Oyfter shells Hartshorn Coral Oriental Bezoar Occidental Bezoar Quick Lime Slacked Lime	297 202 219 236 234 233 108 144 180 216

These experiments do not sufficiently ascertain the point intended by them: in the first fet, the quantity of acid is too vague and indetermined: in the second, we are not told whether the acid was perfectly saturated; and in both, the acids made use of were so very different from any that can be supposed ever to exist in the human body, that little can be concluded from them with regard to the medical effects of these Trial should have absorbents. been made with the mild vegetable acids, as the juices of certain fruits. four fermented liquors; or rather with four milk. Nevertheless these tables, though not fo perfect as could be wished, have their use in the hands of such as can make proper allowances. See the experimental history of the Materia Medica, page 557.

EARTHS NOT DISSOLUBLE in acids, or other liquors.

The earths of this kind may be ranged in two classes.

Class 1. Hard eryftalline earths: as the ruby, garnet, emerald, fapphire, hyacinth, and other precious stones; crystal, flint, &c.

HESE kinds of substances attributed to them, by the superstiwere introduced into medi- tion of the earlier ages. Some of cine, and many fabulous virtues them are still preserved in foreign phar-

sharmacopæias, but at length very uftly expunged from ours, notrithstanding what some late wriers of repute speak of their medical These indissoluble hard irtue. podies are not capable of producng any other effect, than by their igid angular particles (which tho' vigated with the utmost care, the microscope still discovers in them) to offend or wound the intestines. . In levigation, they wear off fo much from the hardest marble instruments, as will equal or exceed their own weight: from this circumstance we may account for their having sometimes appeared to act as absorbents. Some of these flones, exposed to a vehement fire, become in some measure friable; but nevertheless remain indissoluble. Most of the coloured

ones by this treatment lose their colour; and in this state, prove nearly of the fame quality with common crystal: such are, the sapphire, emerald, amethyst, and cornelian. Others melt into a blackish vitreous matter, from which a portion of iron is obtainable by proper fluxes; as the hyacinth and garnet. Geoffroy concludes from hence, that these stones really possess some medical virtues, depending upon their metallic part; but the quantity of metallic matter, sufficient to give them a confiderable tinct, is so exceedingly small, and so inclosed in a stony matter not at all foluble by any of the known menstrua, as scarce to admit of any possibility of its acting in the human body.

## Class 2. Softer earths; the talky, gypseous, and argillaceous.

The tales and gypsums have rarely been used as medicines. Some of the former, from their unctuous softness and filver hue, stand recommended externally as cosmetics; and some of the latter, on little better foundation, internally as astringents. But they have long been deservedly rejected by the judicious practitioners. They seem to possess the ill qualities of the alkaline earths (concreting with the mucus of the stomach, &c.) without any of their good ones.

Several of the clays, boles, and terræ figillatæ were highly celebrated by the ancients as aftringents and alexipharmacs, and some of them still continue in esteem; though it is certain they have no great claim to the virtues that have been attributed to them. Their real effects are, to give a greater degree of consistency to the shuids

in the first passages, and in some measure defend the solids from their acrimony.

Most of these bodies contain. besides the tenacious indissoluble earth, which is their principal characteristic, (1) a portion of an earth soluble in acids, similar to those of the first section; (2) of acid, separable by distillation in a strong fire; this acid is always of the fame nature with that obtained from vitriol, fulphur, and alum (3). The coloured ones contain likewife small quantities of iron, reducible, by inflammable fluxes, into its metallic form. In confequence of the first of these ingredients, these earths may be looked upon in some measure as absorbent; the acid appears to be united with a part of the absorbent earth, into a faline compound. approaching to an aluminous nature; whence they have some degree of astringency; whether they receive any peculiar virtue from the iron, is greatly to be doubted; fince it is in a very crude state, and in quantity extremely small.

These earths unite with water into a turbid liquor, slippery and smooth to the touch, and remain for some time suspended; the sand, grit, or other grosser matters which are often found naturally mingled with them, subsiding. They may be freed by means of acids from their alkaline earth;

by coction in water, from their faline matter; and the coloured ones from their iron by digeftion in aquaregis, the only menstruum we are acquainted with that will extract the ferrugineous matter of argillaceous and bolarearths. Thus purified, they have all nearly the same appearance and qualities. Exposed to a strong sire, they lose their soft glutinous quality, and are reduced into hard masses indiffoluble as at first.

GLUTINOUS, vegetable, and animal substances.

Class 1. Vegetable.

Pure gums:
Tragacanth,
Senica,
The gums of cherry, plum,
and other European trees.

UMS and mucilages are glutinous vegetable productions, of no particular tafte or smell, soluble in water, but not in vinous spirits, or in oils: see-p. 13. They differ from one another, only in degree of tenacity: the more tenacious are called gums; those which are less so, mucilages. The first naturally exude from certain trees and shrubs; the latter are extracted by art. Almost all vegetable substances contain some por-

Vegetables abounding with mucilage:
Orchis roots.
Althæ root,
Quince feeds, &c.

tion of these, which after the resinous part has been extracted by spirit, may be separated from the remaining matter by means of water.

The general virtues of these kinds of substances are, to thicken the sluids, and defend the solids from them, when grown sharp or corrosive. Hence their use in a thin acrimonious state of the juices, and where the natural mucus of the intestines is abraded.

#### Class 2. Animal.

Most animal substances (the fat excepted) contain a viscous matter, in many respects similar to the foregoing, and capable of being extracted by strong coction in water.

Animal glues and gellies have the general qualities of the vegetable gums and mucilages; with this difference, that the former are more nutrimental, and apt to run into a putrid state. Considered as the subjects of chemistry, the difference betwixt them is very great: those of the animal kind are changed by fire into a volatile alkaline salt, and a fetid oil; the vegetable into an acid liquor, and a very small portion of oily matter, considerably less fetid than the former.

## Soft uncruous substances.

Class 1. Insipid wegetable oils; and substances abounding with them, as almonds, and the kernels of most fruits; linseed, and the medullary part of sundry other seeds.

## Class 2. Animal fats; as spermaceti.

Nctuous vegetables unite with water, by trituration, into a milky liquor: and give out their oil upon expression.—These kinds of oils, and animal fats, dissolve not in any menstruum except alkaline ones; which change their quality, and reduce them into a soap, dissoluble in water, but more perfectly in vinous spirits: from this compound, the oil may, by a skilful addition of acids, be recovered in a purer state than before, and rendered soluble, like essential oils, in spirit of wine: see p. 10.

The medical virtues of these subflances are, to obtund acrimonious humours, and to soften and relax

the folids: hence their use internally, in tickling coughs, heat of urine, pains and inflammations; and externally in tension and rigidity of particular parts. The milky folutions, commonly called emulfions, though much less emollient than the oils themselves or animal fats, have this advantage, that they may be given in acute or inflammatory distempers, without danger of the ill consequences which the others might fometimes produce: fats and oils, kept in a degree of heat no greater than that of the human body, foon become rancid and acrimonious; whilst emulfions tend rather to grow four.

#### Astringents.

Galls, Tormentil root, Bistort root,

A Stringent substances are diflinguished by a rough austere taste; and changing solutions of iron, especially those made in the vitriolic acid, of a dark purple or black colour.

Aftringents yield their virtues by infusion both to water and vinous spirits, generally in greatest perfection to the former. Oils extract nothing from them: nor do they give over any of their virtue in distillation: nevertheless, their aftringency is considerably abated Balaustines, Terra Japonica, Acacia, &c.

by evaporating decoctions of them to the confifence of an extract; and totally destroyed by long keep-

The medical effects of these kinds of substances are, to constring the sibres, and incrassate, or lightly thicken the juices. Their more experienced use is in disorders proceeding from a debility, or slaccid state, of the folids; in hæmorrhages, from a thinness of the blood, laxity or rupture of the vessels; in preternatural F 2

discharges of other kinds, after the offending matter has been duly corrected, or evacuated; and in external relaxations.

In some cases, they produce the effects of aperients; the vessels, constringed and strengthened by them, being enabled to protrude the circulating juices with greater sorce.

A good deal of caution is requifite in the use of these medicines, especially those of the more powerful kind. In plethoric habits, inveterate obstructions, critical evacuations, and in all kinds of fluxes in general before the morbific matter has been expelled, or where there is any stricture or spasmodic contraction of the vessels; aftringents prove eminently hurtful. Where critical dysenteries or diarrheas are restrained by styptics, the acrimonious matter, now confined in the intestines, corrodes or instames them; and sometimes occasions a gangrene of the parts.

SWEETS.

Sugar, Honey, Raisins, Liquorice, &c.

wery numerous tribe; almost every plant that has been examined, discovering, in some of its parts, a saccharine juice. The bottoms of slowers, and most kinds of seeds and grain when they begin to vegetate, are remarkably sweet.

Vegetable sweets are extracted both by water and vinous spirits, most readily by the first, but in greatest persection by the latter. Nothing of their taste arises in distillation with either of these liquors: nevertheless, by long boiling with water they become somewhat less agreeable; but are not much injured by being treated in the same manner with rectified spirit.

The purer sweets, as sugar, promote the union of distilled oils with watery liquors, and prevent the separation of the butyraceous part from milk: from this quality they are supposed to unite the unctuous part of the food with the animal juices. Hence some have concluded, that they increase fat: others, that they have a contrary

effect, by preventing the separation of the unctuous matter which forms the fat, from the blood; and others, that they render the juices thicker and more fluggish, retard the circulation and cuticular excretion, and thus bring on a variety of disorders. But sweets have not been found to produce any of these effects, in any remarkable degree: common experience shews, that their moderate, and even liberal use, is at least innocent; that they reconcile, not only to the palate, but the flomach also, substances of themselves disguftful to both; and thus render falutary what would otherwise be injurious to the body.

The unctuous and mucilaginous fweets, as the impure fugars, liquorice, &c. have a confiderable degree of emollient and lubricating virtue.—Those, accompanied with a manifest acid, as in the juices of most sweet fruits, are remarkably relaxing; and if taken immoderately, occasion diarrhæz and dysenteries, which sometimes

have proved fatal.

ACRIDS.

#### ACRIDS.

Crids are substances of a penetrating pungency. Applied to the skin, they inflame or exulce- ed up the nose, provoke sneezing.

rate it: chewed, they occasion a copious discharge of saliva: and snuff-

These substances, considered as the subjects of pharmacy, may be divided into three classes.

yielding their acrimony

( 1. In distillation with water : as horse-radish. mustard, scurvy-grass, &c.

2. By infusion only: as the greater celandine, pyrethrum, &c.

3. Neither to infusion, nor distillation: as arum and dracunculus.

The general effects of acrid medicines are, to stimulate the vessels, and dissolve tenacious juices. In cold leucophlegmatic habits, stagnations of the fluids, and where the contractile power of the folids is weak, they prove powerful expectorants, deobstruents, diurerics, and emmenagognes; and if the patient is kept warm, sudorifics. In hot bilious constitutions, plethoric habits, inflammatory distempers, where there is already a degree of irritation, where the juices are too thin and acrimonious, or the viscera unsound; these stimulating' medicines prove highly prejudicial, and never fail to aggravate the disease.

Certain acrid substances have been lately recommended in dry convultive althmas: of the efficacy of the squill in particular, for the cure of this disorder, several instances are related in the Commercium Literarium of Norimberg for the years 1737 and 1739. theuser thinks, that not the asthma itself, but a particular effect of it, was removed by this medicine. He observes, that in all asthmas, the free circulation of the blood through the pulmonary vessels, is impeded; and hence, during every paroxysm, the lungs are in a kind of cedematous state: that if this ædema, becoming habitual, remains after the fit is over, it is either perpetually occasioning fresh ones, or gives rise to a dropfy of the breast: that acrid medicines, by removing the cedema, remove what was originally an effect of the asthma, and in time a cause of its aggravation.

# AROMATICS.

Romatics are substances of a 🖊 warm pungent taste, and a more or less fragrant smell. Some of the spices are purely aromatic. as cubebs, pepper, cloves; some substances have a sweetness mixed with the aromatic matter, as angelica root, aniseed, fennel seed;

fome an astringency, as cinnamon; some a strong mucilage, as casialignea: fome a bitterness, as orange peel. The aromatic matter itself. contained in different subjects, differs also not a little in its pharmaceutic properties. It is extracted from all by rectified spirit of wine; F 3 from from some in great part, from others scarcely at all, by water. The aromatic matter of some subjects, as of lemon peel, rises wholly in distillation both with spirit and water; that of others, as cinnamon, rises wholly with water, but scarcely at all with spirit; while that of others, as pepper, is in part left behind after the distillation of water itself from the spice.

With regard to the general vir-

tues of aromatics, they warm the stomach, and by degrees the whole habit, raise the pulse, and quicken the circulation. In cold languid cases, phlegmatic habits, and a weak flaccid state of the solids, they support the vis vitæ, and promote the salutary secretions. In hot bilious temperaments, plethoric habits, inflammatory indispositions, dryness and strictures of the sibres, they are generally hurtful.

#### BITTERS.

Gentian root, Hops,

Lesser centuary, Carduus, &c.

BItters for the most part yield their virtue both to watery and spirituous menstrua: some more persectly to one, and others to the other. None of the substances of this class give over any thing considerable of their taste in distillation, either to water or to spirit; their bitterness remaining entire, and frequently improved, in the extracts. Such as are accompanied with slavour, as wormwood, may by this process be reduced into simple slavourless bitters.

These substances participate of the virtues of astringents and aromatics. Their general effects are, to constringe the fibres of the stomach and intestines, to warm the habit, attenuate the bile and juices in the first passages, and promote the natural evacuations, particularly of sweat and urine. In weakness of the stomach, loss of appetite, indigestion, and the like dis-

orders, proceeding from a laxity of the folids, or cold fluggish indisposition of the juices, these kinds of medicines do good service. Where the fibres are already too tense and rigid, where there is any immoderate heat or inflammation. bitters very sensibly increase the distemper; and if their use is continued, communicate it to the kidneys: hence the urine becomes high coloured, small in quantity, and at length suppressed; a dropsy foon fucceeding. If the kidneys were before fo lax, as to remain now uninjured, yet the other vifcera become gradually more and more rigid, and a tabes is at length brought on.

Bitter substances destroy insects, and prevent putresaction. Hence they are recommended as anthelmintic; and externally as antisep-

tics.

## EMETICS and CATHARTICS.

Hellebore, Jalap, Ipecacoanha,

THESE substances consist of a refinous part, in which the purgative or emetic quality reColocynth, Scammony, Gamboge, &c.

fides; and a gummy faline one, which acts chiefly as a diuretic. The first is extracted or dissolved by

by vinous spirits: the latter by water. Nothing arises in distillation from either.

The acrid refins, exhibited by themselves, tenaciously adhere to the coats of the intestines, by their stimulating power irritate and inflame them, and thus produce fundry violent disorders. Hoffman relates, that he has sometimes obferved convulsions, and a paralysis of both fides, from their use.

These inconvenienciès may be avoided, by previously triturating them with substances capable of dividing their tenacious texture, and preventing their adhesion; by this means, they become mild and safe, operate without disturbance, and at the same time more effectually answer the purposes intended

by them.

Some have endeavoured to correct the ill quality of the refinous purgatives, by the addition of acids and aromatic oils. Acids weaken their power, but have no other effect than what a diminution of the dose would equally answer. The purgent essential oils may serve to warm the stomach, make the medicine fit easier, and thus prevent the nausea, which sometimes happens; but as soon as the resin begins to exert itself in the intestines, these oils, instead of correcting, increase its virulence; being themfelves apt to occasion the inconveniencies which they are here intended to prevent, an irritation and

inflammation of the bowels. kaline salts or soaps have a better effect; as they dispose the resin to folution, and promote its operation.

The medicines of this class seem to act by liquefying the juices, and stimulating the coats of the stomach and intestines. If the irritation is ftrong and sudden, their action is quick and upwards: if flower, downwards. Cathartics given in a liquid form, or in very fensible habits, often prove emetic; and emetics, where mucus abounds. cathartic. They operate more violently in robust constitutions, than in those of a contrary temperament; the vessels being in the former more tense and rigid, and consequently less capable of bearing an equal degree of irritation.

The action of these medicines is extended beyond the primæ viæ: This appears evident from the increase of the pulse which always accompanies their operation; and from the common observation of children being purged by the milk, if the nurse has taken a cathartic. Some of them, particularly hellebore, are faid to purge, if only applied externally in iffues .- Purgatives, even of the more powerful kind, exhibited in suitable small doses, in conjunction with the milder aperients, may be introduced into the habit, so as to prove notable deobstruents, diuretics, and diaphoretics, without acting fenfibly by stool.

THE foregoing observations are inserted, not with any view to a method of simples, but to give a general idea of the virtues of such medicinal substances as are possessed of the qualities which make the objects of the respective articles. I shall dwell no longer on general reflections, but proceed to an account of each of the simples separately.

ABIETIS lignum, summitates, coni: maris C. B. vel Abietis tenuiore folio Abietis conis sursum spectantibus sive frustu deorsum spectante Tourn.

filver and red fir; their wood, tops, and cones.

These are large ever-green trees, frequent in the northern climates. The first is said to be found wild in some parts of England, and the fecond on the hills of Scotland. From these trees, in different parts of Germany, the Strasburgh turpentine is extracted, of which here-The wood, and the fruit or cones gathered about the end of autumn, abound with refinous matter; and yield, in distillation with water, an effential oil, not greatly different from that obtained by the same means from turpentine - The wood and tops of the fir trees, on account of their refinous juice, are fometimes employed in decoctions and diet drinks, for promoting urine and fweat, purifying the blood and juices, and cleanfing and healing internal ulcerations, particularly those of the urinary passages. See the article Terebinthina.

ABROTANI MARIS folia:
Abrotani maris angusti folii majoris
C. B. Southernwood; the leaves

[E.]

This is a shrubby plant, clothed with very finely divided leaves of a greyish green colour; the slowers, which are very small and yellowish, hang downwards, several together, from the middle of the branches to the top. It is a native of the warmer countries; in this it is cultivated in gardens: the leaves fall off every winter: the roots and stalks abide many years.

Southernwood has a strong, not very disagreeable smell; and a nauseous, pungent, bitter taste; which is totally extracted by rectified spirit, less persectly by watery liquors. It is recommended as an anthelmintic; and in cold

leucophlegmatic habits, as a fimulant, detergent, aperient, and sudorific. The present practice has almost entirely confined its use to external applications. The leaves are frequently employed in discutient and antisceptic fomentations: and have been recommended also in lotions and unguents for cutaneous eruptions, and the falling off of the hair.

ABROTANI FEMINÆ folia:
Abrotani fæminæ foliis teretibus C.
B. Lavender-cotton; the leaves
[E]

This plant is all over white and hoary: the leaves are composed of small knows set in rows along a middle rib; the slowers stand upright on the tops of the stalks. It is raised in gardens, slowers in June and July, and holds its leaves

all the winter.

The abrotanum femina is supposed to possess the same virtues with the mas; but in a less degree. For external purposes, the medical difference betwixt them is not very great; hence in fomentations (which is the principal intention they are usually applied to) the London college allows either to be taken instead of the other.—The abrotanum femina is recommended by fome in hysteric and other female complaints: it has been customary among the common people to use a decoction of it in milk against worms.

ABSINTHII VULGARIS folia: Al finthii vulgaris majoris J. B. Common wormwood; the leaves

[L.E.]

The leaves of this fort of wormwood are divided into roundish fegments, of a dull green colour above, and whitish underneath. It grows wild in several parts of England; about London, large quan-

tities

tities are cultivated for medicinal use: it flowers in June and July; and after having ripened its seeds, dies down to the ground, excepting a tust of the lower leaves, which generally abides the winter.

Wormwood is a strong bitter; and was formerly much used as fuch, against weakness of the stomach, and the like, in medicated wines and ales. At present it is rarely employed in these intentions, on account of the ill relish and offensive smell which it is These it may accompanied with. be in part freed from by keeping, and totally by long coction, the bitter remaining entire. An extract made by boiling the leaves in a large quantity of water, and evaporating the liquor with a strong fire, proves a bitter sufficiently grateful, without any disgustful flavour .- An oil distilled from this plant [L. E.] and an extract [E.] are kept in the shops.

# ABSINTHII MARITIMI fummitates: Abfintbii marini albi

Gerard. Sea Wormwood, commonly, but falfely, called Roman wormwood; the tops [L].

The leaves of sea wormwood are much smaller than those of the common, and hoary on the upper side, as well as the lower; the stalks also are hoary all over. It grows wild about our falt marshes, and in several parts about the sea coasts.—In taste and smell, it is weaker and less unpleasant than the common wormwood. The virtues of both are supposed to be of the same kind, and to differ only in degree.

The tops enter three of our diftilled waters, and give name to a conferve [L.] They are an ingredient also in the common fomention and green oil [L.]

ABSINTHII ROMANI folia:

Absinthii pontici tenuifolii incani C. B. Roman wormwood; the leaves and tops [E.]

This species is very different in appearance from the two foregoing: it is in all its parts smaller than either; the leaves are divided into fine filaments, and heary on the lower fide; the flalks, either entirely or in part, of a purplish hue. It is a native of the warmer countries, and at present difficultly procurable in this, though as hardy and as eafily raifed as any of the other forts. Sea wormwood has long supplied its place in the markets, and been in general miftaken for it.

Roman wormwood is less ungrateful than either of the others: its smell is tolerably pleasant: the taste, though manifestly bitter, scarce disagreeable. It appears to be the most eligible of the three as a stomachic; and is likewise recommended by some in dropsies.

ACACÍA [L. E.]: the inspissed juice of the unripe fruit of a large prickly tree, called by Caspar Bauhine, Acacia foliis scorpioidis leguminose.

The juice is brought to us from Egypt, in roundish masses, wrapt up in thin bladders. It is outwardly of a deep brown colour, inclining to black; inwardly of a reddish or yellowish brown; of a firm consistence, but not very dry. It soon softens in the mouth, and discovers a rough, not disagreeable taste, which is followed by a sweetish relish. This inspissated juice entirely dissolves in watery liquors; but is scarce sensibly acted on by rectified spirit.

Acacia is a mild affringent medicine. The Egyptians give it in spitting of blood, in the quantity of a dram, dissolved in any conve-

nient

nient liquor; and repeat this dose occasionally: they likewise employ it in collyria for strengthening the eyes, and in gargarisms for quinseys. Among us, it is little otherwise used than as an ingredient in mithridate and theriaca [L.] and is rarely met with in the shops. What is usually sold for the Egyptian acacia, is the inspisated juice of unripe sloes: this is harder, heavier, of a darker colour, and somewhat sharper taste, than the true sort.

ACANTHI folia: Acanthi fativi vel mollis Virgilii C. B. Brankurfine; the leaves.

This is a beautiful plant, growing naturally in Italy, and other warm climates: from its leaves, the ancients took the patterns of their foliage works. All the parts of it have a foft sweetish taste, and abound with a mucilaginous juice: its virtues do not seem to differ from those of althea and other mucilaginous plants.

ACETOSÆ vulgaris, sive oxalidis, folia& radix: Acetosæ arvensis C. B. Oxalidis vulgaris foliolongo J. B. Common forrel; the

roots and leaves [E.]

Sorrel grows wild in fields and meadows throughout England. The leaves have a restringent acid tafte, without any fmell or particular flavour: their medical effects are, to cool, quench thirst, and promote the urinary discharge: a decoction of them in whey affords an useful and agreeable drink in febrile or inflammatory diforders: and is recommended by Boerhaave to be used in the spring as one of the most efficacious aperients and detergents. Some kinds of scurvies have yielded to the continued use of this medicine: the Greenlanders, who are very fubject to this distemper, are said to employ, with good success, a mixture of the jucices of sorrel and of scurvygrass. The only officinal preparation of this plant, is an essential salt from the juice of the leaves [E]

The roots of forrel have a bitterish austere taste, without any acidity: they are said to be deobstruent and diuretic; and have fometimes had a place in aperient apozems, to which they impart a

reddish colour.

The feeds of this plant were formerly used in diarrhœas and dysenteries, but have long been strangers to the shops, and are now justly expunged both from the London and Edinburgh pharmacopæias: they have no remarkable smell, and scarcely any taste.

ACETOSELLA [E.] vide Lu-

ACETUM [L. E.] Vinegar: an acid produced from fermented vinous liquors by a fecond fermen-

tation. See page 6.

Wine vinegar is confiderably purer than that prepared from malt liquors; the latter, however acid and fine, contains a large portion of a viscous mucilaginous substance: as is evident from the ropyness and slimyness which this kind of vinegar is very much fubject to: the stronger and more spirituous the wine, the better and stronger vinegar it yields. French vinegars are faid by Geoffroy to faturate above one-thirtyfifth their weight of fixt alkaline falt, and some of them no less than one-twelfth; the best of the German vinegars little more than one fortieth.

Vinegar is a medicine of excellent use in all kinds of inflammatory and putrid disorders, either internal internal or external: in ardent, bilious fevers, pestilential, and other malignant distempers, it is recommended by Boerhaave as one of the most certain sudorifics, (see the section of acids, page 61.) Weakness, fainting, vomiting, hysterical, and hypochondriacal complaints, have been frequently relieved by vinegar applied to the mouth and nose, or received into the stomach.

ACORUS, vide CALAMUS AROMATICUS.

ADIANTHI VERI feu capilli Veneris folia: Adianthi folio coriandri C. B. True maidenhair; the leaves.

This is a low evergreen herb, and one of those which, from the slenderness of their stalks, are called capillary. It is a native of Italy, and the southern parts of France; from whence the leaves are sometimes brought to us. These have an agreeable, but very weak, smell; and a mucilaginous somewhat roughish taste, which they readily impart to boiling water.

Maidenhair has been greatly celebrated in diforders of the breaft, proceeding from a thinness and acrimony of the juices; and likewife for opening obstructions of the viscera, and promoting the expectoration of tough phlegm. But modern practice pays little regard to it: nor is it often to be met with in the shops; the TRICHOMANES, or English maidenbair, which is of the same quality, generally supplying its place.

AERUGO [L. E.] Verdegris. This is a preparation of copper, made chiefly at Montpellier in France, by firatifying copper plates with grape stalks that have been

impregnated with a fermented vegetable acid: in a few days, the plates are found covered with a pale green downy matter, which is scraped off from the copper, and the process again repeated.

Verdigris, as it comes to us, is generally mingled with stalks of the grape; these may be separated, in pulverization, by discontinuing the operation as soon as what remains seems to be almost entirely composed of them.

Verdegris is rarely or never used internally. Some writers greatly extol it as an emetic, and fay, that a grain or two being taken. acts as foon as received into the stomach; but its use has been too often followed by dangerous consequences. (See the article Cu-RPUM.)-Verdegrisapplied externally, proves a gentle detergent and escharotic, and serves to take down fungous flesh arising in wounds. In these intentions, it is an ingredient in the mel appptiacum, unquentum bafilicum viride [L.] and bulfamum viride [E.]

AGALLOCHUM seu lignum aloes. Aloes wood.

There have been different conjectures concerning this wood; but no fatisfactory account of it has hitherto appeared. Authors diftinguish several forts of agallochum, most of which are strangers to Europe. That which comes to us is in little hard ponderous pieces, of a yellowish brown co. lour, with several black or purplish veins. It has a bitterish aromatic taste; and a fragrant fmell, especially if reduced to powder, or set on fire. Distilled with water, it affords a very fragrant essential oil, but in small quantity; digested in rectified spirit, it yields an elegant tincture, which loses nothing valuable in being evaporated to the confishence of an extract.

Agallochum is at present of verylittle use in medicine, and rarely to be met with in the shops: if it could be easily procured, at least the better sort of it, it bids fair to be a very useful cordial; Hossman greatly recommends, in this intention, the distilled oil and spirituous tincture; and esteems a mixture of this last with tincture of steel an excellent corroborant.

AGARICUS: Agaricus five fungus laricis C. B. Agaric; a fungus growing on old larch trees

This fungus is an irregular fpongy substance, extremely light, and of an uniform snowy white-ness (except the cortical part, which is usually taken off before the agaric is brought into the shops). It cuts freely with a knife, without discovering any hardness or gritiness, and readily crumbles betwixt the singers into a powder. It has no remarkable smell; its taste is at first sweetish; but on chewing for a little while, proves acrid, bitter, and nauseous.

Agaric was formerly in great efteem as a cathartic; but the present practice has almost entirely rejected its use. It operates exceeding flowly, infomuch that fome have denied it to have any purga-Given in subtive virtue at all. stance, it almost always occasions a nausea, not unfrequently vomiting, and fometimes excessive tormina of the bowels; these effects are attributed to its light farinaceous matter adhering to the coats of the intestines, and producing a The best preconstant irritation. paration of agaric seems to be an extract made with water, in which fixt alkaline salt has been dissolved; or with vinegar or wine; the first is faid by Boulduc, and the two latter by Newmann, to prove effectual and safe purgatives. Nevertheless this is at best a precarious medicine, which we stand in no manner of need of; hence the college have justly rejected it from all the compositions which it formerly had a place in, except the mithridate and theriaca [L.]

AGARICUS penis equini facie Tourn. Female agaric, or agaric of the oak, called, from its being very easily inflammable, touchwood, or spunk.

This fungus is frequently met with, on different kinds of trees. in England; and is faid to have been fometimes brought into the shops mixt with the true agaric of the larch; from this it is easily distinguishable by its greater weight, dusky colour, and mucilaginous taste, void of bitterness. The medullary part of this fungus, beaten soft, and applied externally, has been of late greatly celebrated as a flyptic, and faid to restrain not only venal but arterial hæmorrhages, without the use of ligatures. It does not appear however to have any real flyptic power, or to act any otherwise than dry lint, sponge, or other soft. fungous applications.

AGERATI folia et flores: Agerati foliis ferratis C. B. Ptarmice lueæ fuaveolentis Tourn. Maudlin; the leaves and flowers.

This is a flender plant, clothed all over with narrow ferrated leaves. It is a native of Italy, and other warm countries; with us, it is raifed in gardens, and flowers in July and August.

Maudlin has a light agreeable fmell; and a roughish, somewhat warm and bitterish taste. These qualities point out its use as a mild

-01109

corroborant; but it has long been a stranger to practice, and is now omitted both by the London and Edinburgh colleges.

AGNI CASTI femen: Agni folio non ferrato J. B. The chaste tree; its seeds.

This is a small tree, or rather shrub, growing spontaneously in Italy, &c. and raised with us in gardens. Its fruit, which is about the size of a pepper corn, contains four longish seeds, which are said to be of an aromatic smell, and an acrid bitterish taste, but which are found on examination to be almost inodorous and insipid. These seeds have been celebrated as antaphrodisacs; but experience does not warrant their having any such virtues.

AGRIMONIÆ folia: Eupatorii weterum feu agrimoniæ C. B. Agrimony; the leaves [E.]

This is a common plant in hedges, and the borders of fields. The leaves have an herbaceous, fomewhat acrid, roughift tafte, accompanied with an aromatic flavour. Agrimony is faid to be aperient, detergent, and to strengthen the tone of the viscera: hence it is recommended in scorbutic differences, in debility and laxity of the intestines, &c Digested in whey, it affords an useful dietdrink for the spring season, not ungrateful to the palate or stomach,

## ALCANNA, vide Anchusa.

ALCEÆ folia: Alceæ vulgaris majoris C B. Malvæ verbenacæ Ger. Vervain Mallow; the leaves.

This is easily distinguishable from the common and marshmallow, by its leaves being jagged or cut in about the edges. It grows in hedges, and slowers greatest

part of the fummer. Alcea agrees in quality with the ALTHEA and MALVA VULGARIS; but appears to be less mucilaginous than either.

ALCHIMILLÆ folia: Alchimilla vulgaris C. B. Ladies man-

tle; the leaves [E.]

This grows wild in many parts of England, but is rarely met with about London: the leaves feem as if plaited or folded together, so as to have given occasion to the English name of the plant. The leaves of alchimilla discover to the taste a moderate aftringency, and were formerly much esteemed in some semale weaknesses, and in sluxes of the belly. They are now rarely made use of; though both the leaves and roots might doubtless be of service in cases where mild aftringents are required.

ALCIS UNGULA: Elks hoof. The elk is a large animal of the flag kind, met with in Muscovy, and other cold countries. The hoof of one of the hinder feet has been celebrated against epilepsies, from a ridiculous opinion, that the elk is himself subject to disorders of this kind, and prevents or removes them by scratching his ear with his hoof.

ALKEKENGI feu balicacabi fructus: Solani weficarii C. B. Winter cherry: the fruit [E.]

This is a low, branched shrub, bearing leaves like those of night-shade; with white slowers, which stand single at the joints. The slower cup changes into a membraneous cover, which at length bursts and discovers a fruit of a sine red colour, about the size of a common cherry. The stuit ripens in October, and continues frequently to the middle of December. This plant grows wild in

fome parts of France, Germany, &c. the beauty and lateness of its fruit has gained it a place in

our gardens.

Winter cherries are said by most writers to be extremely bitter: but, as Haller justly observes, the cherry itself, if carefully freed from the cover (which is very bitter and pungent) has merely a subacid taste. They stand highly recommended as detergent, aperient, diuretic, and for expelling gravel: four, five, or more of the cherries are directed for a dose. or an ounce of the expressed juice. Mr. Ray tells us of a gouty person who was cured and kept free from returns of his disorder, by taking eight of these cherries at each change of the moon; these occafioned a copious discharge of extremely fetid urine.

ALLIARIÆ folia: Hesperidis allium redolentis Tourn. Sauce alone, or jack by the hedge; the

leaves [E.]

This is common in hedges and hady waste places, flowering in May and June. The leaves have a bitterish acrid taste, and, when rubbed betwixt the fingers, a strong fmell, approaching to that of garlick. They are recommended internally as sudorifics and deobstruents, formewhat of the nature of garlick, but much milder; and externally as antifeptics in gangrenes and cancerous ulcers. Hildanus used to gather the herb for these last purposes in the spring, and expose it for a day to the action of a dry air in a shady place: being then committed to the press, it yielded a juice possessing the smell and tafte of the alliaria; this, he informs us, with a little oil on the surface, keeps in perfection for years; whereas the herb in

fubstance soon loses its virtue in keeping.

ALLIUM: radix Allii sativi C. B. Garlick; the roots [L. E.] These roots are of the bulbous kind, of an irregularly roundish shape, with several fibres at the bottom: each root is composed of a number of leffer bulbs, called cloves of garlick, inclosed in one common membranous coat, and easily separable from one another. All the parts of this plant, but more especially the roots, have a strong offensive smell, and an acrimonious almost caustic taste. The root applied to the skin inflames, and often exulcerates the part. Its fmell is extremely penetrating and diffusive; when the root is applied to the feet, its fcent is foon discoverable in the breath; and when taken internally, its fmell is communicated to the urine, or the matter of an issue, and perspires through the pores

of the Ikin. This pungent root warms and stimulates the solids, and attenuates tenacious juices. Hence in cold leucophlegmatic habits, it proves a powerful expectorant, diuretic, and emmenagogue; and if the patient is kept warm, fudorific. In humoural afthmas. and catharrhous disorders of the breast, in some scurvies, flatulent colics, hysterical and other diseases proceeding from laxity of the folids, and cold fluggish indispofition of the fluids, it has generally good effects: it has likewife been found serviceable in some hydropic cases. Sydenham relates, that he has known the dropfy cured by the use of garlick alone; he recommends it chiefly as a warm strengthening medicine in the beginning of the disease.

The liberal use of garlick is apt to occasion headachs, statulencies, thirst, sebrile heats, instammatory distempers, and sometimes discharges of blood from the hæmorrhoidal vessels. In hot bilious constitutions, where there is already a degree of irritation, where the juices are too thin and acrimonious, or the viscera unsound; this stimulating medicine is manifestly improper, and never fails to aggravate the distemper.

The most commodious form for the taking of garlick, a medicine to most people not a little unplea-fant, is that of a bolus or pill. Infusions in spirit, wine, vinegar, and water, although containing the whole of its virtues, are so acrimonious, as to be unsit for general use. A syrup and oxymel of it are kept in the shops.

Garlick made into an unguent with oils, &c. and applied externally, is faid to resolve and discuss cold tumors, and has been by fome greatly esteemed in cutaneous diseases. It has likewise sometime been employed as a revellent: Sydenham affures us, that among all the substances which occasion a derivation or revulsion from the head, none operate more powerfully than garlick applied to the foles of the feet : hence he was led to make use of it in the confluent small pox: about the eighth day after the feet began to swell, the root cut in pieces, and tied in a linen cloth, was applied to the foles, and renewed once a day till all danger was over.

ALNI VULGARIS folia & cortex Alat rotundifoliæ glutinosæ wiridis C B. The leaves and bark of the elder tree. These have a bitter styptic disagreeable taste. The bark is recommended by some in intermittent severs; and a decoction

of it, in gargarisms, for inflammations of the tonfils.

ALINI NIGRÆ feu frangulæ cortax: Alni nigræ bacciferæ J. B. The black or berry-bearing alder; its bark [E.]

This tree is common in moist woods in divers parts of England. The internal bark of the trunk os root of the tree, given to the quantity of a dram, purges violently, occasioning gripes, naufeæ, and vomiting. These may be in good measure prevented by the addition of aromtics; but as we have plenty of safer and less precarious purgatives, practitioners have deservedly rejected this.

ALOE. Aloes is the inspissated juice of certain plants of the same name. The ancients distinguished two forts of aloes; the one was pure and of a yellowish colour, inclining to a red, resembling the colour of a liver, and thence named hepatic; the other was full of impurities, and hence supposed to be only the drofs of the better At present, various sorts are met with in the shops; which are distinguished either from the places, from the species of the plants, or from some differences in the juices themselves. These may be all ranged in three classes.

(1) ALOE SOCOTORINA [L.E.] Socotorine aloes, brought from the island Socotora in the Indian ocean, wrapt in skins; it is obtained from the aloe Succotorina angustifolia spinosa, store purpureo Breyn, & Commelen.—This fort is the purest of the tree: it is of a glossy surface, clear, and in some degree pellucid; in the lump, of a yellowish red colour, with a purple cast; when reduced to powder, of a bright golden colour. It is hard

hard and friable in the winter, fomewhat pliable in summer, and grows soft betwixt the singers. Its taste is bitter, accompanied with an aromatic slavour, but insufficient to prevent its being disagreeable; the smell is not very unpleasant, and somewhat resembles that of myrrh.

- (2) ALOB HEPATICA [E] Hepatic, Barbadoes, or common aloes; the juice of the Aloe C. B. aloe vera vulgar is Munting .-- Hepatic aloes is not so clear and bright as the foregoing fort: it is also of a darker colour, more compact texture, and for the most part dryer. Its smell is much stronger and more disagreeable: the taste intensely bitter and nauseous, with little or nothing of the fine aromatic flavour of the Socotorine.- The best hepatic aloes comes from Barbadoes in large gourd shells; an inferior fort of it (which is generally foft and clammy) is brought over in casks.
- (3) ALOE CABALLINA. Fetid, caballine, or horse aloes; the produce of the aloe Guineensis caballina vulgari similis sed totu maculata Commelin. This fort is cafily diftinguished from both the foregoing, by its strong rank smell; although, in other respects, it agrees pretty much with the hepatic, and is not unfrequently fold in its stead.-Sometimes the caballine aloes is prepared so pure and bright, as not to be distinguishable by theeye even from the Socotorine; but its offensive smell, which it cannot be divested of, readily betrays it.

All the forts of aloes dissolve in pure spirit, proof spirit, and proof spirit diluted with half its weight of water; the impurities only being lest. They dissolve also by the assistance of heat in water alone;

but as the liquor grows cold, the resinous part subsides, the gummy remaining united with the water. The hepatic aloes is found to contain more resin, and less gum than the Socotorine, and this than the The refins of all the caballine. forts, purified by spirit of wine, have little fmell; that obtained from the Socotorine has scarce any perceptible talle; that of the hepatic, a slight bitterish relish, and the refin of the caballine, a little more of the aloetic flavour. gummy extracts of all the forts are less disagreeable than the crude aloes: the extract of Socotorine aloes has very little smell, and is in taste not unpleasant; that of the hepatic has a somewhat stronger fmell, but is rather more agreeable in tafte than the extract of the Socotorine: the gum of the caballine retains a confiderable share of the peculiar rank smell of this fort of aloes, but its tafte is not much more unpleasant than that of the extracts made from the two other forts.

Aloes is a stimulating cathartic bitter: if given in so large a dose as to purge effectually, it often occasions an irritation about the anus, and sometimes a discharge of blood. Small doses of it frequently repeated, not only cleanse the primæ via, but likewise attenuate and dissolve viscid juices in the remoter parts, warm the habit, quicken the circulation, and promote the uterine and hæmorrhodial fluxes. This medicine is particularly ferviceable to persons of a phlegmatic temperament and sedentary life, and where the stomach is oppressed and weakened: in dry bilious habits, aloes prove injurious, immoderately heating the blood, and inflaming the bowels.

I his juice is likewise, on account of its bitterness, supposed to kill

worms,

worm, either taken internally, or applied in plasters to the umbilical region. It is also celebrated for restraining external hæmorrhages, and cleansing and healing wounds and ulcers.

The ancients gave aloes in much larger doses than is customary at present. Dioscorides orders half a dram or a dram for gently loosening the belly; and three drams when intended to have the full effect of a cathartic. But modern practice rarely exceeds a scruple, and limits the greatest dose to two scruples. For the common purposes of this medicine, ten or twelve grains suffice; taken in these or less quantities, it acts as a gentle stimulating eccoprotic, capable of removing, if duly continued, very obstinate obstructions.

Some are of opinion, that the purgative virtue of aloes refides entirely in its refin; but experience has shewn, that the pure refin has little or no purgative quality; and that the gummy part separated from the refinous, acts more powerfully than crude aloes. aloes indeed be made to undergo long coction in the preparation of the gummy extract, its cathartic power will be confiderably lessened, not from the separation of the refin, but from an alteration made in the juice itself by the heat. The strongest vegetable cathartics become mild by a like treatment, without any remarkable separation of their parts.

Socotorine aloes, as already obferved, contains more gummy matter than the hepatic; and hence it is likewise found to purge more, and with greater irritation. The first fort therefore is most proper where a stimulus is required, as for promoting or exciting the menstrual flux; whilst the latter is better calculated to act as a common purge. It is supposed that the vulnerary and balsamic virtues of this juice reside chiefly in the refin; and hence that the hepatic aloes, which is most resinous, is most serviceable in external application.

The Edinburgh college directs the hepatic aloes in the balfamum traumaticum and tindura myrrhæ et aloes, defigned for external use; and the Socotorine in those preparations or compositions which are to be taken internally, as the tindura facra, elixir facrum, pulvis hieræ picræ, pilulæ aloeticæ, pilulæ Rusi, pilulæ stomachicæ, pilulæ cocciæ, &cc.

The London college uses the Socotorine only. In the winum aloeticum, tinctura sacra, elixir aloes, balsamum traumaticum, pilulæ aromaticæ, and the other pills wherein aloes is an ingredient, the Socotorine kind in substance is directed. In the powder of biera picra, only the pure gummy part of the Socotorine aloes is employed, the separation of which from the resinous matter is given in a distinct process.

ALSINES folia: Alfines vulgaris five morfus gallinæ 7. B. Chick-

weed; the leaves [E.]This plant was employed by the ancients externally against erysipelatous, and other inflammatory diforders. Later times have given it internally in hæmoptues, as a restorative in atrophies and confumptions, and likewise as an antipilen-Some recommend for these purposes the expressed juice to be taken to the quantity of an ounce: others the dried leaves, in the dose of a dram, and others a water diftilled from them. But if any real benefit is expected from alfine, it ought to be used liberally as food: though even then, its effects would not perhaps be superior to those of more approved culinary herbs.

AMMONIACUM **GUMMI** [L. E.] Ammoniacum is a concrete gummy refinous juice, brought from the East Indies, usually in large masses, composed of little lumps or tears, of a milky colour, but foon changing, upon being exposed to the air, of a yellowish hue. We have no certain account of the plant which affords this juice; the feeds usually found among the tears resemble those of the umbelliserous Such tears as are large, dry, free from little stones, seeds, or other impurities, should be picked out and preferred for internal use; the coarser kind is purified by solution and colature, and then carefully inspissating it; unless this be artfully managed, the gum will lose a considerable deal of its more volatile parts. There is often vended in the shops, under the name of strained gum ammoniacum, a composition of ingredients much inferior in virtue.

Ammoniacum has a nauseous fweet taste, followed by a bitter one; and a peculiar smell somewhat like that of galbanum, but more grateful; it softens in the mouth, and grows of a whiter colour upon being chewed. Thrown upon live coals, it burns away in flame: it is in some measure soluble in water and in vinegar, with which it assumes the appearance of milk; but the refinous part, amounting to about one half, subfides on sland-

Ammoniacum is an useful deobstruent; and frequently prescribed for opening obstructions of the abdominal viscera, and in hysterical disorders occasioned by a deficiency of the menstrual evacuations. is likewife supposed to deterge the pulmonary vessels, and proves of considerable service in some kinds of afthmas, where the lungs are oppressed by viscid phlegm; in this intention, a folution of gum ammoniacum in vinegar of squills proves a medicine of great efficacy, though not a little unpleasant. In long and obstinate colics proceeding from viscid matter lodged in the intestines, this gummy-refin has produced happy effects, after purges and the common carminatives had been used Ammoniacum is most in vain. commodiously taken in the form of pills: about a scruple may be given every night, or oftener. Externally it softens and ripens hard tumors: a folution of it in vinegar stands recommended by some for resolving even scirrhous swellings.

In the shops is prepared a solution of it in pennyroyal water, called, from its milky colour, lac ammoniaci [L.] It is an ingredient also in the pettoral oxymel and pills, in the deobstruent and gum pills [E.] and in several plasters [L. E.]

AMOMI VERI semen: Amount nacemos C. B. The feeds of the true amomum brought from the

East-Indies [L.] The true amomum is a round

fruit, about the fize of a middling grape; containing, under a membranous cover, a number of small rough angular feeds, of a blackish brown colour on the outfide, and whitish within: theseeds are lodged in three distinct cells; those in each cell are joined closely together, so as that the fruit, upon being opened, appears to contain only three feeds. Ten or twelve of these fruits grow together in a cluster, and adhere, without any pedicle, to a woody stalk about an inch long: each fingle fruit is surrounded by fix leaves, in form of a cup; and the part of the stalk void of fruit is clothed with leafy scales.

The husks, leaves, and stems have a light grateful fmell, and a modemoderately warm aromatic tafte: the feeds freed from the husks, are in both respects much stronger; their smell is quick and penetrating, their tafte pungent, approach. ing to that of camphor. Notwithstanding amomum is an elegant aromatic, it has long been a

stranger to the shops.

It is directed as an ingredient in the theriaca: the college of Edinburgh has expunged that composition, and as the true amonum is not at present to be procured in this country, they have dropt its name: that of London allows the feeds of the following plant of our own growth to be substituted to those of the oriental amomum.

AMOMI VULGARIS semen: Sisonis quod amomum officinis nostris C. B. Sii aromatici Tourn. feeds of the common amomum; or bastard stone parsley [L. E.]

These are very different in their appearance and manner of growth from the foregoing: they stand in form of umbels, and are joined two together without any common covering: they are fmall, striated. of an oval figure and brown colour. Their taste is warm and aromatic. but considerably different from that of the amomum verum, and very far weaker. Wafer extracts little of their flavour by infusion, but elevates the whole in distillation; rectified spirit extracts the whole, but elevates very little: hence the watery extract has no tafte or fmell of the feeds; whilft the spirituous posselles their flavour in great perfec-It is observable that the tincture drawn from them with pure spirit is of a green colour. These seeds have been recommended as carminative, aperient, diuretic and emmenagogue; but they are at present little regarded in practice.

AMYGDALÆ AMARÆ DULCES. Sweet and bitter almonds [L. E.].

The almond is a flattish kernel. of a white colour, covered with a thin brownish skin; of a fost sweet taste, or a disagrecable bitter one. The skins of both forts are unpleafant, and covered with an acrid powdery substance: they are very apt to become rancid on keeping. and to be preyed on by a kind of insect, which eats out the internal part, leaving the almond to appearance entire. To these circum. stances regard ought to be had in the choice of them.

The fruit which affords these kernels, is the produce of a tree greatly resembling the peach, called by C. B. amygdalus sativa. The eye distinguishes no difference betwixt the trees which produce the sweet and bitter, or betwixt the kernels themselves; it is said that the same tree has, by a difference in culture, afforded

both.

Both forts of almonds yield on expression, a large quantity of oil, which has no smell or any particular taste: this oil separates likewise upon boiling the almonds in water, and is gradually collected on the furface; but on triturating the almonds with water, the oil and water unite together, by the mediation of the other matter of the kernel, and form an uncluous milky liquor.

Sweet almonds are of greater use in food than as medicines; but they are reckoned to afford little nourishment, and when eaten in substance are not easy of digestion, unless thoroughly comminuted. They are supposed, on account of their foft unctuous quality, to obtund acrimonious juices in the primæ viæ: peeled sweet almonds, eaten fix or eight at a time, sometimes

G 3

give present relief in the heart-

Bitter almonds have been found poisonous to dogs, and sundry other animals; and a water distilled from them, when made of a certain degree of strength, has had like effects. Nevertheless, when eaten, they appear innocent to men, and have been not unfrequently used as medicines; Boerhaave recommends them, in substance, as diuretics which heat but moderately, and which may therefore be ventured upon in acute diseases.

The oils obtained by expression from both forts of almonds are in their sensible qualities the same. The general virtues of these oils are, to bluntacrimonious humours, and to soften and relax the folids: hence their use internally, in tickling coughs, heat of urine, pains and inflammations: and externally in tension and rigidity of particular

parts

The milky folutions of almonds in watery liquors, commonly called emulfions, contain the oil of the subject, and participate in some degree of the emollient virtue thereof; but have this advantage above the pure oil, that they may be given in acute or inflammatory disorders, without danger of the ill' effects which the oil might sometimes produce; fince emulsions do not turn rancid or acrimonious by heat, as all the oils of this kind in a little time do. Several unctuous and refinous substances, of themfelves not miscible with water, may by trituration with almonds be eafily mixed with it into the form of an emuliion; and are thus excellently fitted for medicinal use. In this form, camphor and the refinous purgatives may be commodi-The only officinal oully taken. preparations of almonds are the expressed oil and emulsion.

ANACARDIA. Anacardium, or malaca bean.

This is the fruit of a tree growing in Malabar and other parts of the East Indies. It is of a shining black colour, of the shape of a heart statened, about an inch long, terminating at one end in an obtuse point, and adhering by the other to a wrinkled stalk, it contains, within two shells, a kernel of a sweetish taste: betwixt the shells is lodged

a thick acrid juice.

The medicinal virtues of anarcadia have been greatly disputed: many have attributed to them the faculty of comforting the brain and nerves, fortifying the memory, quickening the intellect; and hence a confection made from them has been dignified with the title of confectio sapientum: others think it better deserves the name of confedio fullorum, and mention instances of its continued use having rendered people maniacal. But the kernel of anacardium is not different in quality from that of almonds. The ill effects attributed to this fruit belong only to the juice contained betwixt the kernels, whose acrimony is so great, that it is faid to be employed by the Indians as a caustic This juice is recommended externally for tetters, freckles, and othercutaneous deformities; which it removes only by ulcerating or excoriating the part, fo that a new Skin comes underneath,

ANAGALLIDIS folia: Anagallidis phæniceo flore C. B et Anagallidis flore cæruleo C. B. Common male and female pimpernel; the leaves.

Pimpernel is a low plant, in appearance refembling chick weed but easily distinguishable by its leaves being spotted underneath, and joined immediately to the stalk. The male and semale pimpernels differ no otherwise than in the colour of

theix

their flowers: they are both found wild in the fields, but the male or red flowered fort is most common.

Both the pimpernels have an herbaceous, roughish taste, with little or no fmell. Many extraordinary virtues have been attributed to them. Geoffroy efteems them cephalic, fudorific, vulnerary, antimaniacal, antepileptic, and alexe-Tragus, Caspar, Hoffman, terial. Michaeli, and others, are also very liberal in their praises: one of these gentlemen declares, that he has known numerous instances of the fingular efficacy of a decoction and tincture of pimpernel, in maniacal and melancholic deliria. But later practitioners have not been so happy as to meet with the like fuccels. Pimpernelis not unfrequently taken as food; it makes no unpleasant falad; and in some parts of this kingdom, is a common pot-herb. A spirituous tincture of it contains nothing valuable: the only preparation that promifes any utility, is an extract made with water; or the expressed juice depurated and inspissated.

ANAGALLIS AQUATICA, vide Becabunga.

ANCHUSÆ radix: Buglossi radice rubra Tourn. Alkanet root | E]

Alkanet is a rough hairy plant, much resembling the vipers bugloss: its chief difference from the common buglosses consists in the colour of its roots; the cortical part of which is of a dusky red, and imparts an elegant deep red to oils, wax, and all unstuous substances, but not to watery liquors. This plant is a native of the warmer parts of Europe: it is sometimes cultivated in our gardens; but the greatest quantities are raised in Germany and France, particularly about Montpelier, from whence

the dried roots are usually imported to us. The alkanet root produced in England is much inferior in colour to that brought from abroad; the English being only lightly reddish, the others of a deep purplish red: this has induced some to suspect that the foreign roots owe part of their colour to art; but we think without sufficient soundation.

Alkanet root has little or no smell: when recent, it has a bitterish astringent taste; but when dried scarce any. As to its virtues, the present practice expects not any from it. Its chief use is for colouring oils, unguents, and plasters. As the colour is confined to the cortical part, the small roots are best, these having proportionably more bark than the large.

ANETHI semen: Anethi bortenfis C. B. Dill seed [L. E.]

Dill is an umbelliferous plant, cultivated in gardens, as well for culinary as medical use. The feeds are of a pale yellowish colour, in shape nearly oval, convex on one fide, flat on the other. Their tafte is moderately warm and pungent, their smell aromatic, but not of the most agreeable kind. feeds are recommended as a carminative, in flatulent colics proceeding from a cold cause or a viscidity of the juices. The most efficacious preparations of them are, the distilled oil, and a tincture or extract made with rectified spirit. The oil and simple water distilled from them are kept in the shops  $\{L_i\}$ 

ANGELIC Æ radix, folia, semen:
Angelicæ sativæ C. B. imperatoriæ
sativæ Tourn. Garden angelica;
the roots, leaves, and seeds [L E.]

This is a large umbelliferous plant, growing spontaneously in the northern climates: for the use of the shops, it is cultivated in gardens, gardens, in the different parts of Europe. Bohemia and Spain are faid to produce the best. The London collegedirect the roots brought from Spain to be alone made use of. Angelica roots are apt to grow mouldy, and be preyed upon by insects unless thoroughly dried, kept in a dry place, and frequently aired. We apprehend that the roots which are subject to this inconvenience might be preserved, by dipping them in boiling spirit, or exposing them to its steam, after they are dried.

All the parts of angelica, especially the roots, have a fragrant aromatic smell; and a pleasant bitterish warm talle, glowing upon the lips and palate for a long time after they have been chewed. The flavour of the feeds and leaves is very perishable, particularly that of the latter, which on being barely dried, lose greatest part of their tafte and smell: the roots are more tenacious of their flavour, though even these lose part of it upon keeping. The fresh root, wounded early in the spring, yields an odorous, yellow juice, which flowly exficcated, proves an elegant gummy-refin, very rich in the virtues of the angelica. On drying the root, this juice concretes into diflinct moleculæ, which on cutting it longitudinally, appeardistributed in little veins: in this state they are extracted by pure spirit, but not by watery liquors.

Angelica is one of the most elegantaromarics of European growth, though little regarded in the present practice. The root, which is the most efficacious part, is rarely met with in prescription, and does not enter any officinal composition. The leaves are ingredients in the three alexeterial waters [L.]: the seeds, in the compound anised water [L.], plague water, aqua mi-

rabilis, and aromatic tincture [E] The stalks make an agreeable sweetmeat.

ANGUILLÆ HEPAR. The liver of the cel.

The liver and gall of the eel are extremely acrid. They have been held a specific in difficult births; and enter the principal compositions for that intention in foreign pharmacopæias; although it appears, that in most cases of this kind, acrid irritating medicines are really injurious. Boerhauve obferves, that no fish has a more acrid gall than the eel; and fays, that with pills made of the gall of the eel and pike, he has cured pale rickety children with swelled bellies; the gall powerfully promoting urine, and occasioning the belly to subfide.

ANIME; [E.] a refin exuding from the trunk of a large American tree, called by Pifo jetaiba, by the Indians courbaril.

This refin is of a transparent amber colour, a ligh: agreeable smell, and little or no taste. It dissolves entirely, but not very readily, in rectified spirit of wine; the impurities, which a e often in large quantity, remaining behind. The Brazilians are said to employ anime in sumigations for pains and aches proceeding from a cold cause; with us, it is rarely, if ever, made of for any medicinal purposes.

ANISI semen: Apii anisi disti semine sua veolente Tourn. Anise, the seed [L. E]

Anife is an annual umbelliferous plant, growing naturally in Crete, Syria, and other places of the Eat. It is cultivated in some parts of France, Germany, and Spain, and may be raised also in England: the seeds brought from Spain, which

are fmaller than the others, are

preferred.

Anifeeds have an aromatic smell, and a pleasant warm taste, accompanied with a degree of sweetness. Water extracts very little of their slavour; rectified spirit the whole.

These seeds are in the number of the four greater hot seeds; their principal use is in cold statulent disorders, where tenacious phlegm abounds, and in the gripes, to which young children are subject. Frederick Hossman strongly recommends them in weakness of the stomach, diarrhæas, and for the stomach, diarrhæas, and for strengthening the tone of the viscera in general; and thinks they well deserve the appellation given them by Helmont, intestinorum solamen.

The officinal preparations of these seeds are an effectial oil L. E.] and a spirituous compound water [L. E.] They are ingredients in mitbridate and theriaca; and the essential oil in the paregoric elixir [L]

ANONIS, vide Ononis.

ANSERINA, vide ARGENTINA.

ANTIMONIUM [L E.] fii-

Antimony is a ponderous brittle mineral, composed of long shining streaks like needles, intermingled with a dark lead coloured substance; of no manifest taste or smell. There are several mines of it in Germany, Hungary, and France; and some likewise in England. The English seems to be of all the others the least proper for medicinal use, as frequently containing a portion of lead. The substances sound mixed with the foreign forts are generally of the unfusible stony kind, from which the antimony is melted out in vessels,

whose bottom is perforated with small holes, and received in conical moulds: in these, the lighter and more drossy matter arises to the surface; whilst the more pure and ponderous subsides to the bottom: hence the upper broad part of the loaves is considerably less pure than the lower.

The goodness of antimony is judged of from its weight; from the loaves not being spongy or blebby; from the largeness of the striæ; and from the antimony totally evaporating in a strong sire.

Antimony was employed by the ancients in collyria against inflammations of the eyes; and for staining the eyebrows black. ternal use does not seem to have been established till towards the end of the fifteenth century; and even at that time it was by many looked upon as poisonous. But experience has now fully evinced, that pure antimony, in its crude state, has no noxious quality; that some of the preparations of it are medicines of great efficacy; and that though many of them are most violently emetic and cathartic, yet even these, by a slight alteration or addition, lose their virulence. and become mild in their opera-

This mineral appears from chemical experiments to confift of a metal, un:ted with common fulphur, and separable in its metallic form by the same means whereby other metallic bodies are extracted from their ores.

The pure metal operates, in a very minute dose, with extreme vehemence, as a purgative and emetic: when combined with sulphur, as in the crude mineral, its power is restrained: divested of the inflammable principle which it has in common with all persectly metallic bodies, it becomes an indo-

lent

lent calx. See the preparations of antimony in the third part of this work.

ANTHOR Æ frve antithoræ redix: Aconiti falutiferi C. B. Aconiti foliorum laciniis linearibus, ubique ejufdem latitudinis Linnæi Wholefome wolf's bane; the roots.

This plant may be distinguished from the poisonous aconites by its leaves being more finely divided, and not at all bright or shining: it grows wild on the Alps. root has been supposed useful against poisons, particularly that of the thora, (whence its name.) Some nevertheless look upon this pretended antidute itself as unfafe: Fred. Hoffman says it is cathartic, and has produced dangerous diforders of the stomach, accompanied with heat, thirst, and anxiety. On the other hand Geoffroy relates, that he has never observed any purgative quality in this root, or any ill consequence from its use; that he has frequently exhibited it, and always with good success, against worms, and in malignant fevers, especially such as were occasioned by viscidities in the stomach and intestines; the dose from a scruple to a dram. A competency of experiments to fully determine this point, is as yet wanting, the root never having come into general practice. Its taste is acrid and bitter.

APARINES folia: Aparines oulgaris C. B. Goolegrass, or clivers; the leaves [E.]

This is a flender, rough plant, common in hedges, &c. It is recommended as an aperient, but practice has little regard to it.

APES. Bees; their bodies, honey, and wax [E.]
Bees, dried and pulverized, are

faid to cure the alopecia, and, given internally, to promote urine; but they have been for a long time strangers to the shops. The honey and wax shall be treated of under the respective heads.

APII feu eleofelini radix : Apii foliis caulinis cuneiformibus Linnai.

Smallage, the roots [E.]

This plant is larger than the garden apium (parsley), of a darker green colour, and of a stronger and more unpleasant flavour. The roots are in the number of the five called opening roots, and have been sometimes prescribed as an ingredient in aperient apozems and diet drinks; but are at present disre-The feeds of the plant garded. are moderately aromatic, and were formerly used as carminatives; in which intention they are, doubtless, capable of doing service, though the other warm feeds, which the shops are furnished with, render these unnecessary; and accordingly the Edinburgh college, which retains the roots, has expunged the feeds.

APIUM HORTENSE, vide Petrosolinum.

AQUILEGIÆ folia; stores, femen: Aquilegiæ store simplici Raii Syn. Columbines; the leaves, stowers, and seeds.

This plant grows wild in woods, but is not very common. It has been looked upon as aperient; and was formerly in great effect among the common people for throwing out the small-pox and measles. A distilled water, medicated vinegar, and conserve, were prepared from the flowers; but they have long given place to medicines of greater efficacy.

ARANEARUM TELÆ [E.] Cobwebs.

Thefe

These are never met with in prescription; but are sometimes applied by the common people to stop the bleeding of slight wounds: this they feem to effect by adhering to the part, fo as to close the orifices of the veffels, and prevent the effusion of their contents.

ARESTA BOVIS, vide Ono-NIS.

ARGENTINÆ, potenti/læ, anserinæ folia: Pentaphylloidis minoris fupini, seu procumbentis, foliis alatis urgenteis et serratis, flore luteo Mor. Hist. Ox. Silverweed, or wild

tansey; the leaves [E.]

This plant grows wild about the fides of rivulets and other moitt places; it has no stalk, the leaves lying flat on the ground. The writers on the Materia Medica in general look upon argentina as a very frong aftringent; misled probably by its agreement in botanic characters with tormentil, which is known to be a powerful flyptic. The fenfible qualities of argentina promise no great virtue of this kind; for to the taste it discovers only a flight roughishness, from whence it may be presumed to be entitled to a place only among the milder corroborants. As the aftringency of tormentil is confined chiefly to its root, it might be thought that the argentina also has an astringent root: the root of this plant, however, is found to have no other than a pleasant sweetish tatte, like that of parinips, but not fo itrong.

ARGENTUM. Silver [L. E.] Abundance of virtues have been attributed to crude filver by the Arabians, and by some also of later times, but on very little founda-This metal, taken in its crude flate, has no effect in the

body: combined with a small quantity of the nitrous acid, it proves a powerful, though not always a fafe, hydragogue; with a larger, a strong caustic. The nitrous acid is the only one that perfectly diffolves this metal; on adding to this folution a minute portion of marine acid, or substances containing it, the liquor turns milky. and the filver falls to the bottom in form of a white calx: hence we are furnished with a method of discovering marine falt in waters, &c. See the preparations of filver in the third part.

ARGENIUM VIVUM. Hydrargyrus: Mercurius. Mercury, or

quickfilver [L.E.]

Mercury is an opake filver-coloured mineral fluid; appearing to the eye like tin or lead when melted: it is heavier than any other fluid, and than most of the metallic bodies: it does not congeal in the greatest degree of natural cold hitherto known; in the fire it proves totally volatile. This mineral is either met with in its fluid form, in the earth, or extracted by art from certain ores. are confiderable mines of it in Hungary and Spain; but the greatest quantities come from the Eatt Indies.

The use of mercury in medicine feems to have been little known before the fifteenth century. The ancients looked upon it as a corrouve poison, though of itself perfectly void of acrimony, talleand smell: there are examples of its having been ledged for years in cavities both of bones and fleshy parts, without its having injured or affected them. Taken into the body in its crude state, and undivided, it passes through the intestines onchanged, and has not been found to produce any confiderable

effect.

effect. It has indeed been recommended in afthmas and diforders of the lungs; but the virtues attributed to it in these cases have not been warranted by experience.

Notwithstanding the mildness and activity of crude quicksilver undivided; when resolved by fire into the form of sume, or otherwise divided into very minute particles, and prevented from re-uniting by the interposition of proper substances; or combined with mineral acids, it has very powerful effects; affording the most violent possons, and the most excellent remedies that we are acquainted with.

The mercurial preparations, either given internally or introduced into the habit by external application, seem to liquefy all the juices of the body, even those in the minutest and most remote vessels; and may be fo managed as to promote excretion through all the Hence their comemunctories. mon use in inveterate chronic disorders proceeding from a thickness and fluggishness of the humours, and obstinate obstructions of the execretory glands; in scrophulous and cutaneous diseases; and in the venereal lues. If their power is not restrained by proper additions to certain emunctories, they tend chiefly to affect the mouth; and after having fuled the juices in the remoter parts, occasion a plentiful evacuation of them from the salival glands

The falutary effects of mercurials do not depend on the quantity of scnsible evacuation. This medicine may be gradually introduced into the habit, so as, without occasioning any remarkable discharge, to be productive of very happy effects. To answer this purpose, it should be given in very small doses, in conjunction with such substances as determine its

action to the kidneys or the pores of the fkin By this method inveterate cutaneous and venereal diftempers have been cured, without any other fensible excretion than a gentle increase of perspiration or urine. Where there are ulcers in any part, they discharge for some time a very fetid matter, the quantity of which becomes gradually lefs, and at length the ulcer kindly heals. If the mercury should at any time, from cold or the like, affect the mouth (which I have rarely found to happen) it may be restrained by omitting a dose, and by warmth or fuitable medicines promoting the perspiration.

ARISTOLOCHIA. Birthwort. Three roots of this name are directed for medicinal use:

- (1) ARISTOLOCHIA LONGA [L.E] Long birthwort. This is a tuberous root, sometimes about the fize of the finger, sometimes as thick as a man's arm, and a foot in length: it is nearly of an equal thickness all over, or a little thicker in the middle than at the ends: the outside is of a brownish colour; the inside yellowish.
- (2) ARISTOLOGHIA ROTUNDA [E.] Round birthwort. This has scarce any other visible difference from the foregoing than its roundish shape.

(3) ARISTOLOCHIA TENUIS
[L.] Slender birthwort. This is a
long and flender root, rarely exceeding the thickness of a goose quill.

These roots are the produce of Spain, Italy, and the southern parts of France. Their smell is somewhat aromatic; their taste warm and bitterish. Authors in general represent them as extremely hot and pungent: some say they are the

hottest

hottest of all the aromatic plants; but as usually met with in the shops, they have no great pungency. The long and round forts. on being first chewed, scarce discover any taste, but in a little time prove nauseously bitterish; the long fomewhat the least fo. The other fort instantly fills the mouth with an aromatic bitterness, which is not ungrateful. Their medical virtues are, to heat, stimulate, attenuate viscid phlegm, and promote the fluid secretions in general: they are principally celebrated in suppressions of female evacu-The dose in substance is from a scruple to two drams. The long fortisrecommended externally for cleanfing and drying wounds and ulcers, and in cutaneous difeases-The aristolochia tenuis, is an ingredient in theriaca; and in want of this species, the longa is allowed to be fubflituted to it by the London college.

ARMORACIA. vide RAPHA-NUS RUSTICANUS.

ARNICA, vide Doronicum.

ARSENICUM. Arsenic.

Arfenic is contained, in greater or less quantity, in most kinds of ores, particularly in those of tin and bilmuth, in the white pyrites, and the mineral called cobalt; from which last greatest part of the arsenic brought to us is extracted by a kind of sublimation: the arsenic arises at first in form of greyish meal, which more carefully resublimed, concretes into transparent masses, the while arsenic of the shops.

Arfenic sublimed with one-tenth its weight of sulphur, unites therewith into a bright yellow mass, in some degree transparent; the common yellow arsenic. On doubling

the quantity of sulphur, the compound proves more opake and compact; of a deep red colour, resembling that of cinnabar, but with this difference, that it loses of its beauty upon being reduced into powder, whilst that of cinnabar is improved by this means: this is the common red arsenic, By varying the proportions of arsenic and sulphur, sublimates may be obtained of a great variety of shades of yellow and red.

Natural mixtures of arfenic and falphur resembling the foregoing preparations, are not unfrequently met with in the earth. The fossil red arsenic is the sandaracha of the Greeks, the realgar and rifigal of the Arabians. Both the red and yellow, when of a smooth uniform texture, are named zamichs: and when composed of small scales or leaves, auripigmenta, or orpiments: these last are the only substances to which the Greeks gave the name That the zarnichs and orpiments really contain arfenie (contrary to the opinion of some late writers) is evident from fundryexperiments, whereby aperfect arfenic, and in notable quantity, is The comobtainable from them. pilers of the preceding edition of the Edinburgh dispensatory therefore very justly gave fandaracha Gracorum as a synonymon to red arsenic and auripigmentum to the yellow.

The pure or white arsenic has a penetrating cosrosive taste; and taken into the body proves a most violent poison. Besides the effects which it has in common with other corrosives, it remarkably attenuates the coats of the stomach, occasions a swelling and sphaselation of the whole body, and a funden putresaction after death, particularly, as is said, of the genitals in men. Where the quantity is so very small as not to prove

fatal

fatal, tremors, palfies, and lingering hectics succeed. The remedies recommended against this poison are, milk and oily liquors immediately and liberally drank.

The red and yellow arfenics, both native and factitious, have little taste, and are much less virulentin their effects than the foregoing. Sulphur, which restrains the power of mercury and the antimonial metal, remarkably abates the virulence of this poisonous mineral also. Such of these substances as participate more largely of sulphur, seem to be almost innocent: the factitious red arsenic and the native orpiments, have been given to dogs in confiderable quantity, without their being productive of any apparent ill consequences.

ARTEMISIÆ folia: Artemisiæ vulgaris majoris C.B. Mugwort;

the leaves. [L. E]

This plant grows plentifully in fields, hedges, and waste places, throughout England; and flowers in June. In appearance it somewhat refembles the common wormwood: the difference most obvious to the eye is in the flowers, those of wormwood hanging downwards, whilst the flowers of mugwort sland erect.

The leaves of this plant have a light aromatic smell, and an herbaceous bitterish taste. They are principally celebrated as uterine and antihysteric: an infusion of them is fometimes drank, either alone or in conjunction with other substances, in suppressions of the mentiqual evacuations. dicine is certainly a very mild one, and confiderably less hot than most others to which these virtues are attributed: in some parts of this kingdom, mugwort is of common use as a pot-herb.

ARTHANITÆ five cyclaminis radix: cyclaminis or biculato folio inferne purpurascente C. B.

bread; the root [E.]

This plant is met with in the gardens of the curious. The root has, when fresh, an extremely acrimonious burning tafte, which it almost entirely loses on being dried. It is recommended as an errhine: in cataplasms for scirrhous and scrophulous tumours; and internally as a cathartic, detergent, and aperient; it operates very flowly, but with great virulence, inflaming the fauces and inteslines; and hence is deservedly rejected from the London dispensatory, though retained in that of Edinburgh.

ARI radix: Ari maculati maculis nigris C. B. Wake-robin:

the root [L. E.]

This plant grows wild under hedges, and by the fides of banks, in most parts of England. It sends forth in March, three or four triangular leaves, which are followed by a naked stalk, bearing a purplish pistil inclosed in a long sheath: this is succeeded, in July, by a bunch of reddish berries. In some plants, the leaves are spotted with black, in others with white spots, and in others not spotted at all; the black spotted fort is supposed to be the most efficacious, and hence is expressly directed by the London college.

All the parts of arum, particularly the root, have an extremely pungent, acrimonious taile: if the root be but lightly chewed, it continues to burn and veilicate the tongue for some hours, occasioning at the same time a considerable thirst: these symptoms are alleviated by butter, milk, or oily Dried and kept for fome liquors. time, it loses much of its acrimony, and becomes at length an

almost insipid farinaceous sub-

This root is a powerful stimulant and attenuant. It is reckoned a medicine of great efficacy in some cachectic and chlorotic cases, in weakness of the stomach occasioned by a load of viscid phlegm, and in fuch disorders in general as proceed from a cold fluggish indisposition of the folids and lentor of the fluids. I have experienced great benefit from it in rheumatic pains, particularly those of the fixt kind, and which were seated deep. these cases I have given from ten grains to a scruple of the fresh root twice or thrice a day, made into a bolus or emulfion with uncluous and mucilaginous substances, which cover its pungency, and prevent its making any painful impression on the tongue. It generally excited a slight tingling sensation through the whole habit, and, when the patient was kept warm in bed, produced a copious sweat.

The only officinal preparation, in which this root is an ingredient, is a compound powder; in which form, its virtues are very precarious. Some recommend a tincture of it drawn with wine; but neither wine, water, nor spirit, ex-

tract its virtues.

ASAFOETIDA. Afafetida [L.E.] the concrete juice of a large umbelliferous plant growing in Persia.

This juice exudes (from wounds made in the root of the plant) liquid, and white like milk: on being exposed to the air, it turns of a brownish colour, and gradually acquires different degrees of consistency. It is brought to us in large irregular masses, composed of various little shining lumps or grains, which are partly of a whitish colour, partly reddish, and partly of a violet hue. Those masses are accounted the best which are clear,

of a pale reddish colour, and variegated with a great number of elegant white tears.

This drug has a strong setid smell, somewhat like that of garlick; and a bitter, acrid, biting taste. It loses with age of its smell and strength, a circumstance to be particularly regarded in its exhibition. It consists of about one-third part of pure resin, and two-thirds of gummy matter; the former soluble in rectissed spirit, the other in water. Proof spirit dissolves almost the whole into a turbid liquor; the tincture in rectissed spirit is transparent.

Asafetida is the strongest of the fetid gums, and of frequent use in hysteric and different kinds of nervous complaints. It is likewise of considerable efficacy in statulent colics; and for promoting all the studies fecretions in either sex. The ancients attributed to this medicine many other virtues, which are at present not expected from

ıt.

This gummy-resin is an ingredient in the officinal gum pills, compound powder of myrrh, setid tincture, tincture of soot, setid volatile spirit [L.] and antihysteric plaster [E.]

ASARI folia, radix: Afari C. B. Afarabacca; the roots and leaves.

—The London college directs only the leaves: the Edinburgh both leaves and root.

Asarum is a very low evergreen plant, growing naturally in France, Italy, and other warm countries: the dried roots have been generally brought from the Levant; those of our own growth being supposed weaker.

Both the roots and leaves have a nauseous, bitter, acrimonious, hot taste; their smell is strong and not very disagreeable. Given in substance from half a dram to a dram,

they .

they evacuate powerfully both upwards and downwards. It is faid. that tinctures made in spirituous menstrua, possess both the emetic and cathartic virtues of the plant: that the extract obtained by inspiffating these tinctures, acts only by vomit, and with great mildness: that an infusion in water proves cathartic, rarely emetic: that aqueous decoctions made by long boiling, and the watery extra t, have no purgative or emetic quality, but prove notable diaphoretics, diuretics, and emmenagogues.

The principal use of this plant among us is as a sternutatory. The root of asarum is perhaps the strongest of all the vegetable errhines, white hellebore itself not excepted. Snuffed up the nose, in the quantity of a grain or two, it occasions a large evacuation of mucus, and raises a plentiful spitting. The leaves are confiderably milder, and may be used, to the quantity of three, four, or five grains. Geoffroy relates, that after fuuffing up a dose of this errhine at night, he has frequently observed the discharge from the nose to continue for three days together; and that he has known a paralyfis of the mouth and tongue cured by He recommends this medicine in stubborn disorders of the head, proceeding from viscid tenacious matter, in palsies, and in suporific distempers. The leaves are an ingredient in the pulvis flernutatorius of the shops [L. E.]

ASCLEPIAS, vide Vince-

ASELLI, vide MILLEPEDÆ.

ASPALATHUS, vide Rho-

ASPARAGIradix: Asparagi sa-

This plant is cultivated in gardens for culturary use. The roots have a bitterish mucilaginous taste; inclining to sweetness, the fruit has much the same kind of taste; the young shoots are more agreeable than either. Asparagus promotes appetite, but affords little nourishment. It gives a strong ill fmell to the urine in a little time after eating it, and for this reason chiefly is supposed to be diuretic: it is likewise esteemed aperient and deobstruent; the root is one of the five called opening roots. Some suppose the shoots to be most efficacious; others the root; and others the bark of the root. is of opinion, that none of them have any great share of the virtues usually ascribed to them. Asparagus appears from experience to contribute very little either to the exciting of urine when suppressed, or increasing its discharge; and in

tiwi C B. Asparagus; the root [E.]

ASPERULÆ flores: Aspergulæ aut aspergulæ adoratæ mostratis Lab. Woodroof; the slowers.

cases where aperient medicines ge-

nerally do service, this has little

or no effect.

This is a low umbelliferous plant, growing wild in woods and copfes, and flowering in May. It has an exceeding pleasant smell, which is improved by moderate exsiccation; the taste is subfaline, and somewhat austere. It imparts its slavour to vinous sliquors. Asperula is supposed to attenuate vikid humours, and strengthen the tone of the bowels: it is recommended in obstructions of the liver and biliary ducts, and by some in epilepsies and passes: modern practice has nevertheless rejected it.

ASPHALTUS, vide BITUMEN JUDAICUM.

ASPLE-

ASPLENIUM, vide CETERACH.

A FRIPLICIS OLIDÆ folia:
Atriplicis fætidæ C. B. Chenopodii
fætidi Tourn. Stinking orach, or
arrach; the leaves [L.]

This is a low plant, sprinkled all over with a kind of whitish clammy meal: it grows about dunghills, and other waste places. The leaves have a strong fetid smell, which the hand, by a light touch, becomes so impregnanted with, as not to be eafily freed from. fmell has gained it the character of an excellent antihysteric; and this is the only use it is applied to. Tournefort recommends a spirituous tincture, others a decoction in water, and others a conferve of the leaves, as of wonderful efficacy in uterine disorders.

ATRIPLEX SATIVA. Gardenorrach. or arrach.

The garden oraches (which are either of a pale greenish, or purplish red colour, and hence named atriplex alba and rubra) are chiefly employed for culinary purposes. They are cooling, and gently laxative; a decoction of the leaves is recommended in costiveness, where the patient is of a hot bilious disposition.

AVENA [E.] Oats.

This grain is an article rather of food than of medicine. It is sufficiently nutritive and easy of digestion. The gruels made from it have likewise a kind of soft mucilaginous quality; by which they obtund acrimonious humours, and prove useful in instammatory disorders, coughs, hoarseness, roughness and exulcerations of the fauces.

AURANTIORUM HISPA-LENSIUM succus et cortex: Fructus Mali auraniæ majoris C. B. Seville oranges; the juice and yellow rind [L.E.] The Edinburgh college uses also the flowers of the tree.

The orange is a beautiful evergreen tree, or rather shrub, bearing flowers and fruit all the year: it is a native of the warmer climates, and does not well beat the winters of this.

The flowers are highly odoriferous, and have been for some time past of great esteem as a perfume: their tafte is somewhat warm, accompanied with a degree of bitterness. They yield their flavour by infusion to rectified spirit, and in distillation both to spirit and water: the bitter matter is dissolved by water, and, on evaporating the decoction, remains entire in the extract. The distilled water was formerly kept in the shops, but on account of the scarcity of the flowers is now laid aside: it is called by foreign writers aqua nepbæ. oil distilled from these slowers is brought from Italy under the name of oleum or essentia neroli.

The outer yellow rind of the fruit is a grateful aromatic bitter, and, in cold phlegmatic constitutions, proves an excellent stomachic and carminative, promoting appetite, warming the habit, and strengthening the tone of the viscera. Orange peel appears to be very confiderably warmer than that of lemons, and to abound more with essential oil: to this circumstance therefore due regard ought to be had in the use of these medicines. The flavour of the first is likewise supposed to be less perishable than that of the other, hence the college employ orange peel in the spirituous bitter tincture, which is defigned for keeping, whilst in the bitter watery infusion, lemon peel is preferred. A fyrup and two di-

flilled

filled waters are for the fame reason prepared from the rind of oranges in preference to that of lemons.

The juice of oranges is a grateful acid liquor, of confiderable use in febrile or inflammatory distempers, for allaying heat, abating exorbitant commotions of the blood, quenching thirst, and promoting the falutary excretions: it is likewise of great use in scurvies, especially when given in conjunction with the cochlearia, nasturium, or other acrid antiscorbutics, as in the succi scorbutici of the shops.

AURANTIA CURASLAVEN-SIA. Curaffao oranges [E.]

These are the small young fruit of the Seville orange dried. Their sirstappearance in a public pharmacopœia is in the present edition of the Edinburgh; in which they are made an ingredient in the stomachic tincture and elixir. They appear very well adapted to that intention, being moderately warm bitterish aromatics, of a slavour sufficiently agreeable.

AURICULA JUDÆ: Fungus suricuta Judæ, coloris ex cineraceo nigricantis, perniciofus, in fambuci caudice nascens J. B. Jews-ear, a fungus growing on old elder trees. This fungus is said by some to be a strong purgative; by others an astringent. The more judicious medical writers have declared its internal use dangerous: and hence at the late reformation, the Edinburgh as well as London college have rejected it.

AURICULÆ MURIS folia: Pilosellæ majoris repentis hirsutæ C. B. Mouse-ear; the leaves.

This is a low creeping plant, covered with a kind of hairs: it grows wild in dry pasture grounds.

and flowers in June and July. The leaves have a fomewhat rough bitterish taste: they are recommended as aftringents, but practice pays no regard to them.

AURIPIGMENTUM. Orpiment; a mineral composed of sulphur and arsenic. See Arsenicum.

AURUM. Gold.

This metal was introduced into medicine by the Arabians, who esteemed it one of the greatest cordials and comforters of the nerves. From them Europe received it withoutany diminution of its character; in foreign phamacopæias it is still retained, and even mixed with the ingredients from which simple waters are to be distilled. But no one, it is presumed, at this time expects any fingular virtues from it, fince it certainly is not alterable in the human body. Geoffroy, though unwilling to reject it from the cordial preparations, honestly acknowledges, that he has no other reason for retaining it, than complaisance to the Arabian schools. The chemists have endeavoured, by many elaborate processes, to extract what they call a fulphur or anima of gold: but no method is as yet known of separating the component parts of this metal; all the tinctures of it and aurum potabile's, which have hitherto appeared, are real folutions of it in aqua regis, diluted with spirit of wine or other liquors, and prove injurious to the body rather than beneficial.

AXUNGIA. Fat.

A great variety of fats were introduced into medicine by the Arabians, and recommended as possessing distinct virtues. The college of Wirtemberg, in the last edition of their dispensatory, published in

1741. directs no less than twentyeight different fats, to be kept in the shops: some of these, they informus, are attenuating and resolvent; such are those of the heron, wild cat, stork, partridge, coney, hare, fox, Alpine mouse, the badger, boar, wolf, ferpents, and vipers: others are heating, detergent and feptic; those of the eel. the pike, and the umber: a third class is emollient; the fat of the ox, the deer, and the goat: and a fourth, emollient, digerent, and lenient; this last comprehends the fats of the duck, goose, dog, ca. pon, beaver, horse, hen, and human fat. Experience, however, does not countenance these different virtues ascribed to different fats. They have all one common emollient quality, relax the part to which they are applied, and prevent perspiration: these effects, with the consequence of them, may be expected in a greater or less degree, from fats of every kind. The London college has therefore retained only three fats, of different confistences, for different mixtures, viz. viper's fat, hog's lard, and mutton fuet: to which the Edinburgh college adds goat's fuet. These are certainly sufficient for answering all the intentions that substances of this kind are employed for.

BALAUSTIA: Flores balausties flore pleno majore C. B. Balaustines: the flowers of the balaustine or double-flowered pomgranate tree [L. E.]

The balaukine is a low tree, or rather shrub, growing wild in Italy, &c. The slowers are of an elegant red colour, in appearance refembling a dried red rose. Their taste is bitterish and astringent.

Balaustines are recommended in diarrhoxas, dysenteries, and other cases, where astringent medicines are proper. They are rarely directed in extemporaneous prescription, and enter only one officinal composition, the pulvis e succino compositus. [L.]

BALSAMITÆ MARIS sive costi bortorum folia: Menthæ hortensis corymbisere C. B. Costma-

ry; the leaves [E.]

This was formerly a very common garden plant, and of frequent use both for culinary and medicinal purposes; but is at present very little regarded for either; though it should feem, from its sensible qualities, to be equal or superior, as a medicine, to some aromatic herbs, which practice has retained. The leaves have a bitterish warm aromatic talle; and a very pleafant fmell, approaching to that of mine or a mixture of mint and maudlin. Water elevates their flavour in difillation; and rectified spirit extracissit by infusion.

BALSAMUM COPAIBA(L.E.)
Balsam of Copaiba: a liquid refinous juice, flowing from incisions made in the trunk of a large tree which grows in the Spanish West-Indies.

This juice is clear and transparent, of a whitish or pale yellowish colour, an agreeable (mell, and a bitterish pungent taste. It is usually about the confistence of oil, or a little thicker: long kept, it becomes nearly as thick as honey, retaining its clearness; but has not been observed to grow dry or solid, as most of the other refinous juices We fometimes meet with a thick fort of balfam of Copaiba, which is not at all transparent, or much less so than the foregoing, and generally has a portion of turbid watery liquor at the bottom. This fort is probably either adulterated by the mixture of other substances, or has been extract-

H 2 e

ed by coction from the bark and branches of the tree; its smell and taste are much less pleasant than those of the genuine balsam.

Pure balsam of Copaiba dissolves entirely in rectified spirit, especially if the menstruum be previously alkalized: the solution has a very fragrant smell. Distilled with water, it yields a large quantity of a limpid essential oil; and in a strong heat, without addition, a blue oil.

The balfam of Copaiba is an useful corroborant detergent medicine, accompanied with a degree of irritation. It strengthens the nervous system, tends to loosen the belly, in large doses proves purgative, promotes urine, and cleanses and heals exulcerations in the urinary passage, which it is supposed to perform more effectually than any of the other balsams. Fuller observes, that it gives the urine an intensely bitter taste, but not a violet smell as the turpentines do.

This balsam has been principally celebrated in gleets and the fluor albus, and externally as a vulnerary. The author above mentioned recommends it likewise in dysenteries, in scorbutic cachexies, in diseases of the breast and lungs, and in an acrimonious or putrescent state of the juices: he says, he has known very dangerous coughs which manifestly threatened a consumption, cured by the use of this balsam alone; and that notwithstanding its being hot and bitter, it has good effects even in heetic cases.

The dose of this medicine rarely exceeds twenty or thirty drops, though some direct fixty or more. It may be conveniently taken in the form of an elæosaccharum; or in that of an emulsion, into which it may be reduced by triturating it with almonds, or rather with a thick mucilage of gum arabic,

till they are well incorporated, and then gradually adding a proper quantity of water.

The only officinal preparation of this balfam is an empyreumatic oil distilled with the addition of gum guaicum [L.] The balfam itself is an ingredient in the balfamic tincture, and tincture of cantharides [E.]

BALSAMUM GILEADENSE, vide Opobalsamum.

BALSAMUM PERUVIANUM

[L. E.] Balfam of Peru.

The common Peruvian balfam is faid to be extracted by coction in water, from an odoriferous fhrub growing in Peru, and the warmer parts of America. This balfam, as brought to us, is nearly of the confiftence of thin honey, of a reddish brown colour, inclining to black, an agreeable aromatic fmell, and a very hot biting taste. Distilled with water, it yields a small quantity of a fragrant essential oil of a reddish colour; and in a strong fire, without addition, a yellowish red oil.

Balfam of Peru is a very warm aromatic medicine, confiderably hotter and more acrid than Copaiba. Its principal effects are, to warm the habit, to strengthen the nervous system, and attenuate viscid humours. Hence its use in some kinds of asthmas, gonorrhœas, dysenteries, suppressions of the uterine discharges, and other disorders proceeding from a debility of the folids, or a sluggishness and inactivity of the juices. It is also employed externally, for cleaning and healing wounds and ulcers; and fometimes against palsies and rheumatic pains.

This balfam does not unite with water, milk, expressed oils, animal fats, or wax: it may be

mingled

mingled in the cold with this last, as also with the sebaceous substance called expressed oil of mace; but if the mixture be afterwards liquested by heat, the balsam separates and falls to the bottom. It may be mixed with water into the form of an emulsion after the same manner as the balsam of Copaiba. Alkaline lixivia dissolve great part of it; and rectified spirit the whole.

This balsam is an ingredient in the balsamum guaiacinum, pilulæ aromaticæ [L. tinctura balsamica, elixir petorale, balsamum ceptalicum, and

balsamum Locatelli [E.]

This is another fort of balfam of Peru, of a white colour, and confiderably more fragrant than the former. This is very rarely brought to us. It is faid to be the produce of the fame plant which yields the common or black balfam; and to exude from incifions made in the trunk.

BALSAMUM TOLUTANUM [L. E.] Balfam of Tolu.

This flows from a tree of the pine kind, growing in Tola in the Spanish West-Indies; from whence the balsam is brought to us in little goard shells. It is of a yellowish brown colour, inclining to red; in confistence thick and tenacious: by age it grows hard and brittle, without suffering any great loss of its more valuable parts. The imell of this balfam is extremely fragrant, somewhat resembling that of lemons; its taste warm and fweetish, with little of the pungency, and nothing of the nauseous relish, which accompany the other balsams. It has the same general virtues with the foregoing: but is much milder, and for some purpofes, particularly as a corroborant in gleets and seminal weaknesses,

is supposed to be more efficacious.

It is an ingredient in the wulnerary

balfam [L] the balfamic tincure, and the pectoral pills and elixir [E] A syrup also is impregnated with it in the shops.

BARDANÆ MAJORIS feu lappæ majoris, radix et femen: Lappæ majoris, arcii Dioscordis C. B. Burdock; the roots and

feeds [E.]

This is a common plant about way fides, fufficiently known from its fealy heads, or burs, which flick to the clothes. - The feeds have a bitterish subacrid taste: they are recommended as very efficacious diuretics, given either in the form of emulsion, or in powder, to the quantity of a drachm .- The roots tafte sweetish, with a flight aufterity and bitterishness: they are esteemed aperient, diuretic, and sudorific: and said to act without irritation, so as to be fasely ventured upon in acute disorders. Decoctions of them have of late been used in rheumatic, gouty, and other disorders: and preferred by some to those of sarsaparilla.

BDELLIUM [L. E.] Bdellium. Bdellium is a gummy-refinous concrete juice brought from Arabia and the East-Indies, in glebes of different figures and magnitudes. It is of a dark reddish brown colour, and in appearance somewhat resembles myrrh: upon cutting a piece, it looks somewhat transparent, and as Geoffroy justly observes, like glue. It grows foft and tenacious in the mouth, flicks to the teeth. has a bitterish taste, and not a disagreeable smell. Bdellium is recommended as a sudorific, diuretic, and uterine; and in external applications for maturating tumors, &c. In the present practice, it is scarce otherwise made use of than as an ingredient in the theriaca.

H 3 BECA-

BECABUNGE, seu Anagallidis aquaticæ folia: Veronicæ aquaticæ folio subrolundo Moris. bis. Brooklime; the leaves [L. E.]

This is a low plant, common in little rivulets and ditches of standing water: the leaves remain all the winter, but are in greatest perfection in the spring. Their prevailing taste is an herbaceous one, accompanied with a very light bit-

terishness.

Becabunga has been fupposed to have a saponaceous detergent virtue, and to attenuate viscid humours without pungency or irritation: hence it has been directed in the species of scurvy called hot, where the cochlearie, and other acrid antiscorbutics, were supposed to be less proper. It is now used only in composition with those plants, as in the succi scorbutici [L. E.] but does not perhaps add much to their efficacy. If any virtue is expected from becabunga, it should be used as food.

BELLIDIS MAJORIS folia: Bellidis majoris sylvestris caule folioso C. B. Greater or ox-eye daisy;

the leaves [E.]

This plant is frequent in fields, and among corn, flowering in May and June. The leaves have a mucilaginous, subsaline, rough-They are said to be deish taste. tergent, resolvent, aperient, and also moderately astringent. Geoffroy relates, that the herb, gathered before the flowers have come forth, and boiled in water. imparts an acrid tafte, penetrating and subtile like pepper; and that this decoction is an excellent vulnerary and diuretic: but this account feems to belong more properly to the following plant.

BELLIDIS MINORIS frue confolidæ minimæ foliu: Bellidis minoris sylvestris C. B. Common daisy; the leaves [E.] This is common almost every where, and slowers early in the spring.—The leaves have a subtile subacrid taste, and are recommended as vulneraries, and in asthmas and hectic severs, and such disorders as are occasioned by drinking cold liquors when the body has been much heated. Ludovici prefers the bellis minor to the plants commonly used as antiscorbutics, and resolvents of coagulated blood in hypochondriacal disorders.

BENZOINUM [L. E.] Benzoine.

Benzoine is a concrete refinous juice, obtained from a large tree growing naturally in both the Indies, and hardy enough to bear the winters of our own climate. The refin is brought from the East-Indies only; in large masses, composed of white and light brown pieces, or yellowish specks, breaking very easily betwixt the hands: such as is whitest, and siee from impurities, is most esteemed.

This refin has very little tafte, impressing only a light sweetness on the tongue; its smell is extremely fragrant and agreeable, especially when heated. Committed to the fire in proper vessels, it yields a considerable quantity of a white saline concrete, called flowers, of an acidulous taste and grateful odour, soluble in rectified spirit, and by the affishance of heat

in water.

The principal use of benzoine is in perfumes, and as a cosmetic; it is rarely met with in extemporaneous prescription, and enters in substance only one officinal composition, the balfamum traumaticum, designed chiefly for external use. It should nevertheless seem applicable to other purposes, and to have no ill title to the virtues of storax and balfam of Tola.

Tolu, at least in a subordinate degree. The slowers are recommended in disorders of the breast; and in this intention they are made an ingredient in the paregoric elixir [L.] in the pestoral elixir, electuary, and pills, and in the traches of fulphur [E.]

BERBERIS, seu oxyacanthae Galeni, cortex et frustius: Berberis dumetorum C. B. Barberry; the bark and fruit.

The barberry is a small tree, or rather a large bush, covered with an ash-coloured bark, under which is contained another of a deep yellow: the berries are of an elegant red colour, and contain each two hard brown seeds. It grows wild on chalky hills in several parts of England; and is frequently planted in hedges and in gardens.

The outward bark of the branches, and the leaves, have an aftringent acid tafte; the inner yellow bark, a bitter one; this last is said to be serviceable in the jaundice; and by some, to be an

nieful purgative. The berries, which to the tafte are gratefully acid, and moderately refiringent, have been given with good success in bilious fluxes, and diseases proceeding from heat, acrimony, or thinness of the Among the Egyptians, barberries are employed in fluxes, and in malignant fevers, for abating heat, quenching thirst, raising the firength, and preventing putrefaction; the fruit is macerated for a day and night, in about twelve times its quantity of water, with the addition of a little fennel feed, or the like, to prevent offence to the flomach; the liquor strained off, and sweetened with fugar, or fyrup of citrons, is riven the patient liberally to drink. Prosper Alpinus (from whose treatise De medicina Egyptiorum this account is extracted) informs us, that he took this medicine himself, with happy success, in a pestilential sever, accompanied with an immoderate bilious diarrhœa. A gelly of the fruit is directed by the Edinburgh college as an officinal.

BET A folia: Bete albe vel pallescentis que sicula et cicla officinarum Mor. et Bete rubre vulgaris C. B. et Bete rabre radice rape C. B. White and red beets; and the turnep-rooted red beet, or beetrave [E.]

These plants are cultivated in gardens, chiefly for culinary use. The eye distinguishes little other difference betwixt them, than that expressed in their titles. Decoctions of beets gently loosen the belly; hence they have been ranked among the emollient herbs: the plants remaining after the boiling are supposed to have rather a contrary effect. They afford little nourishment, and are faid by some to be prejudicial to the stomach. The juice expressed from the roots, is a powerful errhine.

BETONICÆ folia: Betonicæ purpureæ C. B. Common or wood betony; the leaves [E.]

Betony is a low plant, growing in woods and shady places, in several parts of England: the slowers come forth in June and July: they are of a purplish colour, and stand in spikes on the tops of the stalks. The leaves and slowers have an herbaceous, roughish, somewhat bitterish taste, accompanied with a very weak aromatic slavour. This herb has long been a savourite among writers on the Materia Medica, who have not been wanting to attribute to it abundance of good qualities. Expe-

rience does not discover any other virtue in betony, than that of a mild corroborant; as such, an infusion or light decoction of it, may be drank as tea, or a faturated tincture in recined spirit given in fuitable doses, in laxity and debility of the viscera, and disorders proceeding from thence. powder of the leaves, snuffed up the nose, provokes sneezing; and hence betony is sometimes made an ingredient in sternutatory powders; this effect does not feem to be owing, as is generally supposed, to any peculiar stimulating quality in the herb, but to the rough hairs which the leaves are covered with. The roots of this plant differ greatly in quality from the other parts: their taste is bitter and very nauseous: taken in a small dose, they vomit and purge violently, and are supposed to have somewhat in common with the roots of hellebore. It is pretty fingular, if true, that belony affects those who gather any confiderable quantity of it with a disorder resembling drunkenness; as affirmed by Simon Paulli and Bartholinus.

BETONICA AQUATICA, vide Scrophularia aquatica major.

BETONICA PAULI, vide VERONICA MAS.

BETULÆ cortex et lachryma: Betulæ C.B. The birch tree: the

bark and sap. [E.]

This tree grows wild in moift woods; its bark confifts of a thick brittle substance of a brownish red colour; and of several very thin, smooth, while, transparent membranes. These last are highly inflammable, and appear to abound with resinous matter, though scarcely of any particular smell or

tafle; the thick brittle part is less resinous, and in taste roughish: of the medical virtues of either, little or nothing is known with certain-

tv.

Upon deeply wounding or boring the trunk of the tree in the beginning of spring, a sweetish juice issues forth, sometimes, as is said, in so large quantity as to equal in weight the whole tree and root; one branch will bleed a gallon or more in a day. This juice is chiefly recommended in scorbutic disorders, and other soulnesses of the blood: its most sensible effect is to promote tht urinary discharge.

BEZOAR lapis. Bezoar stone. The bezoar stone is a calculous concretion found in the stomach of certain animals which are said to be of the goat kind. It is composed of concentrical coats surpunding one another, with a little cavity in the middle, containing a bit of wood, straw, hair,

or the like substances. The shops distinguish two forts of bozoar, one brought from Persia and the East Indies, the other from the Spanish West-Indies. The first. or best fort, called oriental bezoar, is of a shining dark green or olive colour, and an even imooth furface; on removing the outward coat, that which lies underneath it appears likewife smooth and shining. The occidental has a rough furface, and less of a green colour than the foregoing; it is likewise much heavier, more brittle, and of a loofer texture; the coats are thicker, and on breaking exhibit a number of firiz curiously interwoven. The oriental is generally less than a walnut; the occidental for the most part larger, and sometimes as big as a goofe egg. The first is universally most esteemed, and is

the only fort now retained by the London college: the Edinburgh, in the edition of their pharmacopœia preceding the present, directed both; but they now seem to allow them to be used promiscuously, retaining in their catalogue only the name bezoar lapis.

Kæmpler (in whose Amanitates ex tice, a full account of the bezoar animal may be feen) informs us, that this stone is in high escem among the Persians, and even of greater value than in Europe; this, with fundry other circumstances needless to relate here, has given occasion to many to suspect, that the true bezoar is never brought to us. Some authors relate with great confidence, that all the stones commonly fold under this name are artificial compositions. That some of them are fo. is evident; hence the great differences in the accounts which different persons have given of their qualities : the stones examined by Slare as oriental bezoar did not dissolve in acids; those which Grew and Boyle made trial of did: those employed by Geoffroy (in fome experiments related in the French memoirs 1710) did not feem to be acted on by rectified spirit; whilst some of those examined by Neumann at Berlin almost totally dissolved therein. The common mark of the goodness of this stone, is its striking a deep green colour on white paper that has been rubed with chalk.

Bezoar was not known to the ancient Greeks, and is first taken notice of by the Arabians, who extol it in a great variety of disorders, particularly against poisons. Later writers also bestow extraordinary commendations on it as a sudorific and alexipharmac; yirtues to which it certainly has no pretence. It is a morbid concre-

tion, much of the fame nature with the human calculus, of no fmell or talle, not digestible in the flomach of the animal in which it is found, and scarce capable of being acted on by any of the juices of the human body. It cannot be confidered in any other light than as an absorbent; and is much the weakest of all the common substances of that class. It has been given to half a dram, and sometimes a whole dram, without any sensible effect; though the general dose (on account of its great price) is only a few grains.

BISMALVA, vide ALTHEA.

BISMUTHUM. Bismuth. Bismuth is a ponderous brittle metal, resembling in appearance the antimonial regulus and zinc. but greatly differing from them in quality. It dissolves with vehemence in the nitrous acid, which only corrodes the regulus of antimony: and is scarce at all soluble in the marine acid which acts strongly on zinc. A calx and flowers of this semimetal have been recommended as similar in virtue to certain antimonial preparations but are at present of no other use than as a pigment or cosmetic. See Part III. chap. ix.

BISTORT E radix: Bistorte majoris radice minus intorta C. B. Bittort, or snakeweed; the root [L E.]

This plant grows wild in moift meadows, in feveral parts of England; but is not very common about London. The root is about the thickness of the little finger, of a blackish brown colour on the outside, and reddish within: it is writhed or bent vermicularly (whence the name of the plant) with a joint at each bending, and

full of bushy fibres; the root of the species here intended has, for the most part, only one or two bendings; others have three or more.

All the parts of bistort have a rough austere taste, particularly the root, which is one of the Arongest of the vegetable astringents. It is employed in all kinds of immoderate hamorrhages and other fluxes, both internally and externally, where aftringency is the only indication. It is certainly a very powerful flyptic, and is to be looked on fimply as fuch; the fudorific, antipestilential, and other like virtues attributed to it, it has no other claim to, than in confequency of its aftringency, and of the antiseptic power which it has in common with other vegetable flyptics. The largest dose of the root in powder is one dram. It enters only one officinal composition, he species e scordio [L.]

BITUMEN JUDAICUM [L.]

Asphaltus Jews pitch.

I his is a light, folid bitumen, of a dusky colour on the outside, and a deep shining black within; of very little taste, and scarcely any smell, unless heated, when it emits a strong pitchy one. It is found plentifully in the earth, in several parts of Egypt, and on the surface of the Dead Sea; but is very rarely brought to us. In its room, the shops employ other bituminous substances found in France, Germany, and Switzerland: these have a much stronger, pitchy fmell; but in other respects agree pretty much with the true afphaltus. Sometimes pitch itself, or the caput mortuum remaining after the distillation of amber, are substituted. Abundance of virtues are attributed to this bitumen, as refolvent, discutient, glutinant, su-'dorific, emollient, emmenagogue,

•

&c. but it has not for a long time been any otherwise used, than as an ingredient in theriaca. The Edinburgh college having now expunged the theriaca, have expunged also the bitumen Jadaicum.

BOLI. Boles are viscid earths, less coherent, and more friable than clay, more readily uniting with water, and more freely subsiding from it. They are soft and unctuous to the touch, adhere to the tongue, and by degrees melt in the mouth, impressing a light sense of astringency. A great variety of these kinds of earths have been introduced into medicine; the principal of which are the following.

- (1) BOLUS ARMENA. Armenian bole, or bole-armenic [L. E.]
  Pure Armenian bole is of a bright red colour, with a tinge of yellow; it is one of the hardest and most compact of the bodies of this class, and not smooth or glossy like the others, but generally of a rough dusty surface. It raises no effervescence with acids.
- (2) BOLUS GALLICA. French bole [L. E.] The common French bole is of a pale red colour, variegated with irregular specks or veins of white and yellow. It is much softer than the foregoing; and slightly effervesces with acids.
- (3) BOLUS BLESENSIS. Bole of Blois. This is a yellow bole, remarkably lighter than the former, and than most of the other yellow earths. It effervesces strongly with acids.
- (4) Bolus Bohemica. Bohemian bole. This is of a yellow colour, with a cast of red, generally of a staky texture. It is not acted on by acids.

- (5) TERRA LEMNIA. Lemnian earth. This is a pale red earth; slightly effervescing with acids.
- (6) TERRA SILESIACA. Silefian earth, is of a brownish yellow colour: acids have no sensible effect upon it. These and other earths, made into little masses, and stamped with certain impressions, are called terra sigillata.

The boles of Armenia and Blois, and the Lemnian earth, are rarely met with genuine in the shops; the coarser boles, or white clay coloured with ochre, caput mortuum of vitriol, &c. frequently supply their place. The genuine may be distinguished by their subsiding uniformly from water, without any separation of their parts; the genuine yellow boles retain their colour, or have it deepened, in the fire; whilst the counterseit forts burn red.

These earths have been recommended as astringent, sudorisic, and alexipharmac; in diarrhœas, dysenteries, hæmorrhages, and in malignant and pestilential distempers. In intestinal sluxes, and complaints in the first passages from thin acrimonious humours, they may, doubtless, be of some use; but the virtues ascribed to them in the other cases appear to have no foundation.

In the London pharmacopæia bole is an ingredient in the pulvis e bolo, e feordio, tabellæ cardialgicæ, theriaca, and in one composition for external use, viz. the lapis medicamentosus. No earth of this kind is employed in any of the compositions of the Edinburgh pharmacopæia.

BOMBYX, vide Sericum.

BONI HENRICI five lapathi unduofi folia: Lapathi unduofi olidi perennis spinachiæ facie Moris. English herb mercury; the leaves [E.]

This herb is met with by road fides, and in uncultivated places. It is ranked among the emollient herbs, but rarely made use of in practice. The leaves are applied by the common people for healing slight wounds, cleansing old ulcers, and other like purposes.

BORRAGINIS flores: Boraginis flore caruleo J. B. Borage; the flowers [E]

This is a rough plant, clothed with small prickly hairs: it grows wild in waste places, and apon old walls. An exhilarating virtue has been attributed to the slowers of borage, which are hence ranked among the so called cordial flowers: but they appear to have very little claim to any virtue of this kind, and seem to be altogether insignificant.

BORAX [L. E.] Tincar, or Borax.

This is a faline fubstance, brought from the East-Indies in great masses, composed partly of large crystals, but chiefly of smaller ones, partly white and partly green, joined together as it were by a greasy yellow substance, intermingled with sand, small stones, and other impurities: the purer crystals, exposed to the fire, melt into a kind of glass, which is nevertheless dissoluble in water.

This falt dissolved and crystallized, forms small transparent masses: therefiners have a method of shooting it into larger crystals; but these differ in several respects from the genuine salt, insomuch that Cramer calls them not a purified, but adulterated borax. The origin of this salt is as yet unknown, and its composition is known only in part. Thus much experiments

have

have clearly shewn, that it consists of a fixt alkaline falt, the same with the basis of sea salt, in some degree neutralized by a smaller proportion of another faline fubflance, which has no where, that I know of, been yet discovered but in borax itself.

Nor have the medical virtues of borax been sufficiently ascertained by experience: it is supposed to be, in doses of half a dram or two scruples, diuretic, emmenagogue, and a promoter of delivery; the only officinal composition it is employed in is the pulvis ad partum of the Edinburgh pharmacopoia. Mr. Biffet, in an essay on the medical constitution of Great Britain, recommends a solution of this falt in water as the most powerful dissolvent yet known of aphthous crusts in the mouth and fauces of children. There are strong reasons to believe, that the virtues of borax are much greater than they are in general supposed to be.

BOTRYOS folia: Chenopodii embrofioidis folio finucio Tourn. Atriblicis odora seu suaveolentis Morif. lerusalem oak : the leaves [E.]

This plant is cultivated in gardens. In has a strong not disagreeable smell; and a warm somewhat pungent taste. It is recommended as a carminative and pectoral. Infusions of it may be drank in tea.

BRASSICA SATIVA: Braffica eapitata alba C. B. et Brassica capitata rubra C. B. et Braffica rubra C. B. et Brossica alba capite oblongo non penitus claufo C. B. Brafsica subauda Ger. et Park. et Brasfica cauliflora C. B. White and red cabbages, coleworts, Savoy cabbages, and cauliflower [E.]

. These are cultivated in gardens rather for culinary than medicinal They are all supposed to be hard of digestion, to afford little nourishment, and to produce flatulencies: though probably on no very good foundation. They tend strongly to putrefaction, and run into this state sooner than almost any other vegetable; when putrefied, their smell is likewise the most offensive, greatly resembling that of putrefied animal substances. A decoction of them is faid to loosen the belly. Of all these plants, cauliflower is reckoned the easiest of digestion. The white is the most fetid: and the red most emollient or laxative: a decoction of this last is recommended for foftening acrimonious humours in some disorders of the breast, and in hoarseness.

BRASSICÆ MARINÆ fes foldanella folia: Convolvuli maritimi soldanellæ ditti Raii Sea coleworts, Scotch scurvy grafs, or foldanella; the leaves [E.]

This is a trailing plant, growing on the fea beach in many parts of the north of England. The root, leaves, and stalks, yield a milky

Soldanella is a strong cathartic, operating very churlishly, and hence deservedly rejected from practice. Those who recommend its use differ considerably with regard to the dose; some direct half a dram, others three drams, and others a whole handful.

BRITANNICA, vide LAPA-THUM.

BRUNELLA, vide PRUNELLA.

BRUSCUS, vide Ruscus.

BRYONLE ALBÆ radix: Bryoniæ asperæ søve albæ baccis rubris C. B. White bryony, or wild vine: the roots [E.] This This is a rough plant, growing on dry banks under hedges, and climbing upon the bushes. The roots are large, sometimes as thick as a man's thigh: their smell, when fresh, is strong and disagreeable; the taste nauseously bitter, acrid, and biting: the juice is so sharp, as in a little time to excoriate the skin; in drying, they lose great part of their acrimony, and almost the whole of their scent.

Bryony root is a strong irritating cathartic; and as such has sometimes been successfully exhibited in maniacal cases, in some kinds of dropties; and in feveral chronical disorders, where a quick solution of viscid juices, and a sudden Rimulus on the folids, were required. An extract prepared by water, acts more mildly, and with greater fafety than the root in substance; given from half a dram to a dram, it is faid to prove a gentle purgative, and likewise to operate powerfully by urine.

Bryony root, applied externally, is faid to be a powerful discutient: it enters a cataplasm for that intention in the Edinburgh pharmacopoeia.

BUFO. The toad.

This animal has been generally looked upon as poisonous, particularly its faliva, and a certain acrid liquor, supposed to be the urine, which it throws out, when irritated, to a confiderable diffance. It was first introduced into medicine upon occasion of a cure performed on an hydropic person, to whom powdered toads were given in order to dispatch him, but who voided a large quantity of urine after taking them, and soon recovered of his disorder: since this time, the toad dried by a gentle heat and pulverized, has been greatly effeemed as a diuretic. This preparation is faid likewise, applied externally to the navel, to restrain hæmorrhages, particularly those from the uterus. The Edinburgh college, in the preceding edition of their pharmacopeais, estained the toad in their catalogue of simples, and gave likewise the process of drying it, but have now wholly rejected this loathsome animal.

BUGLOSSI radix, folia, flores: Buglossi augustifolii majoris C. B. Garden bugloss; the roots, leaves,

and flowers [E.]

This is a rough, hairy plant, refembling borage, but less prickly: a wild fort is commonly met with in hedges and among corn, which differs from the garden only in being smaller. Bugloss has a slimy sweetish taste, accompanied with a kind of coolness: the roots are the most glutinous, and the flowers the least so. These qualities point out its use in hot bilious or inflammatory distempers, and a thin acrimonious state of the sluids. The flowers are one of the four called cordial flowers: the only quality they have that can entitle them to this appellation, is, that they moderately cool and foften, without offending the palate or ftomach; and thus in warm climates, or in hot diseases, may in some measure refresh the patient.

RUGULÆ five confolidæ mediæ foliu: Bugulæ fylvæticæ vulgæris cæruleæ Morison. Bugle or middle consound; the leaves [E.]

This grows wild in woods, hedges, and moist meadows. The leaves have at first a sweetish taste, which gradually becomes bitterish and roughish. They are recommended as vulnerary medicines, and in all cases where mild astringents.

aftringents or corroborants are proper.

BUNIAS, vide NAPUS.

BURSÆ PASTORIS folia: Thlaspis fatui, bursæ pastoris didi Raij. Shepherdspurse; the leaves.

This plant is common in waste places; and is found in flower all the fummer. Shepherdspurse has long been celebrated as an astringent, and strongly recommended in diarrheas, dysenteries, uterine fluors, and in general in all diseases where aftringents of any kind can avail. Some have esteemed it so powerful a styptic, as scarce to be safely exhibited internally. Others have thought it to be of a hot fiery nature, and supposed it to stop sluxes and hæmorrhges, by coagulating the juices like alcohol, and burning or fearing the orifices of the vessels. The fenfible qualities of shepherdspurse discover little foundation for either of these opinions; it has no perceptible heat, acrimony, pungency, and scarcely any attringency: the talle is almost merely harbaceous, fo as sufficiently to warrant the epithet given this plant by Mr. Ray, Fatuum.

BUXI lignum [L.] et folia [E.]: Buxi arborescentes C. B. The box tree: the leaves and wood.

The box is a small tree, growing wild in some parts of Kent and Surry. The wood is of a yellow colour, more solid, compact, and ponderous than any other of the European woods. The leaves have a strong nauseous taste, and when fresh, a fetid smell: they are said to purge violently, in the dose of a dran. A decoction of the wood is recommended by some as powerfully sudorisic, preferable even to guaiacum: but the taste readily discovers that it wants the qualities

of that wood. Neither the wood nor leaves of the box tree are at present employed for any other medicinal purpose than for the distillation of an empyreumatic oil [L.] and an oil of nearly the same quality is obtainable by the same treatment from almost all woods.

CACAO [E] Chocolate nuts.
These are the fruit of an American tree resembling the almond.
The principal use of these nuts is for the preparation of the dietetic liquor chocolate. This is a mild, unctuous, nutritious sluid, capable of softening acrimonious humours, and of great service in consumptive disorders; especially if made with milk, and with only a small proportion of aromatics.

CALIMINARIS LAPIS [L. E.] Calamy or calamine stone.

This mineral is found plentifully in England, Germany, and other countries, either in dittina mines, or intermingled with the ores of different metals. usually of a greyish, brownish. yellowish, or pale reddish colour: confiderably hard, though not fufficiently so to strike fire with steel. It has been looked upon by fome as a fimple earth, by others as an iron ore: later experiments have discovered it to be an ore of zinc. Calamine is generally roafted or calcined before it comes into the shops, in order to separate some fulphureous or arienical matter which the crude mineral is funposed to contain, and to render it more easily reducible into a fine powder. In this state, it is employed in collyria against defluxions of thin acrid humous upon the eyes; for drying up moif, running ulcers; and healing excoriations. It is the basis of an officinal epulotic cerate.

CALA-

CALAMINTHÆ folia: Calaminthæ puligii odore seu nepetæ C. B. calaminthæ foliis ovætis, obtusis, caule procombente Halleri. Field calamint; the leaves [L.]

This is a low plant, growing wild about hedges and highways, and in dry fandy foils. The leaves have a quick warm tafte, and fmell ftrongly of pennyroyal: as medicines, they differ little otherwife from spearmint, in than being somewhat hotter, and of a less pleasant odour; which last circumstance has procured calamint the presence in hysteric cases.

CALAMINTHÆ MONTANÆ folia: Colaminthæ flore magno vulgaris J. B. Common calamint; the leaves [E.]

This plant, notwithstanding its name, is, among us, much less common than the former, which has generally supplied its place in the markets: hence the London college have now dropt this montana, and received the other. The calamintha montana is also less efficacious than the foregoing fort: the taste is weaker: the smell approaches to that of the wild mints, without any thing of the strong pennyroyal slavour of the other.

CALAMI AROMATICI radix:

Actori weri five calami aromatici officinarum C. B. Sweet scented flag;
the roots [L. E.]

This flag resembles, as to its leaves, the common iris, but in other respects differs greatly from it: the stalk growsatalittle distence from the leaves: the lower half, up to where the slowers came forth, is roundish; the part above this broad like the other leaves; the slowers are very small, whitish, and stand in a kind of head about the size of a singer. This plant grows plentifully in rivulets and marshy

places, about Norwich and other parts of England; in the canals of Holland; in Switzerland; and in other countries of Europe. The shops have been usually supplied from the Levant with dried roots, which do not appear to be superior to those of our own growth.

The root of acorus is full of ioints, crooked, fomewhat flatted on the fides, internally of a white colour, and loofe spongy texture : its smell is strong; the taste warm, acrid, bitterifb, and aromatic; both the fmell and tafte are improved by exficcation. This root is generally looked upon as a carminative and stomachic medicine, and as such is sometimes made use of in practice. It is said by some to be superior in aromatic flavour to any other vegetable that is produced in these northern climates: but fuch as I have had an opportunity of examining, fell short, in this respect, of several of our common plants. It is, nevertheless, a sufficiently elegant aromatic. It is an ingredient in the mithridate and theriaca of the London pharmacopæia, and in the aromatic and stomachic tinctures, and compound arum powder, of the Edinburgh. The fresh roots candied after the manner directed in our dispensatory for candying eryngo root, is faid to be employed at Constantinople as a preservative against epidemic diseases. leaves of this plant have a sweet fragrant smell, more agreeable, though weaker, than that of the roots.

CALENDULÆ flores: Calendulæ flore fativæ Rait: Calendulæ flore fimplici J. B. Garden marigold; the flowers [E.]

This herbis common in gardens, where it is found in flower greatest part of the summer. Marigold flowers are supposed to be aperient

BADO

and attenuating; as also cardiac. alexipharmac, and fudorific: they are principally celebrated in uterine obstructions, the jaundice, and for throwing out the small-pox. Their sensible qualities give little foundation for these virtues: they have scarcely any taste, and no consider-The leaves of the able fmell. plant discover a viscid sweetishness. accompanied with a more durable faponaceous pungency and warmth: these seem capable of answering some useful purposes, as a stimulating, aperient, antiscorbutic medicine.

CALX VIVA [L E] Quicklime. Quicklime is usually prepared among us, by calcining certain stones of the chalky kind. All chalks and marbles, and in general all the mineral earths that dissolve in acids, burn intoquicklime; with this difference, that the more compact the stone, generally the stronger is the lime. In maritime countries, in desect of the proper stones, sea shells are made use of, which asford a calx agreeing in most respects with the stone limes.

All these limes are, when fresh buint, highly acrimonious and corrosive. In this state they are emplyed in some external applications as a depilatory; for rendering sulphur soluble in water [L]; and for encreasing the power of fixt alkames falts either for the purposes of a caustic [L.E.] or to enable them more readily to dissolve oils for making soap [L.] If the lime be exposed for a length of time to the air, it falls by degrees into a powder, and loses greatly of its acrimony.

Water poured directly upon quicklime, takes up a confiderable portion of it: the solution has a strong taste, somewhat styptic, drying the mouth, and accompa-

nied with a kind of sweetishness. This liquor does not effervesce either with acids or alkalies, but is rendered by the latter turbid and milky: it prevents the coagulation of milk, and hence is sometimes made use of along with milk diets: agitated with expressed oils, it unites with them into a thick compound, recommended by Dr. Slare against burns and inflammations. Both the simple folution of the lime, and the folution impregnated with other materials, are directed as officinal, under the title of fimple and compound lime waters [L. E.]

Lime water drank to the quantity of a quarter of a pint three or four times aday, and continued for a length of time, has been found ferviceable in scrophulous cases. and other oblinate chronic diforders It generally promotes urine, and not unfrequently the cuticuler discharge: for the most part it binds the belly, and sometimes produces troublesome costiveness, unless this effect be occasionally provided against, by the interposition of proper medicines. It does good service in debility and laxity of the viscera in general; in those of the uterine and seminal vessels it is particularly recommended. most he had not to use this medieine too liberally in hot bilious confitutions, or where the patient is much emaciated, or the appetite weak, or at the time of any critical or periodical evacuation. Its principal use is in cold, moitt, fluggift, and corpulent habits.

This liquor has lately been found an efficacious dissolvent of the human calculas: the lime water prepared from calcined oyster shells proves, for this purpose, a more powerful menstruum than that made from the stone limes, the dissolving power of the former being more than double to that of the latter. See a paper on this subject in the Edinburghessays, vol. v. art. 69. Abridg. vol. i. p. 471.

CAMPECHENSE LIGNUM, vide LIGNUM CAMPECHENSE.

CAMPHORA [L. E.] Camphor is a folid concrete, extracted from the wood and roots of a tree growing in Japan, by a process fimilar to that whereby effential oils are obtained. As it first sublimes from the wood it appears brownish, composed of semipellucid grains mixed with dirt: in this state it is exported by the Dutch, and purified by a second sublimation: after which, it is reduced into loaves (in which it is brought to us) probably by fusion in close veffels; for it does not assume this form in fublimation.

Pure camphor is very white, pellucid, somewhat uncluous to the touch; of a bitterish, aromatic, acrid tafte, yet accompanied with a sense of coolness; of a very fragrant smell, somewhat like that of rosemary, but much stronger. It is totally volatile, and inflammable; foluble in vinous spirits, oils, and the mineral acids; not in water, alkaline liquors, or the acids of the vegetable kingdom. This concrete-is effeemed one of the most efficacious diaphoretics; and has long been celebrated in fevers, malignant and epidemical diffempers: in deliria, where opiates fail of procuring sleep, and oftentimes aggravate the symptoms, this medicine frequently fucceeds.

Frederick Hoffman has wrote an express differtation De camphoræ ulu interno securissimo et præstantissimo: the substance of his observations is, that camphor seems to penetrate very quickly through the whole body, and notably increase

perspiration: that though given to the quantity of half a dram, diffolved in spirit of wine, and duly diluted, it does not raise the pulse, or occasion any heat, but rather causes a sense of coolness about the præcordia: that on continuing its use for some time, the blood became sensibly more fluid, and the quantity of watery ferum, which the habit before abounded with, was notably diminished: that in malignant fevers, and all disorders. whether acute or chronical, proceeding from an acrid or putrefcent state of the juices, camphor has excellent effects, correcting the acrimony, expelling the putrid morbific matter through the cutaneous pores, and preventing an inflammation or sphacelus, where there is previously any disposition thereto: that, by strengthening the vessels. it restrains hæmorrhages happening in acute fevers, and promotes critical and periodical evacuations that it expels even the venereal virus; and that he has known examples of the lues being cured by camphor alone, a purgative only being premised: and that in recent infections he has found no medicine equal to it in efficacy. In inflammatory cases, where there is a tendency to mortification, intense heat, thirst, or where the skin is dry and parched, whether before or after a delirium has come on, small doses of camphor joined with nitre produced happy effects, almost immediately relieving the fymptoms, occasioning a calm sleep and plentiful sweat, without fatiguing the patient. He farther observes. that this simple, by its antiphlogistic quality, prevents the ill effects of the more irritating medicines: that cantharides, and the acrid stimulating cathartics and diutetics, by the admixture of a small proportion of camphor, become

much more mild and fafe in operation.

The common dose of camphor is from one grain to ten. Its officinal preparations are, a julep [L.] and emulsion [E.] for internal use; and a solution in rectified spirit [L. E.] and in expressed oil [E.] for external applications. It is an ingredient also in the paregoric elixir, camphorated vitriolic water, [L.] camphorated white cintment, and saponaceous liniment [L E.]

CANCRORUM CHELÆ[L.E] Crabs claws: the black tips of the claws of the common sea crab, or sancer marinus.

CANCRORUM OCULI didi [L. E.] Crabs eyes fo called: stony concretions found in the head, or rather stomach, of the astacus

fluviarilis, or craw fish.

The only virtue of these simples is to absorb acidities in the primæ viæ. The claws enter an officinal lozenge, and give name to a powder, for this intention. They are ingredients also in some other officinal compositions, in which they do not seem to be of much advantage; viz. the compound arum powder, contrayerva powder, and cordial confection.

Crabs eyes are said by most writors on the Materia Medica to be frequently counterfeited with tobacco-pipe clay, or compositions of chalk withmucilaginous substances. This piece of fraud, if really practised, may be very easily discovered; the counterfeits wanting the leafy texture which is observed upon breaking the genuine; more readily imbibing water; adhering to the tongue; and dissolving in vinegar, or the stronger acids diluted with water, either entirely, or not at all, or by piece-meal, whilst the true crabs eyes, digested in these liquors, become foft and transparent, their original form remaining the fame: this change is owing to the earthy part, on which depended their opacity and hardness, being dissolved by the gentle action of the acid, which leaves the conglutinating matter unhurt.

CANELLA ALBA: Cinnamomum five canella tubis minoribus alba C. B. Cortex Winteranus falso dictus Park Canella alba.

This is a bark rolled up into long quills, thicker than cinnamon, and both outwardly and inwardly of a whitish colour, lightly inclining to yellow. It is the produce of a tall tree growing in great plenty in the low lands in Jamaica, and other American islands, called by Sir Hans Sloane arbor baccifera laurifelia aromatica, fructa viridi calycu-lato racemoso. The canella is the interior bark, freed from an outward thin rough one, and dried in the shade. The shops distinguish two forts of canella, differing from one another in the length and thickness of the quills; they are both the bark of the same tree, the thicker being taken from the trunk, and the thinner from the branches. This bark is a warm pungent aromatic, not of the most agreeable kind: nor are any of the preparations of it very grateful. It is lately fometimes met with in extemporaneous prescription, and is an ingredient in the officinal hiera picra and tindura facra [L.] and in the aqua raphani and pulvis ari compositus [E.]

CANNABIS semen: Cannabis sativa C. B. Hemp; the seed [E]

This plant, when fresh, has a rank narcotic smell: the water in which the stalks are soaked, in order to facilitate the separation of the tough rind for mechanic uses, is said to be violently poisonous, and to produce its effects almost

as soon as drank. The seeds also have some smell of the herb; their taste is unctuous and sweetish; on expression they yield a considerable quantity of insipid oil: hence they are recommended (boiled in milk, or triturated with water into an emulsion) against coughs, heat of urine, and the like. They are also said to be useful in incontinence of urine, and for restraining veneral appetites; but experience does not warrant their having any virtues of this kind.

CANTHARIDES [L. E.] Spanish flies. These insects are of a shining green colour, intermingled with more or less of a blue and a gold yellow. They are found adhering to different kinds of trees and herbs, in Spain, Italy, and France; the largest and most esteemed come from Italy.

Cantharides are extremely acrimonious: applied to the skin, they first inflame, and afterwards excoriate the part, raising a more perfect blifter than any of the vegetable acrids, and occasioning a more plentiful discharge of serum. All the bliftering compositions have See Part cantharides for the basis. The external ap-IV. chap. x. plication of cantharides is often followed by a strangury, accompanied with thirst and feverish heat: this inconvenience may be remedied by foft unctuous or mucilaginous liquors liberally drank.

Cantharides taken internally, often occasion a discharge of blood by urine, with exquiste pain: if the dose is considerable, they seem to inflame and exulcerate the whole intestinal canal; the stools become mucous and purulent; the breath fetid and cadaverous; intense pains are selt in the lower belly; the patient saints, grows giddy, raving mad, and dies. All these terrible

consequences have sometimes happened from a few grains. Herman relates, that he has known a quarter of a grain instance the kidneys. and occasion bloody urine with violent pain. There are nevertheless cases in which this stimulating fly. given in larger doses, proves not only safe, but of singular efficacy for the cure of diseases that vield little to medicine of a milder class. In cold phlegmatic fluggish habits, where the vifcera are overloaded. and the kidneys and ureters obstructed with thick viscid mucus matter, cantharides have excellent effects; here the abounding mucous defends the folids from the acrimony of the fly, till it is itself expelled; when the medicine ought to be discontinued. Groenvelt employed cantharides with great fuccess in dropsies, obstinate suppresfions of urine, and ulcerations of the bladder; giving very considerable doses made into boluses with camphor; and interposing large draughts of emulsions, milk, or other emollient liquids: by this means, the excessive irritation. which they would otherwise have occasioned, was in great measure The camphor did not prevented. perhaps contribute fo much to this effect as is generally imagined: fince it has no fensible quality that promises any considerable abates ment of the acrimony of cantharides; nitre would answer all that the camphor is supposed to perform: this, with milk, or emollient mucilaginous liquors, drank in large quantity, are the best correctors. Cantharides, in very small doses, may be given with safety also in other cases. Dr. Mead observes, that the obstinate gleetings which frequently remain after the cure of venereal maladies, and which rarely yield to balfamic medicines, are effectually remedied by cantharides; and that no one remedy is more efficacious in leprous disorders; in which last, proper purgatives are to be occasionally taken during the use of the cantharides. The best and safest preparation of cantharides for these purposes, is a spirituous tincture [L. E.] and indeed in all cases, the tincture is far preferable, for internal use, to the sly in substance.

The virtues of cantharides are extracted by rectified spirit of wine, proof spirit, and water; but do not arise in distillation. The watery and spirituous extracts blister as freely as the fly in substance; whilst the fly remaining after the several menstrua have performed their office, is to the taste insipid, and does not in the least blister, or instance the skin.

CAPILLUS VENERIS, vide ADIANTHUM.

CAPPARIS radicis cortex, et florum gemmæ: Capparis spinosæ fructu minore, folio rotundo C. B. Caper bush; the bark of the root, and buds of the flowers.

This is a low prickly bush, found wild in Italy, and other countries: it is raised with us by sowing the seeds upon old walls, where they take root betwirt the bricks, and

endure for many years.

The back of the root is pretty thick, of an ash colour, with several transverse wrinkles on the surface; cut in slices and laid to dry, it rolls up into quills. This bark has a bitterish acrid taste: it is reckoned aperient and diuretic; and recommended in several chronic disorders, for opening obstructions of the viscera.

The buds, pickled with vinegar, &c. are used at table. They are supposed to excite appetite, and promote digestion; and to be par-

ticularly useful, as detergents and aperients, in obstructions of the liver and spleen. Their taste and virtues depend more upon the saline matter introduced into them, than on the caper buds.

CAPRIFOLII folia et flores: Periclymeni non perfoliati Germanici C. B. Woodbind, or honeyfuckle; the leaves and flowers.

This is a climbing fhrub, common in hedges, the beauty of its flowers has gained it a place also in gardens. The leaves have a disagreeable smell; the flowers a very pleasant one; the taste of both is herbaceous and roughish. They are said to be diuretic and aperient; but practice has not for a long time paid any regard to them.

CAPSICUM, vide PIPER In-

CARABE, vide Succinum.

CARANNA [E.] Caranna.
This is a refinous substance brought from New Spain, and other parts of America, in little masses, rolled up in leaves of slags: it is said to exude from a species of palm tree. This refin is very rarely made use of in medicine, or met with in the shops; whence the London college have rejected it from their catalogue, though it is still retained by the Edinburgh.

CARDAMOMI MAJORIS femen. Greater cardamom seed.

The greater cardamom is adried fruit or pod, about an inch long, containing under a thick skin two rows of small triangular seeds of a warm aromatic slavour.

CARDAMOMI MINORIS femen. Lesser cardamom [L. E.]

This fruit is scarce half the length of the foregoing; the seeds

are confiderably stronger both in fmell and tafte. Hence this fort has long supplied the place of the other in the shops, and is the only one now directed.

Cardamom seeds are a very warm, grateful, pungent aromatic, and frequently employed as fuch in practice: they are faid to have this advantage, that notwithstanding their pungency, they do not, like those of the pepper kind, immoderately heat or inflame the bowels. Both water and rectified spirit extract their virtues by infusion, and elevate them in distillation; with this difference, that the tincture and distilled spirit are considerably more grateful than the infusion and distilled water: the watery infusion appears turbid and mucilaginous: the tincture made in spirit, limpid and transparent. The husks of these seeds, which have very little fmell or tafte, may be commodioully separated, by committing the whole to the mortar, when the feed will readily pulverize, so as to be freed from the shell by the fieve: this should not be done till just before using them: for if kept without the husks, they soon lose confiderably of their flavour. -The officinal preparations of these seeds are a spirituous water and tincture : they are employed also as a spicy ingredient in several of the officinal compositions.

CARDIACÆ folia: Marubii cardiacæ di&i, forte primi Theophrasti C. B. Motherwort; the leaves.

This plant is common in waste places, and found in flower greatest part of the summer. The leaves have a bitter talle, and a pretty firong smell; they are supposed to be useful in hysteric disorders, to strengthen the stomach, to promote urine: and indeed it may be judged from their smell and taste,

that their medical virtues are confiderable, though they are now rejected both from the London and Edinburgh pharmacopæias.

CARDUI BENEDICTI folia semen: Cnici sylvestris birsutioris sive cardui benedicti C. B. Carduii lutei procumbentis, sudorofici et amari Morison. Bleffed thiftle; the leaves

[L. E.] and feed [E.]

Thisis an annual plant, cultivated in gardens: it flowers in June and July, and perfects its feeds in the The herb should be gaautumn. thered when in flower, dried in the shade, and kept in a very dry airy place, to prevent its rotting or growing mouldy, which it is very apt to do. The leaves have a penetrating bitter tafte, not very ftrong, or very durable; accompanied with an ungrateful flavour, which they are in great measure freed from by keeping. Water extracts, in a little time, even without heat, the lighter and more grateful parts of this plant: if the digestion is continued for some hours, the disagreeable parts are taken up: a strong decoction is very nauseous and offensive to the stomach. Rectified spirit gains a very pleasant bitter taste, which remains uninjured in the extract.

The virtues of this plant seem to be little known in the present practice. The nauseous decoction is fometimes used to provoke vomiting; and a strong infusion to promote the operation of other emetics. But this elegant bitter, when freed from the offensive parts of the herb, may be advantageously applied to other purposes. I have frequently experienced excellent effects from a light infusion of carduus in loss of appetite, where the stomach was injured by irregularities. Astronger infusion made in cold or warm water, if drank freely, and the 13

patiem

patient kept warm, occasions a plentiful sweat, and promotes all

the fecretions in general.

The seeds of this plant are also confiderably bitter, and have been fometimes used in the same intention as the leaves. The Edinburgh college makes them an ingredient in the stomachic tincture.

CARICÆ [L.E.]Figs; the dried fruit of the ficus communis C. B.

The principal use of these is as a fost, emollient sweet; in this intention they enter the pectoral decoction and lenitive electary of the shops. They are also esteemed by some as suppuratives, and hence have a place in the maturating cataplaím.

CARLINÆ, seu Chamæleontis albi radix: Carlinæ acqulæ magno flore albo C. B. Carline thistle; the

root [E.]

This is a very prickly fort of thistle, growing spontaneously in the fouthern parts of France, Spain. Italy, and the mountains of Swifferland; from whence the dried roots are brought to us. This root is about an inch thick, externally of a pale rufty brown colour, corroded as it were on the furface. and perforated with numerous small holes, appearing when cut as if worm eaten. It has a strong smell, and a subacrid, bitterish, weakly aromatic taste. Carlina is looked on as a warm diaphoretic and alexipharmac; and has been for some time greatly esteemed by foreign physicians, but never came much into use among us: the present practice has entirely rejected it: nor is it often to be met with in the shops. Frederick Hoffman, the elder, relates, that he has observed a decoction of it in broth to occasion vomiting.

CARPOBALSAMUM: Fruc-

tus balsami Syriaci rutæ folio. C. B.

Carpobalfam [L.]

This is the fruit of the tree that yields the opobalfam, or balm of Gilead. It is about the fize of a pea, of a whitish colour, inclosed in a dark brown wrinkled bark. This fruit, when in perfection, has a pleasant warm glowing taste, and a fragrant smell, resembling, that of the opobalsamum itself. It is very rarely found in the shops ; and fuch as we now and then do meet with, has almost entirely lost its fmell and tafte. It is of no other use in this country than as an ingredient in the mithridate and theriaca, in both which the college direct cubebs as a substitute to it.

CARTHAMI semen: Cartami officinarum flore croceo Tourn. tard faffron, or fafflower; the feeds.

The bastard saffron is a soft kind of thistle, with only a few prickles about the edges of the leaves. It is cultivated in large quantity in some places of Germany; from whence the other parts of Europe are supplied with the flowers as a colouring drug, and the feeds as a medicinal one. The flowers, well cured, are not easily distinguishable by the eye from faffron; but their want of smell readily The feeds are discovers them. white, fmooth, of an oblong roundish shape, yet with four senfible corners, about a quarter of an inch in length, so heavy as to fink in water; of a viscid sweetish taste, which in a little time becomes acrid and nauseous. feeds have been celebrated as a cathartic: they operate very flowly, and for the melt part dilorder the bowels, especially when given in substance; triturated with aromatic distilled waters, they form an emulsion less offensive, yet inferior in efficacy to more common purgatives.

CARUI, carvi, seu cari, semen: Cumini pratensis carui ossicinarum C. B. Caraway; the seeds [L. E.]

Caraway is an umbelliferous plant, cultivated with us in gardens, both for culinary and medi-The feeds have an arocinal use. matic fmell, and a warm pungent taste. These are in the number of the four greater hot feeds; and frequently employed as a stomachic and carminative in flatulent colics. and the like. Their officinal preparations are an effential oil [L. E.] and a spirituous water [L.] they are ingredients also in the compound juniper water, tincture of fena, stomachic tincture, oxymel of garlic, electary of bayberries and of scammony, philonium, and the cummin-seed plaster [L.]

## CARYOPHYLLA AROMA-TłCA [L. E.] Cloves.

Cloves are the flower-cups (not as is generally supposed the fruit) of a bay-like tree, growing in the East-Indies. In shape, they somewhat resemble a short thick nail: in the inside of each clove are found a stylus and stamina with their apices, as in other flowercups: at the larger end shoot out from the four angles four little points like a star, in the middle of which is a round ball, composed of four little leaves, which are the unexpanded petala of the slower.

Cloves have a very strong agreeable aromatic smell, and a bitterish pungent taste, almost burning the mouth and sauces. The Dutch, from whom we have this spice, frequently mix it with cloves which have been robbed of their oil: these, though in time they regain from the others a considerable share both of taste and smell, are

easily distinguishable by their weaker flavour and lighter colour. Cloves, considered as medicines, are very hot stimulating aromatics, and possess in an eminent degree the general virtues of substances of this class. An extract made from them with rectified spirit is excesfively hot and pungent: the distilled oil has no great pungency; an extract made with water is nauseous and fomewhat flyptic. The only officinal preparation of them is the essential oil [L. E.] Both the cloves themselves and their oil are ingredients in many officinal compolitions.

## CARYOPHYLLA RUBRA: Flores Caryophylli altilis majoris C. B. Clove July flowers [L. E.]

A great variety of these flowers are met with in our gardens: those made use of in medicine ought to be of a deep crimson colour, and a pleasant aromatic smell, somewhat like that of cloves: many forts have scarce any smell at all. The carrophylla rubra are said to be cardiac and alexipharmac: Simon Paulli relates, that he has cured many malignant fevers by the use of a decoction of them; which he fays powerfully promotes sweat and urine, without greatly irritating nature, and also raises the spirits, and quenches thirst. present these flowers are chiefly valued for their pleasant flavour, which is entirely loft even by light coction; hence the college directs the fyrup, which is the only officinal preparation of them, to be made by infusion.

CARYOPHYLLAT E radix: Caryophyllatæ vulgaris flore parwo luteo J. B. Avens, or herb benet: the root.

Avens is a rough plant found wild in woods and hedges.

The root has a warm, bitterish, aftringent taste, and a pleasant fmell, fomewhat of the clove kind. especially in the spring, and when produced in dry warm foils. kinfor observes, that such as is the growth of moist foils has nothing of this flavour. This root has been employed as a stomachic, and for strengthening the tone of the vifcera in general: it is still in some esteem in foreign countries, though not taken notice of among us. It yields on distillation an elegant odoriferous essential oil, which concretes into a flaky form.

CASCARILIA, vide ELEU-

CASIA FISTULARIS [L. E.] the fruit of an oriental tree, re-

sembling the walnut.

This fruit is a cylindrical pod, starce an inch in diameter, a foot or more in length: the outlide is a hard brown bark: the infide is divided by thin transverse woody plates, covered with a foft black pulp of a sweetish taste, with some There are degree of acrimony. two forts of this drug in the shops: one brought from the East-Indies, the other from the West: the canes or pods of the latter are generally large, rough, thick-rinded; and the pulp nauseous: those of the former are less, smoother, the pulp blacker, and of a sweeter taste; this fort is preferred to the other. Such pods should be chosen as are weighty, new, and do not make a rattling noise (from the seeds being loofe within them) when fhaken. The pulp should be of a bright, shining black colour, and a sweet taste, not harsh (which happens from the fruit being gathered before it has grown fully ripe) or fourish (which it is apt to turn upon keeping:) it should nei-

ther be too dry, nor too moist, nor at all mouldy, which, from its being kept in damp cellars, or moistened, in order to increase its weight, is very subject to be. Greatest part of the pulp dissolves both in water and in rectified spirit; and may be extracted from the cane by either. The shops employ water, boiling the bruised pod therein, and afterwards evaporating the solution to a due consistence.

The pulp of casia is a gentle laxative medicine, and frequently given, in a dose of some drams, in costive habits. Some direct a dose of two ounces or more as a cathartic, in inflammatory cases, where the more acrid purgatives have no place: but in these large quantities it generally nauseates the stomach, produces flatulencies, and sometimes gripings of the bowels, especially if the casia is not of a very good kind; these effects may be prevented by the addition of aromatics, and exhibiting it in a liquid form. Geoffroy fays, it does excellent fervice in the painful tension of the belly, which fometimes follows the imprudent use of antimonials; and that it may be advantageously acuated with the more acrid purgatives, or antimonial emetics, or employed to abate their force. Vallisnieri relates, that the purgative virtue of this medicine is remarkably promoted by manna; that a mixture of four drams of casia, and two of manna, purges as much as twelve drams of casia, or thirty-two of manna alone. Sennertus observes, that the urine is apt to be turned of a green colour by the use of casia; and sometimes, where a large quantity has been taken, This drug gives name blackish to an officinal electary, and is an ingredient also in another.

CASIA LIGNEA: the bark of an Indian tree called by Breynius arbor canellifera Indica, cortice acrimo visicido seu mucilaginoso, qui

cassia lignea officinarum.

This bark, in appearance and aromatic flavour, approaches to cinnamon; from which it is eafily distinguishable by its remarkable viscidity: chewed, it seems to disfolve in the mouth into a flimy fubstance; boiled in water, it gives out a strong mucilage, the aromatic part exhaling; the water obtained by distillation, unless drawn with great care, has an unpleasant fmell, somewhat of the empyreumatic kind: nevertheless the distilled oil proves nearly of the same quality with that of cinnamon. Cassia possesses the aromatic virtues of cinnamon; but in an inferior degree; and its effects are less durable. Its elutinous quality renders it useful in some cases where simple aromatics are less proper.

CASTOREUM [L. E.] Caftor Castor is the inguinal glands of the beaver; a four footed amphibious animal, frequent in several parts of Europe and America. The best comes from Russia: this is in large round hard cods, which appear, when cut, full of a brittle red liver-coloured substance, interspersed with membranes and fibres exquisitely interwoven. inferior fort is brought from Dantzick, this is generally fat and The worst of all is that of New England, which is in longish thin cods.

Russia castor has a strong not agreeable smell, and an acrid, bitting, bitterish nauseous taste. Water extracts the nauseous part, with little of the finer bitter; rectified spirit extracts this last, without much of the nauseous; proof spirit both: water elevates the whole

of its flavour in distillation; rectified spirit brings over nothing.

Castor is looked upon as one of the capital nervine and antihysteric medicines: some celebrated practi-, tioners have nevertheless doubted its virtues: and Neumann and Stahl declare it infignificant. Experience, however, has shewn, that the virtues of castor are considerable, though they are certainly far less than they have been generally supposed to be. Its officinal preparations are a simple water [L.] a spirituous tincture [L, E.] and a tincture in the volatile oily spirit [E.] It is an ingredient in fundry other compositions. as the compound elixir and powder of myrrh [L.] the feric pills, gumpills, and powder for promoting delivery [E.]

CASUMUNAR [L. E.]

This is a tuberous root, an inch or more in thickness, marked on the surface with circles or joints like galangal, of a brownish or ash colour on the outside, and a dusky yellowish within: it is brought from the East-Indies, cut in transverse slices: what kind of plant it produces, is not known.

Casumunar has a warm bitterish taste, and an aromatic smell, somewhat resembling that of ginger. It has been celebrated in hysteric cases, epilepsies, passes, loss of memory, and other disorders: the present practice sometimes employs it as a stomachic and carminative; but it is not so much used or known

as it deserves to be.

CAUDA EQUINA, seu Equisetum: Equisetum palustre longioribus setis C B Horsetail; the herb [E.]

This plant is common in watery places. It is faid to be a very strong astringent: it has indeed a manifest astringency, but in a very low degree.

CENTAŬ-

CENTAURII MAJORIS seu Rhupontici vulgaris radix: Gentaurii majoris folio in lacinias plures diviso C. B. Greater centaury; the root.

The greater centaury is a large plant, cultivated in gardens. The root has a rough fomewhat acrid taste, and abounds with a red viscid juice; its rough taste has gained it some esteem as an astringent; its acrimony as an aperient; and its glutinous quality as a vulnerary; the present practice takes little notice of it in any intention.

CENTAURII MINORIS fummitates: Centaurii minoris flore purpureo J. B. Lesser centaury; the tops [L. E.]

This grows wild in many parts of England, in dry pasture grounds and amongst corn. The tops are an useful aperient bitter; the Edinburgh pharmacopæia directs an extract to be prepared from them, and employs them as an ingredient in the bitter insusion and stomachic tincture.

CENTINODIUM: Polygonum latifolium C. B. Knotgrass; the herb.
This is said to be vulnerary and aftringent, but on no very good foundation.

CEPA [L. E.]: Cepa vulgaris C. B. Onions.

Onions differ from other bulbous rooted plants, in having single roots or such as cannot be parted so as to increase the plant. These roots are considered rather as articles of food than of medicine: they are supposed to afford little or no nourishment, and when eaten liberally produces flatulencies, occasion thirst, headachs and turbulent dreams: in cold phlegmatic habits, where viscid mucus abounds, they doubtless have their use; as by their simulating qua-

lity they tend to excite appetite, attenuate thick juices, and promote their expulsion; by some they are strongly recommended in suppressions of urine and in dropsies. The chief medicinal use of onions in the present practice is in external applications, as a cataplasm for suppurating tumours, &c. They are an ingredient in the cutaplasma suppurans [E.]

CERA FLAVA [L. E.] Yellow bees wax.

This is a folid concrete obtained from the honeycombs after the honey is got out, by heating and pressing them betwixt iron plates. The best fort is of a lively yellow colour, and an agreeable smell, somewhat like that of honey; when new it is toughish, yet easy to break; by age becomes harder and more brittle, lose, its fine colour, and in great measure its smell.

CERA ALBA [L. E.]—White wax is prepared from this yellow, by reducing it into thin flakes, and exposing it for a length of time to the air; when sufficiently bleached, it is melted, and cast into cakes. The best fort is of a clear and almost transparent whiteness, and of a light agreeable smell like that of the yellow wax, but much weaker.

The chief medical use of wax is in cerates, plasters, unguents, &c. as an emollient for promoting suppuration, &c. It readily unites with oils and animal fats, but not with watery or spirituous liquors. It is given also internally in diarrheas, dysenteries, &c. either mixed with oily substances, as in the balsamum Locatelli [L. E.] or divided by earthy powders, as in the pulvis restaceus ceratus. [E.]

CERACA: Fruitus Ceraci majoris

of Sylvessis, fruitu Subdulci, nigro
colore

colore inficiente C. B. et Cerasi sativæ, frustu rotundo rubro et acido Tourn et Cerasa acidissima sanguineo succo C. B. The sweet cherry with a black juice: the pleafantly sourish cherry, with a colourless juice; and the very sour cherry, with a blood red juice; commonly called black, red, and morello cherries.

These fruits, especially the acid sorts, are very useful and agreeable coolers and quenchers of thirst; and are sometimes directed in this intention, in hot bilious, or febrile distempers. Boerhaave was extremely fond of these and the other fruits called boræi, as aperients in some chronic cases; and declares himself persuaded, that there is no kind of obstruction of the viscera capable of being removed by medicine, which will not yield to the continued use of these.

CERUSSA [L. E.] Cerusse, or white lead

I his is prepared by exposing lead to the steam of vegetable acids till it is corroded into a white powdery substance. It is sometimes adulterated with a mixture of common whiting; this, if in any considerable quantity, may be easily discovered by the specific lightness of the compound: the fort called stake lead is not subject to abuse. See the article Plumbum; and Cerussain the third part.

CETERACH: Ceterach officinarum C. B. Spleenwort, or miltwast.

This is a small bushy plant growing upon rocks and old walls. It has an herbaceous, somewhat mucilaginous, roughish taste: it is recommended as a pectoral and for promoting urine in nephritic cases. The virtue which it has been most celebrated for, is that which it has the least title to, diminishing the spleen.

CHÆREFOLII folia: Chærophylli fativi C. B. Chervil; the leaves [E.]

This is a low annual plant, somewhat like parfley, commonly cultivated in gardens for culinary purpoles. This plant is grateful both to the palate and flomach gently aperient, and diuretic. Geoffroy affores us, that he has found it from experience to be of excellent service in dropsies: that in this disorder, it promotes the discharge of urine when suppressed, renders it clear when feculent and turbid, and when high and fiery of a paler colour: that it acts mildly without irritation, and tends rather to allay than excite inflammation. He goes fo far as to fay, that dropsies which do not yield to this medicine, are scarce capable of being cured by any other. He directs the juice to be given in the dose of three or four ounces every fourth hour, and continued for some time, either alone, or in conjunction with nitre and fyrup of the five opening roots.

CHALYBS, vide FERRUM.

CHAMÆCYPARISSUS, vide ABROTANUM FOBMINA.

CHAMÆDRYOS, seu Trissaninis folia et summitates cum semine: Chamædryos minoris repentis C. B. Germander; the leaves [E.] and tops with the seed [L.]

This is a low shrubby plant, cultivated in gardens. The leaves, tops, and seeds have a bitter taste, with some degree of astringency and aromatic slavour. They are recommended as sudorific, diuretic, andemmenagogue, and for strengthening the stomach and viscera in general. With some they have been in great esteem in intermittent servers; as also in scrophulous and other chronic disorders.

CHAMÆ-

CHAMÆLEON ALBUS, vide CARLINA.

CHAMÆMELI folia, flores: Chamæmeli mbilis feu leucanthemi odoratioris C. B. Single flowered chamomile (the trailing fort with larger leaves and flowers, and the disk of the flower not very convex) the leaves and flowers [L. E.]

These have a strong not ungrateful, aromatic smell, and a very bitter nauseous taste. They are accounted carminative, aperient, emollient, and in some measure anodyne: and fland recommended in flatulent colics, for promoting the uterine purgations, in spasmodic pains, and the pains of childbed women: fometimes they have been employed in intermittent fevers, and the nephritis flowers are frequently also used externally in discutient and antisceptic fomentations, and in emollient glysters: they enter the fotus communis, decoctum commune pro clystere and oleum viride of our dispensatory: an essential oil [L] simple water and extract [ E. ] are likewise prepared from them in the shops.

CHAMÆMELUM flore multiplici C. B. Double-flowered chamomile; the flowers.

These dister from the foregoing in having several rows of the white petala set thick together about the middle disk, which is much smaller. In this disk the medicinal qualities of the slower chiefly reside; and hence the double or small disked fort is inserior in essicacy to the single.

CHAM ÆPITYOS sive Ivæ arthriticæ folia: Chamæpityos luteæ vulgaris sive folio trisido C. B. Ground pine; the leaves.

This is a low hairy plant clammy to the touch, of a strong aromatic refinous smell, and a bitter roughish taste. It is recommended as an aperient and vulnerary, as also in gouty and rheumatic pains.

CHEIRI feu Leucoii lutei flores: Leucoii lutei vulgaris C. B. Wall-flower.

This grows upon old walls and among rubbish, in several parts of England. The slowers have a pleasant smell, and a subacrid, bitterish, not agreeable taste: they are said to be cordial, anodyne, aperient, and emmenagogue, but are wholly neglected by practice.

CHELIDONII MAJORIS folia, radix: Chelodonii majoris vulgaris C. B. Celandine; the leaves and root [E.]

This plant grows upon old walls, among rubbish, and in waste shady places. The herb is of a blueish green colour; the root of a deep red; both contain a gold coloured juice: their smell is disagreeable; the taste somewhat bitterish, very acrid, biting and burning the mouth; the root is the most acrid. Juice of celandine has long been celebrated in disorders of the eyes; but it is greatly too sharp, unless plentifully diluted, to be applied with fafety to that tender organ. It has been fometimes used, and it is said with good fuccess, for extirpating warts, cleansing old ulcers, and in cataplasms for the herpes miliaris. This acrimonious plant is rarely given internally: the virtues attributed to it are those of a stimulating aperient, divretic, and sudorific: it is particularly recommended in the flow kind of jaundice, where there are no symptoms of inflammation. and in dropfies: fome suppose the root to have been Helmont's specific in the hydrops ascites. Half a dram or a dram of the dry root is directed for a dose; or an infusion

in wine of an ounce of the fresh root. The root and leaves are ingredients in the icterial decoction of the Edinburgh pharmacopæa.

CHELIDONII MINORIS folia, radix: Chelidoniæ rotundifoliæ minoris C. B. Pilewort; the leaves

and root [E.]

This is a very small plant, found in moist meadows and by hedge-sides: the roots consist of slender sibres, with some little tubercles among them, which are supposed to resemble the hæmorrhoids; from whence it has been concluded, that this root must needs be of wonderful efficacy for the cure of that diftemper: to the taste, it is little other than mucilaginous.

## CHERMES, vide KERMES.

CHINÆ radix, China root [E.] There are two forts of this root in the shops, one brought from the East Indies, the other from the West. They are both longish, full of joints, of a pale reddish colour: of no smell, and very little taste: the oriental, which is the most esteemed, is confiderably harder and paler coloured than the other. Such should be chosen as is fresh, close, heavy, and upon being chewed appears full of a fat unctuous juice. China root was either unknown or difregarded by the ancient physicians. It was first introduced into Europe about the year 1535, with the character of a specific against venereal and cutaneous disorders, and as such was made use of for some time, but at length gave place to medicine of a more powerful kind. is generally supposed to promote insensible perspiration and the urinary discharge: and by its uncluous quality to obtund acrimonious juices.

CHINA CHINÆ, vide PERU-VIANUS CORTEX, CICERIS RUBRI femen: Ciceris floribus et semenibus ex purpura rubescentibus C. B. Red chiches, or chich peas.

This is a fort of pulse cultivated in the warmer climates, where our finer peas do not thrive so well. They are a strong flatulent sood, hard of digestion. Lithontriptic and diuretic virtues are attributed to them on no good soundation.

CICHOREI folia, radix: Cichorei fylwestris siwe officinarum C. B. Wild succory; the root and leaves [E.]

The root has a moderately bitter tafte, with some degree of roughness: the leaves are somewhat less bitter: the roots, stalks, and leaves yield, on being wounded, a milky faponaceous juice. By culture this plant loses its green colour and its bitterness, and in this state is employed in falads: the darker coloured and more deeply jagged the leaves, the bitterer is their tafte. Wild succery is an useful detergent, aperient, and attenuating medicine; acting without much irritation, tending rather to cool than to heat the body, and at the same time corroborating the tone of the intestines. The juice taken in large quantities, fo as to keep up a gentle diarrhoea, and continued for some weeks, has been found to produce excellent effects in scorbutic and other chronical disorders.

CICUTÆ folia: Cicutæ majoris C. B. Hemlock; the leaves [E.] This is a large umbelliferous plant, common about the fides of fields, under hedges, and in moist shady places: the leaves are winged, divided into a great number of small fern-like sections of a dark or black-ish green colour, and appearing as it were rough: the stalk is hollow (as is likewise great part of the root after-

after the stalk has arisen) and spotted with several blackish, red, or purple spots. Hemlock is sometimes applied externally as a discutient, and in this intention is an ingredient in one of the plasters of the Edinburgh pharmacopœia. With regard to its virtue when taken internally, it has been generally accounted poisonous, which it doubtless is, in a high degree, when ased in any considerable quantity. But Dr. Storck has lately found, that in certain small doses it may be taken with great safety, and that, without at all disordering the constitution, or even producing any fensible operation, it sometimes proves a powerful resolvent in many obstinate disorders. See Extractum cicutæ in the third part of this work.

CINERES RUSSICI. Ruffia

potash [L.]

Potash is an impure alkaline salt, produced from vegetable matters by burning. The strongest is brought from Russia, in dark coloured very hard masses, which do not soon deliquiate in the air, like the purer alkaline salts. This fort is said to be prepared by burning wood with a close imothering heat, and making the ashes, with a ley drawn from the coarfer part of them, into a paste, which is afterwards stratified with fome of the more inflammable kinds of wood, and burnt a fecond time: by this means the falt melts, and concretes with the earthy matter of the ashes into hard cakes: but it appears from experiment, that this kind of potash contains, besides the vegetable earth, a large proportion of quicklime. A purer and whiter falt is brought to us from Germany, under the name of pearl ashes: this is extracted from wood ashes by means of water, and afterwards reduced into a dry form

by evaporation. These salts are liable to great abuses from fundry admixtures, and therefore should never be employed for medicinal purposes, without due purification: this may be effected by folution in cold water, filtration, and exficcation. See Part III. chap. viii.

CIMOLIA ALBA, seu Argillæ Tobacco-pipe clay [E.]

This is a pure white clay, nearly of the same general nature with the boles already spoken of, but more viscous when moistened with water, and hence probably more effectual for obtunding and incrasfating acrimonious thin humours. It is scarcely ever used for any medicinal purpose.

CIMOLIA PURPURASCENS. Fullers earth [E.]

This earth is more viscous than the boles, and less so than the tobaccopipe clay. It is wholly neglected in practice, though retained in the Edinburgh pharmacopæia.

CINABARIS NATIVA. Native

cinnabar [E.]

This is a ponderous mineral of a red colour, found in Spain, Hungary, and feveral other parts of the The finest fort is in pretty world. large masses, both externally and internally of an elegant deep red colour, which greatly improves upon grinding the mass into fine powder; this is imported by the Dutch from the East-Indies. There is another fort of a good colour, in roundish drops, smooth without, and striated within.

This mineral appears from chemical experiments to be composed of mercury and fulphur, in fuch a manner, that the quantity of the former is commonly above fix times greater than that of the latter: the iner the colour of the cinnabar, the

more mercury it is found to hold. Native cinnabar has been by many preferred as a medicine to that made by art: but there does not appear to be any just foundation for this preference. The native has sometimes been observed to occasion nausex, vomiting, and anxiety: these probably proceeded from an admixture of some arsenical particles which it could not be freed from by repeated ablution. When pure, it has no quality or medical virtue diftinct from those of the artificial cinnabar, like which it is not disfoluble in the animal fluids, and is commonly found of little activity. See Part III. chap. ix. sect. 7.

CINNAMOMUM; Cinnamoanum five canella zeylanica C. B.

Cinnamon [E.]

This is a light, thin bark, of a reddish colour, rolled up in long quills or canes; of a fragrant, delightful smell, and an aromatic, fweet, pungent taste, with some degree of aftringency. It is generally mixed with the cassa bark: this last is easily distinguishable by its breaking over smooth, whilst cinnamon sp!inters; and by its slimy mucilaginous taile, without any thing of the roughness of the true cinnamon. Cinnamon is a very elegant and useful aromatic, more grateful both to the palate and stomach, than most other substances of this class: by its astringent quality it likewise corroborates the viscera, and proves of great service in several kinds of alvine fluxes, and immoderate discharges from the uterus. An essential oil, a simple and spirituous distilled water, and a tincture of it, are kept in the Thops: it is likewise employed as a foicy ingredient in a great number of compositions.

CITREORUM cortex et succus:

Frudus mali medica C. B. Citrons; the yellow rind and juice [E.]

The citron is an evergreen tree or shrub, of the same genus with the orange and lemon: it was sirst brought from Assyria and Media (whence the fruit is called mala Assyria, mala Medica) into Greece, and thence into the southern parts of Europe, where it is now cultivated. Citrons are rarely made use of among us; they are of the same quality with lemons, except that their juice is somewhat less acid.

CITRULLI semen: Anguriæ citrulli didæ C. B. Citruls; the

feed [E]

This plant is rarely met with among us unless in botanic gardens. The seeds are in the number of the sour greater cold seeds, and agree in quality with the others of that class.

CNICUS, vide CARTHAMUS.

COCCINELLA, seu Cochinilla.

Cochineal [L. E.]

This is a small irregular roundish body of a dark red colour on the outfide, and a deep bright red within: it is brought from Mexico and New Spain. This fubstance has long been supposed to be the feed of a plant: it appears from chemical experiments to be an animal, and from the accounts of the more celebrated naturalists, an infect, which breeds on the American prickly pear-tree, and adheres thereto without changing its place. Cochineal has been strongly recommended as a fudorific, cardiac, and alexipharmac: but practitioners have never observed any considerable effects from it. its greatest confumption is among the scarlet dyers; and in medicine its principal use is a colouring drug: both watery and spirituous liquors extract

extract its colour. In the London pharmacopæia three tinctures, in the Edinburgh eight, with two decoctions, an infusion, and a confection, receive from this drug a fine ted colour.

· COCHLEÆ TERRESTRES, vide Limaces terrestres.

COCHLEARIÆ HORTEN-SIS folia: Cochleariæ folio subrosundo C. B. Gardea scurvy-grass; the leaves [L. E]

COCHLEARIÆ MARINÆ folia: Cochleariæ foliosin unto C. B. Sea scurvy-gras; the leaves.

These plants have little other difference, as to their external appearance, than that expressed in their titles: in taste and medical virtue, the first is considerably stronges; and hence is alone retained, both by the London and Edinburgh colleges.

Scurvy-grass is a pungent stimulating medicine; capable of diffolving viscid juices, opening obstructions of the viscera and the more distant glands, and promoting the fluid fecretions: it is particularly celebrated in scurvies, and is the principal herb employed in these kinds of dilorders in the northern countries. Its officinal preparations are a conferve [L. E.] and fpirit  $[E_{\bullet}]$ : it is an ingredient also in the scorbutic juices and compound horseradish water [L. E.] and its spirit is used for drawing a tincture from gum lac [E.]

COFFEA [E] Coffee: the fruit of an oriental thrub called by Justien jasminum Arabicum lauri solio, cujus semen apud nos casse dicitur.

This fruit is employed rather as food than as medicine. The medical effects expected from it, are to affift digeftion, promote the natural fecretions, and prevent or remove a disposition to sleepiness.

COLOCYNTHIDIS medulla. Coloquintida, or bitter apple; the medullary part of the dried fruit [L. E.]

This is the produce of a plant of the gourd kind, growing in Turkey. The fruit is about the fize of an orange; its medullary part, freed from the rind and seeds, is alone made use of in medicine: this is very light, white, spongy, composed of membranous leaves; of an extremely bitter, nauseous, acrimonious taile. Colocynth is one of the most powerful and most violent cathartics. Many eminent phyficians condemn it as dangerous, and even deleterious: others recommend it not only as an efficacious purgative, but likewise as an alterative in obstinate chronical disorders. Thus much is certain, that colocynth in the dole of a few grains, acts with great vehemence, disorders the body, and sometimes occasions a discharge of bloods Many attempts have been made to correct its virulence by the addition of acids, astringents, and the like; these may lessen the force of the colocynth, but no otherwise than might be equally done by a reduction of the dose. The best method of abating its virulence, without diminishing its purgative virtue, seems to be by triturating it with gummy farinaceous substances, or the oily seeds, which without making any alteration in the colocynth itself, prevent its refinous particles from cohering, and sticking upon the membranes of the intestines, so as to irritate, inflame, or corrode them. It is an ingredient in some of the purgative pills, and the cathartic extract of the shops.

CONSO-

CONSOLIDÆ MAJORIS, feu Symphyti majoris radiz: Symphyti confolidæ majoris C. B. Comfry; the root [E.]

This is a rough hairy plant, growing wild by river-fides and in watery places. The roots are very large, black on the outfide, white within, full of a viscid glutinous juice, of no particular tafte. They agree in quality with the roots of althæa; with this difference, that the macilage of confolida is somewhat stronger bodied. Many ridiculous histories of the confolidating wirtues of this plantare related by authors. It is an ingredient in the compound white decoction of the Edinburgh pharmacopœia.

CONSOLIDA MEDIA, vide Bugula.

CONSOLIDA MINIMA, vide Bellis minor.

CONTRAYERVA [L. E.]

This is a knotty root, an inch or two in length, about half an inch thick, of a reddish brown colour externally, and pale within: long, tough, flender fibres shoot out from all sides of it; these are generally loaded with finall round knots. This root is of a peculiar kind of aromatic fmell, and a somewhat oftringent, warm, bitterish taffe, with a light and sweetish kind of acrimony when long chewed: the Sbres have little taste or smell; the tuberous part therefore should be alone chosen. Contrayerva is one of the mildest of those substances called alexipharmacs: it is indifputably a good and vieful diaphoretic, and may be fafely given in much larger doses than the common practice is accustomed to exhibit it in. Its virtues are extracted both by water and rectified spirit. and do not srife in eva with either: the spirituous and extract taste stronger root than the aqueous of gives name to an officinal and is an ingredient in the burgh therizes.

COPAL [E.] a refin of from feveral forts of lar growing in New Spain. In is brought to us in i lumps, fome transparent, lowish or brown colour, of mitransparent and whitish never come into use as a nand is rarely met with in the state of t

CORALLINA: Muscu mus sive corallina officinari Coralline, or sea moss [E

This is a branched fle stance of a white colour, on rocks, and sometime shells of sistes. It is co as a vermisuge, on what son I know not: to the traintirely insipid.

CORALLIUM RU Red coral [L. E.]

This is also a marine jon, of the same nature foregoing. It cannot rebe looked upon in any ot than as a mere absorbeat it enters the officinal crowder, and is sometimes tice directed by itself.

CORIANDRI semen: majoris C. B. Coriander [L. E.]

Coriander is an enb plant, differing from all t of that class in producing seeds. These, when fres strong disagreeable smel improves by drying, and sufficiently grateful; the commended as carminativ machic. They are an ingredient in the officinal compound limewater and electary of bayberries [L.]

CORNU CERVI[L E.] The

stag or harts horns.

Many extraordinary virtues have been attributed to these horns, and to all the parts of the animal in general; but experience gives no countenance to them; nor do they seem to have any other foundation than the great timidity of the hart, the annual renewal of his horns, and an opinion of his extraordinary longwity; from these circumstances it was inferred, that all the parts of him must be proper for intimidating the enraged Archeus, renewing health and strength, and prolonging life.

The horns boiled in water give out an emollient nutritious gelly [E.] Burnt to whiteness, they yield an absorbent earth, purer from gelatinous matter than the natural testaceous absorbents, but which appears to be weaker in its absorbent power. This earth is employed in the officinal white decoction

 $\{L. E.\}$ 

CORNI fructus: Corni bortenfit maris C. B. The cornel tree; its fruit.

This fruit is moderately cooling and astringent, but not regarded as an article of the Materia Medica, and hence rejected both from the London and Edinburgh pharmacopæias.

COSTUS [L. E.] Coftus: a root brought from the East-Indies.

Authors mention two forts of costus, sweet and bitter: in the shops we seldom meet with any more than one, the costus dulcis officinarum C. B. The root is about the size of the singer; and consists of a yellowish woody part inclosed

within a whitish bark: the former is very tough, of no fmell, and very little tafte; the cortical part brittle, of a warm, bitterish, aromatic tafte, and an agreeable smell, somewhat refembling that of violets or Florentine orris. Costus is said to attenuate viscid humours, to promote expectoration, perspiration, and urine. At present it is rarely met with in prescription, and not often in the shops; in mithridate, theriaca, and the confectio pauling, the only officinal compositions it is directed in, zedoary supplies its place.

COSTUS HORTORUM, vide BALSAMITA MAS.

COTULÆ FOETIDÆ folia: Chamæneli fætidi C. B. Mayweed, or wild chamomile; the leaves [E.]

This plant is common among corn, and in waste places. In appearance it resembles some of the garden chamomiles, but is easily distinguishable from them by its strong settle scent. It is rarely or never used in the present practice.

CRASSULÆ five Telephii folia: Telephii vulgaris C. B. Or-

pine; the leaves [E.]

This is a very thick leaved juicy plant, not unlike the houseleeks. It has a mucilaginous roughish taste, and hence is recommended, as emollient and astringent, but has never been much regarded in practice.

CREPITUS LUPI, vide Ly-

CRETA [L. E.] White chalk. This is a pure alkaline earth, totally foluble in vinegar, and the lighter acids, so as to destroy every sensible mark of their acidity. This earth is one of the most useful of the absorbents, and is to be looked

upon

upon fimply as such: the aftringent virtues which some attribute to it, have no foundation, unless in so far as the earth is satiated with acid, with which it composes a faline concrete manifestly subastringent. It gives name to an official julep [L.] and decoction [E] and is an ingredient in the cardialgic troches. It is employed also for extricating the volatile salt of sal ammoniac [L.]

CRITHMI folia: Crithmi frue famiculi maritimi minoris C. B. Samphire; the leaves [L.]

This plant grows wild on rocks, and in maritime places: the leaves are somewhat like those of sennel, but the segments much thicker and shorter: their smell resembles that of smallage; the taste is warm, bitterish, not agreeable. They are said to be stomachic, aperient, and diuretic.

CROCUS: Crocus fativus C. B. Saffron; the chives or fleshy capillaments growing at the end of the pistil of the flower, carefully picked and pressed together into cakes [L. E.]

There are three forts of faffron met with in the shops, two of which are brought from abroad, the other is the produce of our own country: this last is greatly superior to the two former, from which it may be distinguished by its blades being broader. When in perfection, it is of a fiery orange red colour, and yields a deep yellow tincture: it should be chosen fresh, not above a year old, in close cakes, neither dry, nor yet very moif, tough and firm in tearing, of the same colour within as without, and of a ftrong, acrid, diffutive smell.

Saffron is a very elegant and useful aromatic: besides the virtues which it has in common with all the bodies of that class, it remarkably exhilarates, raises the spirits, and is deservedly accounted one of the highest cordials: taken in large doses, it is said to occasion immoderate mirth, involuntary laughter, and the ill effects which follow from the abuse of spirituous li-This medicine is particuquors. larly ferviceable in hysteric depresfions proceeding from a cold cause or obstructions of the uterine secretions, where other aromatics, even those of the more generous kind, have little effect. Saffron imparts the whole of its virtue and colour to rectified spirit, proof spirit, wine, vinegar, and water: a tincture drawn with vinegar, loses greatly of its colour in keeping: the watery and vinous tinctures are apt to grow four, and then lose their colour also: that made in pure spirit keeps in perfection for many years. Its officinal preparations are, a spirituous tincture [E.] a vinous tincture, and a fyrup [L.] It is an ingredient in the cordial confection [L.] the sudorific tincture, the sectoral and paregoric elixirs, the powder for promoting delivery [E.] and feveral of the aloetic compositi-

CUBEBÆ [L. E.] Cubebs.

Cubebs are a fruit brought from the East Indies. This fruit has a great resemblance to pepper. The principal difference distinguishable by the eye, is, that each cubeb is furnished with a long slender stalk (whence they are called by some piper caudatum.) In aromatic warmth and pungency, cubebs are far inserior to pepper. They are an ingredient in mithridate and theriaca, [L.] and in the compound spirit of lavender [E.]

CUCUMERIS HORTENSIS K 2 femen. Semen. Garden cucumbers; the

feeds [E.]

These are in the number of the four greater cold feeds; they are less apt to grow rancid in keeping than the others of that class.

CUCUMERIS **AGRESTIS** fructus: Cucumeris sylvestris afinini ditti C. B. Wild cucumber; the

fruit [ L. E. ]

This plant, found wild in foreign countries, is with us cultivated in gardens. Its principal botanic difference from the former, is the smallness of its fruit, which is no bigger than a Spanish olive: when ripe, it burits on a light touch, and sheds its seeds with violence, and hence was named by the Greeks elaterium. This name was applied likewise to the inspissated juice of the fruit, the only preparation of the plant made use of in medicine. Blaterium is a strong cathartic, and very often operates also upwards. Two or three grains are accounted in most cases a sufficient dose. Simon Paulli relates fome instances of the good effects of this purgative in dropsies: but cautious practitioners not to have recourse to it till after milder medicines have proved ineffectual; to which caution we heartily subscribe. cines indeed in general, which act with violence in a small dose, require the utmost skill to manage them with any tolerable degree of fafety: to which may be added, that the various manners of making these kinds of preparations, as practifed by different hands, must needs vary their power.

CUCURBITÆ semen: Cucurbit a oblonga, flore albo, folio molli C. B. The gourd; its feeds [E.]

These are in the number of the . four greater cold feeds. They unite with water by trituration into an

emulfion, and yield to the press a foft infipid oil, and possess the general virtues of uncluous subftances.

CUMINUM. vide Cyminum.

CUPRESSI frudus. The cy-

press tree; its fruit. I his is a tall tree growing wild

in the warmer climates. fruit is a strong astringent; and in fome places frequently used as such: among us it is very rarely employed, and not often met with in the shops.

CUPRUM [L. E.] Copper.

The preparations of copper are violently enletic, and therefore very rarely exhibited internally. Some have ventured upon a folution of a grain or two of the metal in vegetable acids, and observe, that it acts almost as soon as received into the stomach, so as to be of good use for occasioning poisonous substances that have been swallowed to be immediately thrown up again. Boerhaave recommends a faturated folution of this metal in volatile alkaline spirits, as a medicine of great service in disorders proceeding from an acid, weak, cold, phlegmatic cause: if three drops of this tincture be taken every morning with a glass of mead, and the dole doubled every day to twenty-four drops, it proves (he fays) aperient, attenuating, warming, and diuretic: he affures us, that by this means he cared a confirmed afcites, and that the urine run out as from an open pipe; but at the fame time acknowledges, that upon trying the fame medicine on others, it failed him. He likewise recommends other preparations of copper, as of wonderful efficacy in certain kinds of ill habits, weaknefs of the stomach, &c. but we CARBOT

cannot think the internal use of this metal commendable, or even safe. physicians in general seem to be agreed, that it has really a virulent quality; and too many examples are met with of satal consequences ensuing upon eating food that had been dressed in copper vessels not well cleaned from the rust which they had contracted by

lying in the air.

Great care ought to be had that acid liquors, or even water defigned for internal use, be not suffered to stand long in vessels made of copper; otherwise they will dissolve so much of the metal as will give them disagreeable qualities. Hence in the distillation of simple waters with copper stills, the last runnings, which are manifestly acid, have frequently proved eme-It is romarkable, that whilst weak acid liquors are kept boiling in copper vessels, they do not seem to dissolve any of the metal; but if suffered to remain in them for the same length of time without boiling, they become notably impregnated with the copper. Hence the confectioners, by skilful management, prepare the most acid fyrups in copper vessels without giving them any ill tafte from the metal.

CURCUMA [E.] Turmeric.

Turmeric is a root brought from the East-Indies. It is internally of a deep lively yellow, or fasfron colour, which it readily imparts to watery liquors. It has an agreeable, weak smell, and a bitterish, somewhat warm taste. Turmeric is esteemed aperient and emmeagogue, and of singular essicacy in the jaundice. It is an ingredient in the icteric decoction of the Edinburgh pharmacopeeia. It tinges the urine of a fasfron colour.

CUSCUTA. Dodder.

This is of the class of plants called parasitical, or which grow out from the body of others: it has no leaves, confilling only of a number of juice alaments matted together. There are two forts of it, cuscuta major C. B. which grows commonly in heaths on furzes, nettles, &c. and likewise in fields of flax, and other manured plants and the cufcuta minor, or epithymum of the same author, so called from its being found only upon thyme. This last is preferred for medicinal use, and is usually brought from Leghorn and Turkey, with tops and stalks of thyme amongst it. Epithymum has a pretty ftrong fmell, and roughish somewhat pungent subtile taste. Its virtues remain as yet to be determined: the ancients ranked it among cathartics; but those who have given it in that intention have been generally disappointed.

CYANI flores: Cyani segetum C. B. Bluebottle; the flowers.

This is a common weed among corn. The flowers are of an elegant blue colour, which, if carefully and haftily dried, they retain for a conderable time. As to their virtues, the present practice expects not any from them; notwithstanding they have been formerly celebrated against the bites of poisonous animals, contagious diseases, palpitations of the heart, and many other distempers.

CYCLAMEN, vide ARTHA-

CYDONIA MALA, corumque femina: Fructus Cotoneæ mali J. B. The quince-tree; the fruit and its feeds [L. E.]

Quinces have a very auftere acid tafte: taken in small quantity, they are supposed to restrain vomiting,

K 3 and

and alvine fluxes; and more libe- fylvefiris vulgaris flore odorate incarrally to loofen the belly. The feeds abound with a mucilaginous substance, of no particular taste, which they readily impart to watery liquors: an ounce will render three pints of water thick and ropy like the white of an egg. A fyrup [L.] and jelly [E.] of the fruit and mucilage of the feeds [L.] are kept in the shops.

CYMINI semen: Cymini semine longiore C. B. Faniculi orientalis cumini dicti Tourn. Cummin; the

feeds [L. E.]

This is an umbelliferous plant, in appearance resembling fennel, but much smaller; the seeds are brought chiefly from Sicily and Malta. Cummin seeds have a bitterish warm taste, accompanied with an aromatic flavour, not of the most agreeable kind. are accounted good carminatives, but not very often made use of. An effential oil of them is kept in the fhops, and they give name to a plaster and cataplasm [L.]

CYNOGLOSSI radix: Cynogloss majoris vulgaris C. B. Hounds-

tongue; the root.

The leaves of this plant are in shape thought to resemble a tongue, whence its name; they are clothed with a whitish down: it grows wild in shady lanes. The roots have a rank disagreeable fmell, and rough bitterish taste, covered with a glutinous sweetish-The virtues of his root are very doubtful: it is generally furposed to be narcotic, and by some to be virulently fo: others declare, that it has no virtue of this kind. and look upon it as a mere glutinous astringent. The present practice takes no notice of it in any intention.

CYNOBASTI frudus: Rose nato C. B. The wild brian, dog rose, or hip tree; its fruit [L. E.]

This bush grows wild in hedges throughout England. The flowers have a pleasant smell; but so weak, that Parkinson and others have named the plant rosa sylvestris inodora: a water distilled from them fmells agreeably. The fruit or hips contain a fourish sweetish pulp, with a rough prickly matter inclosing the feeds, from which the pulp ought to be carefully separated before it is taken internally: the Wirtemberg college observes, that from a neglect of this caution, the pulp of hips fometimes occasions a proritus, and uneasiness about the anus; and I have known the conserve of it to excite violent vomiting. The conferve is the only officinal preparation of this fruit.

CYPERI LONGI radix: Crperi odorati radice lunga, sive cyperi officinarum C. L. Long cyperus; the root.

This is a plant of the graminifolious kind: it is sometimes found wild, in marshy places in England; the roots have been generally brought to us from Italy. This root is long, slender, crooked, and full of knots; outwardly of a dark brown, or blackish colour, inwardly whitish; of an aromatic smell, and an agreeable warm taile: both the taile and finell are improved by moderate exficcation. Cyperus is accounted a good stomachic and carminative, but at present very little regarded.

DACTYLI: Frudus Palme majoris C. B. Dates [E]: a half dried fruit, about the shape of an acorn, but generally larger, confisting of a sweet pulpy part and a hard stone: the best are brought from Tunis. They were formerly used in pectoral decoctions, and supposed, besides their emollient and incrassating virtue, to have a slight astringency.

DAUCI CRETICI femen: Dauci foliis fæneculi tenuissimis C.B. Candy carrot, or carrot of Crete;

the seeds [L. E.]

This is an umbelliferous plant growing wild in the Levant, and the warmer parts of Europe. The feeds, which are brought from Crete, have a warm biting tafte, and a not difagreeable aromatic fmell. They are carminative, and faid to be diuretic; but at prefent little otherwise used than as ingredients in the mithridate and theriaca.

DAUCI SYLVESTRIS femen: Pastinacæ sykvestris tenuifoliæ Diofcoridis, svel dauci officinarum C. B. Wild carrot; the seed [E.]

This is common in pasture grounds and fallow fields throughout England. The feeds possess the virtues of those of the daucus Creticus, in an inferior degree: and have often supplied their place in the shops; and been themselves supplied by the seeds of the garden carrot: these last are, in warmth and slavour, the weakest of the three: the seeds of the Candy carrot are much the strongest.

DENTIS LEONIS five Taraxaci, radix et folia: Dentis leonis latiore folio, et angustiore folio C. B. Dandelion; the root and herb [E.]

This plant is common in fields, and uncultivated places; it has feveral narrow dentated leaves lying on the ground, with a flender naked stalk sustaining a yellow slower. The root, leaves, and stalk, contain a bitter milky juice: they

promise to be of use as aperient and detergent medicines; and have sometimes been directed in this intention with good success. Boerhaave esteems them capable, if duly continued, of resolving almost all kinds of coagulations, and opening very obstinate obstructions of the viscera.

DIAPENSIA, vide Sanicula.

DICTAMNUS ALBUS, vide FRAXINELLA.

DICTAMNI CRETICI folia: Origani Cretici latifolii tomentofi Tourn. Dittany of Crete [L. E.]

This is a kind of origanum, said to grow plentifully in the island of Candy, in Dalmatia, and in the Morea: it has been found hardy enough to bear the ordinary winters of our own climate. The leaves, which are the only part in use with us, come from Italy. best fort are well covered over with a thick white down, and now and then intermixt with purplish slowers. In smell and taste they fomewhat refemble lemonthyme: but have more of an aromatic flavour, as well as a greater degree of pungency; when fresh they yield a confiderable quantity of an excellent effential oil. are ingredients in the pulvis e myrrha, species e scordio, mithridate, and tberiaca [L.]

DIGITALIS folia: Digitalis purpureæ folio aspero C. B. Fox-

glove; the leaves.

This grows wild in woods, and on uncultivated heaths: the elegant appearance of its purple flowers (which hang in spikes along one side of the stalk) has gained it a place in some of our gardens. The leaves have been strongly recommended, externally against

against scrophulous tumours; and likewise internally, in epileptic disorders: what service they may be capable of doing in these cases, we have no experience. Several examples are mentioned by medical writers of their occasioning violent vomiting, hypercatharsea, and disordering the whole constitution: infomuch that Boerhaave accounts them poisonous. Their taste is bitter and very nauseous.

DORONICI ROMANI radix; Derenici radice scorpii C. B. Roman wolfsbane; the root.

This grows spontaneously on the Alps, and in fundry places of Germany. It has been greatly disputed whether this plant is to be ranked among the poisonous or falutary ones: we shall not here enter into this controversy, obferving only, that all the inten-tions, it has been recommended for, may certainly be answered by other medicines of no less efficacy, and known to be innocent: and that therefore the use of doronicum may be very reasonably laid aside: in this we are warranted by common practice, which has not, for a long time, paid any regard to it.

DORONICI GERMANICI, feu Arnicæ, folia et radix: Doroniei plantaginis folia a'terins C. B. German leopardíbane; the leaves and soot [E.]

This plant is a native of Germany. It has been effected a specific for resolving coagulated blood; but operates so violently, that it is rarely used. The dose is said to be one or two pugils of the leaves, and, in some cases, of the roots.

DRACONTIUM: Dracuncu-

lus polyphillus C. B. Arum polyphillum Revini. Dragons, or the munyleaved arum.

This is cultivated in gardens. It has scarce any other medical difference from the common arum, than being in all its parts somewhat more pungent and acrimonious.

DRAKENA, vide Contra-

DULCAMARÆ, seu Amaradulcis, solani lignosi, herba, radix z Solani scandentis seu dulcamaræ C. B. Bittersweet, or woody nightshade; the herb and roots se.

This plant grows wild in moist hedges, and climbs on the bushes with woody brittle stalks. The taste of the twigs and roots, as the name of the plant expresses, is both bitter and sweet; the bitterness being first perceived, and the sweetness afterwards. They are commended as deobstruents, for resolving coagulated blood, &cc. and are said to occasion generally some considerable evacuation by sweat, urine, or stool, particularly the latter.

EBULI folia, cortex, radix: Sambuci humilis frue ebuli C. B. Dwarf elder, or danewort; the root, bark, and leaves [E.]

This plant grows wild in some counties of England; but about London is rarely met with, unless in gardens: the eye distinguishes little difference betwixt it and the elder tree, except in the size; the elder being a pretty large tree, and the dwarf elder only an herb three or sour seet high. The leaves, roots, and bark of chalus have a nauseous, sharp, bitter taste, and a kind of acrid ungrateful smell it they are all strong cathartics, and as such are recommended in drong for

sies, and other cases where medicines of that kind are indicated. The bark of the root is said to be strongest; the leaves the weakest. But they are all too churlish medicines for general use: they sometimes evacuate violently upwards, almost always nauseate the stomach, and occasion great uneasiness of the bowels. By boiling they become (like the other draftics) milder, and more safe in operation. Fernelius relates, that by long coction they entirely lose their purgative The berries of this plant are likewise purgative, but less virulent than the other parts. A rob prepared from them, may be given, to the quantity of an ounce, as a cathartic; and in imaller ones as an aperient and deobstruent in chronic disorders: in this last intention, it is said by Haller to be frequently used in Switzerland, in the dole of a dram.

ELATINES folia: Linariæ segetum nummulariæ solio non villoso Tourn. Fluellin, or semale speedwell: the leaves.

This is a low creeping plant, growing wild in corn-fields. The leaves have a very bitter, roughish tafte. They were formerly accounted excellent vulneraries, and of great use for cleansing and healing old ulcers and spreading cancerous fores: some have recommended them internally in leprous and scrophulous disorders; as also in hydropic cases. It gives name to one of the officinal honeys [L.]; but the plant itself is never used in the present practice, and this preparation of it is in no great esteem.

ELEMI [L. E.] Gum elemi.
This is a refin brought from the
Spanish West-Indies, and sometimes from the East-Indies, in

long roundish cakes, generally wrapt up in flag leaves. The best fort is softish, somewhat transparent, of a pale whitish yellow colour, inclining a little to greenich, of a strong not unpleasant smell. It almost totally dissolves in pure spirit, and sends over some part of its fragrance along with this menstruum in distillation : distilled with water, it yields a confiderable quantity of a pale-coloured, thin, fragrant essential oil. This resin gives name to one of the officinal unguents, and is at present scarce any otherwise made use of; though it is certainly preferable, for internal purpoles, to some others which are held in greater efteem.

ELEOSELINUM, vide APIUM.

ELEUTHERIÆ feu Cascarillæ; a bark said to be imported into Europe from one of the Bahama islands called Elatheria, in curled pieces, or rolled up into short quills, about an inch in width, pretty much resembling in appearance the Peruvianus cortex, but of a paler brown colour on the inside, less compact, and more friable.

Its take is bitterer, yet less disagreeable and less rough than that of the Peruvian bark; with a considerably greater share of aromatic pungency and heat: the thin outward skin, which is of a whitish colour, has no taste. It is easily stammable, and yields, whilst burning, a very fragrant smell: this peculiar property distinguishes the eleutheria from all other known barks.

Stifferus seems to have been the first that employed the certex eleutheriæ as a medicine, in Europe; he relates (in his Att. laborat, chym. published in the year 1693) that he received this asomatic bark from England;

England; and that some time after it was fold at Brunswick for Peruvian bark; that a tincture of it in alkalized vinous spirits, or dulcified alkaline ones, proved carminative and diuretic, and did confiderable fervice in arthritic. fcorbutic, and calculous cases; and that if taken immediately after meals, it affected the head a little. theria was foon after employed by Apinus in an epidemic fever which raged in some parts of Norway in 1694 and 1695: this disease, which at first had the appearance of an ordinary intermittent, at length was accompanied with petechial spots. The common alexipharmacs and sudorifics were found ineffectual: but the powder or extract of this bark, joined with them, proved successful, even after petechiæ had come forth: dyfenteries succeeding the sever were removed by the same medicine. During the use of the eleutheria, the patient generally sweated plentifully, without loss of strength, or other inconvenience: the belly was likewise kept open; those who did not sweat, had three or four flools a day: where the menfirmal or hæmorrhoidal fluxes were fuppressed at the beginning of the disorder, they generally, upon the use of this medicine; re-appeared. Among the Germans, the eleutheria is at present in very great esteem, and frequently employed against common intermittents, in preference to the Peruvian bark, as being less subject to some inconveniences which the latter, on account of its greater aftringency, is apt to occasion: it is also given, with good success, in flatulent colics, internal hæmorrhages, dyfenteries, the diarrhese of acute fevers, and other like disorders. The gentlemen of the French academy found this bark of excellent

fervice in an epidemic dyfentery in the year 1719; in which ipecacoanha proved ineffectual. Mr. Boulduc observed, that this last left a lowness and weakness of stomach, which continued for a long time, whilst eleutheria soon raised the strength, and promoted appetite. Among us the use of this bark is not yet so general as it seems to deserve: insusions of it are sometimes directed for promoting expectoration.

ENDIVIÆ radix, folia: Intybi sativæ latifoliæ C. B. Endive, the roots and leaves (E.)

Endive is raised in gardens for culinary use. It is a gentle cooler and aperient, nearly of the same quality with the cicborium. The feeds are ranked among the four lesser cold seeds.

ENULÆ CAMPANÆ seu Helenii radix: Asteris omnium maximi Tourn. Elecampane; the root

[L. E.]

This is a very large downy plant, sometimes found wild in moift rich foils. The root, especially when dry, has an agreeable aromatic smell : its taste, on first chewing, is glutinous, and as it were somewhat rancid; in a little time it discovers an aromatic bitterness, which by degrees becomes confiderably acrid and pungent. Elecampane root possesses the general virtues of alexipharmacs, it is principally recommended for promoting expectoration in humoural asthmas and coughs, in which intention it enters the pectoral oxymel of the Edinburgh pharmacopœia: liberally taken, it is faid to excite urine, and loosen the belly. In some parts of Germany, large quantities of this root are candied, and used as a stomachic, for strengthening the tone of

the viscera in general, and for attenuating tenacious juices. Spirituous liquors extract its virtues in greater perfection than watery ones: the former scarce elevate any thing in distillation; with the latter an essential oil arises, which concretes into white flakes: this possesses at first the flavour of the elecampane, but is very apt to lose it in keeping. An extract made with water (a preparation now kept in the shops) possesses the bitterness and pungency of the root, but in a less degree than one made with spirit.

EQUISETUM, vide CAUDA EQUINA.

BRIGERI, seu Senecionis folia: Senecionis minoris vulgaris C. B. Groundsel; the leaves,  $\{E.\}$ 

This is a common weed, which, notwithstanding its being annual, is met with at all times of the year. The juice, or an infusion of it in ale, is generally faid to be a mild and safe emetic; but unless taken in very large quantity, it has no effect this way. The fresh herb, beaten into a very coarse pulp, and applied externally, cold, to the pit of the stomach, is said to have occasioned throng vomiting: but, as Haller juftly suspects, there was probably some fallacy in the observation.

ERUCÆ semen: Erucæ latifoliæ albe, fative Dioscordis C. B. Rocket; the seeds [E.]

This was formerly much cultivated in gardens for medicinal use, and for falads: but is at present less common. In appearance, it resembles mustard, but is easily distinguishable by the smoothness of its leaves, and its disagreeable finell. The feeds have a pungent taste, of the mustard kind, but weaker: they have long been celebrated as aphrodifiacs, and may, probably, have in some cases a title to this virtue, in common with other acrid plants.

ERVUM, vide OROBUS.

ERYNGII radix: Eryngii maritimi C. B. Erynga; or sea-holly;

the loot [L. E.]

This plant grows plentifully on fome of our fandy and gravelly shores; the roots are flender, and very long; of a pleasant sweetish tafte, which, on chewing them for some time, is followed by a light degree of aromatic warmth and acrimony. They are accounted aperient and diuretic, and have also been celebrated as aphrodifiac; their virtues however are too weak to admit them under the head of medicines. The candied root is ordered to be kept in the shops.

ERYSIMI folia: Eryfimi vulgaris C. B. Hedge multard; the

leaves [E.]

This is a low hairy plant, common in waste places, and by way-The leaves are faid to promote expectoration, excite urine, and the other fluid feeretions, attenuate and dissolve viscid juices, &c. This they are supposed to perform by an acrimonious stimulating quality; but the taste discovers in them only an herbaceous foftness void of acrimony: the feeds indeed are confiderably pungent, and the roots in some small degree.

ESULA MAJOR et MINOR, vide TITHYMALUS.

EUPATORII CANNABINI folia: Eupatorii cannabini C. B. Hemp agrimony, water agrimony, or water hemp; the leaves.

This plant is found wild by the fides of rivers and ditches. It has an acrid smell, and a very bitter tafte.

taste, with a considerable share of pungency. The leaves are greatly recommended for strengthening the tone of the viscera, and as an aperient; and said to have excellent effects in the dropsy, jaundice, cachexies, and scorbutic disorders. Boerbaave informs us, that this is the common medicine of the turf-diggers in Holland, against scurvies, foul ulcers, and swellings in the seet, which they are subject to. The root of this plant is said to operate as a strong cathartic.

EUPATORIUM MESUES, vide Ageratum.

EUPATORIUM GRÆCO-RUM, vide AGRIMONIA.

EUPHORBIUM [E.] a gummy refin exuding from a large oriental shrub. It is brought to us immediately from Barbary, in drops of an irregular form; fome of which, upon being broken, are tound to contain little thorns, small twigs, flowers, and other vegetable matters: others are hollow, without any thing in their cavity: the tears in general are of a pale yellow colour externally, somewhat white withinfide: they eafily break betwixt the fingers. Lightly applied to the tongue, they affect it with a very sharp biting taste; and upon being held for some time in the mouth, prove vehemently acrimomious, inflaming, and exulcerating the fauces, &c. Euphorbium is extremelytroublesome to pulverize; the finer part of the powder, which flies off, affecting the head in a vio-lenr manner. The acrimony is fo great as to render it absolutely un-fit for any internal use: several correctors have been contrived to abate its virulence; but the best of them are not to be trusted to; and as there seems to be no real uccafion for it, unless for some external purposes, we think, with Hoffman and others, that it dught to be expunged from the catalogue of internal medicines.

EUPHRASIÆ folia: Euphrafiæ officinarum C. B. Eyebright; the leaves [E.]

This is a very low plant, growing wild in moit fields. It was formerly celebrated as an ophthalmic, both taken internally, and applied externally. Hildanus fays, he has known old men of feventy, who had loft their fight, recover tagain by the use of this herb: later practitioners, however, have not been so happy as to observe any such good effects from it. At present it is totally, and not unjustly, disregarded.

FABÆ flores et semen: Fabæ flore candido lituris migris conspicuo Tourn. Garden beans; the flowers and seed [E.]

Beans are of greater use for culinary than medicinal purposes: they are a strong statulent food, sufficiently nutritious, but not easy of digestion; especially when growing old. A water distilled from the slowers has been celebrated as a cosmetic, and still retains its character among some female artists.

FARFARA, vide Tussilago.

FERRUM et CHALYBS [L.E.] Iron and steel [L. E.]

Steel is accounted less proper for medicinal use than the softer iron, as being more difficultly acted upon by the animal juices and the common menstrea: iron dissolves readily in all acids, and rusts freely in the air, especially if occasionally moistened with water; steel requires a longer time for

its folution, and does not ruft fo

eafily.

The general virtues of these metals, and the several preparations of them, are, to confiringe the fibres, to quicken the circulation, to promote the deficient secretions in the remoter parts, and at the same time repress inordinate discharges into the intestinal tube. After the use of them, if they take effect, the pulse is very senfibly raised; the colour of the face, though before pale, changes to a florid red; the alvine, urinary, and cuticular excretions, are increased. Nidorous ernelations, and the fæces voided of a black colour, are marks of their taking due effect.

An aperient virtue is usually attributed to some of the preparations of iron, and an astringent to others; but in reality, they all produce the effects both of aperients and astringents, and seem to differ only in degree Those distinguished by the name of astringent sometimes occasion a very copious discharge of urine, or a diarrhæa; whilst those called aperient frequently stop these evacuations.

Where either a præternatural difcharge, or suppression of natural fecretions, proceed from a languor and sluggishness of the sluids, and weakness of the solids; this metal, by increasing the motion of the former, and the strength of the latter, will suppress the slux, or remove the suppression: but where the circulation is already too quick, the solids too tense and rigid, where there is any stricture or spasmodic contraction of the vessels; iron, and all the preparations of it, will aggravate equally both distempers.

Though the different preparations of iron act all in the same manner, yet they are not equally proper in all constitutions. Where

acidities abound in the first passages, the crude filings, reduced into a fine powder, prove more serviceable than the most elaborate preparation of them. On the other hand. where there is no acid in the primæ viæ, the metal requires to be previously opened by faline menstrua: hence a folution of iron in acid liquors, has, in many cases, excellent effects, where (as Boerhaave observes) the more indigestible preparations, as the calces made by fire, have scarce any effect at all. If alkalescent juices are lodged in the stomach, this metal, though given in a liquid form, proves at least useless; for here the acid folvent is absolved by the alkaline matters which it meets with in the body, so as to leave the iron reduced to an inactive calx.

Chalybeate medicines are likewise supposed to differ, independent of differences in the constitution, according to the nature of the acid united with the metal: vegetable acids superadd a detergency and aperient virtue; combined with the vitriolic, it acts in the first passages powerfully as an aperient; whilst the nitrous renders it extremely styptic; and the marine still more so. For the different preparations of iron, see the third part of this work.

FILIPENDULÆ radix: Filipendulæ vulgaris, an Moli Plinit C. B. Dropwort; the root

This plant grows wild in fields and chalky grounds: the root confifts of a number of tubercles, fattened together by flender strings; its taste is rough and bitterish, with a slight degree of pungency. These qualities point out its use in a slaccid state of the vessels, and a sluggishness of the juices: the natural evacuations are in some measure restrained or promoted by

is, where the excess or deficiency proceed from this cause. Hence some have recommended it as an astringent in dysenteries, immoderate uterine fluors, &c. others as a diuretic; and others as an aperient and deobstruent in scrophulous habits. At present it is wholly difregarded.

FILICIS MARIS radix: Filicis non ramosæ, dentatæ C. B. Common male fern; the root [E.]

FILICIS FŒMINÆ radix: Filicis ramosæ majoris pinnulis obtufis non dentatis C. B. Female fern,

or brakes; the root [E.]

FILICIS FLORIDÆ, seu Osmunde regalis radix: Felicis ramose non dentase, floride C. B. Ofmund royal, or the flowering

fern; the root [E.]

The roots of these plants (which are the only part directed for medicinal use) have, when first chewed, somewhat of a sweetish glutinous take, which soon becomes bitterish, subastringent, and nauseous. They are said to be aperient and anthelmintic. Simon Paulli tells us, that they have been the grand fecret of some empirics against the broad kind of worms called tenia; and that the dose is one, two, or three drachms of the powder. The third fort is supposed to be the weakest, and the second the strongest; this therefore has been generally made choice of; practice has, however, at length expunged them all, though the college of Edinburgh still retains them in their catalogue of fimples.

FŒNICULI DULCIS semen Fæniculi dulcis C. B. Sweet fen-

nel; the feeds [L. E.]

FŒNICULI VULGARIS folia, radix, semen: Fæniculi vulgaris germanici C. B. Common fennel; the feeds, roots, and leaves

The sweet fennel is smaller in all its parts than the common, except the feeds, which are, confiderably larger. The feeds of the two forts differ likewise in shape and colour: those of the common are roundish, oblong, flattish on one fide, and protuberant on the other, of a dark almost blackish colour: those of the sweet are longer. narrower, not fo flat, generally crooked, and of a whitish or pale vellowish colour. Both forts are cultivated in our gardens: the common is a perennial plant: the fweet perishes after it has given feed; nor do its feeds come to fuch perfection in this climate as those which we receive from Germany.

The feeds of both the fennels have an aromatic smell, and a moderately warm, pungent tafte: those of the fæniculum dulce are in flavour most agreeable, and have also a confiderable degree of sweetishness: hence our college have directed the use of these only. They are ranked among the four greater hot feeds; and not undeservedly looked upon as good stomachics and carminatives. An effential oil [E.] and fimple water [L.] are prepared from them in the shops; they are ingredients also in the compound juniper water, garlic oxymel, mithridate, theriaca, and decoction for glyf-

ters [L.]

The root is far less warm. but has more of a sweetish taste. than the feeds; it is one of the five roots called openers; and has sometimes been directed in apperient apozems. Boerhaave says, that this root agrees in tafte, smell, and medical qualities, with the celebrated ginfeng of the Chinese; from which, however, it

appears

appears to be very considerably different.

The leaves of fennel are weaker than either the roots or feeds, and have very rarely been employed for any medicinal use. A simple water is directed to be prepared from them in the Edinburgh pharmacopæia.

FŒNI GRÆCI femen: Fæni græci fativi C. B. Fænugreck; the feeds [L. E.]

This plant is cultivated chiefly in the fouthern parts of France, Germany, and in Italy; from whence the feeds are brought to us. They are of a yellow colour, a rhomboidal figure; a difagreeable ftrong fmell, and a mucilaginous tafte. Their principal use is in cataplasms, fomentations, and the like, and in emollient glysters. They enter the oleum a mucilaginibus of the shops; to which they communicate a considerable share of their smell.

FOLIUM INDUM, vide Ma-LABATHRUM.

FORMICÆ. Ants; their bo-

dies and eggs.

These insects are at present of no use with us in medicine, though formerly much celebrated for aphrodifiac virtues, and still employed in the aque magnanimitatis, and other like compositions of foreign dispensatories. It is remarkable, that these animals contain a truly acrid juice, which they shed in small drops upon being irritated: by infusing a quantity of live and vigorous ants in water, an acid liquor is obtained nearly as strong as good vinegar. Neumann abserves, that on distilling them either with water or pure spirit, a clear limpid oil arises, which has scarce any taste, or at

least is not hot or pungent like the effential oils of vegetables.

FRAGARIÆ folia, fruetus: Fragariæ ferensis fragra rubra J. B.
The strawberry bush; its leaves and fruit [E.]

The leaves are somewhat styptic, and bitterish; and hence may be of some service in debility and laxity of the vifcera, and immoderate secretions, or a suppresfion of the natural evacuations depending thereon: they are recommended in hæmorrhages and fluxes; and likewise as aperients, in suppressions of urine, obstructions of the viscera, in the jaundice, &c. The fruit is in general very grateful both to the palate and stomach: like other fruits of the dulco-acid kind, they abate heat, quench thirst, loosen the belly, and promote urine; but do not afford much nourishment. Geoffroy observes; that the urine of those who eat liberally of this fruit, becomes impregnated with its fragrant smell.

FRANGULA, vide Alnus nigra.

FRAXINELLÆ seu Dictamni albi, radix: Dictamni vulgo, seve fraxinellæ C. B. White or bastard

dittany; the root [E.]

This plant grows wild in the mountainous parts of France, Italy, and Germany; from whence the cortical part of the root, dried and rolled up into quills, is sometimes brought to us. This is of a white colour, a weak, not very agreeable smell; and a durable bitter, lightly pungent tasse. It is recommended as an alexipharmac; but not regarded by common practice, nor often kept in the shops.

4

FRAXINI cortex et semen: Fraxini excessionis C. B. Fraxini vulgationis J. B. The ash tree;

its bark and seeds [E.]

The bark of this tree is moderately aftringent, and as such has sometimes been made use of: the seeds, which are somewhat acrid, have been employed as aperients. There are so many other medicines more agreeable, and more efficacious for these intentions, that all the parts of the ash tree have long been neglected.

FULIGO lignorum combustorum.

Wood foot [L. E.]

This concrete is of a shining black colour, a disagreeable smell, and an acrid bitter nauseous taste. Its chief use is in hysteric cases, in which it is sometimes given in conjunction with the fetid gums: it gives name to a tincture of this kind in the shops. Its virtues are extracted both by watery and spirituous liquors, each of which, if the foot is of a good kind, dissolve about one fixth. Soot is faid to differ greatly in quality according to the wood it was produced from: the more refinous the wood, the more the foot abounds with oily matter.

FUMARIÆ fola: Fumariæ
officinarum et Diascordis C. B.
Fumitory; the leaves [E.]

This is a common weed in shady cultivated grounds, producing spikes of purplish slowers in May and June. It is very juicy, of a bitter taste, without any remarkable smell. The medical effects of this herb are, to strengthen the tone of the bowels, gently loosen the belly, and promote the urinary and other natural secretions. It is principally recommended in melancholic, scorbutic, and cutaneous disorders;

for opening obstructions of the viscera, attenuating, and promoting the evacuation of viscid juices. Frederick Hossman had a very great opinion of it as a purifier of the blood; and assure us, that in this intention scarce any plant exceeds it. Both watery and spirituous menstrua extract its virtues.

GALANGÆ MINORIS radix [E.] Galangal; a root brought from Chinz.

This root comes to us in pieces fearce an inch long, and not half for thick, full of joints, with feveral circular rings on the outside; of an aromatic smell, and a bitterish, hot, biting taste. Galangal is a warm stomachic bitter; it has been frequently prescribed in bitter insusions, but the slavour it gives is not agreeable. The London college has rejected it from the officinal simples.

GALBANUM [L. E.]

This is the concrete juice of an African plant of the ferulaceous kind. The juice, as brought to us, is semipellucid, soft, tenacious; of a strong, and to some unpleasant smell, and a bitterish warm tafte: the better fort is in pale coloured masses, which, on being opened, appear composed of clear white tears. Geoffroy relates, that a dark greenish oil is to be obtained from this fimple by distillation, which, upon repeated rectifications, becomes of an elegant sky blue colour. The purer forts of galbanum are faid by some to dissolve entirely in wine, vinegar, or water; but thefe liquors are only partial menstrua with regard to this drug; nor do spirit of wine, or oils, prove more effectual in this respect: the best diffolvenz . dissolvent is a mixture of two parts spirit of wine, and one of water. Galbanum agrees in virtue with gum ammoniacum; but is generally accounted less efficacious in asthmas, and more so in hysterical complaints. It is an ingredient in the gum pills, species e scordio, mithridate, theriaca, consectio Paulina, maturating cataplasm [L.] and antihysteric plaster [E.]

GALEGÆ folia: Galegæ vulgaris floribus ceruleis C. B. Goats

rue; the herb.

This is celebrated as an alexipharmac; but its fensible qualities discover no foundation for any virtues of this kind: the taste is merely leguminous; and in Italy (where it grows wild) it is said to be used as food.

GALLÆ [L. E.] Galls.

There are excrescences found, in the warmer countries, upon the oak tree: they are produced by a kind of infect (which wounds the young buds or branches) and afterwards ferve as a lodgement for its eggs: the animal within the gall eats its way through; those which have no hole are found to have the infect remaining in them. The best galls come from Aleppo: these are not quite round and smooth, like the other forts. and have several tubercles on the furface. Galls have a very auftere flyptic tafte, without any fmell: they are very firong astringents, and as fuch have been fometimes made use of both internally and externally, but are not much taken notice of by the present practice.

GALLII folia: Gallii lutei C. B. Ladies bedfiraw, or cheese-rennet; the herb [E.]

This herb has a subacid taste, with a very faint, not disagreeable smell: the juice changes blue vegetable insusions of a red colour, and coagulates milk, and thus discovers marks of acidity. It stands recommended as a mild styptic; but has never been much in use.

GAMBOGIA [L. B.] Gamboge; a folid concrete juice, brought from the East-Indies, in large cakes or rolls. The best fort is of a deep yellow or orange colour, breaks thining and free from drofs: it has no smell, and very little tafte, unless kept in the mouth for some time, when it impresses a slight sense of acrimony. It immediately communicates to spirit of wine a bright golden colour, and almost entirely dissolves in it; Geosfroy says, except the fixth part: alkaline falts enable water to act upon this fubstance powerfully as a menstruum: the folution made by their means is fomewhat transparent, of a deep blood-red colour, and passes the filtre: the dulcified spirit of sal ammoniac readily and entirely dissolves it, and takes up a confiderable quantity; and what is pretty remarkable, this folution mixes either with water or spirit, without growing turbid.

Gamboge evacuates powerfully both upwards and downwards a Hoffman and fome others condemn it as acting with too great violence, andoccasioning a dangerous hypercatharses; whilst others are of a contrary opinion. Geoffroy seems particularly fond of this medicine, and informs us, that he has frequently given it, from two to four grains, without its proving at all emetic: that from four to eight grains, it both vomits and purges, without violence; that its operation is

foon

foon over; and that if given in a liquid form, and sufficiently diluted, it stands not in need of any corrector; that in the form of a bolus or pill, it is most apt to prove emetic, but very rarely has this effect if joined along with mercurius dulcis. He nevertheless cautions against its use where the patient cannot easily bear vomiting. It gives name to a pill in the Edinburgh pharmacopæia, but is not used in any composition of the London.

GENISTÆ folia, flores, semen: Cytiso genistæ scopariæ vulgaris store luteo Tourn. Broom; the leaves, flowers, and feeds [E]

The leaves of this shrub have a nauseous bitter taste: decoctions of them loofen the belly, promote urine, and stand recommend-

ed in hydropic cases.

The flowers are faid to prove cathartic in decoction, and emetic in substance, though in some places, as Lobel informs us, they are commonly used, in large quantity, in falads, without producing any effect of this kind. qualities of the seeds are little better determined: some report, that they purge, almost as strongly as hellebore, in the dose of a drachm and balf; whilst the author above-mentioned relates, that he has given a decoction of two ounces of them as a gentle emetic.

GENTIANÆ radix: Gentianæ majoris luteæ C. B. Gentian;

the root [L. E.]

This plant is found wild in fome parts of England: but the dried roots are most commonly brought from Germany, &c. they should be chosen fresh, and of a yellow or bright gold colour within. This root is a strong bitter, and as such, very frequently made use of in practice: in taste it is less exceptionable than most of the other substances of this class: infusions of it, flavoured with orange peel, are sufficiently grateful. It is the capital ingredient in the bitter wine, tincture, and infusion of the shops. An extract made from it is likewise an officinal preparation.

A poisonous root was some years ago discovered among some of the gentian brought to London: the use of which occasioned violent disorders, and in some instances death. This is easily distinguishable by its being internally of a white colour, and void of bitterness. This poisonous fimple feems to be the root of the thora valdenfis of Ray, the atonitum primum pardalianches of Geiner; a plant which Lobel informs us the inhabitants of some parts of the Alps used formerly to empoison darts with.

GERANII BATRACHOIDIS folia. Crowfoot cranesbill; the leaves.

GERANII ROBERTIANI folia. Herb Robert; the leaves.

These plants are found wild, the first in hedges, the second in moist meadows. The leaves have an austere taste, and have hence been recommended as astringents; but they have long been difregarded in practice, and have therefore been omitted at the late reformation both of the London and Edinburgh pharmacopæias.

GINSENG [E.] a finall root brought from North America, and fometimes from China; an inch or two in length, taper, finely striated, of a whitish or yellowish colour. It has a very sweet tafte, accompanied with a flight bitterishness and warmth.

The

The Chinese are said to have a very extraordinary opinion of the virtues of this root, and to look upon it as an universal restorative, in all decays, from age, intem-The great perance, or disease. value, there set upon it, has prevented its being exported from thence into other countries, and its discovery in North America is but of late date, fo that among us it has hitherto been very rarely made use of; although, from what can be judged of it from the tafte, it seems to deserve some regard, especially as it is now procurable in plenty.

GITH, vide NIGELLA.

GLASTI folia: I fatidis fative wel latifolia C. B. Wood; the leaves. This plant is cultivated for the use of the dyers; but is never employed for any medicinal purposes. The virtues attributed to it are those of an astringent.

GLADIOLI LUTE! radix: Iridis palustris lutea, five acori adulterini J. B. Acori vulgaris pharm. August. et Wirt. Yellow water-flag, bastard acorus, or water flower-de-luce: the roots [L.]

This grows common by the brinks of rivers and in other watery places. The root has a very acrid tafte, and proves, when fresh, a strong cathartic: its expressed juice, given to the quantity of eighty drops every hour or two, and occasionally increased, has occasioned a plentiful evacuation, after jalap, gamboge, &c. had proved ineffectual. (See the Edinburgh effays, vol. v. art. 8. Abridg. vol. i. page 202.) By drying, it lofes its acrimony and purgative virtue. The pulvis ari of our difpenfatory contains about one-fifth of the dry root; the Edinburgh uses in its place the acerus werus, or calamus aromaticus.

GLYCYRRHIZÆ radix: Glycyrrbizæ filiquosæ vel Germanicæ C.

B. Liquorice; the root [L. E.]

This is produced plentifully in all the countries of Europe: that which is the growth of our own is preferable to fuch as comes from abroad; this last being generally mouldy, which this root is very apt to become, unless kept in a dry place. The powder of liquorice usually fold is often mingled with flower, and I fear too often with substances not quite so wholesome: the best fort is of a brownish yellow colour (the fine pale yellow being generally fophisticated) and of a very rich fweet taste, much more agreeable than that of the fresh root. Liquorice is almost the only sweet that quenches thirst; whence it was called by the Greeks adipson. Galen takes notice, that it was employed in this intention in hydropic cases, to prevent the necesfity of drinking. Mr. Fuller, in his Medicina gymnastica, recommends this root as a very vseful pectoral, and fays it excellently foftens acrimonious humours, at the same time that it proves gently detergent: and this account is warranted by experience. It is an ingredient in the pectoral syrup, pectoral troches, the compound lime waters, decoction of the woods, compound powder of gum tragacanth, lenitive electary, and theriaca. An extract is directed to be made from it in the shops, but this preparation is brought chiefly from abroad, though the foreign extract is not equal to fuch as is made with proper care among ourselves.

L 2

GRAMINIS CANINI radix: Graminis canini arvensis, sive graminis Dioscoridis C. B. Quickgrass; the roots.

Grass roots have a sweet roughish taste. They are principally recommended in aperient spring drinks, for what is called purifying and sweetening the blood.

GRANA PARADISI: Cardamomum majus semini piperato Geoffroii. Grains of paradise; a fruit, brought from the East Indies [E.]

This fruit is about the fize of a fig, divided internally into three cells, in each of which are contained two rows of small seeds like cardamoms. These seeds are somewhat more grateful, and confiderably more pungent, than the common cardamoms, approaching in this respect to pepper, with which they agree also in their pharmaceutical properties; their pungency residing, not in the distilled oil, as that of cardamom seeds does, but in the resin extracted by spirit of wine.

GRANATI cortex: Frudus Mali punice fative C. B. The rind of the pomegranate, called

malicorium [L. E.]

The pomegranate tree is fometimes met with in our gardens; but the fruit, for which it is chiefly valued, rarely comes to fuch perfection as in warmer climates. This fruit has the general qualities of the other fweet fummer fruits, allaying heat, quenching thirst, and gently loosening the belly. The rind is a strong astringent, and as such is occasionally made use of.

GRATIOLÆ folia: Gratiolæ centaurioidis C. B. Hedge hyfop; the leaves [E.]

This is a small plant, met with,

among us, only in gardens. The leaves have a very bitter, difagreeable tafte; an infusion of a handful of them when fresh, or a drachm when dried, is faid to operate strongly as a cathartic. Kramer reports (Tentam. botanic. p. 18.) that he has found the root of this plant a medicine similar in virtue to ipecacoanha.

GUAIACI lignum, cortex, gummi: Guaiaci Americani primi fructus aceris, five legitimi Breyn. prodr. Guaiacum, a tree growing in the warmer parts of the Spanish West-Indies; its wood, bark, and refin called gum guaiacum [L. E.]

The wood is very ponderous, of a close compact texture; the outer part is of a yellow colour, the heart of a deep blackish green, or variegated with black, green, pale and brown colours: the bark is thin, smooth, externally of a dark greyish hue: both have a lightly aromatic, bitterish, pungent taste; the bark is somewhat the weakest. The refin (which exudes from incisions made in the trunk of the tree) is brought to us in irregular masses, usually friable, of a dusky greenish, and sometimes of a reddish cast, with pieces of the wood among them: its tafte is more acrid and pungent than that of the wood or bark.

Their general virtues are those of a warm, stimulating medicine: they strengthen the stomach and other viscera; and remarkably promote the urinary and cuticular discharge: hence in cutaneous defedations, and other disorders proceeding from obstructions of the excretory glands, and where sluggish serous humours abound, they are eminently useful: rheumatic and other pains have often been relieved by them. The resin is the most active of these drugs;

and

and the efficacy of the others depends upon the quantity of this part contained in them: the refin is extracted from the wood in part by watery liquors, but much more perfectly by spirituous ones; the watery extract of this wood, kept in the shops, proves not only less in quantity, but confiderably weaker than one made with spirit. This last extract is of the same quality with the native refin, and differs from that brought to us only in being purer. The gum, or extracts, are given from a few grains to a scruple or half a drachm: which last dose proves for the most part confiderably purgative. The officinal preparations of guaiacum are an extract of the wood [L.] a folution of the gum in rectified spirit of wine [L.] and a folution in volatile spirit [L. E.] as also an empyreumatic oil distilled from the wood. The wood is an ingredient in the compound lime water [L. E.] and in the compound tinctures of jalap and fena [E.] of which it increases the purgative virtue; the gum, in the aromatic pills [L.] ecphractic pills [L. E.] mercurial and ethiopic pills, Edinburgh theriaca [E.] and the compound oil of balfam of Copaiba [L.]

GUMMI AMMONIACUM, vide Ammoniacum.

GUMMI ARABICUM [L. E.] Gum Arabic; a concrete gum, exuding from the Egyptian acacia tree. This is brought to us from Turkey, in small irregular masses or strings, of a pale yellowish colour. The true gum Arabic is rarely to be met with in the shops; gum senega or senica, which comes from the coasts of Guinea, being usually sold for it: this greatly resembles

the other, and perhaps, as Dale conjectures, exudes from a tree of the same kind: it is generally in large pieces, rough on the outfide; and in these circumttances possibly consists the only difference betwixt the two; although the former is held to be the purer and finer gum, and therefore preferred for medicine; and the latter the strongest, most substantial, and cheapest, and consequently more employed for mechanic uses. The virtues of this gum are the same with those of gummy and mucilaginous substances in general: it is given, from a scruple to two drachms, in hoarsenesses, a thin acrimonious state of the juices. and where the natural mucus of the intestines is abraded. It is an ingredient in the white decoction, chalk julep, the com-pound powders of bole, fcordium, amber, gum tragacanth, the common emulsion, mithridate, theriaca, and some of the troches.

GUMMI CERASORUM. Cherry-tree gum.

There is not any medical difference betwirt this and the preceding. Some have supposed that all the gum brought to us from the East, under the name of Arabic, is no other than the gum of cherry, plum, and other trees common among ourselves. This opinion is nevertheles erroneous; for these trees, as Geoffroy observes, do not grow in the countries from whence gum Arabic is brought; whilst the acaciae are very common there.

GUMMI ELEMI, vide E-

GUMMI TRAGACANTHÆ
[L. E.] The gum of the tragacanth,
a thorny bush growing in Crete,
L 3
Asia,

Asia, and Greece. This gum is of a much stronger body than either of the foregoing, and does not so persectly dissolve in water. A drachm will give a pint of water the consistence of a syrup, which a whole ounce of gum Arabic is scarce sufficient to do. Hence its use for forming troches, and the like purposes, in preference to the other gums. It gives name to an officinal powder, and is an ingredient in the compound powders of cerus and amber.

GUTTA GAMBA, vide GAM-

HÆMATITES lapis [L. E.] Hematites, or bloodstone.

This is an elegant iron ore, extremely hard, of a dark reddift or yellowish colour: it is found either along with other ores of iron, or in distinct mines by itself. With regard to its medical virtues, we conceive they do not vary from those experienced from rust, and the common croci of iron, notwithstanding the extraordinary opinion which many, even of the late practitioners, have entertained of it; as of its curing ulcers of the lungs, which Geoffroy says the hæmatites dries and heals.

HALICACABUM, vide Alkekengi.

HEDERÆ ARBOREÆ folia, Baccæ, et gummi seu resina; Hederæ communis majoris Raii. Ivy; the leaves, berries, and resin called gum hederæ [E.]

This is a climbing shrubby plant, growing commonly from the trunks of trees, or on old walls. The leaves have very rarely been given internally, notwithstanding they are recommended (in the Ephem. natur. suriof. vol. ii, obs. 120.) against

the atrophy of children: their tafte is nauteous, acrid, and bitter. Externally they have fomerimes been employed for drying and healing ichorous fores, and likewife for keeping issues open. The berries were supposed by the ancients to have a purgative and emetic quality: later writers have recommend. ed them in small doses, as diaphoretics and alexipharmacs; and Mr. Boyle tells us, that in the London plague the powder of them was given with vinegar, with good fuccess, as a sudorific. It is probable the virtue of the composition was rather owing to the vinegar than to the powder. The refin was ranked by the ancients (if their diagon To mood was the same with our gummi bederæ) among the depilatories; from this class, which it certainly had no title to, it has fince been removed to that of conglutinaters of wounds, to which it has no very just one.

HEDER & TERRESTRIS folia; Heder & terrestris vulgaris C. B. Ground-ivy; the leaves [E.]

Ground-ivy is a low plant, frequent in hedges and shady places. It has an aromatic, though not veryagreeable smell; and a quick, bitterish, warm taste. This herb is an useful corroborant, aperient, and detergent; and hence stands recommended against laxity, debility, and obstructions of the viscera: seme have had a great opinion of it for cleanfing and healing ulcers of the internal parts, even of the lungs; and for purifying the blood. It is customary to infuse the dried leaves in malt liquors: a practice not to be commended, though it readily communicates its virtue, and likewise helps to fine them down: scarce any other herb has this effect more remarkably than ground-ivy.

HELE-

HELENIUM, vide Enula CAMPANA.

HELLEBORI ALBI radix: Hellebori albi flore fubviridi C. B. White hellebore; the root [L. E.]

This plant grows spontaneously in Switzerland, and the mountainous parts of Germany. The root has a nauseous, bitterish, acrid taste. burning the month and fauces: wounded when fresh, it emits an extremely acrimonious juice, which mixed with the blood, by a wound, is said to prove very dangerous: the powder of the dry root, applied to an issue, occasions violent purging: inuffed up the noie, it proves a strong, and not always a fafe sternutatory. This root. taken internally, acts with extreme violence as an emetic, and has been observed, even in a small dose, to occasion convulsions, and other terrible disorders. The ancients sometimes employed it in very obstinate cases, and always made this their last resource. Modern practice seems to have almost entirely rejected its internal use; though I am informed that some have lately ventured upon so large a dose as a fcruple, in maniacal cases, and found good effects from it after the stronger antimonial preparations had been given in vain. A tincture and honey of it are kept in the shops [L.]

HELLEBORI NIGRI radix: Hellebori nigri flore roseo C. B. Black hellebore; the roots [L. E.]

This plant grows wild in the mountainous parts of Switzerland, Austria, and Stiria: the earliness of its flowers, which sometimes appear in December, has gained it a place in our gardens.

In some parts of Germany, a species of black hellebore has been made use of, which not unfrequent-

ly produced violent, and fometimes deleterious effects: this the Wirtemberg college particularly caution against, though without mentioning any marks by which it may be distinguished, or even giving the precise name of the plant. It appears to be the fetid black hellebore of C. B. called in England, where it grows wild, setterwort, settlewort, or bastard hellebore: the roots of this may be distinguished from the officinal fort by their being less black. roots of the poisonous aconites refemble in appearance those of the black hellebore: and in the Breslaw collections we find fome instances of fatal effects occasioned by mistaking the former for the latter: these also are happily discoverable by their colour; the aconitum being lighter coloured than even the paleft of the black hellebores. faculty of Paris, by allowing the use of one of the paler hellebores (the green-flowered, which grows wild in England, and is called by our farriers, peg-root) have in some measure deprived the shops of the benefit of this criterion: but the London college have directed the darkest coloured of all the roots of this class. Since therefore the two noxious roots which the buyer is most apt to mistake for this, are distinguishable from it by their colour, but have no other external mark by which they may be with certainty known, particular regard ought to be had to this circumstance; only the deepest black being chosen, and all the paler roots rejected.

The tafte of Hellebore is acrid and bitter. Its acrimony, as Dr. Grew observes, is first felt on the tip of the tongue, and then spreads immediately to the middle, without being much perceived on the intermediate part: on chewing it

OL

for a few minutes, the tongue feems benumbed, and affected with a kind of paralytic stupor, as when burnt by eating any thing too hot: the fibres are more acrimonious than the head of the root which they issue from. Black hellebore root, taken from fifteen grains to half a drachm, proves a strong cathartic; and as fuch has been celebrated for the cure of maniacal, and other disorders, proceeding from what the ancients called atra bilis: in which cases, medicines of this kind are doubtless occasionly of use, though they are by no means possessed of any specific power. It does not however appear, that our black hellebore acts with fo much violence as that of the ancients: whence many have supposed it to be a different plant: and indeed the descriptions which the ancients have left us of their heliebore, do not agree to any of the forts usually taken notice of by modern botanists. Another species has been discovered in the eastern countries, which Tournefort diftinguishes by the name of belleborus niger orientalis, amplissimo folio, caule præalto, flore purpurascente, and supposes to be the true ancient hellebore, from its growing in plenty about mount Olympus, and in the island Anticyra, celebrated of old for the production of this antimaniacal drug: he relates, that a scruple of this fort, given for a dose, occasioned convulfions.

Our hellebore is at present looked upon principally as an alterative, and in this light is frequently employed, in small doses, for attenuating viscid humours, promoting the uterine and urinary discharges, and opening inveterate obstructions of the remoter glands; it often proves a very powerful emmenagogue in plethoric habits, where seed

is ineffectual or improper. An extract made from this root with water, is one of the mildest, and for the purposes of a cathartic, the most effectual preparation of it; this operates sufficiently, without occasioning the irritation which the pure resin is accompanied with. A tincture drawn with proof spirit, contains the whole virtue of the hellebore, and feems to be one of the best preparations of it when defigned for an alterative: this tincture, and the extract, are kept in the shops. The college of Edinburgh makes this root an ingredient in the purging cephalic tingture, and compound tincture of jalap; and its extract in the purging deobstruent pills, gamboge pills, the laxative mercurial pills, and Rudius pills, or the compound cathartic extract.

HELXINE, vide PARIETA-

HEPATICÆ NOBILIS folia: Ranunculi tridentati verni, flore simplici cæruleo Tourn. Noble liverwort; the leaves [E.]

This herb has a place in our gardens on account of the beauty and early appearance of its flowers. It is a cooling, gently refiringent herb; and hence recommended in a lax state of the sibres as a corroborant.

HEPATICA TERRESTRIS, vide Lichen.

HERBÆ PARIS folia et frudus; Solani quadrife'ii bacciferi C. B. Herb Paris, truelove, or one-berry; the leaves and fruit.

This is a low plant, growing wild in shady woods. It is said, but on no good grounds, to be alexipharmac: Gesner relates, that its juice has killed poultry; and its smell and tafte manifelly agree with except that the hippofelinum is somethose of the narcotic herbs.

HERMODACTYLUS. Hermodactil; a root brought from Turkey. It is of the shape of a heart flatted, of a white colour, compact, yet easy to cut or powder; of a viscous sweetish taste, with a light degree of acrimony.

Hermodactils were of great repute among the ancients as a cathartic; but those we now meet with in the shops have very little purgative virtue; Neumann declares he never found them to have any effect at all.

HERNIARIÆ folia: Polygoni minoris hwe millegranæ majoris glabræ C. B. Rupture-wort; the leaves.

This is a low herb, growing wild in fandy and gravelly grounds. It is a very mild restringent, and may, in some degree, be serviceable in diforders proceeding from a weak flaccid state of the viscera: the virtue which it has been most celebrated for, it has little title to, that of curing bernias.

HIPPOGLOSSI folia: Rusci angustifolii, frudu folio innascente Tourn. Double tongue; the leaves.

This is met with only in garden. where plants are cultivated for curiofity. It has rarely been taken notice of by medicinal writers, though retained in our pharmacopœias till the late emendation.

HIPPOSELINI, seu Smyrnii, folia, rudix, semen: Hipposelini Theophrasti, vel Smyrnii Dioscoridis, C. B. Alexanders; the leaves, root, and feeds [E.]

This is an umbelliferous plant, differing from the others of that class, in bearing a large tumid black feed: it grows by the fea fide, upon rocks. In medical qualities it agrees with apium (smallage) what stronger.

HIRUNDINARIA, vide VIN-CETOXICUM.

HORDEI semen: Hordei distichi, quod spica binas ordines habeat Plinie C. B. Common barley  $\{L. E.\}$ 

HORDEUM GALLICUM hive MUNDATUM. French barley, or the common barley freed from the fhell.

HORDEUM PERLATUM dictum [L.] Pearl barley; prepared in Germany and Holland, by grinding the shelled barley into little round granules, which appear of a kind of pearly whiteness.

Barley, in its several states, is more cooling, less glutinous, and less nutritious than wheat or oats; among the ancients, decoctions of it were the principal aliment, and medicine, in acute diseases. The London college direct a decoction of pearl barley; and both the London and Edinburgh make common barley an ingredient in the pectoral decoction.

HORMINI SATIVI, seu Sclareæ, folia, semen: Hormini sclareæ diti C. B. Garden clary; the leaves and seeds [E.]

These have a warm, bitterish pungent tafte; and a strong, not very agreeable smell: the touch discovers in the leaves a large quantity of glutinous or refinous matter. They are principally recommended in the fluor albus, and other female weaknesses, in hysteric diforders, and in flatulent colies.

HYBERNICUS LAPIS: Tegula vel ardesia Hybernica [E.] Lish

This a blackish sossile stone brought from Ireland. It feems to confift of an argillaceous or bolar warth, earth, flightly impregnated with fulphur and iron; and may be prefumed to possess in a low degree the virtues of the other ferugineous minerals.

HYDRARGYRUS, vide Argentum vivum.

HYDROLAPATHUM, vide LAPATHUM.

HYOSCYAMI folia: Hyoscyami a'bi majoris wel tertii Dioscoridis et quarti Plinii G. B. White henbane; the leaves [E.]

This is met with only in botanic

gardens.

HYOSCYAMI NIGRI folia: Hyoscyami vulgaris vel nigri C. B. The common wild, or black hen-

bane; the leaves.

These plants stand recommended for fundry external purpoles; and by some likewise internally against dyfentries and hæmorrhages; but these are so many examples of their pernicious effects, that common practice has very deservedly rejected them. They are strong and virulent narcotics, greatly disorder the fenses, occasioning deliria and madness, either deadly, or of long duration. Haller tells us of one who eat of all the poisons of the physic garden, the napelli, apocyna, bella donna, without injury, but was mastered by this; that after its common effects as a narcotic had abated, a paralysis of one of the legs remained; and that Boerhaave had his senses disordered by only making a plaster from this plant. There are other examples also, though from less unexceptionable authorities, of henbane proving narcotic, though none of it was received into the body.

HYPERICI folia, flores, femen: Hyperici wulgaris, C. B. St. John's

wort; the leaves, flowers [L. E.] and feeds [E.]

This plant grows wild in woods and uncultivated places throughout England. Its tafte is rough and bitterish; the smell disagreeable. Hypericum has long been celebrated as a corroborant, diuretic, and vulnerary; but more particularly in hysterical and maniacal disorders: it has been reckoned of such efficacy in these last, as to have thence received the name of fuga dæmonum. It is observable, that the flowery tops tinge expressed oils of a red colour (which very few vegetable substances will do) and communicate a blood red to rectified spirit. The oil tinged by them is kept in the shops [L.]

HYPOCISTIDIS succus inspissatas: Hypocistidis sub cisto C. B. Juice of hypocistis [L. E.]

Hypocifits is a flethy production, growing in the warmer climates from the roots of different kinds of cifti. Its inspissated juice is an astringent, similar to acacia, but somewhat stronger. At present it is scarce otherwise made use of, than as an ingredient in some of the old compositions, viz. mithridate, theriaca, and the compound powder of amber [L.]

HYSSOPI folia: Hysfopi officinarum, carulea frue spicata C. B. Hysfop; the leaves [L. E.]

The leaves of hysfop have an aromatic smell, and a warm pungent taste. Besides the general virtues of aromatics, they are particularly recommended in humoral asthmas, coughs, and other disorders of the breast and lungs; and said to notably promote expectoration. A simple water is prepared from them in the Edinburgh pharmacopæia.

JACOBÆÆ folia: Jacobææ vilgaris laciniatæ C. B. Ragwort,

or feggram: the leaves.

This ragged-leaved plant grows wild by road fides, and uncultivated places. Its tafte is roughish, bitter, pungent, and extremely unpleasant: it stands strongly recommended by Simon Paulli against dysenteries; but its forbidding taste has prevented its coming into practice.

JALAPIUM [L. E.] Jalap.

Jalap is the root of an American convolvulus, brought to us in thin transverse slices, from Xalapa, a province of New Spain. Such pieces should be chosen as are most compact, hard, weighty, dark coloured, and abound most with Slices of black circular striæ. bryony root are faid to be sometimes mixed with those of jalap: these may be easily distinguished by their whiter colour, and less compact texture. This root has no fmell, and very little taffe upon the tongue; but when swallowed, it affects the throat with a sense of heat, and occasions a plentiful difcharge of saliva.

Jalap in substance, taken in a dole of about half a drachm (less or more, according to the circumstances of the patient) in plethoric, or cold phiegmatic habits, proves an effectual, and in general a fafe purgative, performing its office mildly, feldom occasioning nausea or gripes, which too frequently accompany the other strong cathartics. In hypochondriacal disorders, and hor bilious temperaments, it gripes violently, if the jalap be good: but rarely takes due effect as a purge. An extract made by water purges almost universally, but weakly; and at the same time, has a confiderable effect by urine: the root remaining after this process, gripes violently. The pure resin, prepared by spirit of wine, occasions most violent gripings, and other terrible symptoms, but scarce proves at all cathartic: triturated with fugar, or with almonds into the form of an emuliion, or disfolved in spirit and mixed with fyrups, it purges plentifully in a fmall dose, without occasioning much disorder: the part of the jalap remaining after the separation of the refin, yields to water an extract, which has no effect as a cathartic, but operates powerfully by urine. Its officinal preparations are an extract made with water and fpirit [L. E.] a refin [E.] a fimple tincture [L. E.] and a compound tincture [E.] The extract is the basis of one of the purgative pills

Frederick Hoffman particularly cautions against giving this medicine to children, and affures us, that it will destroy appetite, weaken the body, and perhaps occasion even death. In this point, this celebrated practitioner was probably deceived: children, whose vessels are lax, and the food soft and lubricating, bear these kinds of medicines, as Geosfroy observes,

better than adults.

JAPONICA TERRA five catechu [L. E] Japan earth, improperly so callea; being neither an earth, nor the produce of Japan; but an inspissated vegetable juice, prepared in the East-Indies from the fruit, as is supposed, of the areca palm tree. It is dry and pulverable, outwardly of a reddish colour, inwardly of a thining dark brown, almost black. with some cast of red. When pure, it dissolves totally in water, and almost totally in rectified spirit : as we usually meet with it, a confiderable quantity of fandy matter is left by both these menstrua. This medimedicine is a mild aftringent, and frequently employed as fuch in alvine fluxes, uterine profluvia, in laxity and debility of the viscera in general, and in coughs proceeding from thin acrid defluxions. Its taile is more agreeable than that of most other substances of this class: chewed for some time, it leaves a kind of sweetishness in the mouth. The troches and tineture, kept in the shops, are very elegant preparations of it. It gives name to an officinal confection [E.] and is an ingredient in the compound powder of amber, mithridate, and theriaca [L.]

JASMINI flores: Jasmini vulgatioris flore albo C. B. Jasmine; the flowers.

This is a small tree, commonly planted in our gardens. The flowers have a strong smell, which is liked by most people, though to fome disagreeable: expressed oils extract their fragrance by infusion; and water elevates somewhat of it in distillation, but no essential oil has hitherto been obtained from them; the distilled water, kept for a little time, loses its odour. to their medical virtues, the present practice expects not any from them, notwithstanding they have been recommended forpromoting delivery, curing ulcerations of the uterus, &c.

IBERIDIS foliu: Lepidii gramineo fulio five iberidis Tourn. Sciatica cresses; the herb.

This is met with in botanic gardens: in taste, smell, and medical virtues, it agrees with the nasturtium. It has been particularly recommended in external applications against the sciatica, whence the English name of the plant.

ICHTHYOCOLLA [E] Fish-glue, or ifing-glass.

This is a folid glutinous substance. obtained from a large kind of fish, caught in the feas of Muscovy. The skin, and some other parts of the animal are boiled in water, the decoction inspissated to a proper consistence, and then poured out so as to form thin cakes; these are either farther exsiccated till perfectly dry, or cut whilst soft into slices, which are afterwards bent, or rolled up into spiral, horseshoe, and other shapes. This glue is more employed for mechanic purpofes than in medicine. It may be given in a thin acrimonious state of the joices, after the same manner as the vegetable gums and mucilages; regard being had to their different disposition to putrescence.

IMPERATORIÆ, seu Magistrantiæ radix: Imperatoriæ majoris C.B. Masterwort; the root [E.]

This is a native of the Alps and Pyrenean mountains, and some parts of Germany, from whence we are supplied with roots superior in aromatic flavour to those raised in our gardens. The smell of this root is very fragrant; its taste bitterish, warm, and pungent, glowing in the mouth for a long time after it has been chewed. fimple, though undoubtedly an elegant aromatic, is not regarded in the present practice: it is scarcely ever directed in extemporaneous prescription, and the only officinal composition it has a place in, is the plague water of the Edinburgh pharmacopæia. Its flavour is fimilar to that of angelica, but stronger.

IPECACOANHA [L. E.] a root brought from the Spanish West-Indies.

It is divided into two forts, Peruvian and Brazilian: but the eye dislinguishes three, ash coloured or

grey,

grey, brown, and white. ' The ash coloured, or Peruvian ipecacoanha of the shops, is a small wrinkled root, bent and contorted into a great variety of figures, brought over in short pieces, full of wrinkles, and deep circular fissures, quite down to a small white woody fibre that runs in the middle of each piece: the cortical part is compact, brittle, looks smooth and refinous upon breaking: it has very little finell; the tafte is bitterish and fubacrid, covering the tongue, as it were, with a kind of mucilage. The brown is small, and somewhat more wrinkled than the foregoing, of a brown or blackish colour without, and white within; this is brought from Brazil. The white fort is woody, has no wrinkles, and no perceptible bitterness in tafte. The first fort (the ash coloured or grey ipecacoanha) is that usually preferred for medicinal use. The brown has been sometimes observed, even in a small dose, to produce violent effects. The white, though taken in a large one, has scarce any effect at all: Mr. Geoffroy calls this fort bastard ipecacoanha, and complains that it is an imposition upon the public. what species of plant the ipecacoanha belongs, has not as yet been determined. Geoffroy, Neumann, Dale, and Sir Hans Sloane, inform us, that the roots of a kind of apocynum (dogs bane) are too frequently brought over inflead of it: and inflances are given of ill confequences following from the use of these roots: if the marks above laid down, particularly the ash colour, brittleness, deep wrinkles, and bitterish tafte, be carefully attended to, all mistakes of this kind may be prevented.

Ipecacoanha was first brought into Europe about the middle of last century, and an account of it published about the same time by Piso: but it did not come into general use till about the year 1686, when Helvetius, under the patronage of Lewis XIV. introduced it into practice. This root is one of the mildest and safest emetics we are acquainted with; and has this peculiar advantage, that if it should not operate by vomit, it passes off by the other emunctories. It was first introduced among us with the character of an almost infallible remedy in dysenteries, and other inveterate fluxes; as also in diforders proceeding from obstructions of long standing: nor has it lost much of its reputation by time. In dysenteries, it almost always produces happy effects, and often performs a cure in a very short space of time. In other fluxes of the belly, in beginning dysenteries, and fuch as are of a malignant kind, or where the patient breathes a tainted air, it has not been found equally successful: in these cases it is necessary to continue the use of this medicine for several days, and to join with it opiates, diaphoretics, This root, given in and the like. fubstances, it as effectual, if not more so, than any of the preparations of it: the pure refin acts as a strong irritating emetic, but is of little service in dysenteries; whilst an extract prepared with water is almost of equal service in these cases with the root itself, though it has little effect as an emetic. Geoffroy concludes from hence, that the chief virtue of ipecacoanha in dysenteries depends upon its gummy fabitance, which lining the intestines with a fost mucilage, when their own mucus has been abraded, occasions their exulcerations to heal, and defends them from the acrimony of the juices: and that the refinous part in which the emetic quality resides, is required,

required, where the morbific marters is lodged in the glands of the stomach and intestines. But if the virtues of this root were entirely owing to its mucilaginous, or gummy parts, pure gums, or mucilages, might be employed to equal advantage. Water, assisted by a boiling heat, takes up from all vegetables a confiderable portion of refinous along with the gummy matter: if the ipecacoanha remaining after the action of water be digested with pure spirit, it will not yield half so much resin as at first: so that the aqueous extract differs from the crude root only in degree, being proportionably less refinous, and having less effect, both as an emetic, and in the cure of dysenteries. The virtues of ipecacoanha, in this diforder, depends upon its promoting perspiration, the freedom of which is here of the utmost importance, and an increase of which, even in healthful persons, is generally obferved to suppress the evacuation by stool. In dysenteries, the skin is for the most part dry and tense, and perspiration obstructed: the common diaphoretics pals off without effect through the intestinal canal: but ipecacoanha, if the patient, after a puke or two, be covered up warm, brings on a plentiful sweat. After the removal of the dysentery, it is necessary to continue the use of the medicine for some time longer, in order to prevent a relapse: for this purpose, a sew grains, divided into feveral doses, so as not to occasion any fensible evacuation, may be exhibited every day; by this means the cure is effectually established. And indeed small doses given, even from the beginning, have been often found to have better effects in the cure of this disease than larger ones. Geoffroy informs us, from his own experience, that he has observed ten grains of the powder to act as effectually as a scruple or two; and therefore confines the doses betwixt six and ten grains: it has lately been found, that even smaller doses prove sufficiently emetic. The only officinal preparation of this root is a tincture made in wine [L.E.]

IRIDIS FLORENTINÆ radix: Iridis Florentinæ albæ C. B.
Florentine orris; the root.

IRIDIS PURPUREÆ NOS-TRATIS radix: Iridis vulgaris Germanicæ five fykvestris C. B. Flower-de-luce; the root [E.]

Both these appear to be the same species of plant: several varieties of it are cultivated in our gardens on account of the elegance of their flowers. The roots, when recent, have a bitter, acrid, nauseous taste: and taken into the body, prove strongly cathartic; and hence the juice is recommended in dropfies, in the dose of three or four scruples. By drying they lose this quality, yet still retain a somewhat pungent, bitterish taste: their smell in this flate is of the aromatic kind; those produced in the warmer climates have a very grateful flavour, approaching to that of March violets: hence the use of the Florentine iris in perfumes, and for flavouring liquors: the shops employ it in the white pectoral troches [L.E.] and as an ingredient in the theriaca [L.] pectoral decoction, and pectoral oxymel [E.]

IVA ARTHRITICA, vide CHAMÆPITYS.

JUGLANDIS cortex et fruëtus, The walnut, and its outer shell.

The kernal of the fruit is fimilar in quality to almonds: the shell is astringent; aftringent; but neither of them are at present employed in emdicine.

JUJUBÆ, Jujubes; a half-dried fruit brought from France.

Jujubes have a pleasant sweet taste. They are recommended in an acrimonous state of the juices; in coughs from thin sharp defluxions; and in heat of urine: but they are at present, among us, a stranger to medicinal practice, and to the shops.

JUNCUS ODORATUS: Juncus odoratus five aromaticus C. B. Sweet rush, or camels hay [L.]

This, is a dry smooth stalk. brought to us along with the leaves, and sometimes the flowers. from Turkey and Arabia, tied up in bundles about a foot long. The stalk, in shape and colour, fomewhat resembles a barley straw: it is full of a fungous pith, like those of our common rushes: the leaves are like those of wheat, and furround the stalk with feveral coats, as in the reed: the flowers are of a carnation colour, flriped with a lighter purple. The whole plant, when in perfection, has a hot bitterish, not unpleasant, aromatic taste: and a very fragrant smell; by long keeping, it loses greatly of its aromatic fla-Distilled with water, it Your. yields a confiderable quantity of essential oil. It was formerly often used as an aromatic, and in ob-Rructions of the viscera, &c. but at present is scarce otherwise employed than as an ingredient in mithridate and theriaca.

JUNIPERI bacca, lignum, gummi: Juniperi vulgaris fruticofa C. B. Juniper; the berries [L. E.] wood, and the refin (improperly called gum) which exudes from it in the warmer climates [E.] This is an evergreen shrub, growing upon heaths and hilly grounds in all the parts of Europe: the wood and resin are not at present made use of for medicinal purposes: the berries are brought from Holland, where this

thrub is very plentiful.

Juniper berries have a strong, not disagreeable smell; and a warm, pungent sweet taste, which, if they are long chewed, or previously well bruised, is sollowed by a bitterish one. The pungency feems to refide in the bark; the fweet in the juice; the aromatic flavour in oily vesicles, spread through the substance of the pulp. and distinguishable even by the eye; and the bitter in the feeds: the fresh berries yield, on expresfion, a rich, sweet, honey-like, aromatic juice; if previously pounded so as to break the seeds, the juice proves tart and bitter.

These berries are useful carminatives and stomachies: for these purposes, a spirituous water [L.] and essential oil distilled from them [L. E.] are kept in the shops: they are ingredients also in the compound horseradish water, tincture of jalap, tincture of sena [E.] mithridate and theriaca [L.] the liquor remaining after the distillation of the oil, passed through a strainer, and gently exhaled to the confistence of a rob, proves likewife a medicine of great utility, and in many cases is perhaps preferable to the oil, or berry itself: Hoffman is expressly of this opinion, and strongly recommends it in debility of the Romach and intestines, and says it is particularly of service to old people who are subject to these disorders, or labour under a difficulty with regard to the urinary excretion: this rob is of a dark, brownish yellow colour, a balsamic sweet taste,

with a little of the bitter, more or less, according as the feeds in the berry have been more or less bruised.

KALI folia: Kali mujoris cochleato semine C. B. Glaffwort ; its leaves, and the alkaline falt called cineres clavellati, or potash [E.] which used formerly to be prepared from this plant only, but now from fundry forts of woods, and other vegetable matters indifferently (see the article CINE-RES RUSSICIA) Several forts of these salts, differing in degree of purity and strength, are to be met with in the shops of the dry-salter: they are rarely to be found under this denomination in those of the apothecary or druggift.

KERMES [L E.] a round grain about the bulk of a pea, found (in Spain, Italy, and in the fouthern parts of France) adhering to the branches of the ilex aculeata

eocciglandifera C. B.

I hese grains appear, when fresh, full of small, reddish ovula, or animalcules, of which they are the nidus. On expression, they yield a red juice, of a bitterish, iomewhat rough and pungent tafte, and a not unpleasant smell: this is brought to us from the fouth of The grains themselves France. are cured by sprinkling with vine. gar before exiccation: this prevents the exclusion of the ova, and kills such of the animals as are already hatched; otherwise, they change into a winged infect, leaving the grain an empty hulk.

Kermes, confidered as a medicine, is a grateful, very mild refiringent, and corroborant. In this light it was looked upon by the Greeks: the Arabians added a cordial virtue: European writers also have in general recommended it

for exhilarating the spirits, and against palpitations of the heart: but more particularly for promoting birth, and preventing abortion. I have known, fays Geoffroy, many women, who had never reached the end of pregnancy, made joyful mothers by the use of pills composed of kermes, germin. ovor. exficcat. and confectio de by. acintho (a composition, containing some vegetable aftringents and aromatics, together with gold and filver leaf, four precious stones, and other ingredients of less value:) three of these pills must be taken for the first dose, and this repeated three times, at the interval of twice three hours; after which three pills more are to be taken every morning on the three last days of the moon in every month till delivery. Notwithstanding this affertion, we conceive our readers will with us believe, that neither the kermes, or its auxiliaries, are to be much depended on. kermes gives name to an officinal confection, which appears to be greatly superior to the above composition.

LABDANUM [L. E.]

This is a refinous substance exuding upon the leaves of the ciftus ladanifera cretica flore purpures Tourn. This refin is said to have been formerly collected from the beards of goats, who brouzed the leaves of the ciftus: at present, a kind of rake, with several straps or thongs of fkins fixed to it, is drawn lightly over the shrub, so as to take up the unctuous juice. which is afterwards scraped off with knives. It is rarely met with pure, even in the places which produce it; the dust, blown upon the plant by the wind, mingling with the tenacious juice: the inhabitants are also said to mix with

it a certain black fand. In the shops two forts are met with: the best (which is very rare) is in dark coloured, almost black maffes. of the confistence of a soft plaster, which grows still fofter upon being handled; of a very agreeable smell, and of a light pungent bitterish taste: the other fort is harder, not so dark coloured, in long rolls coiled up: this is of a much weaker smell than the first, and has a large admixture of a fine fand, which, in the labdanum examined by the French academy, made up three-fourths of the mass. Rectified spirit of wine almost entirely dissolves pure labdanum, leaving only a small portion of gummy matter which has no tafte or smell: and hence this resin may be thus excellently purified for internal purposes. It is an useful ingredient in the stomachic and cephalic plasters of the shops.

LAC [E.] Milk.

Milk appears to be a vegetable juice, with little or nothing of an The quality and animal nature. use of this fost nutritious liquor are in general well known: we shall therefore, in this place, only give an account of some experiments, pointing out the alterations it undergoes from different admixtures,

and the difference in quality of the milk of different animals.

New milk mixes uniformly with common water, the mineral chalybeat waters, wines, and malt liquors that are not acid, weak vinous spirits, solutions of sugar, foaps, and neutral falts; but not with oils expressed or distilled. Acids both mineral and vegetable coagulate it; as also do fixt and volatile alkalies, and highly rectified spirit of wine: the curd made by acids is in part resolved again by alkaline liquors; as that made by alkalies likewife is by acids. Neutral falts, nitre in particular, preserve it from coagulating spontaneously; and likewise render it less easily coagulable by acids.

The human milk is the sweetest of these liquors, and that of asses next to it: this last is the most dilute of them all: on fuffering it to coagulate spontaneously, the curd scarce amounted to two drachms from twelve ounces, whilst that of cows milk was five times as much: the coagulum of affes milk, even when made by acids, forms only into fine light flakes. which swim in the serum; that of goats milk concretes into more compact masses, which sink.

Upon evapo- rating twelve ounces of		From which water extracted a fwee t faline substance, amounting, when exficcated, to drachms.
Cows milk	13	1½
Goats milk	12 <u>1</u>	1½
Human milk	8	6
Asses milk	8	6

The saline substance obtained from affes milk was white, and brown or yellow, and confiderably

less sweet; that of cows milk, the least sweet of all. It appears, therefweet as sugar; those of the others . fore, that asses milk contains more ferum, and much more of a fac-M

charine faline matter, than those of cows and goats; and that the two latter abound most with unctuous gross matter: hence these are found to be most nutritious, whilst the first proves most effectual as an

aperient and detergent.

The inspissated residuum of milk, digested with about as much water as was wasted in the evaporation, yields an elegant kind of whey, more agreeable in tatte, and which keeps better than that made in the common manner. This liquor promotes the natural secretions in general, and if its use is duly continued, does good service in scorbutic, and other disorders, proceeding from thick phlegm and obstructions of the viscera.

There are considerable differences in the milk of the same animal, according to its different aliment. Dioscorides relates, that the milk of goats, who feed on the scammony plant and spurges, proved cathartic: and examples are given in the Acta Haffniensia of bitter milk from the animal having eaten wormwood. It is a common observation. that cathartics and spirituous liquors given to a nurse, affect the child: and that the milk of animals feeding on green herbs, is much more dilute than when they are fed with dry ones. Hoffman, from whom most of the foregoing observations are taken, carries this point fo far, as to direct the animal to be dieted according to the disease which its milk is to be drank for.

LACCA, gummi refina [E.] Lac,

improperly called gum lac.

This is a fort of wax of a red colour, collected in the East-Indies, by certain infects, and deposited on Ricks fallened for that purpose in the earth. It is brought over, either adhering to the sticks, or in small transparent grains, or

in semi-transparent flat cakes: the first is called stick lac, the second feed lac, and the third shell lac. On breaking a piece of stick lac, it appears composed of regular cells like the honeycomb, with small corpufcles of a deep red colour lodged in them; these are the young infects, and to these the lac owes its tincture; for when freed from them its colour is very dilute. The shell and seed lacs, which do not exhibit any infects or cellular appearance upon breaking, are supposed to be artificial preparations of the other: the feed fort is faid to be the stick lac bruised and robbed of its more foluble parts; and the shell to be the feed lac, melted and formed into cakes. The flick lac therefore is the genuine fort, and ought alone to be employed for medicinal purposes. This concrete is of great effeem in Germany. and other countries, for laxity, and sponginess of the gums, proceeding from cold, or a scorbutic habit: for this use the lac is boiled in water, with the addition of a little alum, which promotes its folution: or a tincture is made from it with rectified spirit. This tincture is recommended also internally in the fluor albus and in rheumatic and scorbutic disorders: it has a grateful smell, and a not unpleasant, bitterish, astringent taste: in the Edinburgh pharmacopœia, a tincture is directed to be made with spirit of scurvy-grass. The principal use of lac among us is in certain mechanic arts as a colouring drug, and for making fealing wax.

LACTUCÆ folia, semen: Lactucæ sativæ C. B. Garden lettuce; the leaves and seeds.

The feveral forts of garden lettuces are very wholesome, emollient, cooling sallad herbs, easy of digestion, and somewhat loosening

the

the belly. Most writers suppose that they have a narcotic quality; and indeed, in many cases, they contribute to procuse rest; this they effect by abating heat, and relaxing the fibres. The seeds are in the number of the sour lesser cold seeds.

There are two wild forts of lettuce, not unfrequently met with under hedges, &c. One of these differs greatly in quality from the foregoing; as may be judged from its strong soporific smell; it is called by Morison, lactuca sylvestris opii odore vehement soporifero et vireso. The upper leaves of this are jagged about the edges, the lower ones not. All the leaves of the other wild fort are very deeply jagged: hence this is by the same author distinguished by the name lactuca sylvestris laciniata.

LAMII ALBI folia, flores: Lamii albi non fatentis folio oblongo C. B. White archangel, or dead nettle: the leaves [E.] and flowers [L. E.]

This grows wild in hedges; and flowers in April and May. The flowers have been particularly celebrated in uterine fluors, and other female weaknesses, as also in disorders of the lungs; but they appear to be of very weak virtue.

LAPATHUM. Dock; the roots.

We have ten or eleven docks growing wild in England, the roots of most of which are brought to market promiscuously; though two have been generally directed by physicians in preference to the others: these are

OXYLAPATHUM: Lapathum folio acuto plane C. B. The dock with long, narrow, tharp-pointed leaves, not curled up about the edges [E.]

HYDROLAPATHUM, feveHerba Britannica: Lapathum aquaticum folio cubitali C. B. The great water dock [E:]

The feaves of the docks gently loosen the belly, and have sometimes been made ingredients in decoctions for removing a custive ha-The roots are celebrated for the cure of scorbutic and cutaneous disorders, both exhibited internally, and applied externally in ointments, cataplasms, and fomentations. Muntingius published a treatise on these plants in the year 1681, in which he endeavours to prove, that our great water dock is the berba Britannica of the ancients: and indeed the description which Dioscorides gives of the latter, does not ill agree to the former. This author therefore attributes to the hydrolapathum all the virtues ascribed of old to the Britannica, particularly recommending it in the scurvy and all its symptoms. Where this disorder is of very long standing, so as not to yield to the hydrolupathum alone, he directs a composition, by the use of which, he fays, even the venercal lues will, in a short time, be effectually cured. Six ounces of the roots of the water dock, with two of faffron; and of mace, cinnamon, gentian root, liquorice root, and black pepper, each three ounces, (or, where the pepper is improper, fix ounces of liquorice) are to be reduced into coarse powder, and put into a mixture of two gallons of wine, with half a gallon of strong vinegar, and the yolks of three eggs; and the whole digested, with a moderate warmth, for three days, in a glazed vessel, close stopt: from three to fix ounces of this liquor are to be taken every morning on an empty stomach, for fourteen or twenty days, or longer.

M 2 LAPA-

LAPATHUM UNCTUO-SUM, vide Bonus Henricus.

LAPIS BEZOAR, CALAMINARIS, HÆMATITES, LAZULI; vide Bezoar, Calaminaris, &c.

LAPPA MAJOR, vide BAR-

LAVENDULÆ flores: Lavendulæ angustifoliæ C. B. Common, or narrow leaved lavender, or spike; the flowers [L.]

LAVENDUI Æ flores: Lavendulæ latifoliæ C. B. Greater, or broad-leaved lavender; the flow-

ers [E.]

These plants have a fragrant fmell, to most people agreeable; and a warm, pungent, bitterish tafte: the broad leaved fort is the strongest in both respects, and yields in distillation thrice as much essential oil as the other: its oil is also hotter, and specifically heavier; hence in the fouthern parts of France, where both kinds grow wild, this only is made use of for the distillation of what is called oil The narrow-leaved is of spike. the fort commonly met with in our gardens, and therefore alone directed by the London college.

Lavender is a warm stimulating aromatic. It is principally recommended in vertigoes, palsies, tremors, suppression of the menstrual evacuations; and in general in all disorders of the head, nerves, and uterus, proceeding from a weakness of the folids, and lentor or sluggishness of the juices. It is sometimes also used externally in somentations for paralytic limbs. The distilled oil is particularly celebrated for destroying the pediculi inguinalis, and other cutaneous infects: if soft spongy paper, dipt

in this oil, either alone, or mixed with that of almonds, be applied at night to the parts infested by infects, they will certainly, says Geoffroy, be all found dead in the morning. The officinal preparations of lavender, are the effential oil, a spirit [L.E.] and a conserve [L.]: the flowers in substance are an ingredient in the sternutatory powder [L.] and the oil in the cephalic balsam and cephalic plaster [E.]

LAUREOLÆ folia, baccæ: Laureolæ semper virentis slore viridi, quibus dam laureolæ maris C. B. Spurgelaurel; the leaves and berries.

This is a small shrub, growing wild in some of our woods. The leaves berries, and bark, both of the stalks and roots, have an extremely acrid, hot taste, which last for along time, burning and instaming the mouth and sauces. Taken internally they operate with great violence by stool, and sometimes by vomit; so as scarce to be exhibited with any tolerable degree of safety, unless their virulence be previously abated by boiling.

LAURI folia, baccæ: Lauri vulgaris C. B. The bay tree; its léaves and berries [L. E.]

These are generally brought from the Streights, though the tree bears the colds of our own climate. They have a moderately strong aromatic smell, and a warm, bitterish, pungent tafte: the berries are thronger in both respects than the leaves, and afford in distillation a larger quantity of aromatic effential oil; they yield also an almost infipid oil to the press, in confequence of which they prove unctu-ous in the mouth. These simples are warm carminative medicines, and fometimes exhibited in this intention against flatulent colics; and likewise in hysterical disorders.

Their principal use in the present practice is in glysters, and some external applications. The leaves enter our common somentation: and the berries, the plaster and cataplasm of cummin; they also give name to an electary, which is little otherwise used than in glysters.

LAZULI lapis; [E.]-a compact ponderous fossil, of an opake blue colour, met with in the eastern countries, and in some parts of Germany. It is a strong emetic, rarely or never used in the present practice.

LENTIS VULGARIS semen: Lentis vulgaris semine subruso C. B. Lentile; the seed.

This is a strong, flatulent food, very hard of digestion: it is never, at least with us, used for any medicinal purpose.

LENTISCUS: Lentiscus verus ex insula Chio, cortice et soliis suscis Commelin. The lentisc, or mastich tree; the wood [E.]

This tree or shrub is a native of the warm climates, but bears the common winters of our own. The wood is brought to us in thick knotty pieces, covered with an ash coloured bark, and white within, of a rough, somewhat pungent taste, and an agreeable, though faint smell; the smaller tough sprigs are both in taste and smell the strongest. This wood is accounted a mild balsamic restringent; a decoction of it is, in the German ephemerides, dignified with the title of vegetable aurum potabile, and strongly recommended in catarrhs, nauseæ, and weakness of the stomach: for strengthening the tone of the viscera in general, and promoting the urinary secretion.

This is the tree, which in the island Chio affords the resin, call-

ed mastich. See the article Mas-

LEPIDII folia: Lepidii latifolii C. B. Common broad ditander, pepperwort, or poor man's pepper; the leaves [E.]

This plant is sometimes found wild by the fides of rivers, and in other moist places. The leaves have an aromatic, pungent, biting tafte, fomewhat approaching to that of pepper, but going off fooner than that of most other substances; They are very rareof this class. ly employed in medicine, though strongly recommended as antiscorbuties, and for promoting the urinary and cuticular secretions; virtues, which they have undoubtedly a good title to.

LRUCOIUM LUTEUM, vide CHEIRI.

LEVISTICI, feu Ligustici radix femen [E.] Angelicæ montanæ perennis, paludapii folio Tourn. Lovage; the root and feed [E.]

This is a large umbelliferous plant, cultivated with usin gardens. The root nearly agrees in quality with that of angelica: the principal difference is, that the lovage root has a stronger smell, and a somewhat less pungent taste, accompanied with a more durable sweetness: the seeds are eather warmer than the root. There simples, though certainly capable of being applied to useful purposes, are not at prefent regarded: neither of them is directed in extemporaneous prefcription, and the root enters no officinal composition; the seeds are an ingredient in the compound valerian water and troches of myrrh of the Edinburgh pharmacopæia, in both which they excellently coincide with the other ingredients. LICHEN:

LICHEN: Licben petræus cauliculo pileolum suffinente C. B. Liverwort; the herb.

This grows wild in moist shady places, and by the sides of rivers. It has a faint, not disagreeable smell; and an herbaceous, roughish, and somewhat bitterish taste. Great virtues have been attributed to this simple, in obstructions of the liver, jaundice, &c. which practitioners do not now expect from it.

LICHEN CINEREUS TER-RESTRIS: Lichen terrestris cinereus Raii. Ash-coloured ground liverwort [L. E.]

This confifts of pretty thick digitated leaves, flat above, of a rericular texture underneath, and faftened to the earth by small fibres: the leaves when in perfection are of an ash colour; by age they become darker coloured or reddish. It is met with on commons and open heaths, where it quickly spreads on the ground. Dr. Mead informs us, that this plant grows in all countries, and has been brought over from America along with the Peruvian bark: that it is found at all times, but ought to be gathered from autumn to winter, as being then in its freshest vigour.

This simple is said to be a warm diuretic; but the taste discovers in it little or no warmth. It is chiefly celebrated for its virtue in the cure of the disorders occasioned by the bite of a mad dog. An account of the remarkable effects in these cases of a powder composed of the dried leaves and pepper, was communicated to the Royal Society by Mr. Dampier, and published in the Philosophical Transactions, No 2 37. This powder was afterwards inferted (in the year 1721) into the London pharmacopæia, under the title of pulvis antilyffus, at the defire of an eminent physician, who

had great experience of its good effects. Some years after, the fame gentleman published and difperfed a paper containing the method of cure, which he had, in a great number of instances, constantly found successful. In this paper, the directions were to the following effect: " Let the patient be blood-" ed nine or ten ounces; and af-" terwards take a drachm and a " half of the powder every morn-" ing tasting, for four mornings fucceffively, in half a pint of " cows milk warm. After these " four doses are taken, the pa-" tient must go into the cold bath, " or a cold spring or river, every " morning fasting for a month: " he must be dipped all over, but " not stay in (with his head above " water) longer than half a minate, if the water be very cold: " after this he must go in three " times a week for a fortnight "longer." In the year 1745, the world was favoured with a new edition of the mechanical account of poisons, in which we find the same method of cure again recommended, as having, in a course of thirty years experience, never failed of fuccess, where it had been. followed before the hydrophobia begun. It is greatly to be with ed, that the efficacy of this medicine, in preventing these terrible disorders, was absolutely certain; and proved by incontestible facts. Instances have been produced of its proving unfuccefsful; and the many examples of the fatality of the disease which continually occur, feem arguments either of the inefficacy of the medicine, or a strange negligence in applying it. We shall only farther observe, that Boerhaave, who is in general sufficiently liberal in the commendation of remedies, ranks this among those infignificant trisles, which whoever

. ... . .

whoever depends upon, will find himself deceived.

LIGNUM ALOES, vide A-GALLOCHUM.

LIGNUM RHODIUM [L. E.] et ASPALATHUS [E.] Rosewood, a wood or root, brought from the Canary Islands: and afpalathus, a simple of considerable esteem among the ancients, but which has not come to the know-

ledge of later times.

The writers on botany, and the materia medica, are much divided about the lignum rhodium, not only with regard to the plant which affords it, but likewise in their accounts of the drug itself, and have described, under this name, simples manifestly different. This confusion seems to have arisen from an opinion, that the rhodium and aspalathus are the fame; whence different woods brought into Europe for the unknown aspalathus were sold again by the name of rhodium.

As to aspalathus, the ancients themselves disagree; Dioscorides requiring by this appellation the wood of a certain shrub freed from the bark, and Galen the bark of a root. At present, we have nothing under this name in What was heretofore the shops. fold among us as aspalathus, were pieces of a pale coloured wood brought from the East-Indies, and more commonly called calambac.

The lignum rhodium of the shops is usually in long crooked pieces, full of knots, which when cut, appear of a yellow colour like box, with a reddish cast: the largest, fmoothest, most compact, and deepest coloured pieces, should be chosen; and the small, thin, or pale ones, rejected. The taste of this wood is lightly bitterish, and somewhat fores: Lilii convallium albi C. B.

pungent; its smell very fragrant. resembling that of roses: long kept, it seems to lose its smell; but on cutting, or rubbing one piece against another, it smells as well as at first Distilled with water, it yields an odoriferous ettential oil, in very small quantity. Rhodium is at present in esteem only upon account of its oil, which is employed as an high and agreeable perfume in scenting pomatums, and the like. But if we may reason from analogy, this oderiserous fimple might be advantageoully applied to nobler purpoles: a tincture of it in rectified spirit of wine, which contains in a small volume the virtue of a confiderable deal of the wood, bids fair to prove a ferviceable cordial, not inferior perhaps to any thing of this kind.

LIGNUM TINCTILE CAM-PECHENSE [L. E]: Lignum Brafilio simile, carules tingens J. B. Campeachy or logwood; a wood brought from Campeachy in the. bay of Honduras.

This is usually in large logs, very compact and hard, of a red colour, and an aftringent sweet tafte. It has been for a long time used by the dyers; but not till very lately as a medicine: a decoction of it, and the extract, are in use in our hospitals, and said to have proved very serviceable in diarrhœas. The extract is now received into the

LILII ALBI radix flores: Lilii albi flore eredo et vulgaris C. B. White lily; the roots and flowers [E.]

shops. See part III. chap. vi.

This is cultivated in gardens, more for the beauty of its flowers,

than medicinal use.

LILII CONVALLIUM radix. Lily Lily of the valley, or May lily; the roots and flowers [E.] This grows wild in woods and shady

places, flowering in May.

The flowers of these plants are faid to be cephalic and nervine. They have a pleasant sweet smell, which they impart by infusion to expressed oils, and give over in distillation both to water and spirit; but no essential oil has been hitherto obtained from them. Ettmuller fays, that the distilled spirit is more fragrant than the water. The roots of the garden lily abound with a foft mucilage, and hence they have been used externally in emollient and maturating cataplasms: they are an ingredient in the suppurating cataplasm of the present Edinburgh pharmacopæia. Those of the wild lily are very bitter: dried, they are faid to prove a gentle errhine; as also are the flowers.

LIMONUM fuccus, cortex; Fractús mali limoniæ accidæ C. B. Lemons; their juice, yellow rind [L. E.] and its effential oil, called

csence of lemons [L]

The juice of lemons is fimilar in quality to that of oranges, from which it differs little otherwife than in being more acid. The yellow peel is an elegant aromatic, and is frequently employed in fromachic tinctures and infusions: it is considerably less hot than orange peel, and yields in distillation with water a less quantity of effential oil: its flavour is nevertheless more perishable, yet does not arise so readily with ipirit of wine; for a spirituous extract made from lemon peel possessies the aromatic taste and imell of the subject in much greater perfection than an extract prepared in the same manner from the speels of oranges. In the shops, a syrup is prepared from the juice, and the peel is candied; the peel is an ingredient in the bitter insusions, bitter wine, and both the peel and juice in one of the insusions of sena; the essential oil in the volatile aromatic spirit, saponaceous pills, and ointment of sulphur [L.]

LINARIÆ folia: Linariæ vulgaris luteæ florc majore C. B. Toad-

flax; the leaves [E.]

This grows wild upon banks and about the fides of fields. It is faid by some to be a powerful diuretic; whence it is named by Tragus herba urinalis; by others, to be a strong cathartic, insomuch that Brunfelsius has called it by a German name, expressing this quality, scheisskraut. Experience scarcely warrants either of these appellations; nor does common prastice take any notice of the plant.

LINGUÆ CERVINÆ. seu Scolopendrii solia: Linguæ cervoinæ officinærum C. B. Harts-tongue; the leaves [E.]

This plant confills of a number of long narrow leaves, without any stalk: it grows upon rocks and old walls, and remains green all the year. The leaves have a roughish, somewhat mucilaginous tafte, like that of the maidenhairs, but more disagreeable. They are recommended in obstructions of the viscera, and for strengthening their tone; and have sometimes been made use of for these intentions, either alone, or in conjunction with maiden-hair, or the other plants called capillary.

LINI CATHARTICI folia; Lini pratenfis flosculis exiguis C. B. Purging Purging flax, or mill-mountain; the leaves [E.]

This is a very small plant, not above four or five inches high, found wild upon chalky hills, and in dry pasture grounds. Its virtue is expressed in its title: an insusion in water or whey of a handful of the fresh leaves, or a drachm of them in substance when dried, are said to purge without inconvenience.

LINI VULGARIS femen: Lini fativi C. B. Common flax; the feed, called linfeed [L. E.]

Linseed yields to the press a confiderable quantity of oil; and boiled in water, a strong mucilage: these are occasionally made use of for the same purposes as other substances of that class: and fometimes the feeds themselves in emollient and maturating cataplasms. They have also been employed in Asia, and, in times of scarcity, in Europe, as food; but are not agreeable, or in general wholesome. Tragus relates, that those who sed on these seeds, in Zealand, had the hypochondres much diffended, and the face and other parts swelled, in a very short time; and that not a few died of these complaints. The expressed oil is an officinal preparation [L. E.]: the cake, remaining after the expression of the oil, is an ingredient in the suppurating cataplaim, and the feeds themselves in the nephritic decoction [E.]

LIQUIDAMBRA [E.] Liquidambar; a refinous juice which flows from a large tree growing in New Spain, Virginia, and other provinces of South America. This juice is at first about the confishence of turpentine, but by long keeping hardens into a refin: it is of a yellow colour inclining to red,

a warm taste, and a fragrant smell not unlike that of storax heightened with a little ambergris. It was formerly of great use as a persume, but is at present a stranger to the shops.

LITHARGYRUS [L. E.] Litharge; a preparation of lead, usually in form of fost flakes, of a yellowish reddish colour. If calcined lead be urged with a hafty fire, it melts into the appearance of oil, and on cooling concretes into litharge. Greatest part of the litharge met with in the shops, is produced in the purification of filver from lead, and the refining of gold and filver by means of this metal: according to the degree of fire and other circumstances, it proves of a pale or deep colour; the first has been commonly called litharge of filver, the other litharge of gold. See the article PLUMBUM.

LITHOSPERMI, feu Mili folis, femen: Lithospermi majoris eredî C. B. Gromwell; the seed [E.]

This is found wild in dry fields and hedges. Its feeds are roundiff, hard, of a whitish colour, like little pearls; and from these circumstances have been supposed peculiarly ferviceable in calculous disorders. Their taste is merely farinaceous.

LOTI URBANÆ fol a, semen: Loti hortensis odoræ C. B. Sweet trefoil; the leaves and seeds

The flowers of this plant are fironger in smell than the other parts: these have been recommended for diaphoretic, alexipharmac, anodyne, and other virtues: but their effects have not been sound considerable enough to continue them in practice.

LUJULÆ

LUJULÆ folia: Oxyos albæ Gerærd. Wood forrel; the leaves

[L. E.]

This is a small plant, growing wild in woods. In taste and medical qualities, it is similar to the common forrel (see the article Acetosa), but considerably more grateful, and hence is preserved by the London college. Boiled with milk, it forms an agreeable whey; and beaten with sugar, a very elegant conserve, which has been for some time kept in the shops, and is now received in the dispensatory.

LUMBRICI et LIMACES TERRESIRES. Earth-worms

and inails [E.]

Both these are supposed to cool and cleanse the viscera. The latter, from their abounding with a viscid glutinous juice, are recommended as a reflorative in consumptions: for this purpose, they are directed to be boiled in milk; and thus managed, they may poffibly be of some service. They give over nothing in distillation either with water or spirit; and hence the distilled waters of them, though formerly in great esteem, are not found to have any of the virtues which the animals themselves are supposed to possess.

LUPINI semen: Lupini vulgaris, semine et slore albo, sativi J. B.

White lupines; the feeds.

These have a leguminous taste, accompanied with a disagreeable bitter one. They are said to be anthelmintic, both internally raken, and applied externally. Caspar Hossman cautions against their internal use, and tells us (from one of the Arabian writers) that they have sometimes occasioned death. Simon Paulli also says, that he saw a boy of eight or ten years of

age, after taking a drachm of thefe feeds in powder, seized with exquisite pains of the abdomen, a difficulty of respiration, and almost total loss of voice; and that he was relieved from these complaints by a glyster of milk and fugar, which brought away a vast quantity of worms. But Mr. Geoffroy observes, very justly, that either these symptoms were owing to the worms, and not to the medicine; or that these seeds, if they have any noxious quality, lose it, with their bitterness, in boiling; fince they were commonly used among the Greeks as food, and recommended by Galen as very wholesome.

LUPULUS: Convolvulus perennis, heteroclitus, floribus berbaceis, capfulis foliaceis strobili infar Morif. Hops; the loose leasy heads which grow on the tops of the stalks.

These are one of the most agreeable of the strong bitters, though rarely employed for any medicinal purposes. Their principal consumption is in malt liquors, which they render less glutinous and dispose to pass off more freely by urine.

LYCOPERDON: Fungus sotundus orbicularis C. B. Yuff-ball, or

dufty mushroom [E.]

This fungus is found in dry pasture grounds. It seems to be nearly of the same quality with the agaric of the oak (see p. 76.) and has, like it, been employed for restraining external hæmorrhages and other fluxions. The fine dust, with which it becomes filled by age, has been applied also in the same intentions.

MACIS [L. E.] Mace; one of the coverings of the nutmeg (see

the article Nux moschata) This spice, considered as the subject both of medicine and of pharmacy, agrees nearly with the nutmeg. The principal difference is, that mace is somewhat less aftringent, yields to the press a more fluid oil, and in distillation a more volatile one: what is called in the shops expressed oil of mace, is prepared, not from this spice, but from the nutmeg. Mace is an ingredient in the officinal steel wine [L.] and aqua mirabilis [E.]; and the expressed oil in the stomachic and cephalic plasters [L.]

MAGISTRANTIA, vide IM-PERATORIA.

MAJORANÆ folia: Majoranæ vulgaris C. B. Sweet marjoram;

the leaves [L. E.]

Marjoram is raised annually in our gardens for culinary, as well as medicinal uses; the seeds are commonly procured from the fouthern parts of France, where the plant grows wild. It is a moderately warm aromatic, yielding its virtues both to aqueous and spirituous liquors by insusion, and to water in distillation. It is principally celebrated in diforders of the head and nerves, and in the humoural ashmas and catarrhs of old people. An effential oil of the herb is kept in the shops. The powder of the leaves proves an agreeable errhine, and enters the officinal sternutatory powder.

MALABATHRUM folium: Folium cinnamomi free canella Malavarica et Javanensis C. B. Indian leaf [L.]

This leaf is of a green colour, firm texture, very smooth on one fide, leafs so on the other, on

which run three remarkable ribs through its whole length. Lemery and Pomet affirm, that these leaves have no perceptible smell or taste; Herman and others, that they have a very great share of both: those met with in our shops have little or no smell till they are well rubbed, when they emit an agreeable spicy odour: on chewing, they are found extremely mucilaginous. This drug is of no farther use in medicine, than as an ingredient in the mithridate and theriaca; and is, when in its greatest perfection, much inferior to the mace, which our college direct as a succedaneum to it.

MALVÆ folia, flores: Mulvæ fylvesstris folio sinua: o C. B. Mallow; the leaves and slowers

[L. E.]

These have a somewhat mucilaginous sweetish taste. The leaves are ranked the first of the four emollient herbs: they were formerly of some esteem, in food, for loofening the belly; at present, decoctions of them are sometimes employed in dysenteries, heat and sharpness of urine, and in general for obtunding acrimonious humours: their principal use is in emollient glysters, cataplasms, and fomentations. The leaves enter the officinal decoction for glysters, and a conserve is prepared from the flowers [L.]

MALA: Fructus mali sativæ

Raii. Apples [E.]

All the forts of apples have the common quality of cooling and abating thirst: the more acid kinds loosen the belly; the austere have rather a contrary effect.

MALA SYLVESTRIA: Frudus mali fylvestris acido frudu Tourn.

Crab apples, or wildings Tourn.

These are so acid as not to be eatable : their juice, called verjuice, has sometimes supplied the place of vinegar, and has been made an ingredient in cooling and refiringent gargarisms. At present, they are scarce ever employed for any medicinal use.

MANDRAGORÆ folia: Mandragoræ fructu rotundo C. B. Man drake: the leaves.

The qualities of this plant are very doubtful; it has a strong difagreeable smell resembling that of the narcotic herbs, to which class it is usually referred. has rarely been any otherwise made use of in medicine, than as an ingredient in one of the old officinal unguents. Both that composition and the plant itself are now rejected from our pharmacopœias.

MANNA [L. E.]; the juice of certain trees of the ash kind (growing in Italy and Sicily) either naturally concreted on the plants, or exficcated and purified by art. There are several forts of manna in the shops. larger pieces, called flake manna, are usually preferred; though the fmaller grains are equally as good, provided they are white, or of a pale yellow colour, very light, of a sweet not unpleasant taile, and free from any visible impurities. Some people injudicioully prefer the fat honey-like manna to the foregoing: this has either been exposed to a moist air, or damaged by fea or other This kind of manna is water. faid to be tometimes counterfeited by a composition of sugar and honey, mixed with a little scammony: there is also a fac-

titious manna, which is white and dry, faid to be composed of sugar, manna, and some purgative ingredient, boiled to a proper confisience; this may be distinguished by its weight, solidity, untransparent whiteness, and by its tafte, which is different from that of manna.

Manna is a mild, agreeable laxative, and may be given with. safety to children and pregnant women: nevertheles, in some particular conflitutions, it acts very unkindly, producing flatulencies, and distension of the vifcera; these inconveniencies may be prevented by the addition of any grateful warm aromatic. Manna operates to weakly as not to produce the full effect of a cathartic, unless taken in large doses, and hence it is rarely given in this intention by itself. It may be commodiously dissolved in the purging mineral waters, or joined to the cathartic falts, sena, rhubarb, or the like. Geoffroy recommends acuating it with a few grains of emetic tartar; the mixture is to be divided into several doses, each containing one grain of the emetic tartar: by this management, he says, bilious serum will be plentifully evacuated, without any naufea, gripes, or other inconvenience. It is remarkable, that the efficacy of this drug is greatly promoted, (if the account of Valisnieri is to be relied on) by a fubitance which is itself very flow of operation, (See the article CASIA) Manna is an ingredient in the electary of casia  $[L, E_{\cdot}]$  and gives name to an officinal lohoch [E.]

MARGARITÆ [L. E.] Pearls; small concretions of a transparent whiteness, found on the inside of the shell of the concha margaritifera or mother-of-pearl fish, as also of certain oysters, muscles, and other shell fishes. The pearls most esteemed are brought from the East and West-Indies, and distinguished by the names of oriental and occidental: the oriental, which are valued most, have a more shining filver hue than the occidental; these last are somewhat milky: a fort inferior to both these is sometimes met with in our own feas, particularly on the coast of Scotland. The coarse, rough pearls. and the very small ones which are unfit for other uses, are those generally employed in medicine. They have been greatly celebrated as cordial, alexipharmac, and comforting the nerves: but the only virtue that can be reasonably expected from them is, that of abforbing acidities in the prime viæ. in which intention they enter three of the officinal powders. comparative strength, with regard to the other absorbents, may be feen among the tables at the beginning of this work: see p. 63.

MARRUBII folia: Murrubii albi vulgaris C. B. White horehound: the leaves [L. E.]

These bave a very strong, not disagreeable smell, and a roughish very bitter talle. Besides the virtues which they possess in common with other firong bitters, they are supposed to be peculiarly serviceable in humoural afthmas and coughs, the vellow jaundice proceeding from a viscidity of the bile, and other chronical disorders. They are doubtless an useful aperient and deobstruent, promote the fluid secretions in general, and liberally taken. loosen the belly. They are an ingredient only in the theriaca [L.]

MARI SYRIACI folia: Mari cortusi J. B. Chamædryos maritimæ

incanæ frutescentis foliis lanceolatis Tourn. Syrian herb mastich; the leaves [L. E.]

This is a finall furubby plant, growing spontaneously in Syria. Candy, and other warm climates, and cultivated with us in gardens. The leaves have an aromatic bitterish taste; and, when rubbed betwixt the fingers, a quick pungent fmell, which foon affects the head. and occasions sneezing: distilled with water, they yield a very acrid. penetrating essential oil, resembling one obtained by the same means from scurvygrass. These qualities fufficiently point out the uses so which this plant might be applied; at present, it is little otherwise employed than in cephalic fnuffs. It is an ingredient in the pulvis sternetatorius of the London pharmacopœia.

MARI VULGARIS folia: Sampfuci seve mari mastichen redolemis C. B. Thymbræ Hispanicæ majoranæ folio Tourn. Herb mastich; the leaves [L. E.]

This pungent aromatic place also is become almost a stranger to practice.

MASTICHE [L. E.] Mastich: a refin exuding from the lentife tree (see Lentiscus) and brought from Chio, in small, yellowish, transparent grains or tears, of an agreeable smell, especially when heated or fet on fire. This refin is recommended in old coughs, dyfenteries, hæmoptoes, weakness of the stomach, and in general in all de- . bilities and laxity of the fibres. Geoffroy directs an aqueous decoction of it to be used for these purpofes: but water extracts little or nothing from this refin; rectified spirit almost entirely dissolves it: the folution taftes very warm and pungent.

MATRI-

MATRICARIÆ folia, flores:
Matricariæ vulgaris seu sativæ C. B.
Common wild feathersew or seversew; the leaves [L. E.] and slow-

ers [E.]

This plant is a celebrated antihysteric. Simon Paulli relates, that he has experienced most happy effects from it in obstructions of the uterine evacuations. I have often seen, says he, from the use of a decoction of matricaria and chamomile flowers with a little mugwort, hysteric complaints instantly relieved, the discharge succeed plentifully, and the patient from a lethargic state, return as it were into life again. Matricaria is likewife recommended in fundry other disorders, as a warm stimulating bitter: all that bitters and carminatives can do, fays Geoffroy, may be expected from this. It is undoubtedly a medicine of fome use in these cases, though not perhaps equal to chamomile flowers alone, with which the matricaria agrees in sensible qualities, except in being weaker.

MECHOACANNÆ radix [E.]; the root of an American convolvulus, brought chiefly from Mechoacan, a province of Mexico, in thin flices like jalap, but larger, and of a whitish colour. It was first introduced among us (about the year 1524) as a purgative universally fafe, and capable of evacuating all morbific humours from the most remote parts of the body. as jalap became known, Mechoacan gradually loft its reputation, which it has never fince been able to retrieve. It is nevertheless, by some still deemed an useful cathartic: it has very little fmell or tafte, and is not apt to offend the stomach: its operation is flow, but effectual and safe. Geoffroy affirms, that there is scarce any purgative

accompanied with fewer inconveniencies. It feems to differ from jalap only in being weaker; the refins obtained from both have nearly the same qualities, but jalap yields five or fix times as much as Mechoacan; hence it is found necessary to exhibit the latter in fix times the dose of the former, to produce the same effects.

MEL[L. E.] Honey.—Honey is a vegetable juice, obtained from the honeycomb, either by feparating the combs, and laying them flat upon a fieve, through which the honey spontaneously percolates; or by including the comb in canvas bags, and forcing the honey out by a press: the first fort is the purest; the latter is found to contain a good deal of the matter of which the comb is formed, and fundry other impurities: there is another fort fill inferior to the two foregoing, obtained by heating the combs before they are put into the press. The best fort is thick, of a whitish colour, an agreeable smell. and a very pleasant taite: both the colour and flavour differ according to the plants which the bees collect it from: that of Narbonne in France, where rolemary abounds, is faid to have a very manifest flavour of that plant, and to be imitable by adding to other honey an infusion of rolemany flowers. Honey, confidered as a medicine, is a very useful detergent and aperient, powerfully dissolving viscid juices, and promoting the expectoration of tough phlegm: in some particular constitutions it has an inconvenience of griping or proving purgative; this is said to be in some measure prevented, by previoully boiling the honey.

MELAMPODIUM, vide Hel-LEBORUS NIGER.

MELI-

MELILOTI folia flores: Trifolii adorati feu meliloti vulganis J. B. Melilot; the leaves and flowers [E.]

This grows wild in hedges and among corn; and has likewise, for medicinal ofes, been cultivated in gardens. The green herb has no remarkable fmell: when dry, a pretty strong one: the taste is roughish, bitter, and if long chewed. nauseous. A decoction of this herb has been recommended in inflammations of the abdomen: and a decoction of the flowers in the fluor albus. But modern practice rarely employs it any otherwise than in emollient and carminative glysters, and in fomentations, cataplasms, and the like; and in these not often. It formerly gave name to one of the officinal plasters. which received from the melilot a green colour, but no particular virtue.

MELISSÆ folia: Melissæ bortensis C. B. Balm; the leaves [L. E.]

This plant, when in perfection, has a pleasant smell, somewhat of the lemon kind; and a weak rough-The young ish aromatic taste. shoots have the strongest slavour; the flowers, the herb itself, when old, or produced in very moist rich foils, or rainy feasons, are much weaker both in smell and taste. Balm is appropriated, by the writers on the Materia Medica, to the head, stomach, and uterus; and in all disorders of these parts, is supposed to do extraordinary service. So high an opinion have some of the chemists entertained of balm. that they have expected to find in it a medicine which should prolong life beyond the usual period. The present practice, however, holds it in no great esteem, and ranks it (where it certainly deserves to be) among the weaker corroborants: in distillation, it yields an elegant ef-

fential oil, but in exceeding small quantity; the remaining decoction tastes roughish. Strong insusions of the herb, drank as tea, and continued for some time, have done fervice in a weak lax state of the viscera: these liquors, lightly acidulated with juice of lemons, turn of a fine reddish colour, and prove an useful, and to many a very grateful drink, in dry parching severs. A simple water of the plant is directed in the Edinburgh pharmacopæia as an officinal.

MELONUM femen. Melons; the feeds. These stand among the four greater cold seeds. They have been sometimes used, with the others of that class, as cooling and emollient: but are at present little taken notice of.

MENTHA CATARIA, vide Nepeta.

MENTHÆ VULGARIS folia: Menthæ angustifoliæ spicatæ C. B. Garden or spearmint; the leaves [L. E.]

The leaves of mipt have a warin. roughish, somewhat bitterish taste. and a strong, not unpleasant, arematic smell. Their virtues are those of a warm stomachic and carminative: in loss of appetite, naufez, continual reachings to vomit, and (as Boerhaave expresses it) almost paralytic weaknesses of the flomach, there are few fimples perhaps of equal efficacy. In colicky pains, the gripes to which children are subject, lienteries, and other kinds of immoderate fluxes, this plant frequently does good fervice. It likewise proves beneficial in sundry hysteric cases, and affords an useful cordial in languors and other weaknesses consequent upon delivery. The best preparations for these purposes are, a strong infufion made from the dry leaves in water (which is much superior to one from the green herb) or rather a tincture or extract prepared with These possess the rectified spirit. whole virtues of the mint: the effential oil and distilled water contain only the aromatic part; the expressed juice only the astringency and bitterishness, together with the mucilaginous substance common to The effential oil, all vegetables. a simple and spirituous water, and a conserve, are kept in the shops: the Edinburgh college directs an infusion of the leaves in the distilled water. This herb is an ingredient also in the three alexitereal waters; and its effential oil in the flomach plaster [L.] and stomachic pills [E.]

MENTASTRI folia: Mentastri spicati folio longiore candicante J. B. Horse mint; the leaves. and several other forts of mint are found wild in moist meadows. marshes, and on the brinks of rivers. They are much less agreeable in smell than spearmint, and have more of a hot unpleasant bitterness.

MENTHÆ PIPERTIDIS folia: Menthæ spicis brevioribus et babitioribus, foliis menthæ fuscæ, sapore servido piperis Raii Sy-Peppermint; the leaves nupf. [L. E.]

This species has been lately introduced into practice, and represent pharmacopoia; very few of the botanical or medical writers make mention of it: it grows wild in some parts of England, in moist watery places, but is much less common than the other forts. The leaves have a more penetrating smell than any of the other mints, and a much warmer, pungent, glowing taste like pepper, finking as it were into the tongue. The principal use of this herb is in flatulent colics, languors, and other like diforders : it feems to act as foon as taken, and extend its effects through the whole fystem. instantly communicating a glow-Water extracts the ing warmth. whole of the pungency of this herb by infusion, and elevates it in distillation. Its officinal preparations are an effential oil, and a simple and spirituous water [L.] The Edinburgh college employ it also in the aqua mirabilis and elixir of vitriol.

MERCURIALIS maris et seminæ folia: Mercurialis testiculatæ sive maris, et spicata se famina Diofcoridis et Plinii C. B. Male and female French mercury; the leaves

These stand among the five emollient herbs; and in this intention are sometimes made use of in glysters. A syrup made from the leaves, given in the dose of two ounces, is faid to prove a mild and useful laxative.

There is another fort of mercurialis growing in woods and hedges, which though recommended by fome botanic writers, as having the same virtues with the foregoing, and as more palatable, has been lately found possessed of noxious qualities. (See Raii Synops. edit. 3. page 138. Phil. Trans. abr. Lowthorp. ii. 640.) This may ceived for the first time in our be distinguished from the foregoing, by its being a perennial plant, larger, having its leaves rough, and the stalk not at all branched. The officinal fort is named by Linnæus mercurialis caule brachiato, foliis glabris; the poisonous mercurialis caule simplicissimo, foliis scabris; it is commonly called dogs mercury.

MER-

TUM VIVUM.

MESPILA: Frudus mespili vulgaris J. B. The medlar-tree; its truit.

Mediars are scarce ever made use of for any medicinal purposes. They have a very austere astringent taste, insomuch as not to be eatable until mellowed by keeping.

MEIATHAMANTICI radix: Mei foliis anethi C. B. Spignel; the root [L. E.]

Spignel is an umbellifetous plant. found wild in Italy, and the warmer parts of Europe, and sometimes also in England. The roots have a pleasant aromatic smell, and a warm, pungent, bitterish taste: in virtue, they are fimilar to the levisticum, from which this root feems to differ only in being weaker, and somewhat more agreeable. It is an useful aromatic and carminative, though at present little regarded.

MEZEREI radix, cortex, bacca: Laureolæ folio deciduo, flore purpureo, officinis laureolæ fæminæ C. B. Mezereon, or spurge-olive; the root, bark, and berries.

The bark and berries are strong purgatives, similar to the laureola, or spurge laurel. The root is sometimes used in diet drinks.

MILII semen: Milii semine luteo C. B. Millet; the seed.

These seeds are frequently employed in food, but hardly ever as medicines: they are sufficiently nutritious, and not difficult of digestion.

MILIUM SOLIS, vide Li-THOSPERMUM.

MILLEFOLII folia: Millefolii vulgaris albi, et Millefolii purpurei

MERCURIUS, vide Argen - C. B. Milfoil, or yatrow; the leaves [E.]

> This grows plentifully about the fides of fields, and on dry commons, flowering greatest part of the summer. The leaves have a rough bitterish taste, and a faint aromatic smell. Their virtues are those of a very mild astringent, and as such they stand recommended in hæmorrhages both internal and external, diarrhoeas, debility and laxity of the fibres; and likewise in spasmodic hysterical affections. In these cases, some of the Germans have a very high opinion of this herb, particularly Stahl, who esteemed it a very effectual aftringent, and, in his language, one of the most certain tonics and fedatives. Its virtues are extracted in great perfection by proof spirit: water takes up its aftringency and bitterness, but little of its aromatic flavour; tinctures made in rectified spirit contain both, though rather weaker than those in proof spirit.

The flowers of milfoil are confiderably stronger in aromatic flavour than the leaves; in distillation they yield a small quantity of essential oil, of an elegant blue colour.

The roots, taken up in the fpring, have an agreeable, warm, pungent tafte. Dr. Grew resembles them to contrayerva, and imagines they might, in some measure, supply its place; this, however, is greatly to be doubted, fince there is such a remarkable difference betwixt the two, that whilst one retains its taste for a length of time after it has been brought to us from America, the tafte of the other is in great meafare lost by drying.

-MILLEPEDÆ [L. B.] Woodlice, hoglice, flaters. Thefe

These insects are found in cellars, under stones, and in cold moift places; in the warmer countries they are rarely met with. Millepedes have a faint disagreeable fmell, and a somewhat pungent, sweetish, nauseous taste. They have been highly celebrated in suppressions of urine, in all kinds of obstructions of the bowels, in the jaundice, weakness of fight, and a variety of other diforders. Whether they have any just title to these virtues, is greatly to be doubted: thus much is certain, that their real effects come far short of the character usually given of them. Their officinal preparations are, the millepedes dried and powdered [L. E.] and an infusion of the live insect in wine [E]: they are an ingredient also in the icteric decoction of the Edinburgh pharmacopæia.

MINIUM [L. E.] Red lead; lead calcined to redness. See the article PLUMBUM.

MORSUS DIABOLI, seu Succise radix, folia: Scabiosa pratensis nostratis præmorsa radice Morison. Devils bit; the leaves and roots. These stand recommended as alexipharmacs, but they have long given place to medicines of greater efficacy.

MORI fructus, et cortex radicis: Mori frudu nigro C. B. The mulberry tree; its fruit [L. E.] and the bark of the roots [E.]

This tree is commonly cultivated on account of its fruit, which is rather eaten for pleasure than used as a medicine: it has the common qualities of the other sweet fruits, abating heat, quenching thirst, and promoting the groffer secretions; an agreeable fyrup, made from the juice, is kept theless strongly impregnated with

in the shops. The bark of the roots has been in considerable esteem as a vermifuge; its taste is bitter, and fomewhat aftringent.

MOSCHUS [L. E.] Mulk.

Musk is a grumous substance like clotted blood, found in a little bag fituated near the umbilical region of a particular kind of animal met with in China, Tartary, and the East-Indies: the best musk is brought from Tonquin, an inferior fort from Agria and Bengal, and a still worse from Russia.

Fine musk comes to us in round. thin bladders; which are generally about the fize of a pigeon's egg, covered with short brown hairs, well filled, and without any appearance of having been opened. The musk itself is dry, with a kind of unctuofity, of a dark reddish brown, or rusty blackish colour, in small round grains, with very few hard black clots, and perfectly free from any fandy or other visible foreign matter. If chewed, and rubbed with a knife on paper, it looks smooth, bright, yellowish, and free from grittiness. Laid on a red-hot iron, it catches flame, and burns almost entirely away, leaving only an exceeding small quantity of light greyish ashes; if any earthy substances have been mixed with the musk, the quantity of the residuum will readily discover them.

Musk has a bitterish subacrid taste: a fragrant smell, agreeable at a distance, but when smelt near to, so strong as to be disagreeable, unless weakened by the admixture of other substances. If a small quantity be infused in spirit of wine in the cold for a few days, it imparts a deep, but not red tincture: this, though it discovers no great smell of the musk, is neverits virtues; a fingle drop of it communicates to a whole quart of wine a rich musky flavour. The degree of flavour which a tincture drawn from a known quantity of musk communicates to vinous liquors, is perhaps one of the best criteria for judging of the goodness of this commodity. Neumann informs us, that spirit of wine dissolves ten parts out of thirty of musk, and that water takes up twelve; that water elevates its smell in distillation, whilst pure spirit brings over nothing.

Musk is a medicine of great esteem in the eastern countries: among us, it has been for fome time pretty much out of use, even as a perfume, on a supposition of its occasioning vapours, &c. in weak females, and persons of a fedentary life. It appears, however, from late experience, to be, when properly managed, a remedy of good fervice even against those disorders which it has been supposed to produce. Dr. Wall has communicated (in the Philosophical Transactions, No. 474) an account of some extraordinary effeets of musk in convulsive and other diteases, which have too often baffled the force of medicine. The doctor observes, that the fmell of perfumes is often of differvice, where the substance taken inwardly, and in confiderable quantity, produces the happiest effects; that two persons, labouring under a subsultus tendinum, extreme anxiety, and want of fleep. from the bite of a mad dog, by taking two doles of mulk, each of which was fixteen grains, were perfectly relieved from their complaints. He likewise observes, that convuluve hiccups, attended with the worst symptoms, were removed by a dose or two, of ten grains: and that in some cases, where this medicine could not, on account of strong convulsions, be administered to the patient by the mouth, it proved of service when injected as a glyster. He likewise adds, that under the quantity of fix grains he never found much effect from it; but that taken to ten grains, and upwards, it never fails to produce a mild diaphoresis, without at all heating or giving any uneafiness; that on the contrary, it eases pain, raises the spirits, and that after the fweat breaks out, the patient usually falls into a refreshing fleep; that he never met with any hysterical person, how averse soever to perfumes, but could take it, in the form of a bolus, without inconvenience. To this paper is annexed an account of some farther extraordinary effects of musk. observed by another gentleman. Repeated experience has fince confirmed its efficacy in these disorders. I have myself frequently given it with remarkable fuccess; and sometimes increased the dose as far as twenty grains every four hours, with two or three spoonfuls of the musk julep between. The julep is the only officinal preparation of it.

MYROBALANI. Myrobalans; dried fruits brought from the East-Indies; their outward part, freed from the stone.

Five kinds of myrobalans were formerly directed as officinals; (1) The yellow, myrobalani teretes citrini C. B. (2) The chebule, myrobalani maximæ oblongæ angulofæ C. B. (3) The Indian or black, myrobalani nigræ ocangularis C. B. (4) The belliric, morobalani rotundæ belliricæ C. B. (5) The emblic, myrobalani emblicæ in fegmentis nucleum habentes, angulofæ J. B.

All the myrobalans have a low degree of purgative virtue. They

have also an astringent quality, discoverable by the taste, from their use among the Indians for tanning leather, and from their striking a black colour with chalybeate folutions: in consequence of this, they are supposed to strengthen the bowels after their operation as a cathartic is over. Nevertheless their purgative virtue is fo inconsiderable, that practitioners have for a long time laid them intirely aside in that intention; and the college of Edinburgh, as well as that of London, have now rejected them from the catalogue of officinal fimples.

MYRRHA [L. E.] Myrrh.

Myrrh is a concrete gummy-refinous juice, brought from the East-Indies, in glebes or drops, of various colours and magnitudes. The best fort is of a brown or reddish yellow colour, fomewhat transparent; of a lightly pungent, bitter tafte, with an aromatic flavour, though not sufficient to prevent its proving nauseous to the palate; and a strong not disagree-able smell. The medical effects of this aromatic bitter are, to warm and strengthen the viscera, and disfolve thick tenacious juices: it frequently occasions a mild diaphorefis, and promotes the fluid secretions in general.

Hence it proves serviceable, in languid cases, diseases arising from a simple inactivity, those semale disorders which proceed from a cold, mucous, sluggish indisposition of the humours, suppressions of the uterine discharges, cachectic disorders, and where the lungs and thorax are oppressed by viscid phlegm. Myrrh is likewise supposed in a peculiar manner to resist putresaction in all parts of the body; and in this light slands recommended in malignant, putrid,

and peffilential fevers, and in the fmall-pox, in which last it is said to accelerate the eruption.

Rectified spirit extracts the fine aromatic flavour and bitterness of this drug, and does not elevate any thing of either in evaporation: the gummy substance left by this menstruum has a disagreeable tafte. with scarce any thing of the peculiar flavour of the myrrh: this part dissolves in water, except some im-In diftillapurities which remain. tion with water, a confiderable quantity of a ponderous essential oil arises, resembling in flavour the original drug. Myrrh is the bafis of an officinal tincture [L. E.] and gives name to a compound tincture E.]elixir, powder[L.] and troches [E.] It is an ingredient in the aloetic wine or elixir proprietatis, the gum pills, Rufus's pills [L E.] stomachic pills [E.] mithridate, theriaca [L.] and theriaca Edinenfis [E.]

MYRRHIDIS folia, femen: Myrrhidis magno femine, longo, fulcato J. B. Sweet cicely; the leaves and feeds.

This plant is cultivated in gardens: it agrees in quality with the charefolium; see p. 123.

MYRTI baccæ: Myrti communis Italicæ C. B. Myrtle; the berries [E.]

This is an evergreen shrub, growing in Italy, and cultivated in our botanic gardens. The leaves and berries have been sometimes made use of as astringents, but are not at present regarded.

NAPI semen: Napi dulcis officinarum: Napi sative C.B. Sweet navew or navew gentle; the seeds [L.]

This is a fort of turnep, fown in fome of our gardens for culinary use: the roots are warmer than the

common turnep. The feeds have a bitterish tasse, accompanied with a faint aromatic slavour: abundance of virtues have been ascribed to them, as attenuating, detergent, alexipharmac, and others; at present, they are of no farther use in medicine, than as an ingredient in the theriaca.

NAPI SYLVESTRIS femen: Napi filvestrii C. B. Rape; the feeds.

This has little other external difference from the foregoing than being smaller: it grows wild upon dry banks and among corn. The feeds of this are warmer and more pungent than those of the garden fort: the only use, however, they are applied to, is the preparation of the oil called rape oil, which is obtained by bruising and pressing the seeds: large quantities of the plant are cultivated for this purpose in the isle of Ely.

NARDUS CELTICA: Radix nurdi Celticæ Dioscoridis C. B. Valerianæ Celticæ Tourn. Celtic nard [L. E.] the root, brought from the Alps, &c.

This root confifts of a number of fibres, with the lower part of the stalks adhering; these last are covered with thin yellowish scales, the remains of the withered leaves.

NARDUS INDICA [L. E]: Nardus Indica, que spica, spicu, nardi, et spica Indica officinarum C. B. Indian nard, or spikenard, brought from the East-Indies.

This is a congeries of small sibres issuing from one head, and matted close together, so as to form a bunch about the size of the singer, with some small strings at the opposite end of the head. The matted sibres (which are the part chosen for medicinal purposes) are supposed by some to be the head or

fpike of the plant, by others the root: they feem rather to be the remains of the withered stalks, or the ribs of the leaves: sometimes entire leaves and pieces of stalks are found among them: we likewise now and then meet with a number of these bunches issuing from one root.

Both the nards have a warm, pungent, bitterish taste; and a strong, not very agreeable smell. They are stomachic and carminative; and said to be alexipharmac, diuretic, and emmenagogue; their only use at present is as ingredients in the mithridate and theriaca.

NASTURTII AQUATICI folia: Nasturtii aquatici supini C. B. Water-cresses: the leaves [L. E.]

This plant grows wild in rivulets, and the clearer standing waters: its leaves remain green all the year, but are in greatest perfection in the spring. They have a quick pungent smell (when rubbed betwixt the fingers) and an acrid tafle, fimilar to that of cochlearia, but weaker. As to their virtues, they are among the milder aperient antiscorbutics. Hoffman has a mighty opinion of this plant, and recommends it as of fingular efficacy for accelerating the circulation, strengthening the viscera, opening obstructions of the glands, promoting the fluid fecretions, and purifying the blood and humours; for these purposes, the expressed juice, which contains the peculiar tafte and pungency of the herb, may be taken in doses of an ounce or two, and continued for a confiderable time. The juice is an ingredient in the fucci scorbutici of the shops.

NASTURTII HORTENSIS folia, femen: Nasturtii vulgariis feu hortensis tenuiter divisi Morison. N 3 GarGarden creffes; the leaves and

feeds [E.]

The leaves of garden creffes make an useful sallad in scorbutic habits: in taste and medical virtues, they are similar to the foregoing, but much weaker. The seeds also are considerably more pungent than the leaves.

NEPETÆ folia: Menthæ catariæ vulgaris et majoris C. B. Nep, or catmint; the leaves [L E.]

This plant is commonly cultivated in our gardens, and is sometimes also sound growing wild in hedges and ondry banks: It is a moderately aromatic plant, of a strong smell, not ill resembling a mixture of mint and pennyroyal; of the virtues of which it likewise participates.

NEPHRITICUM LIGNUM: Lignum peregrinum, aquam caruleam reddens C. B. Nephritic wood.

This is an American wood, brought to us in large, compact, ponderous pieces, without knots, of a whitish or pale yellow colour on the outfide, and dark coloured or reddish within: the bark is usual-This wood imparts to ly rejected. water or rectified spirit a deep tincture, appearing, when placed betwixt the eye and the light, of a golden colour, in other fituations blue; pieces of another wood are fometimes mixed with it, which give only a yellow colour to water. The nephritic wood has scarce any fmell, and very little tafte. It stands recommended in difficulty of urine, nephritic complaints, and all disorders of the kidneys and urinary passages; and is said to have this peculiaradvantage, that it does not, like the warmer diuretics, heat or offend the parts. Practitioners. however, have not found these virtues warranted by experience.

NICOTIAN Æ folia: Nicotianæ

latifoliæ majoris C. B. Tobacco; the leaves [L. E.]

This plant was first brought into Europe, about the year 1560, from the island Tobago in America; and is now cultivated, for medicinal use, in our gardens: the leaves are about two feet long, of a pale green colour whilst fresh, and when carefully dried of a lively yellowish. They have a strong, disagreeable smell, like that of the narcotic plants; and a very acrid burning tafte. Taken internally, they prove virulently cathartic and emetic, occasioning almost intolerable cardialgic anxieties. By boiling in water, their virulence is abated, and at length destroyed: an extract made by long coction is recommended by Stahl and other German physicians, as a fafe and most effectual aperient, expectorant, detergent, &c. but this medicine, which is extremely precarious and uncertain in strength, has never come into esteem among us. Tobacco is fometimes used externally in unguents, for destroying cutaneous infects, cleanfing old ulcers, &c. Beaten into a mash with vinegar or brandy, it has sometimes proved ferviceable for removing hard tumours of the hypochondres; an account is given in the Edinburgh essays of two cases of this kind cured by it.

There is another fort of tobacco found wild on dunghills, in several parts of England: this is calld by C. Bauhine nicotiana minor, by Gerard hypscyamus luteus. It seems to agree in quality with the hypscyamus formerly mentioned, though (as Dale informs us) often substituted in our markets for the true tobacco: from which it may be distinguished by the leaves being much smaller, and the flowers not reddish as those of the officinal fort, but of a yellowish green colour.

NIGELLÆ

NIGELLÆ femen: Nigellæ flore minore fimplici candido C. B. Fennelflower; the feeds.

This plant is fown annually in fome of our gardens; the feeds most esteemed are brought from Italy. They have a strong, not unpleasant smell: and a subacrid, somewhat unctuous disagreeable taste. They stand recommended as aperient, diuretic, &c. but have long been strangers to practice, and are by some suspected to have noxious qualities.

NITRUM [L. E.] Nitre, or faltpetre; a falt extracted, in Perfia and the East-Indies, from certain earths that lie on the sides of hills; and artificially produced in some parts of Europe, from animal and vegetable matters rotted together (with the addition of lime and ashes) and exposed for a length of time to the air, without the access of which, nitre is never generated: the salt extracted from the earths, &c. by means of water, is purified by colature and crystallization.

Pure nitre dissolves in about fix times its weight of water, and concretes again into colourless transparent crystals; their figure is that of an hexagonal prism, terminated by a pyramid of an equal number of fides. It readily melts in the fire; and in contact with fuel deflagrates, with a bright flame and confiderable noise; after the detonation is over, a large quantity of alkaline falt is found remaining. The taste of nitre is sharp, penetrating, and bitterish, accompanied with a certain fenfation of coldness.

Nitre is a medicine of celebrated use in many disorders. Besides the aperient quality of neutral salts in general, it has a manifestly cooling one, by which it quenches thirst. and abates febrile heats and commotions of the blood: it has one great advantage above the refrigerating medicines of the acid kind, that it does not coagulate the animal juices; blood, which is coagulated by all the mineral acids, and milk, &c. by acids of every kind, are by nitre rendered more dilute, and preserved from coagulation: it nevertheless somewhat thickens the thin, ferous, acrimonious humours, and occasions an uniform mixture of them with fuch as are more thick and viscid; by this means preventing the ill consequences which would otherwise ensue from the former, though it has not, as Juncker supposes, any property of really obtunding acrimony. This medicine for the most part promotes urine; sometimes gently loosens the belly; but in cold phlegmatic habits, very rarely has this effect, though given in large doses: alvine fluxes, proceeding from too great acrimony of the bile or inflammation of the intestines, are suppressed by it: in choleric and febrile disorders, it generally excites fweat; but in malignant cases, where the pulse is low, and the strength lost, it retards this falutary evacuation and the eruption of the exanthemata.

Dr. Stahl has written an express treatise upon the medical virtues of nitre; in which he informs us, from his own experience, that this salt added to gargarisms employed in inflammations of the fauces in acute severs, thickens the salval moisture upon the palate and sauces into the consistence of a mucus, which keeps them moist for a considerable time; whereas, if nitre is not added, a sudden dryness of the mouth immediately ensues: that in nephritic complaints, the pru-

den

dent use of nitre is of more service than any of the numerous medicines usually recommended in that disease: that nitre gives great relief in suppression and heat of urine, whether fimple or occafioned by a venereal taint; that it is of great service in acute and inflammatory pains of the head, eyes, ears, teeth, &c. in all eryfipelatous affections, whether particular or universal, and likewise in chronic deliriums; that in diarrhϾ happening in petechial fevers, nitre, mixed with absorbents and diaphoretics, had the best effects, always putting a ftop to the flux, or rendering the evacuation salutary; that in diarrhoeæ happening in the small-pox, it had been employed with the like fuccess, two doses, or three at most (confisting of two, three or four grains each, according to the age, &c. of the patient) given at the interval of two or three hours, putting a stop to the flux, after the bezoardic powders, both with and without opium, had been given without success. The same author recommends this falt likewife as a medicine of fingular fervice in choleras attended with great anxieties and heat of the blood; in the flatulent spasmodic heartburns familiar to hypochondriacal people; and the loss of appetite, nausea, vomiting, &c. which gouty persons are sometimes seized with upon the pains of the feet, &c. fuddenly remitting. In cases of this last kind, the use of nitre furely requires great caution, although the author affures us, that no bad consequences are to be feared from it. Nevertheless he observes, that in a phthisis and ulcerous affections, it has been found to be of no service; and that therefore its use may be superseded in these complaints,

Indeed in disorders of the lungs in general it is commonly reckoned to be rather hurtful than beneficial.

The usual dose of this medicine among us is from two or three grains to a scruple; though it may be given with great safety, and generally to better advantage, in larger quantities: the only inconvenience is its being apt to fit uneasy on the stomach. have affirmed, that this falt loses half its weight of aqueous moifture by fusion, and consequently that one part of melted nitre is equivalent to two of the crystals; but it did not appear, upon several careful trials, to lose so much as one-twentieth of its weight. officinal preparations of nitre are a decoction or folution in water [E.] and troches [L.] A corrofive acid spirit is also extracted from it; see Part II. chap. viii. fect. 3. It is employed likewise in operations on metallic bodies. for promoting their calcination, or burning out their inflammable matter.

NUMMULARIÆ folia: Lyfimachiæ humifusæ, folio rotundiore, flore luteo Tourn. Moneywort, or herb two-pence; the leaves.

This grows spontaneously in moist watery places, and creeps on the ground, with two little roundish leaves at each joint. Their taste is subastringent, and very lightly acid: hence they stand recommended by Boerhaave in the hot scurvy, and in uterine and other hæmorrhages. But their effects are so inconsiderable, that common practice takes no notice of them.

NUX MOSCHATA [L. E. Nux moschata frudu rotundo C. B.]
Nutmegs; the kernel of a round

ich nut which grows in the East-Indies. The outlide covering of this fruit is fost and fleshy, like that of a walnut, and spontaneously opens when the nut grows ripe; immediately under this lies the mace, (see the article MACIS) which forms a kind of reticular covering; through the fiffures whereof appears a hard woody shell that includes the nutmeg. These kernels have long been made use of both for medicinal and culinary purposes, and deservedly looked upon as a warm agrecable aromatic. They are supposed likewise to have an astringent virtue; and are employed in that intention in diarthosas and dysenteries. astringency is said to be increased by torrefaction, but this does not appear to the taste: this treatment certainly deprives the spice of some of its finer oil, and therefore renders it less efficacious to any good purpose; and if we may reason from analogy, probably abates of its aftringency. Nutmegs, distilled with water, afford a large quantity of essential oil, resembling in flavour the spice itself; after the distillation, an infipid sebaceous matter is found fwimming on the water: the decoction, inspissated, gives an extract of an unctuous, very lightly bitterish taste, and with little or no aftringency. Rectified spirit extracts the whole virtue of nutmegs by infusion, and elevates very little of it in distillation: hence the spirituous extract possesses the flavour of the spice in an eminent degree.

Nutmegs yield to the press (heated) a considerable quantity of limpid yellow oil, which in cooling concretes into a sebaceous consistence. In the shops we meet with three sorts of unctuous sub-

stances, called oil of mace, though really expressed from the nutmeg. The best is brought from the East-Indies, in stone jars; this is of a thick confistence, of the colour of mace, and an agreeable fragrant smell; the second fort, which is paler coloured and much inferior in quality, comes from Holland in solid masses, generally flat and of a square figure: the third, which is the world of all, and usually called common oil of mace, is an artificial composition of serum, palm oil and the like, flavoured with a little genuine oil of the nutmeg. These oils yield all that part in which their aromatic flavour resides, in distillation to water, and to pure spirit by infusion: the distilled liquor and spirituous tincture nearly resemble in quality those prepared immediately from the nutmeg, The officinal preparations of nutmegs are, a spirituous water, esfential oil, and the nutmegs in substance roasted [L.] The nutmeg itself is used in the compound horseradish water, compound spirit of lavender, cordial confection, cardialgic troches, and fyrup of buckthorn [L.]; its essential oil, in the volatile aromatic spirit [L.] and the expressed oil in mithridate and theriaca, stomachic and cephalic plasters [L.] and . cephalic balfam [E.]

NUX PISTACHIA: Nucleus e fructu Pistaciæ Raii. Pistachio [E]

This is a moderately large nut, containing a kernel of a pale greenish colour, covered with a reddish skin. The tree which produces it, grows spontaneously in Persia, Arabia, and several islands of the Archipelago: it bears likewise the colds of our own climate, so as to have produced fruit

not inferior to that which we received from abroad: Pistachio nuts have a pleasant, sweet, unctuous taste, resembling that of almonds. They are ranked amongst the analeptics, and are by some much esteemed in certain weaknesses, and in emaciated habits.

NYMPHÆÆ ALBÆ radix, flores: Nymphæ albæ majoris C. B. White water lily; the root and

flowers [E.]

This grows in rivers and large lakes, flowering usually in June. The roots and flowers have a rough, bitterish glutinous taste; (the flowers are the least rough) and when fresh, a disagreeable fmell, which is in great measure loft by drying: they are recommended in alvine fluxes, gleets, and the like, The roots are supposed by some to be in an eminent degree narcotic, but on no very good foundation. Lindestolpe informs us, that in some parts of Sweden, they were in times of scarcity used as food, and did not prove unwholesome.

OCHRA. Yellow ochre: a foft friable ore of iron, of a yellow colour, dug in several parts of England. It possesses the virtues of the calces of iron and hæmatites; but in so low a degree, that the shops have deservedly rejected it; its principal use is as a pigment.

OCIMI folia: Ocimi vulgatioris C. B. Basil; the leaves [E.] This is a small plant, raised annually in our gardens; it slowers in June and July, and produces its seeds in August, but rarely perfects them in this country. The leaves have a soft, somewhat warm taste; and when rubbed, a strong unpleasant smell, which, by moderate drying, becomes more agreeable. They are faid to attenuate viscid phlegm, promote expectoration, and the uterine secretions; but have not, for a long time, been regarded in practice.

OLIVÆ earumque cleum: Fructus cleæ fativæ C. B. The clive tree; the fruit [E.] and its oil [L. E.]

This tree grows in the fouthern parts of France, in Spain, Italy, and other warm countries: with us it is usually preserved in the green houses of the curious, though it will bear our ordinary winters in the open air, and produce very good fruit. Olives have an acrid. bitter, extremely disagreeable taste: pickled (as we receive them from abroad) they prove less disagree-able; the Lucca olives, which are smaller than the others, have the weakest taste; the Spanish, or larger, the Arongest; the Provence, which are of a middling fize, are generally the most esteemed.

The oil obtained from this fruit has no particular tafte or smell, and does not greatly differ in quality from oil of almonds. Authors make mention of two forts of this oil, one expressed from the olives when fully ripe, which is our common oil olive; the other, before it has grown ripe; this is called oleum immaturum, and omphacinum. Nothing is met with in the shops under this name; and Lemery affirms, that there is no fuch-oil; unripe olives, yielding only a viscid juice to the press. From the ripe fruit, two or three forts are obtained, differing in degree of purity: the pureft runs by light pressure: the remaining magma, heated and preffed more strongly, yields an inferior fort, with fome dregs at the bottom,

called amurca. All these oils contain a confiderable portion of aqueous moisture, and a mucilaginous substance; which subject them to run into a putrid state: to prevent this, the preparers add fome fea falt, which imbibing the aqueous and mucilaginous parts, finks with them to the bottom; by this means, the oil becomes more homogene, and consequently less susceptible of alteration. In its passage to us, some of the falt, thrown up from the bottom by the shaking of the vessel, is fometimes mixed with and detained in the oil, which, in our colder climate, becomes too thick to suffer it freely to subfide; and hence the oil is fometimes met with of a manifestly faline taste Oil olive is used in the simple balfam of fulphur, Locatelli's balsam, and several ointments. is oftner employed in this last intention than the other expressed oils, but more rarely for internal medicinal purposes.

OLIBANUM [L. E.] a gummy-refin, brought from Turkey and the East-Indies, usually in drops or tears, like those of mattich, but larger, of a pale yellowish, and sometimes reddish colour; a moderately warm pungent tafte, and a strong, not very agreeable smell. This drug has received many different appellations, according to its different appearances: the fingle tears are called simply olibanum, or thus: when two are joined together, they have been called thus masculum, and when two were very large, thus famininum: fometimes four or five about the bigness of filberds, are found adhering to a piece of the bark of the tree which they exuded from; these have been named thus corticolum; the finer powder which rubs off from the tears in the carriage, mica thuris; and the coarfer powder, manna thuris. This drug is not however, in any of its states, what is now called thus or frankincense in the shops (see the article 1 Hus.)

Olibanum consists of about equal parts of a gummy and refinous substance, the first soluble in water, the other in rectified spirit. With regard to its virtues, abundance have been attributed to it. particularly in diforders of the head and breast, in hæmoptoes, and in alvine and uterine fluxes: but its real effects, in these cases. are far from answering the promises of the recommenders. verius is said to have had large. experience of the good effects of this drug in plurifies, especially epidemic ones: he directs a scooped apple to be filled with a drachm of olibanum, then covered and roasted under the ashes; this is to be taken for a dose, three ounces of carduns water drank after it. and the patient covered up warm in bed: in a short time, he says, either a plentiful sweat, or a gentle diarrhœa enfue, which carry off the disease. Geoffroy informs us, that he has frequently made use of this medicine, after venæsection, with good success; but acknowledges that it has fometimes failed. Olibanum is an ingredient in the pulvis e succino. theriaca [L.] confectio japonica, pilulæ ex olibano, and emplastrum defensivum [E.]

ONONIDIS, Anonidis five Areftæ bovis radix: Anonidis spinofæ flore purpureo C. B. Restharrow, cammock, or petty-whin; the root [E.]

This plant grows wild in waste grounds, and dry fields. The root has a disagreeable smell, and

a nan-

a nauseous sweetish taste: it stands recommended as an aperient and diuretic; but has never been much regarded among us.

OPHIOGLOSSI folium: Ophioglossi vulgati C. B. Adders

tongue; the leaf.

This plant has only one leaf, with a slender stalk arising from the bottom of it, dented about the edges, and supposed to resemble the tongue of a serpent: it grows wild in moist meadows. any other virtues are attributed to it than those of a vulnerary.

OPIUM [L. E.] Opium; the concrete milky juice of the poppy

(fee PAPAVER.)

This juice has not yet been collected in quantity in Europe. Egypt, Persia, and some other provinces of Asia, have hitherto fupplied us with this commodity: in those countries, large quantities of poppies are cultivated for this The opium prepared about Thebes in Egypt, hence named Thebaic opium, has been usually esteemed the best; but this is not now distinguished from that col-This juice lected in other places. is brought to us in cakes or loaves. covered with leaves and other vegetable matters, to prevent their flicking together: it is of a folid consistence, yet somewhat softish and tenacious, of a dark reddish brown colour in the mass, and when reduced into powder, yellow: of a faint disagreeable smell, and a bitterish taste, accompanied with a pungent heat and acrimony.

The general effects of this medicine are, to relax the folids, and render them less sensible of irritation, to chear the spirits, ease pain, procure sleep, promote perspiration and sweat, but restrain all other

evacuations. When its operation is over, the pain, and other symptoms which it had for a time abated, return; and generally with greater violence than before, unless the cause has been removed by the diaphorefis, or relaxation which it occasioned.

The operation of opium is generally attended with a flow, but strong and full pulse, a dryness of the mouth, a redness and light itching of the skin: and followed by adegree of nausea, a difficulty of respiration, lowness of the spirits,

and a weak languid.pulfe.

The principal indications of opium are, great watchfulness, immoderate evacuations proceeding from acrimony and irritation. cramps or spasmodic contractions of the nerves, and violent pains of almost every kind. In these cases, opiates procure, at least, a temporary relief, and an opportunity for other medicines, properly interposed, to take effect.

Opium sometimes defeats the intention of the physician, and instead of producing rest, occasions vomiting, &c. great anxiety, Taken on a full stomach, it often proves emetic; where the patient is exhausted by excessive evacuations, it occasions generally great lowness. It has been observed to operate more powerfully in perfons of a lax habit, than in the opposite circumstances; whilst it usefully restrains preternatural discharges proceeding from irritation, it proves injurious in those that arise from a contrary cause, as in the colliquative diarrhoea attending hectic fevers. By relaxing, taking off strictures, and occasioning a paralysis of particular parts, it often promotes such evacuation as those parts are concerned in. Boerhaave observes, that it sometimes enables the ureters to allow an easy passage

even to the calculus: but this effect is by no means constant.

With regard to the dose of opium, one grain is generally a fufficient, and often too large a one; maniacal persons, and those who have been long accustomed to take it, require three or more grains to have the due effect. Among the eastern nations, who are habituated to opium, a drachm is but a moderate dose: Garcias relates. that he knew one who every day took ten drachms. Those who have been long accustomed to its use. upon leaving it off, are seized with great lowness, langour, and anxiety; which are relieved by having again recourse to opium, and, in some measure, by wine or

spirituous liquors.

Opium is partially foluble in water, and rectified spirit: proof spirit, wine, and vinegar, totally dissolve it; the impurities only being left. The folutions, in proof fpirit and wine, have the same effects with the juice in substance; with this difference, that they exert themselves sooner in the body, and are less apt to leave a nausea on the stomach. A tincture made in rectified spirit is supposed to operate, in an equal dose, more powerfully than the foregoing liquors: Geoffroy informs us, from his own experience, that whilst the watery and vinous folutions occasioned pleasant quiet sleep, a tincture drawn with pure spirit brought on a phrenfy for a time. Alkaline salts diminish the soporific virtue of this medicine: fixt alkalies render it diuretic, whilst volatile ones determine its action chiefly to the cutaneous pores. Acids almost entirely destroy its Many have endeavoured power. to correct fome imaginary ill qualities of this drug, by toailing it, by fermentation, by long con-

tinued digestions, by repeated diffolutions and distillations. These processes, though recommended by many late writers, do not promise any singular advantage: they may indeed weaken the opium; but by this very means become prejudicial, rendering the medicine more uncertain in its operation, and the dose more undetermined.

Opium applied externally, gives ease in sundry pains, but does not, as some have supposed, stupify the part, or render it insensible of pain: used immoderately, it is said to produce the same ill effects, as when taken to excess internally.

The officinal preparations of opium are, the thebaic extract, or stained opium, and a vinous [L.] and spirituous [E.] tincture. It is a capital ingredient in fundry compositions, as the paregoric elixir [L. E.] sudorific tincture [E.] faponaceous and storax pills [L.] olibanum and pacific pills [E.] the compound powder of bole, scordium and amber, electary of fcordium, confectio Paulina; philonium, mithridate, theriaca [L.] theriaca Edinensis, confectio japonica, and anodyne balfam [E.]

OPOBALSAMUM [L. E.] Bal-Samum Judaicum, Syriacum, e Mecha. Opobaliam, or balm of Gilead: a refinous juice, obtained from an evergreen tree, or shrub growing spontaneously in Arabia. The best fort, which naturally exudes from the plant, is scarce known to Europe; and the inferior kind, faid to be extracted by lightly boiling the leaves and branches in water. are very rarely feen among us. The true opobalsam, according to Alpinus, is at first turbid and white, of a very strong pungent smell, like that of turpentine, but much fweeter.

sweeter, and of a bitter, acrid, astringent taste: upon being kept for some time, it becomes thin, limpid, light, of a greenish hue; then of a gold yellow; and at length of the colour of honey: after this it grows thick like tur-pentine, and loses much of its fragrance. This balfam is of great esteem in the eastern countries, both as a medicine, and as an odoriferous unguent, and cosmetic. Its great scarcity has prevented its coming into use among us: in the mithridate and theriaca, which it is directed as an ingredient in, the London college allows the expresfed oil of nutmegs as a succedaneum to it.

OPOPANAX [L. E.] Opopanax; a concrete gummy refinous juice, obtained from the roots of an umbelliferous plant, panax pastinacæ folio C. B. which grows spontaneously in the warmer countries, and bears the colds of this. The juice is brought from Turkey and the East-Indies, sometimes in round drops or tears, but more commonly in irregular lumps, of a reddish yellow colour on the outfide, with specks of white, inwardly of a paler colour, and frequently variegated with large white pieces. It has a peculiar strong fmell, and a bitter, acrid, somewhat nauseous taste. Its virtues are those of an attenuating and aperient medicine. Boerhaave frequently employed it, along with ammoniacum and galbanum, in hypocondriacal disorders, obstructions of the abdonimal viscera, and suppressions of the menstrual evacuations from a fluggishness of mucous humours, and a want of due elasticity of the folids: in these intentions it is an useful. ingredient in the pilula gummofa and compound powder of myrrh

of the London pharmacopæia; but is not employed in any composition of the Edinburgh. It may be given by itself in the dose of a scruple, or half a drachm: a whole drachm proves, in many constitutions, gently purgative.

ORCHIS, vide SATYRION.

ORIGANI folia: Origani folvestris, cunilæ bubulæ Plinii C. B. Wild marjoram; the leaves [L.E.]

This is met with upon dry chalky hills, and in gravelly foils, in feveral parts of England. It has an agreeable smell, and a pungent taste, warmer than that of the garden marjorum, and much resembling thyme, which it seems to agree with in virtue. An essential oil, distilled from it, is kept in the shops.

There is another fort of origanum, called Creticum, whose flowers or rather flowery tops, are sometimes brought to us from candy: these have an agreeable aromatic flavour, somewhat stronger than

the common fort.

OROBI femen: Orobi filiquis articulatis femine majore C. B. Bitter vetch; the feeds.

This plant is cultivated, though not very often, in our gardens. The feeds have a farinaceous bitterish disagreeable taste: they stand recommended in nephritic complaints, but have long been strangers to practice.

ORYZÉ semen [E.] Rice; the seeds, freed from the outward skin; these are brought chiefly from Carolina, where the plant is cultivated in large quantities. They are sufficiently nutritious, and afford an useful food in diarrhoese, dysenteries, and other disorders from a thin acrimonous state of the juices.

OSTE-

OSTEOCOLLA [E.]

This is a fossil substance, found in many parts of Germany, as also in England, and other coun-It is generally met with in loose sandy grounds, spreading, from near the surface to a considerable depth, into a number of branches, like the roots of a tree: it has a whitish colour, rough on the furface, and for the most part either hollow within, or filled with folid wood, or a powdery woody matter. Sometimes the roots of living trees are found changed into this kind of substance (See Neumann's chemical works, page 11, and the Berlin Memoirs for the year 1748.)

Powdered ofteocolla feparates, on ablution with water, into two diffinet lubstances; the finer matter washed over, burns into quick-lime, and agrees on all trials with powdered limestone: the grosser part which remains is mere fand: the fand and calcareous earth are for the most part nearly in an equal proportion. From this analysis we may easily judge of the virtue which this fossil is celebrated for, that of bringing on a callus in

fractured bones.

OXALIS, vide ACETOSA.

OXYACANTHA GALENI, vide Berberis.

OXYACANTHA VULGA-RIS, vide Spina alba.

OXYLAPATHUM, vide La-

PÆONIÆ radix, flores, semen; Pæoniæ solio nigricante splendido, quæ mas C. B. wel Pæoniæ saminæ store pleno rubro majore C. B. Male and semale peony; the roots, slowers, and seeds [L. E.]

These plants are cultivated in our gardens on account of the beauty of their flowers; the female, which is the largest and most elegant, and for this reason the most common, is the only one which the shops are supplied with. In quality they are scarce sensibly different; and hence the college allow them to be taken promifcuously. The roots and feeds of peony have, when recent, an unpleasant scent, approaching to that of the narcotic plants; and a somewhat glutinous subacrid taste. with a light degree of bitterness and aftringency: the leaves also discover an astringent quality both to the tafte, and by changing chalybeate folutions of a purple colour: the flowers have little tafte. and a very faint, not agreeable fmell. The parts which have chiefly been used for medicinal purposes. are the roots and feeds. Thefe are looked upon as emollient, corroborant, and lightly anodyne; and supposed to be of service in some kinds of obstructions, erosions of the viscera, heat of urine, pains in the kidneys, and the like. virtue they are chiefly celebrated for, is that of curing spasmodic and epileptic complaints; which many have been abfurd enough to believe that the root of this plant would do, by being only worn about the neck. The root is an ingredient in the pulvis ad epileptices of the Edinburgh pharmacopæia.

PALMÆ oleum: Palmæ foliorum pediculis spinosis, frustu prumiformi, luteo, oleoso Sloan. Palmoil [E.]

This oil is obtained from the kernels of the fruit of a species of palm tree, which is a native of the coast of Guinea and Cape Verd islands: from these places it has

been

been transplanted into Jamaica and Barbadoes. The oil, as brought to us, is about the confiftence of an ointment, and of an orange colour; a strong, not disagreeable smell, but very little taste: by long keeping, it loses its high colour, and becomes white, when it ought to be rejected, as no longer fit for use. The inhabitants of the Guinea coast are said to make this oil part of their food, and to employ it for the same puposes as we do butter. With us, it is rarely given inwardly, and used only in some external applications, for pains and weakness of the nerves, cramps, fprains, and the like. The common people apply it to the cure of chilblains, and when early made use of, not without fuccess. It is an ingredient in the emollient ointment and stomach plaster of the Edinburgh pharmacopϒa.

PANICI semen: Panici Germanici, siwe panicula minore C. B.

Panic; the feeds.

This plant is cultivated in some parts of Germany; the seeds have been made use of in sood, but are not regarded as medicines.

PAPAVERIS ALBI capita: Papaveris hortensis semine albo C B. The large garden poppy, with white flowers and seeds; or the white poppy; its heads [L.]

PAPAVER NIGRUM: Papaver hortense nigro semine C. B. The lefter garden poppy, with purple flowers and black seeds; or the black poppy. The college of Edinburgh seems to allow this species to be used promiscuously with the soregoing; having dropt the distinction of white and black, and retained in the catalogue only the title of papaver bortense; of

which they direct the heads, feeds, and leaves, for medicinal use.

The heads and stalks of these plants contain a milky juice; which may be collected in confiderable quantity, by lightly wounding them when almost ripe : this juice, exposed for a few days to the air, thickens into a stiff tenacious mass, agreeing in quality with the opium brought from abroad. (See the article Opium.) The juices of both the poppies appear to be fimilar to one another; the only difference is in the quantity afforded, which is generally in proportion to the fize of the plants: the larger, or white poppy, is the fort cultivated by the preparers of opium in the eastern countries, and for medicinal uses in this.

Poppy heads, boiled in water, impart to the menstruum their narcotic juice, together with the other juices, which they have in common with vegetable matters in general. The liquor, strongly pressed out, suffered to settle, clarified with whites of eggs, and evaporated to a due confistence, yields about one-fifth, or one-fixth the weight of the heads, of extract. This possesses the virtues of opium; but requires to be given in double its dose to answer the fame intention, which it is faid to perform without occasioning a nausea and giddiness, the usual consequences of the other. the Edinburgh esfays abridg, vol. i. pag. 132 and 158.) A strong decoction of the heads, mixed with as much fugar as is fufficient to reduce it into the consistence of a fyrup, becomes fit for keeping in a liquid form; and is the only officinal preparation of the poppy. Both these preparations are very useful ones, though liable to variation in point of strength: nor

does

does this inconvenience feem avoid able by any care in the prescriber, or the operator; fince the poppy heads themselves (according to the degree of maturity, and the soil and season of which they are the produce) contain different proportions of the narcotic matter to the other juices of the plant; as has been observed in the Pharmacopæia Reformata.

The feeds of the poppy are by many reckoned suporific: Juncker fays, they have the same quality with those of hyoscyamus, and Herman looks upon them as a good substitute to opium; misled probably by an observation which holds in many plants, that the feeds are more efficacious than the vessels in which they are contained.

The feeds of the poppy have nothing of the narcotic juice which is lodged in their covering, and in the stalks; an oil expressed from them has been used for the same purposes as oil olive; and the feeds themselves taken as food: their tafte is sweetish and farinaceous.

PAPAVERIS ERRATICI, feu Papaveris rbæados flores: Papaveris erratici majoris C. B. Red poppy or corn role: the greater of the hairy wild poppies, with deep red flowers and dark coloured feeds; its flowers [L. E.]

The flowers of this plant yield upon expression a deep red juice, and impart the same colour by infusion to aqueous liquors. A fyrup of them is kept in the shops: this is valued chiefly for its colour; though fome expect from it a lightly anodyne virtue.

PARALYSIS flores: Verbasculi pratenfis odorati C. B. Primulæ veris majoris Raii. Cowslips: the Mowers [L. E.]

This plant grows wild in marshes

and moist meadows. The flowers appear in April; they have a pleafant sweet smell, and a subacrid, bitterish, somewhat aftringent talte. An infusion of them, used as tea. is recommended as a mild corroborant, in nervous complaints, and in some semale disorders proceeding from a deficiency of the menfirual purgations. A strong infusion of them forms, with proper quantity of fugar, an agreeable fyrup, which has long maintained a place in the shops: by boiling, even for a little time; their fine flavour is destroyed.

PAREIRA BRAVA [E.]

This is the root of an American convolvulus, brought to us from Brazil, in pieces of different fizes, some no bigger than one's finger, others as large as a child's arm: it is crooked, and variously wrinkled on the furface: outwardly of a dark colour, internally of a dull yellowish, and interwoven with woody fibres; fo that, upon a transverse section, a number of concentric circles appear, crossed with fibres, which run from the center to the circumference : it has no smell : the tafte is a little bitterish, blended with a sweetness, like that of liquorice. This root is highly extolled by the Brazilians and Portuguese, in a great variety of diseases, particularly against suppressions of urine, nephritic pains, and the calculus. In the two first, Geoffroy fays he has given it with good success, and that the patient was almost instantly relieved by it, a copious difcharge of urine fuce eding. He likewise observed large quantities of gravel, and even im il flones. voided after its use: this effect he attributes not to any lithontriptic power, but to its disfolving the viscid mucus, by which the sabu-

lobi

lous matter had been detained. He likewise relates, that he has had frequent experience of the good effects of this root in deterging and healing ulcers of the kidneys and bladder, where the urine came away purulent and mucous, and could not be voided at all without extreme pain: by the use of the pareira, the urine foon became clear, and of a due confistence, and was evacuated freely; and by joining to this medicine balsam of Copaiba, the ulcer perfectly healed. The attenuating quality, which he had discovered in this root, induced him to make trial of it in other diseases, proceeding from tenacious juices, and in these likewise it fully answered his expectations: in humoral afthmas, where the lungs were stuffed up, and the patient almost suffocated by thick phlegm, an infusion of pareira, after many other medicines had proved ineffectual, occasioned a plentiful expectoration. and foon completed a cure: in the jaundice, proceeding from thick bile, it did excellent fervice; but in another icterical case, where the liver was swelled and hard, this medicine did no good. His dose of the root in substance is from twelve grains to half a drachin, in decoction two or three drachms.

PARIETARIÆ, seu Helxines i solia: Parietariæ officinarum C. B. Pellitory of the wall; the leaves

[L. E.]

This is a small plant, growing upon old walls; of an herbaceous, subsaline taste, without any smell. It is one of the five emollient herbs, and in this intention is occasionally made use of It is an ingredient in the nephritic decoction of the Edinburgh pharmacopæia. The expressed juice has been given in the dose of three ounces as a diuretic.

PARTHENIUM, vide MA-TRICARIA.

PASTINACA HORTENSIS: Paftinacu latifolia fativa Raii. Garden parsneps.

PASTINACA SILVESTRIS: Puftinaca filvestris latifolia Raij.

Wild parineps.

The roots of the garden parfnep are used as food, and prove sufficiently nutritious. The seeds of both sorts are lightly aromatic; those of the wild are strongest.

PENTAPHYLLI radix: Quinquefolii majoris repentis C. B. Cinquefoil; the root [L.]

plentifully This grows in hedges, and by road-fides. root is moderately aftringent; and as fuch is fometimes given internally against diarrhœas, and other fluxes; and employed in gargarisms for strengthening the gums, &c. The cortical part of the root may be taken, in substance, to the quantity of a drachm: the internal part is confiderably weaker, and requires to be given in double the dose to produce the same effect. It is scarcely otherwise made use of than as an ingredient in the theriaca.

PEPONUM femen: Peponis oblongi C. B. The pumpion; its feeds [E.]

These seeds are very rarely met with in the shops: in quality they are not different from those of cucumbers, melons, and the others called cold seeds.

PERICLYMENUM, vide Ca-

PERSICARIÆ MITIS folia: Perficariæ maculofæ Raii. Spotted arlmart: the leaves.

This grows wild in moist watery places: the leaves somewhat resemble

iembie

femble those of the perfica malus, and have generally a blackish spot in the middle: their taste is roughish and subsaline. This herb is recommended chiefly for external purposes: Tournesort assumed in the Memoirs of the French academy, 1703) that it is one of the best vulneraries and antiseptics he knows, and that a decoction of it in wine stops gangrenes in a surprising manner. The present practice, however, has no dependence on it.

PERSICARIÆ URENTIS folia: Perficariæ vulgaris acris, five hydropiperis Raii. Biting arfmart, lakeweed, or water pepper; the leaves [E.]

This fort is readily distinguishable from the former, by its pungent, biting, pepper-like tafte. Its virtues are those of an acrid stimu-Jating medicine: in phlegmatic habits it promotes the urinary discharge, and has frequently done good service in scorbutic complaints. The fresh leaves are sometimes applied externally for cleanfing old fiftulous ulcers, and confuming funguous flesh: for these purposes they are said to be employed by the arriers, among whom they have been principally made use of.

PERSICÆ MALI flores: Perficæ molli carne, &c. C. B. The peach-tree; its flowers [E.]

Peach flowers have an agreeable fmell, and a bitterish taste; distilled, without any addition, by the heat of a water-bath, they yield one fixth their weight, or more, of a whitish liquor, which, as Mr. Bulduc observes, communicates, to a large quantity of other liquids, a flavour like that of the kernels of fruits. An infusion in water of half an ounce of the fresh gather-

ed flowers, or a drachm of them when dried, sweetened with sugar, proves for children an useful laxative and anthelmintic: the leaves of the tree are, in this intention, somewhat more efficacious, though less agreeable. The fruit has the same quality with the other sweet fruits, that of abating heat, quenching thirs, and gently loosening the belly.

PERUVIANUS CORTEX [L. E.] Peruvian bark; the bark of a tall flender tree, growing in Peru. It is brought to us in pieces of different fizes, sometimes rolled up into short thick quills, and fometimes flat: the outside is brownish, and generally covered in part with a whitish moss; the inside is of a yellowish, feddish, or rufly iron colour. It has a lightly aromatic finell, fomewhat musty, yet not disagreeable; a bitterish, astringent taste, which dwells long upon the tongue, accompanied with a degree of aromatic warmth. The small, thin, flat pieces are by fome accounted the best: by others. the quill fort, with the roughest coat, especially if of a bright cinnamon colour on the infide though the large flat pieces, whether rough or smooth, of lighter or darker colour, are often of equal goodness. The best bark is that which is strongest in smell and taste: this likewise proves friable betwixt the teeth, and does not separate into fibres; it breaks, not shivery, but close and smooth.

The virtues of this bark, as a febrifuge, were discovered by the Indians about the year 1500: Europe did not become acquainted with it till 1649: nor was it received into general practice till feveral years after this; some ill confequences, ensuing from its imprudent use, having brought it, for a

time,

Οá

time, into difrepute. At present, it is looked upon as the most effectual remedy in intermittent fevers of almost every kind, and safe in all ages and constitutions; provided it be judiciously and seasonably administered, and due regard be had to the circumstances of the disease. The modern practice, previous to the use of this medicine, usually gives an emetic at the beginning of a paroxylm: in fome cases a cathartic, and in plethoric habits venæsection, are premised: these render the bark not only more safe, but likewise more certain and speedy in its operation: where these evacuations are neglected, or not sufficiently plentiful. the disease, if of long standing, scarce yields to the cortex; or if it appears at length subdued, yet the patient does not recover his firength, and foon fuffers a relapfe. The use of the bark is begun at the end of a paroxysm, and repeated, in the quantity of half a drachm (more or less, according to the circumstances of the patient) every third or fourth hour during the intermission: where the fever is of the bilious kind, and accompanied with great heat, a little nitre is joined: in all cases, moderate exercile generally promotes its effect. At first, it usually loosens the belly, and sometimes operates as if a cathartic had been taken; and by this means supplies the omission of evacuations before its exhibition: if the purging continues, the medicine does not answer the purposes intended by it; in such case, a little opium is added, which effectually suppresses the flux: if after this the patient continues too coftive, recourse is had to glysters. The loofeness, however, ought not to be stopt too foon: on the contrary, where the bark does not itself produce this effect, it is neces-

fary, as Dr. Mead informs us, to join to it a little rhubarb, so as to occasion for a time two stools a day; by this means the disease is .more effectually cured, and less subject to be followed by a dropsy, or ill habit of body: after a drachm or two of rhubarb have been taken. it is to be discontinued, and the bark exhibited by itself. the fever has been removed, the medicine is continued for some time longer, to prevent a relapse: and evacuations, unless absolutely necessary, abstained from. disease is nevertheless seldom completely cared before fome very confiderable evacuation, either by ftool, urine, or perspiration, ensues: if this does not fucceed spontaneously, cathartics, diuretics, or diaphoretics, are given in conjunction with the bark, otherwise the patient continues weak, and without appetite. till either the disease returns, or changes into one of a different kind.

In fymptomatic agues, hectic and purulent fevers, cacochymic habits, and where the hypochondres are swelled and distended, this medicine is improper, and for the most part prejudicial. Its manifest astringency forbids its use in obstructions of the obdominal viscera, or suppression of any critical evacuation; until the obstruction is suffer removed, or the evacuation had its due course.

In acute, inflammatory, or malignant fevers, the bark does not feem to have any good effect. Nevertheless, in the decline of long nervous fevers, or after a remission, when from bad habit, old age, fatigue, or the like, the patient is extremely weak, and the pulse low, the cortex proves a medicine of excellent service; provided that there is no extravassation, that the vessels remain entire, and pus is not already formed.

Peru. .

Peruvian bark has likewise been found eminently serviceable in gangrenes and mortifications, proceeding either from an internal or external cause. In all the cases of this kind, where it proved successful, it occasioned a kind suppuration, which degenerated when the use of the medicine was discontinued. and again turned kindly upon refuming it. Some have been hence induced to try the cortex in variolous cases, where either the pustules did not rightly suppurate, or petechiæ shewed a disposition to a gangrene; and here likewise it answered expectation: the empty vesicles filled with matter, watery fanies changed into thick white pus, the petechiæ become gradually of a pale colour, and at length disappeared, and the pox began to turn fooner than was expected. See the Edinburgh medical esays.

The bark has been applied likewife, and not without success, to the cure of periodic head-achs, hysteric and hypochondriac fits, and other disorders, which have regular intermissions. By its astringency and aromatic quality, it firengthens the whole nervous system, and proves useful in weakness of the stomach, and fundry chronical disorders. proceeding from too great laxity of the fibres. In obslinate uterine fluxes, and old gleets, bark joined with chalybeats has notable effects.

The virtues of Peruvian bark reside chiesty in a resinous subflance, and hence are extracted in perfection by rectified spirit." By strong coction in water, the refin is melted out, and mingled with the water, which, whilst hot, appears transparent, but in cooling, grows turbid, and deposits great part of the resin to the bottom. Water elevates in distillation the

aromatic part of the bark; pure spirit brings over nothing. Hence an aqueous extract proves not only less in quantity, but likewise inferior in quality to one made with rectified spirit. Proof spirits extract the virtues of this drug, in tolerable perfection, in the cold; heat enables it to take up more than it can retain when cold. Spirit of sal ammoniac, prepared with fixt alkaline falts, gains very little from the cortex, either with or without heat: the spirit prepared with quick-lime, and the dulcified spirit, in a few hours become strongly impregnated with its smell and tafte.

The officinal preparations of bark are, an extract [L. E.] refin [E] spirituous tincture [L, E]tincture in volatile spirit [L.] and compound tincture [E.] it is an ingredient also in the stomachic

tincture [E]

The substances usually joined with bark in prescription seem calculated either to promoté its efficacy, or merely for reducing it into the intended form, without much regard to its agreeableness, and the conveniency of taking it: this is nevertheless a point of great consequence, as its taste, and the quantity which is necessary, make the patient too frequently louth it, before enough has been taken to produce the defired effect. If defigned to be given in the folid form of a bolus, electary, &c. it should be made up, not, as is customary, with fyrups, but with mucilages: with the former, it flicks about the mouth and fauces. whence its take remains for a confiderable time; with the latter, it passes freely, scarce leaving any taste in the mouth. Aromatics do not prevent the tafte of the bark from discovering itself; extract of liquorice very effectually conceals

O 3

it. The extract of logwood alfo, joined to that of bark, and a proper quantity of mucilage, form a very elegant and agreeable composition.

PETASITIDIS radix: Petafitidis majoris et vulgaris C. B. But-

terbur; the root [E.]

This grows wild by the sides of ditches and in meadows: it fends forth short scaly stalks in the spring, bearing spikes of purplish flowers; after this the leaves appear, which are very large, and hollowed in about the middle, so as to resemble a bonnet, or what the Greeks called wireson, whence the name of the plant. The roots have a strong fmell, and a bitterish, aromatic, not very agreeable taile: they have been given in the dole of a drachm or more, as an aromatic, and likewise as an operient and deobstruent; these virtues, however, they possess in so low a degree, as to have lost their reputation in the shops.

PETROLEUM [E.] Rock oil. This is a general name for fundry liquid bitumens, or mineral oils, which spontaneously exude from the earth, or from clests of rocks. These oils are found in almost all countries, but in greatest quantities in the warmer ones: some are met with in different parts of England; and many of our common bituminous minerals, as pit-coal, &c. afford, on distillation, oils not greatly different from them.

The finest fort of this commodity comes from the duchy of Modena in Italy, where three different kinds are found; the best is almost as clear, sluid, and transparent as water, of a highly penetrating, yet not disagreeable smell, somewhat like that of rectified oil of amber; the second fort is of a

clear yellow colour, not fo fluid as the former, less penetrating, and partaking more of the oil of amber imell: the third, or worft, is of a blackish red colour, of a thicker confistence, and more disagreeable than the two foregoing. The first of these is very rarely met with in the shops; the second. mixed with a little of the third, and some subtile oil, is usually sent us instead of it. troleum readily catches fire, and, if pure, burns entirely away: diftilled, it becomes somewhat more pellucid than before (a small quantity of yellowish matter remaining) and lofes greatly of its natural fmell: it unites with the essential oils of vegetables, not at all with vinous spirits: the finer forts are fo light as to swim upon the most highly rectified spirit of wine.

Petroleum is at present very rarely employed as a medicine, though if the finer kinds could be procured genuine, they should seem to deferve fome notice: they are more agreeable than the oil of amber, and milder than that of turpentine; the virtues of both which they partici-They are principally repate of. commended by authors for external purposes, against pains and aches, in paralytic complaints, and for preventing chilblains. For these intentions, some of the more common mineral oils have been made use of with good success; an oil extracted from a kind of stone coal has been cried up among the common people, under the name of British oil, forrheumatic pains, &c. even this is often counterfeited by a small portion of oil of amber added to the common expressed oils.

PETROLEUM BARBADENSE [L.] Barbadoes tar.

This is thicker than the foregoing petrolea, and nearly of the confift-

ence

ence of common tar. It is of a reddish black colour, adisagreeable fmell, less pungent than the other forts. This bitumen is found in feveral of our American islands, where it is esteemed by the inhabitants of great service as a sudorific, and in disorders of the breast and lungs, though in cases of this kind, attended with inflammation, it is certainly improper; they likewise apply it externally as a discutient, and for preventing paralytic difor-Among us it is rarely used, and not often to be met with ge-The London colleged emnuine. ploys it as a menstruum for sulphur in the balsamum sulphuris Barbadense, and directs an oil to be diftilled from it: that of Edinburgh has not yet received it.

PETROSELINI MACEDO-NICI femen: Apii Macedonici C B. Macedonian parsley; the seeds [L.]

PETROSELINI VULGARIS femen, folia, radix: Apii hortenfis seu petroselini vulgo C. B. common parsley; the roots, leaves, [E.] and seeds [L. E.]

The first of these plants is sometimes met with in our gardens; the second is commonly cultivated The feeds for culinary purposes. of both have an aromatic flavour, and are occasionally made use of as carminatives, &c. Those of the Macedonian parsley are the strongest, though generally supplied by the other. The root of parsley is one of the five aperient roots, and in this intention is fometimes made an ingredient in apozems and dietdrinks; if liberally used, it is apt to occasion flatulencies, and thus, by distending the viscera, produces a contrary effect to that intended by it; the taste of this root is somewhat sweetish, with a light degree of warmth and aromatic flavour.

The feeds of the Macedonian parfley are an ingredient in mithridate and theriaca; and those of the common in the electary of bayberries [L.]

PEUCEDANI radix: Peucedani Germanici C. B. Hogs fenuel, or sulphurwort; the root.

This plant grows wild by the fea shores, and in moist shady places. The roots have a strong disagreeable smell, somewhat resembling that of sulphureous solutions; and an unctuous, subacrid, bitterish tase. They are looked upon as stimulating and attenuating, and supposed to promote expectoration and urine: the expressed juice was employed by the ancients as an errhine in lethargic disorders. The present practice pays no regard to them in any intention.

PHU, vide VALERIANA 1YL-

PILOSELLA, vide Auricula muris.

PIMENTA, vide PIPER JA-MAICENSE.

PIMPINELLÆ SANGUISOR-BÆ folia: Pimpinellæ fanguisorbæ minoris hirsutæ et lævis C. B. Burnet; the leaves.

This grows wild upon dry chalky hills: such as is cultivated in gardens, though preserved by some, is inferior in quality to the wild sort. The leaves are mildly astringent, and have been sometimes employed in this intention, in dysenteries and hæmorrhages.

PIMPINELLÆ SAXIFRAGÆ radix, femen, folia. Burnet faxifrage; the root [L. E.] leaves and feeds [E.]

Three forts of this plant are taken notice of by medical writers

1. Pim-

1. Pmpinella faxifraga major, umbella candida C. B. This is the species celebrated by the German writers under the name of pimpinella alba: it is not very common in this country, and therefore our markets have been generally supplied with the following

2. Pimpinella saxif agu minor, foliis sangui/orba Reii Tra, oseli num alterum majus Tourn. This is not unfrequently met with in dry

pasture grounds.

3. Pimpinella faxifraga minor C. B foliis diffectis Hift. Oxon. This fort is the most common in the fields about London: it grows taller than the others, but the leaves are less.

All these plarts seem to be possessed of the same qualities, and to differ only in external appearance: and even in this their difference is so inconsiderable, that Linnæus has joined them into one, under the general name of pimpinella. Our college, instead of the first, which has been generally understood as the officinal fort, allow either of the others (which are more common) to be used promiscuously.

The roots of pimpinella have a grateful, warm, very pungent tafte, which is entirely extracted by rectified spirit: in distillation, the menstruum arises, leaving all that it had taken up from the root, united into a pungent aromatic This root promises, from its sensible qualities, to be a medicine of confiderable utility; though little regarded in common practice: the only officinal composition in which it is an ingredient is the pulvis ari compositus [L] Stahl, Hoffman, and other German phyficians, are extremely fond of it, and recommend it as an excellent flomachic, resolvent, detergent, diuretic, diaphoretic, and alexipharmac. They frequently gave it,

and not without success, in scorbutic and cutaneous disorders, foul-nessofthe blood and juices, tumours and obstructions of the glands, and discases proceeding from a desciency of the sluid secretions in general. Boerhaave directs the use of this medicine in asthmatic and hydropic cases, where the strongest resolvents are indicated: the form he prefers is a watery insusion; but the spirituous tincture possesses the virtues of the root in much greater persection.

There is another species of pimpinella called nigra, from its root being externally of a bright black colour, whilft those of the foregoing forts are whitish: this is remarkable for its yielding an effential oil of a blue colour. It grows wild in some parts of Germany, Swifferland, &c. and is now and then met with in our gardens.

PINUS nuclei et refina: Pinus fative C. B. et Pinus sylvessiris C. B. Pine tree; the kernels of its fruit or cones, and its refin [E]

The pine tree differs from the firs in having its leaves standing in pairs, those of the firs being solitary. The pine abounds with the same kind of resinous juice as the fir trees (see the articles Terebinthina and Thus vulgare.) The kernels have a very pleasant sweet taste, and appear to be nearly of the same quality with sweet almonds: they are considered rather as dietetic than medicinal articles.

PIPER NIGRUM [L. E.] Black pepper; the fruit of a plant growing in Java, Malabar, &c. gathered probably before it is fully ripe, and exficcated in the fun. This is the only fpice which we import directly from the East Indies, all the others coming through the hands of the Dutch.

**PIPER** 

PIPER ALBUM [L. E] White pepper: the fruit of the black pepper plant gathered when ripe, and decorticated by maceration in water. The grains, as brought to us, have fometimes pieces of a dark coloured skin still upon them.

PIPER LONGUM [L. E.]
Long pepper. This is the fruit of
a different plant growing also in
the East-Indies. It is of a cylindrical figure, about an inch and a
half in length; the external surface
appears composed of numerous minute grains disposed round the fruit
in a kind of spiral direction.

All these spices have a pungent fmell, and a very hot biting tafte. The long fort, which is the hottest and strongest, is most frequently made use of formedicinal purposes; the black, as being more grateful, for culinary ones: the white, which is the weakest of the three, is rarely employed for either. The warmth and pungency of these spices refides chiefly in their refinous part; their aromatic odour in an essential oil. The genuine distilled oil fmells strong of the pepper, but has very little acrimony; the remaining decoction, inspissated, yields an extract confiderably pungent. A tincture made in rectified spirit is extremely hot and fiery; a few drops of it let the mouth as it were in a flame.

The white pepper is an ingredient in philonium and mithridate, the black, in the pulvis antilyffus, electary of bayberries, confectio Paulina, and theriaca; the long, in the bitter wine, aromatic tincture, powder and pills, the compound powders of bole and fordium, the confectio Paulina, mithridate, and theriaca [L.]

PIPER JAMAICENSE [L. E.] Pimento, or Jamaica pepper; the

amomum of many of the German writers.

This is the produce of our own plantations; it is the fruit of a large tree, growing spontaneoully in the mountainous parts of Jamaica, called by fir Hans Sloan. myrtus arborea aromatica, laurinis. The fmell of this spice resembles a mixture of cinnamon, cloves, and nutmegs: its tafte approaches to that of cloves, or a mixture of the three foregoing; whence it has received the name of all-spice. The shops have been for some time accustomed to employ this aromatic as a succedaneum to the more colly spices, and from them it has been introduced into our hospitals: the London college have given it a place in their late dispensatory, and direct a fimple water to be distilled from it, which possesses the flavour of the pimento in great perfection. yields a large quantity of a pleafant effential oil, which finks in water: this oil, recommended in the Pharmacopœia reformata, is now received into the Edinburgh Pharmacopæia. Rectified spirit extractsits pungency and flavour, and elevates nothing in distillation.

PIPER INDICUM: Capficum fil quis longis propendentibus Tourn. Guinea pepper, or capficum; the fruit.

This is an annual plant cultivated in our gardens; it ripens its red pods in September or October. The taste of capsicum is extremely pungent and acrimonious, setting the mouth as it were on fire. It is rarely made use of in medicine, being chieshy employed for cultinary purposes: a species of it, called in the West-Indies bird pepper, is the basis of a powder brought us from thence under the name of Cayan pepper.

PISUM

PISUM: Pifum arvense store candido, frudu rotundo ulbo C.B. Peas; the seeds [E.]

These are commonly used in food, but very rarely for medicinal purposes.

PIX LIQUIDA [L. E.] Tar; a thick, black, unctuous substance; obtained from old pines and firtrees, by burning them with a close smothering heat. It differs from the native refinous juice of the trees (see Terebinthina) in having received a disagreeable impression from the fire, and containing a portion of the faline and other juices united with the refinous and oily: by the mediation of these, a part of the terebinthinate oil proves diffoluble in aqueous liquors, which extract little or nothing from the purer turpentines. Water impregnated with the more foluble parts of tar, proves, in consequence of this hot pungent oil, warm and slimulating: it fensibly raises the pulse and quickens the circulation: by these qualities, in cold languid phlegmatic habits, it strengthens the folids, attenuates viscid juices, opens obstructions of the minuter vessels, and promotes perspiration and the fluid secretions in general; whilst in hot bilious temperaments, it disposes to inflammation, and aggravates the complaints which it has been employed to remove.

PIX ARIDA [L. E.] Dry or

stone pitch.

This is the pix liquida, exficcated by heat: in this process, a part of the acid and the more volatile oil are dissipated along with the aqueous moisture; and hence the product proves considerably less active. It is made use of only in external applications, as a warm adhesive, resinous substance.

PIX NAVALIS. This is gene-

rally allowed to be the same with the foregoing dry pitch or inspissated tar: hence in the Edinburgh pharmacopœia the terms pix ficca and pix navalis are made synony-According to Geoffroy, it mous. is compounded of a strange mixture of tallow, and tar, and palimpissa, and an artificial black pitch; which artificial pitch is itfelf composed of tar and palimpissa; and this palimpissa is no other than an inspissated tar: so that notwithstanding this show of composition, the result is only a mixture of pitch with a little tallow.

PIX BURGUNDICA [L. E.] This is of a fo-Burgundy pitch. lid confishence, yet somewhat soft, of a reddish brown colour, and more agreeable in smell than either of the foregoing. Geoffroy relates, that it is composed of gallipot (a folid whitish refin which separates from some of the terebinthinæ as they run from the tree) melted with common turpentine and a little of its distilled oil. Dale informs us, from the relation of a gentleman who saw the preparation of this commodity in Saxony (from whence we are chiefly supplied with it) that it is no more than the common turpentine boiled a little.

All these substances are employed in the shops only in external compositions. The dry pitch and Burgundy pitch are ingredients in several plassers, ointments, and cerates: and tar gives name to one of the ointments.

PLANTAGINIS LATIFOLIÆ folia, femen: Common broad leaved plantane, called feptinervia, from its having seven large nerves or ribs running along each leaf; the narrow leaved fort has only sive ribs, and hence is named quinquenervia: they are both common in fields,

and by road-fides. The leaves are lightly aftringent, and the feeds faid to be so; and hence they stand recommended in hæmorrhages, and other cases where medicines of this kind are proper. The leaves bruised a little, are the usual application of the common people to slight shesh wounds. The Edinburgh college directs an extract to be made from the leaves

PLUMBUM [L. E.] Lead.

This is the heaviest of the metals except gold: it melts in a moderate heat, and if kept in fusion, is soon converted partly into fume and partly into an ash-coloured calx (plumbumuflum;) this exposed to a stronger fire, in such a manner that the flame may play upon its surface, becomes first yellow, and af- . terwards of a deep red, (minium or red lead:) if in this process the fire be suddenly raised to a considerable height, the calx melts, assumes the appearance of oil, and on cooling forms a foft leafy substance of a vellowish or reddish colour (litharge.) The proper menstruum of this metal is aquafortis: the vegetable acids likewise dissolve it, but in very small quantity: a quart of distilled vinegar will not take up a drachm; exposed to the steam of vinegar, it is by degrees corroded intoa white powder (cerusse) which is confiderably more easy of solution. The calces of lead dissolve, by heat, in expressed oils; these mixtures are the basis of several officinal plasters and unguents. Crystals of this metal, made with distilled vinegar (called, from their sweetish taste, sugar of lead) and a tincture drawn from these and green vitriol, are likewise kept in the shops.

Preparations of lead, given internally, are supposed to incrassate the sluids, abate instammations, and restrain venereal desires. The sugar

is a strong astringent, and has been used, it is said, with good success, in hæmorrhages, the fluor albus, feminal gleets, &c. The tincture is recommended for the like purpoles; and for checking immoderate sweats in phthisical cases, whence it has been usually called tincura antiphtbisica. The internal use of this metal is nevertheless full of danger, and ought never to be ventured upon unless in desperate cases, after other medicines have been employed without taking effect: it often occasions violent colics; and though it should not prove immediately hurtful, its ill confequences are fure, though flow: tremors, spasms, or lingering tabes, too frequently follow.

POLII, seu Polii montani summitatis. Poley mountain; the tops [L. E.]

It has been disputed among botanic writers, what species of poley ought to be employed in medicine. The London college allows the promiscuous use of two, the polium maritimum erectum Monspeliacum C. B. and the polium angustifolium Creticum C. B. The first is sometimes cultivated in our gardens, and is the fort which the shops have been generally supplied with. They have both a light aromatic smell, and a bitterish taste; that brought from Crete is the most agreeable. They stand recommended in catarrhs, uterine disorders, &c. but at present are scarce otherwise made use of than as an ingredient in the mithridate and theriaca.

POLYGONATUM, vide SI-GILLUM SALOMONIS.

POLYGONUM, vide Centi-

POLYPODII radix: Filicis polypodii *lypodii dicta Herm.* Polypody; the root  $\{E_n\}$ 

Polypody is a capillary plant, growing upon old walls, thetrunks of decayed trees, &c. that found upon the oak is generally preferred though not fenfibly different from the others. The roots are long and slender, of a reddish brown colour on the outside, greenish within, full of small tubercles, which are resembled to the feet of an insect; whence the name of the plant; the taste of these roots is sweetish and nauseous.

Polypody has been employed in medicine for many ages; nevertheless its virtues remain as yet to be determined. The ancients held it to be a powerful purger of melancholic humours; by degrees, it came to be looked upon as an evacuater of all humours in general: at length, it was supposed only to gently loofen the belly; and afterwards even this quality wa denied it: succeeding physicians declared it to be astringent; of this number is Boerhaave, who esteems it moderately flyptic and antiscorbutic. For our own part, we have had no direct experience of it; nor is it employed in practice: it is probable that (as Juncker supposes) the fresh root may loosen the belly, and that it has not this effect when dry.

POLYTRICHUM, vide TRI-

POMPHOLYX: a calx, or flowers, of zinc, produced in the furnaces where copper is made into brafs by calamine the ore of zinc. It is found adhering to the covers of the crucibles, &c. either in form of thin crufts, or of a light downy matter, generally of a pure white colour, though sometimes yellowish. See Zincum.

POPULI NIGR TE gemme: Populi nigræ C. B. The black popular; its buds [E.]

The black poplar is a large tree, growing wild in watery places; it is easily raised and very quick of The young buds or rugrowth. diments of the leaves, which appear in the beginning of spring, abound with a yellow unctuous, odorous juice. They have hitherto been employed chiefly in an ointment, which received its name from them; though they are certainly capable of being applied to other purposes: a tincture of them made in rectified spirit, yields, upon being inspissated, a fragrant refin, superior to many of those brought from abroad.

PORRI radix: Porri communis capitati C. B. Leeks; the root. This participates of the virtues of garlick, from which it differs chiefly in being much weaker. See the article Allium.

PORTULACÆ semen: portulacæ hortensis latisoliæ J. B. Putslane; the seeds.

This herb is cultivated in gardens for culinary uses. The seeds are ranked among the lesser cold seeds, and have sometimes been employed in emulsions, and the like, along with the others of that class.

POTENTILLA, vide Argen-

PRASIUM, vide MARRUBIUM.

PRIMULÆ VERIS folia, radix: Primulæ veris pallido flore bumilis Tourn. Primrose; the herb and root [E.]

This is a low plant, growing wild in woods and hedges, and producing pale yellow flowers in the spring. The leaves have an herbaceous taste. The roots are lightly bitter, with a kind of aromatic slavour, which some refemble to that of aniseeds; their expressed juice, purished by settling, is sometimes used as a sternutatory. The slowers have an agreeable slavour, but very weak: an infusion of them in wine, and a sprirt distilled from them, are employed in some places as cordial and nervine.

PRUNELLÆ, seu Brunellæ solia: Prunellæ majoris soliis non dissettis C. B. Sels-heal; the leaves

 $\{E\}$ 

This plant grows wild in meadows and pasture grounds, and produces thick spikes of purplish slowers during the latter part of the summer. It has an herbaceous roughish taste: and herbaceous recommended in hamorrhages and alvine sluxes: it has been principally celebrated as a vulnerary, whence its name; and in gargarisms for aphthæ, and inslammations of the sauces.

PRUNUS HORTENSIS. The plum tree. Three forts of plums are looked upon as articles of the Materia Medica. They are all met with in our gardens; but the shops are supplied with the fruit moderately dried from abroad.

PRUNA BRIGNOLENSIA: Pruna ex flavo rufescentia, mixti saporis, gratissima C. B. The Brignole plum, brought from Provence
under the name of prunelloes [E.]

PRUNA GALLICA [L. E.]: Frudus Pruni frudu parvo, dulci, atro-caruleo Tourn. French or common prunes [L. E.] This is the plum called by our gardeners the little black damask.

PRUNA DAMASCENA:
Fructus Pruni fructu magno, dulci,
atro cæruleo Tourn Damascene
plums, or damsons. This is the
fort called the great damask violet
of Tours. It is seldom met with
dry in the shops, and is generally
supplied by the common prune.

The medical effects of the damfon and common prunes are, to abate heat, and gently loofen the belly; which they perform by lubricating the passage, and softening the excrement. They are of confiderable service in costiveness, accompanied with heat or irritation, which the more stimulating cathartics would tend to aggravate: where prunes are not of themselves sufficient, their effects may be promoted by joining with them a little rhubarb or the like : to which may be added some carminative ingredient, to prevent their occasioning flatulencies. Prunelloes have scarce any laxative quality: these are mild grateful refrigerants, and by being occasionally kept in the mouth, usefully allay the thirst of hydropic persons.

PRUNA SILVESTRIA: Fructus pruni filvesfiris C. B. Sloes; the fruit of the common black thorn, or sloe bush [L. E]

These have a very rough austere taste, especially before they have been mellowed by frosts. The juice of the unripe fruit, inspissated to a proper consistence, is called acaci Germanica, and usually sold in the shops for the true Egyptian acacia: it is equally askringent with the Egyptian sort, but has more of a sharp or tartish taste, without any thing of the sweetish relish of the other. The inspissated juice is directed as an officinal by the Edinburgh college, and a conserve of the fruit by the London,

**PSYLLII** 

PSYLLII semen: Psyllii majoris eredi C. B. Fleawort; the seeds.

This is a fort of plantane, growing wild in the warmer climates, and fometimes met with in our gardens: it differs from the common plantanes in having its stalks branched, with leaves upon them; hence it is named by Ray, plantago caulifera. The feeds have been usually brought from the fouth of France; they are small, but supposed to resemble in shape a slea, whence the English name of the These seeds have a nauseous, mucilaginous taste; boiled in water, they yield a considerable quantity of mucilage, which is sometimes made use of in emollient glysters, and the like. Alpinus relates, that among the Egyptians this mucilage is given in ardent fevers, and that it generally either loosens the belly or promotes fweat.

PTARMICÆ radix: Dracunculi pratenfis, serrato folio C. B. Sneezewort, or bastard pellitory: the

root [E.]

This grows wild upon heaths and in moist shady places: the slowers, which areof a white colour, come forth in June and July. The roots have an acrid smell, and a hot biting taste: chewed they occafion a plentiful discharge of saliva; and when powdered and fnuffed up the nose, provoke sneezing. These are the only intentions to which they have been usually applied.

PULEGII folia: Pulegii latifolii C. B. Menthæ aquaticæ seu pulegii vulgaris Tourn. Pennyroyal; the leaves [L. E.]

This plant grows spontaneously in several parts of England upon moist commons, and in watery places; trailing upon the ground, and striking roots at the joints.

Our markets have been for some time supplied with a garden fort, which is larger than the other, and grows upright: this is called by Mr. Dale pulegium erectum.

Pennyroyal is a warm, pungent herb of the aromatic kind, similar to mint, but more acrid and less agreeable: it has long been held in great effeem, and not undeservedly, as an aperient, and deobstruent, particularly in hysteric complaints, and suppressions of the uterine purgations. For these purposes, the distilled water is generally made use of, or what is of equal efficacy, an infusion of the leaves. It is observable, that both water and rectified spirit extract the virtues of this herb by infusion, and likewise elevate greatest part of them in distillation.

In the shops are kept a simple [L. E.] and spirituous [L.] water and essential oil [L. E.] of the plant: this herb is used also in the compound valerian water and troches of myrrh [E.] and its simple water for making the lac ammoniaci [L.] and the camphorated

emultion [E]

PULEGII CERVINI folia: Pulegii angustifolii C. B. Harts pennyroyal; the leaves.

This species is met with, though not very often, in our gardens. It is somewhat stronger, yet rather more agreeable, than the foregoing, both in taste and smell.

PULMONARIÆ MACULO-SÆ folia: Pulmonariæ Italorum al buglossam accedentis J. B. Spotted lungwort, or fage of Jerusalem; the leaves [E.]

This is met with in gardens: the leaves are of a green colour spotted with white; of an herbaceous somewhat mucilaginous taste, without any smell. They stand re-

commended

commended against ulcers of the lungs, phthises, and other like disorders: nevertheless experience gives little countenance to these virtues, nor does the present practice expect them.

PYRETHRI radix: Pyrethri flore bellidis C. B. Pellitory of

Spain; the root [L. E.]

This plant, though a native of the warm climates, bears the ordinary winters of this: and often flowers succeffively, from Christmas to May; the roots also grow larger with us than those which the shops are usually supplied with from abroad.

Pellitory root has no fensible fmell; its taste is very hot and acrid, but less so than that of arum or dracunculus: the juice expressed from it has scarce any acrimony, nor is the root itself so pungent when fresh as after it has been dried. Water, assisted by heat, extracts some share of its taste, rectified spirit the whole; neither of them elevate any thing in distillation. The principal use of pyrethrum in the present practice is as a masticatory, for promoting the falival flux, and evacuating viscid humours from the head and neighbouring parts: by this means it often relieves the tooth-ach, some kinds of pains of the head, and lethargic complaints.

QUERCUS cortex: Quercus cum longis pediculis C. B. Oak tree;

the bark [E.]

This bark is a strong astringent, and hence stands recommended in hæmorrhages, alvine sluxes, and other præternatural or immoderate secretions.

RANARUM SPERMA. Frogs fpawn. This has been celebrated as an excellent cooler for external purposes; but practitioners have not experienced from it any peculiar effects that could deserve its being continued in use, and both the London and Edinburgh colleges have now discarded it.

RAPHANI RUSTICANI radix: Raphani rusticani C. B. Cochsariæ folio cubitali Tourn. Horseradish; the root [L. E.]

This plant is sometimes found wild about river fides, and other moift places; for medicinal and culinary uses, it is cultivated in gardens; it flowers in June, but rarely perfects its feeds in this country. Horse-radish root has a quick pungent smell, and a penetrating acrid taste; it nevertheless contains in certain vessels a sweet juice, which fometimes exudes upon the furface. By drying, loses all its acrimony, becoming first sweetish, and afterwards almost infipid: if kept in a cool place covered with fand, it retains its qualities for a confiderable time. The medical effects of this root are to stimulate the solids, attenuate the juices, and promote the fluid fecretions: it feems to extend its action through the whole habit, and affect the minutest glands. has frequently done good service in some kinds of scurvies and other chronic disorders proceeding from a viscidity of the juices, or obstructions of the excretory ducts. denham recommends it likewise in dropsies, particularly those which fometimes follow intermittent fevers. Both water and rectified spirit extract the virtues of this root by infusion, and elevate them in distillation: along with the aqueous fluid, an essential oil arises, possessing the whole taste and pungency of the horse-radish. The college have given us a very elegant compound water, which takes its name from this root.

RAPI

4

RAPI radix, semen: Rapi sativi rotundi C. B. Turneps; the roots

and feeds [E.]

These roots are accounted a wholesome aperient food: the liquor pressed out from them after boiling has been sometimes used medicinally as a deobstruent and diuretic. The seeds are slightly pungent.

REALGAR, a foffil composed of arsenic and sulphur. Vide Arsenicum.

REGINA PRATI, vide UL-

RHABARBARUM [L. E.] Rhubarb; the root of a plant of the dock kind, which grows spontaneoully in China, and endures the colds of our own climate. forts of rhubarb are met with in the shops. The first is imported from Turkey and Russia, in roundish pieces freed from the bark, with a hole through the middle of each; they are externally of a yellow colour, and on cutting appear variegated with lively reddish The other, which is less esteemed, comes immediately from the East-Indies, in longish pieces, harder, heaviet, and more compact than the foregoing. The first fort; unless kept very dry, is apt to grow mouldy and worm-eaten; the fecond is less subject to these inconveniencies. Some of the more industrious artists are said to fill up the worm holes with certain mixtures, and to colour the outfide of the damaged pieces with powder of the finer forts of rhubarb, and sometimes with cheaper materials: this is often so nicely done, as effectually to impose upon the buyer, unless he very carefully examines The marks of good each piece. rhubarb are, that it be firm and folid, but not flinty; that it be eafily pulverable, and appear, when powdered, of a fine bright yellow colour: that upon being chewed, it imparts to the spittle of a saffron tinge, without proving slimy or mucilaginous in the mouth. It taste is subscrid, bitterish, and somewhat astringent; the smell

lightly aromatic.

Rhubarb is a mild cathartic, which operates without violence or irritation, and may be given with fafety even to pregnant women and children. Besides its purgative quality, it is celebrated for an astringent one, by which it strengthens the tone of the stomach and intestines, and proves useful in diarrhee and diforders proceeding from a laxity of the fibres. Rhubarb in substance operates more powerfully as a cathartic than any of the preparations of it. Watery tinctures purge more than the spirituous ones; whilst the latter contain in greater perfection the aromatic, aftringent, and corroborating virtues of the rhubarb. The dofe, when intended as a purgative, is from a scruple to a drachm or more.

The Turkey rhubarb is, among us, univerfally preferred to the East-India fort, though this last is for some purposes at least equal to the other; it is manifestly more astringent, but has somewhat less of an atomatic flavour. Tinctures drawn from both with reclified spirit, have nearly the same taste: on distilling off the menstruum, the extract left from the tincture of the East-Indiathubarb proved considerably the strongest. They are both the produce of the same climate, and probably the roots of the same plant taken up at different seasons, or cured in a different manner.

The officinal preparations of this drug are, roasted rhubarb [L.] a watery infusion [E.] and vinous and spirituous tinctures [L. E.] It is an ingredient in sundry composi-

tions,

tions, as the fyrup of sena and rhubarb, dysenteric electary, stomachic pills [E.] ecphractic pills [L.] &c.

RHAMNUS CATHARTICUS, vide Spina Cervina.

RHAPONTICI radix: Rhabarbari Dioscoridis et antiquorum Rhapontic; the root of a large roundish-leaved dock, growing wild on the mountain Rhodope in Thrace, from whence it was brought into Europe, about the year 1610, by Alpinus; it bears the hardest winters of this climate, and is not unfrequent in our botanic gardens. The root of this plant (which appears evidently to have been the rhubarb of the ancients) is by fome confounded with the modern rhubarb, though confiderably different both in appearance and quality. The rhapontic is of a dusky colour on the surface, of a loose spongy texture; considerably more astringent, but less purgative, than rhubarb; in this last intention, two or three drachms are required for a dofe.

RHUS OBSONIORUM, vide Sumach.

RIBESIA: frudus Ribes vulgaris frudu rubro Rais. Red cursant bush; the berries [E.]

These have a cool, acidulous, sweet taste, sufficiently agreeable both to the palate and stomach. The college of Edinburgh directs a gelly to be made from them with sugar; but at present they are rather looked upon as a dietetic than a medicinal article.

ROSA DAMASCENA: Rosa purpurea C. B. The damaik rose [L. E.]

This elegant flower is common in our gardens. Its smell is very

pleasant, and almost universally admired; its taste bitterish and subacrid. In distillation with water, it yields a small portion of a butyraceous oil, whose flavour exactly refembles that of the rofes. This oil, and the distilled water. are very useful and agreeable cordials: Hoffman strongly recommends them as of fingular efficacy for raising the strength, chearing and recruiting the spirits, and allaying pain; which they perform without raifing any heat in the constitution, rather abating it when inordinate. Damask roses, besides their cordial aromatic virtue, which refides in their volatile parts, have a mildly purgative one, which remains entire in the decoction left after the distillation: this, with a proper quantity of fugar, forms an agreeable laxative fyrup, which has long kept its place in the shops. The other officinal preparations of this flower, are a solutive honey, and the distilled water, which last is an ingredient in the musk-julep, the confection of kermes, and faponaceous lotion, and is used also in making the fimple ointment called pomatum. [L.]

ROSA RUBRA: Rosa rubra multiplex B. C. The red rose [L. E.]

This has very little of the fragrance of the foregoing pale fort; and instead of its purgative quality, a mild gratefully aftringent one, efpecially before the flower has opened: this is confiderably improved by hafty exficcation; but both the aftringency and colour are impaired by flow drying. In the shops are prepared a conferve, honey. tincture, troches [L.] vinegar and fyrup [E.] of this flower: it is an ingredient also in the compound powder of scordium, the troches of Japan earth, mithridate, and theriaca [L.]

RORIS-

RORISMARINI summitates, et flores anthos dicti: Korismarini hortensis angustiore folio C. B. Rosemary; the tops and flowers [E.]

This is a native of Spain, Italy, and the fouthern parts of France, where it grows in great abundance upon dry gravelly grounds; in the like foils, it thrives best with us, and likewise proves stronger in fmell, than when produced in moist rich ones: this observation obtains in almost all the aromatic

plants.

Rosemary has a fragrant smell, and a warm pungent bitterish taste, approaching to those of lavender: the leaves and tender tops are Arongest; next to these the cup of the flower; the flowers themselves are confiderably the weakest, but most pleasant. Aqueous liquors extract great share of the virtues of rolemary leaves by infusion, and elevate them in distillation; along with the water arises a considerable quantity of effential oil, of an agreeable strong penetrating smell. Pure spirit extracts in great perfection the whole aromatic flavour of the rofemary, and elevates very little of it in distillation; hence the resinous mass, lest upon abstracting the spirit, proves an elegant aromatic, very rich in the peculiar qualities of the plant. The flowers of rolemany give over great part of their flavour in distillation with pure spirit; by watery liquors their fragrance is much injured; by beating destroyed. The officinal preparations of rolemary are an ellential oil from the leaves [L.] or from the herb in flower [E.] a conserve of the flowers [L. E.] and a spirit, called Hungary water, from the flowery tops [L. E.] The tops are used also in the compound spirit of lavender [L. E.] cordial confection [L.] and cephalic tincture[E.] and the effential oil in the cephalic bal-

faponaceous balfam, and ſam, nerve ointment [E.]

RUBIA TINCTORUM: radix Rubiæ tinctorum fativæ C. B. Madder; the root [L. E.]

Madder is raised in some of our gardens for medicinal purposes: it was formerly cultivated among us, in quantity, for the use of the dyers, who are at present supplied from Holland and Zealand. It has little or no smell; a sweetish tafte, mixed with a little bitterness. The virtues attributed to it, are those of a detergent and aperient, whence it has been usually ranked among the opening roots, and recommended in obstructions of the viscera, particularly of the kidneys, in coagulations of the blood from falls or bruises, in the jaundice, and beginning dropfies. It is an ingredient in the icteric decoction of the Edinburgh pharmacopæia.

It is observable, that this root, taken internally, tinges the urine of a deep red colour; and in the Philosophical Transactions, we have an account of its producing a like effect upon the bones of animals who had it mixed with their food: all the bones, particularly the more folid ones, were changed, both externally and internally, to a deep red, but neither the fleshy or cartilaginous parts suffered any alteration: some of these bones macerated in water for many weeks together, and afterwards fleeped and boiled in spirit of wine, lost none of their colour, nor communicated any tinge to the liquors. This root appears therefore to be polfessed of great subtility of parts, whence its medical virtues seem to

deserve inquiry.

RUBI IDÆI frudus : Rubi idei spinosi C. B. The raspberry bush; the fruit [L.]

This

This shrub is common in our gardens; and has likewise, in some parts of England, been found wild: it flowers in May, and ripens its Raspberries have a fruit in July. pleasant sweet taste, accompanied with a peculiarly grateful flavour; on account of which they are chiefly valued. As to their virtues, they moderately quench thirst, abate heat, strengthen the viscera, and promote the natural excretions. An agreeable syrup, prepared from the juice, is directed to be kept in the shops.

RUBI VULGARIS folia, fructus: Rubi vulgaris five rubi frudu nigro C. B. The bramble, or blackberry bush; its leaves and fruit.

This shrub is frequently found wild in woods and hedges. The berries have a faint taste without any thing of the agreeable slavour of the foregoing: the leaves are somewhat astringent.

RUSCI, five Brusci radix: Rusci myrtifolii aculeati Tourn. Butchersbroom, or knee-holly; the root [E.]

This is a small prickly plant, sometimes found wild in woods. The root has a soft sweetish taste, which is followed by a bitterish one: it is one of the five aperient roots; and in this intention is sometimes made an ingredient in apozems and diet drinks, for opening slight obstructions of the viscera, purifying the blood and juices, and promoting the sluid secretions.

RUTÆ folia semen: Rutæ horzensis latisoliæ C. B. Broad leaved rue; the leaves [L. E.] and seeds

This is a small shrubby plant, met with in gardens, where it slowers in June, and holds its green leaves all the winter: we frequently find in the markets a narrow leaved

fort, which is cultivated by fome in preference to the other, on account of its leaves appearing variegated during the winter, with white streaks.

Ruehas aftrong ungrateful smell, and a bitterish, penetrating taste: the leaves, when in full vigour, are extremely acrid, infomuch as to inflame and blifter the skin, if much handled. With regard to their medicinal virtues, they are powerfully stimulating, attenuating and detergent: and hence, in cold phlegmatic habits, they quicken the circulation, dissolve tenacious juices, open obstructions of the excretoryglands, and promote the fluid secretions. Thewriters on the Materia Medica in general have entertained a very high opinion of the virtues of this Boerhaave is full of its praises, particularly of the effential oil, and the distilled water cohobated or re-distilled several times from fresh parcels of the herb: after fomewhat extravagantly commending other waters prepared in this manner, he adds, with regard to that of rue, that the greatest commendations he can bestow upon it fall short of its merit: " What medicine (says he) can be more efficacious for promoting sweat and perspiration, for the cure of the hysteric passion, and of epilepsies and for expelling poison?" Whatever service rue may be of in the two last cases, it undoubtedly has its use in the others: the cohobated water, however, is not the most efficacious preparation of it. Part iii. chap. v.) An extract made by rectified spirit, contains, in a fmall compass, the whole virtues of the rue; this menstruum taking up by infusion all the pungency and flavour of the plant, and elevating nothing in distillation. With water, its peculiar flavour and warmth arise; the bitterness, and a consi-P 2 derable derable share of the pungency

remaining behind.

A watery extract [E.] essential oil [L. E.] and conferve [L.] of This rue are kept in the shops. herb is used also as an ingredient in the electary of bayberries, compound powder of myrrh, and the green oil [L.]

SABINÆ folia seu summitates: Sabinæ f lio tumarisci Dioscoridis C. B. Savin; the leaves or tops L. E.

This is an evergreen shrub, clothed with small, **fomewhat** prickly leaves: it does not produce fruit till very old, and hence has been generally reputed barren. The leaves have a bitter, acrid, biting taste; and a strong disagreeable smell: distilled with water they yield an essential oil, in larger quantity (as Hoffman observes) than any other known vegetable, the turpentine tree alone excepted.

Savin is a warm irritating aperient medicine, capable of promoting sweat, urine, and all the glandular secretions. The distilled oil is one of the most powerful emmenagogues; and is found of good fervice in obstructions of the uterus, or other viscera, proceeding from a laxity and weakness of the vessels, or a cold sluggish indisposition of the juices.

The effential oil [L. E.] and a watery extract [L.] of favin are kept in the shops: the leaves themfelves are an ingredient in the compound valerian water [E.]; the extract in the compound elixir of myrrh [L.]; and the effential oil in the troches of myrrh [E.]

SACCHARUM ALBUM. White or refined fugar [E.]

SACCHARUM PURISSIMUM. Double-refined fugar [L.]

SACCHARUM RUBRUM Brown or unrefined fugar [L. E.]

SACCHARUM CANDUM.

Sugar-candy [E.]

Sugar is the essential salt of the arundo saccharifera, a beautiful large cane growing spontaneously in the East Indies, and some of the warmer parts of the West, and cultivated in great quantity in our American plantations. The expressed juice of the cane is clarified with the addition of lime water (without which it does not assume the form of a true fugar) and boiled down to a due consistence; when, being removed from the fire, the faccharine part concretes from the groffer unctuous matter, called treacle, or melasses. This, as yet impure or brown fugar, is farther purified, in conical moulds, by spreading moist clay on the upper broad furface: the watery moisture, flowly percolating through the mass, carries with it a confiderable part of the remains of the treacly matter. This clayed fugar, imported from America, is by our refiners dissolved in water, the solution clarified by boiling with whites of eggs and despumation, and after due evaporation poured into moulds; as foon as the fugar has concreted, and the fluid part drained off, the surface is covered with moitt clay as before. The fugar, thus once refined, by a repetition of the process, becomes the double refined fugar of the shops. The candy, or crystals, are prepared by boiling down folutions of fugar to a certain pitch, and then removing them into a hot room; with slicks set across the vessel for the sugar to shoot upon: these crystals prove of a white or brown colour, according as the fugar was pure or impure.

The uses of sugar as a sweet, are sufficiently well known. The impure sorts contain an unctuous, or oily matter, in consequence of which they prove emollient and laxative. The crystals are most difficult of solution, and hence are properest where this soft lubricating sweet is wanted to dissolve slowly in the mouth.

SAGAPENUM [L. E.] a concrete juice brought from Alexandria, either in distinct tears, or run together in large masses. It is outwardly of a yellowish colour, internally somewhat paler, and clear like horn, grows soft upon being handled, and sticks to the singers; its taste is hot and biting; the smell disagreeable, by some resembled to that of a leek, by others to a mixture of asafetida and galbanum.

Sagapenum is an useful aperient and deobstruent; and frequently prescribed either alone, or in conjunction with ammoniacum, or galbanum, for opening obstructions of the viscera, and in hysterical disorders arising from a deficiency of the menstrual purga-It likewise deterges the pulmonary vessels, and proves of confiderable service in some kinds of asthmas, where the lungs are oppressed by viscid phlegm. It is most commodiously given in the form of pills: from two or three grains to half a drachm, may be given every night or oftener, and When continued for some time. sagapenum is scarce, the druggists usually supply its place with the larger and darker coloured masses of bdellium, broken into pieces; which are not easily distinguished from it.

Sagapenum is an ingredient in the compound powder of myrrh, gumpills, electary of bay-berries, mithridate and theriaca of the London

pharmacopœia. The college of Edinburgh has no where employed either this gum or opoponax, givt ing the preference to ammoniacum and galbanum.

SAGO [E.] This is the produce of an oriental tree, called by C. Bauhine palmam referens arbor farinifera. The medullary parof the tree is beaten with water, and made into cakes, which are used by the Indians as bread: these reduced into granules, and dried, are the sago brought to us. It is moderately nutritious, though not perhaps superior to our own grain.

SAL AMMONIACUS. Sal ammoniac  $\{L, E\}$ 

This is an artificial saline concrete, faid to be prepared by fublimation from the foot of cow dung. It is brought to us from Egypt in large round cakes, convex on one fide, and concave on the other: and fometimes in conical loaves: on breaking, they appear composed of needles, or firize, running transversely. The best are almost transparent, colourless, and free from any visible impurities: those most commonly met with are of a grey yellowish colour on the outside, and fometimes black, according as the matter is more or less impure. The taste of this falt is very sharp and penetrating. It dissolves in twice its weight, or a little less, of water; and upon evaporating a part of the menstruum, concretes again into long shining spicula, or thin fibrous plates, like feathers.

Sal ammoniac appears from experiments to be composed of marine acid, united with a volatile alkali. If mingled with fixt salts, or absorbeat earths, and exposed to a moderate fire, a large quantity of pure volatile salt sublimes, the acid

P 3 remaining



remaining united with the intermedium; if treated in the same manner with quicklime, an exceeding penetrating volatile spirit arises, but no solid salt is obtained. Exposed alone to a considerable heat, it sublimes entire, without any alteration of its former properties: ground with certain metallic substances, it elevates some parts of them along with itself, and concretes with the remainder into a mass, which readily flows into a liquor in a moist air; this appears in most respects similar to a saturated folution of the metal made di-

rectly in spirit of salt. Pure sal ammoniac is a perfectly neutral falt, capable of attenuating viscid humours, and promoting a diaphoresis, or the urinary discharge, according to certain circumstances in the constitution, or as the patient is managed during the operation. If a drachm of the falt be taken, dissolved in water, and the patient kept warm, it generally proves sudorific; by moderate exercise, or walking in the open air, its action is determined to the kidneys; a large dose gently loosens the belly, and a still larger proves emetic. falt is recommended by many as an excellent febrifuge, and by fome has been held a great secret in the cure of intermittents. It is undoubtedly a powerful aperient, and feems to pass into the minutest vessels; and as such may in some cases be of service, either alone, or joined with bitters, or the bark, where the latter would by itself produce dangerous obstructions, or aggravate those already formed. This falt is sometimes employed externally as an antiseptic, and in lotions and fomentations, for cedematous tumours; as also in gargarisms for inflammations of the tonfils, and for attenuating and dif-

folving thick viscid mucus. It is an ingredient in the discutient cataplasm of the Edinburgh pharmacopæia.

SAL CATHARTICUS AMA-RUS [L. E.] The bitter purging falt; extracted from the bitter liquor remaining after the crystallization of common falt from lea It was first prepared as a water. cheap substitute to the falt of the Epsom, and other purging mireral waters, from which it does not confiderably differ, either in fenfible qualities, or medical effects. We ofually meet with it in minute crystals of a snowy appearance; dissolved in water, and crystallized afresh, it concretes, if properly managed, into larger ones, of a rectangular prismatic figure, refembling those of the artificial cathartic salt of Glauber, to which they are sometimes substituted in the shops.

All these salts have a penetrating bitterish taste: they dissolve in less than an equal weight of water: in a moderate heat they melt, bubble up into blifters, and foon change into a white spongy mass, with the loss of above half their weight: this calx tastes bitterer than the falts did at first, and almost totally dis-The acid folves again in water. of these salts is chiefly the vitriolic: the basis of the natural is a fine abforbent earth; of the artificial, an alkaline falt, the fame with the bafis of sea salt. Hence, upon adding alkaline salts to a solution of the falts of Glauber, no change ensues: whilst the salts obtained from the purging waters, or the bittern of marine waters, grow milky upon this addition, and deposite their earth, the alkaline salt being taken up in its place.

The fal cathacticus is a mild and gentle purgative, operating with

ſuffi-

fufficient efficacy, and in general with ease and fafety, rarely occafioning any gripes, fickness, or the other inconveniencies which purgatives of the refinous kind are too often accompanied with. Six or eight drachms may be dissolved for a dofe in a proper quantity of common water; or four, five, or more, in a pint, or quart of the purging These liquors may likewaters. wife be so managed as to promote evacuation, by the other emunctories; if the patient is kept warm, they increase perspiration; by moderate exercise in a cool air, the urinary discharge.

SAL COMMUNE. Common. This is a neuor alimentary falt. tral falt, differing from most others in occasioning drought when swallowed. It dissolves in somewhat less than three times its weight of water; the solution slowly evaporated, and fet to shoot, affords cubical crystals, which unite together into the form of hollow truncated pyramids. Exposed to the fire, it crackles and flies about, or decrepitates, as it is called; foon after it melts, and appears fluid as A fmall quantity of this water. fair, added to the nitrous acid, enables it to dissolve gold, but renders it unfit for dissolving silver: if a folution of filver be poured into liquors, containing even a minute portion of common falt, the whole immediately grows turbid and white; this phænomenon is owing to the precipitation of the filver.

This falt is either found in a folid form in the bowels of the earth, or diffolved in the waters of

the sea or faline springs.

1. Sal gemmæ [L. E.] Rock falt. I his is met with in feveral parts of the world, but in greatest plenty in certain ceep mines, of prodigious extent, near Cracow in

Poland; fome is likewise sound in England, particularly in Che-shire. It is for the most part very hard, sometimes of an opake snowy whiteness, sometimes of a red, green, blue, and other colours. When pure, it is persectly transparent and colourless; the other forts are purished by solution in water and crystallization, in order to fit them for the common uses of salt.

2. Sal marinus [L. E.] The salt extracted from sea water and saline springs. Sea waters yield from one fiftieth to one-thirtieth their weight of pure falt: several springs afford much larger quantities; the celebrated ones of our own country at Nantwich, Northwich, and Droitwich, yield (according to Dr. Brownrigg) from one fixth to fomewhat more than one-third. There are two methods of obtaining the common falt from these natural folutions of it: the one, a hafty evaporation of the aqueous fluid till the falt begins to concrete, and fall in grains to the bottom of the evaporating pan, from whence it is raked out, and set in proper vessels to drain from the brine or bittern: the other, a more flow and gradual evaporation, tinued no longer than till a faline crust forms on the top of the liquor, which, upon removing the fire, foon begins to shoot, and run into crystals of a cubical figure. In the warmer climates, both these processes are effected by the heat of the fun. The falts obtained by them differ very confiderably: that got by a hasty evaporation is very apt to relent in a moist air, and run per deliquium: an inconvenience which the crystallized salt is not subject to: this last is likewife found better for the preferving of meat, and fundry other

Com-

parpoles.

Common falt, in small quantities, is supposed to be warming, drying, and to promote appetite and digestion: in large doses, as half an ounce, it proves cathartic. It is sometimes used to check the operation of ametics, and make them run off by stool; and as a stimulus in glysters.

SALVIÆ folia: Salviæ majoris C. B. Common fage (the
green and red forts); the leaves
[L. E.]

SALVIÆ hortensis minoris solia, summitates: Salviæ minoris auritæ et non auritæ C. B Small sage or sage of virtue; the leaves and tops [E.]

These plants are common in our gardens, and flower in May and June: the green and red common fages differ no otherwise than in the colour of the leaves; the feeds of one and the same plant produce both; the small fort is a distinct species; its leaves are narrower than the others, generally of a whitish colour, and never red; most of them have at the bottom a piece standing out on each side in the form of ears. Both forts are moderately warm aromatics, accompanied with a light degree of astringency and bitterness; the fmall fort is the strongest, the large most agreeable.

The writers on the materia medica are full of the virtues of fage, and derive its name from its supposed salutary qualities, (Salvia salvatrix, naturæ conciliatrix-Cur moriator bomo, cui salvia crescit in horto, じc.) Its real effects are, to moderately warm and strengthen the vessels; and hence, in cold phlegmatic habits, it excites appetite and proves serviceable in debilities of the nervous system. The best preparation for these purposes is an insusion of the dry lcaves, drank as tea; or a tincture, or extract, made with rectified spirit, taken in proper doses; these contain the whole virtues of the sage; the distilled water and essential oil, only its warmth and aromatic quality, without any thing of its roughness or bitterishness. Aqueous insusions of the leaves, with the addition of a little lemon juice, prove an useful diluting drink in febrile disorders, of an elegant colour, and sufficiently acceptable to the palate.

SALVIÆ SYLVESTRIS folia: Scorodotidis five scordii foliis salvia J. B. Woed sage; the leaves.

This grows wild in woods and hedges. In smell, taste, and medical virtues, it comes nearer to scordium than sage; it is less disagreeable than the former, but more so than the latter.

SAMBUCI folia, flores, bacca, cortex: Sambuci frudu in umbella nigro C. B. Common black berried elder; the leaves, bark, [E.] flowers, and berries [L. E]

This is a large shrub, frequent in hedges, it flowers in May, and ripens its fruits in September. The inner green bark of its trunk is gently cathartic: an infusion of it in wine, or the expressed juice, in the dose of half an ounce, or an ounce, is faid to purge moderately, and in small doses to prove an efficacious deobstruent, capable of promoting all the fluid fecretices. The young buds, or rudiments of the leaves, are strongly purgative, and act with fo much violence as to be deservedly accounted unsafe. The flowers are very different in quality: these have an agreeable aromatic flavour, which they give over in distillation with water, and impart, by infufion, to vinous and spirituous liquors. The berries have a sweetish, not unpleasant ta'e; nevertheles, eaten in substance, they offend the ftomach:

stomach: the expressed juice, infpissated to the consistence of a rob, proves an useful aperient medicine; it opens obstructions of the viscera, promotes the natural evacyations, and if continued for a length of time, does considerable service in sundry chronical disorders. It is observable, that this juice (which in its natural state is of a purplish colour) tinges vinous spirits of a deep red.

A rob prepared from the berries [L. E.] is used for making up the theriaca Edinensis and pectoral electary [E.] with the intention of which it excellently coincides. The flowers are an ingredient in the alexetereal and plague waters, the common decoction for glysters and fomentations, and the discutient cataplasm [E.] An oil of elder is prepared by boiling the flowers in oil olive [L.]; and an ointment, by boiling them in a mixture of oil and suet [L.] An ointment is made also from the leaves and bark [E.]

SAMPSUCHUS, vide Majo-

SANDARACHA, a fosfil composed of arenic and sulphur. Vide ARSENICUM.

SANGUIS DRACONIS [L.E.] Dragons blood, so called; a refin brought from the East Indies, either in oval drops, wrapped up in flag leaves, or in large masses, composed of smaller tears. The writers on the materia medica in general give the preference to the former, though the latter is not unfrequently of equal goodness: the fine dragons blood of either fort breaks smooth, free from any visible impurities, of a dark red colour, which changes upon being powdered into an elegant bright crimson. Several artificial compositions, coloured with the true dragons blood,

or Brazil wood, are !fometimes fold in the room of this commodity: some of these dissolve, like gums, in water; others crackle in the fire, without proving inflammable; whilft the genuine fanguis draconis readily melts and catches flame, and is not affect on by watery liquors. It totally diffolves in pure spirit, and tinges a large quantity of the menstruum of a deep red colour: it is likewise soluble in expressed oils, and gives them a red hue, less beautiful than that communicated by anchusa. This drug, in substance, has no sensible fmell or taste; when dissolved it discovers some degree of warmth and pungency. It is usually looked upon as a gentle aftringent, and sometimes directed as such in extemporaneous prescription, against seminal gleets, the fluor albus, and other fluxes: in the e cases, it produces the general effects of refinous bodies, lightly incrassating the fluids, and somewhat strengthening the folids. It is an ingredient in the styptic powder [L. E.] Locatelli's balfam [E] and strengthening plaster [L.]

SANICULÆ, seu Diapensiæ soliu: Saniculæ officinarum C. B. Sanicle; the leaves [E.]

This plant grows wild in woods and hedges, and flowers in May. The leaves have an herbaceous, roughish taste: they have long been celebrated for fanative virtues, both internally and externally: nevertheless their effects, in any intention, are not considerable enough to gain them a place in the present practice.

SANTALUM ALBUM. White faunders, a wood brought from the East-Indies, in billets, about the thickness of a man's leg, of a pale whitish colour. Greatest part

of it, as met with in the shops, has no smell or talte, or any sensible quality that can recommend it to the notice of the physician.

SANTALUM CITRINUM. [E.] Yellow saunders: a pale yellowish wood brought from the East Indies; of a pleasant smell, and a bitterish aromatic taste, accompanied with an agreeable kind of pun-This elegant wood might gency. undoubtedly be applied to valuable medical purposes, though at pre-Sent very rarely made use of: it is scarcely ever directed in extemporaneous preicription, and the only officinal composition in which it is an ingredient, is the confectio alkermes of the Edinburgh pharmacopæia: the London college have omitted it in their catalogue of fimples Distilled with water, it yields a fragrant effential oil, which thickens, in the cold, into the confiftence of a balfam. Digetted in pure fpirit, it imparts a rich vellow tincture; which being committed to distillation, the spirit arises, without bringing over any thing confiderable of the flavour of the faunders. The residuum contains the virtues of fix times its weight of the wood. Hoffman looks upon this extract as a medicine of fimilar virtues to ambergris; and recommends it as an excellent restorative in great debilities.

SANTALUM RUBRUM [L. E.] Red faunders; a wood brought from the East-Indies, in large billets, of a compact texture, a dull red, almost blackish colour on the outside, and a deep brighter red within. This wood has no manifest smell, and little or no taste. It has been commended as a mild astringent, and a corroborant of the nervous system; but these are qualities that belong only to the yellow fort.

The principal use of red saunders is as a colouring drug; in which intention it is employed in the balfamum Locatelli [L.] and spiritus lavendula compofitus [L. E.] It communicates a deep red to reetified spirit, but gives no tinge to aqueous liquors: a fmall quantity of the refin, extracted by means of spirit, tinges a large one of fresh spirit, of an elegant blood red. 'I here is scarce any oil, that of lavender excepted, to which it communicates its colour. Geoffroy, and others, take notice, that the Brazil woods are fometimes substituted to red faunders; and the college of Brussels are in doubt whether all that is fold among them for faunders is not really a wood of that kind: according to the account which they have given, their saunders is certainly the Brazil wood; the distinguishing character of which is, that it imparts its colour to common water.

SANTONICUM. [E.] Worm-feed; the produce of a plant of the worm wood or mugwort kind growing in the Levant.

It is a small, light, chaffy seed, composed as it were of a number of thin membraneous coats, of a yellowish colour, an unpleasant smell, and a very bitter taste. These seeds are celebrated for anthelmintic virtues (which they have in common with other bitters) and are sometimes taken in this intention, either along with melasses, or candied with sugar: their unpleasant taste renders the form of a powder or decoction inconvenient. They are not very often met with genuine in the shops.

SAPO DORUS [L.]: Sapo albus Hispanicus [E.] White Spanish soap.

SAPO MOLLIS [L.] Common fost soap.

SAPO NIGER, seu Melanosmegma [E.] Black soft soap.

Soap is composed of expressed vegetable oils, or animal sats, united with alkaline lixivia. The first fort, or white hard soap, is made with the finer kinds of oil olive; the common soft sort, with coarser oils, sat, tallow, or a mixture of all these; and the black (as is said) with train oil.

The purer hard foap is the only fort intended for internal use. This. triturated with oily, or refinous matters, renders them foluble in water, and hence becomes an useful ingredient in pills composed of refins, promoting their diffolution in the stomach, and union with the animal fluids: Boerhaave was a great admirer of foap; and in his private practice seldom prescribed any refinous pills without it; unless where an alkalescent, or putrid state of the juices forbad its use. From the same quality, soap likewife feems well fitted for disfolving fuch oily or unctuous matters, as it may meet with in the body, attenuating viscid juices, opening obstructions of the viscera, and deterging all the vessels it passes through. It is likewise a powerful menstruum for the human calculus: a folution of it in lime water is one of the strongest disfolvents that can be taken with fafety into the stomach; the virtue of this composition is considerably greater than the aggregate of the dissolving powers of the foap and lime water when unmixed. See the Edinburgh medical

The foft foaps are more penetrating and acrimonious than the hard. The only medical use of these is for some external purposes.

Hard foap gives name to an officinal plaster  $\{L.E.\}$  liniment  $\{L.\}$  and balsam  $\{E.\}$ : it is joined to opium, to render it more readily foluble in the stomach, in the pilulæ saponaceæ  $\{L.\}$  and to aloes in the pilulæ aloeticæ  $\{E.\}$  Soft soap is an ingredient in the milder common caustic  $\{L.\}$  and black soap in the anodyne plaster  $\{E.\}$ 

SAPONARIÆ folia, radix: Saponariæ majoris lævis C. B. Soapwort, or bruilewoit; the herb and root [E.]

This grows wild, though not very common, in low wet places, and by the fides of running waters; a double-flowered fort is frequent in our gardens. leaves have a bitter, not agreeable tafte; agitated with water, they raife a saponaceous froth, which is said to have nearly the same effects with folutions of foap itself in taking out spots from cloths. and the like. The roots tafte sweetish and somewhat pungent: and have a light smell like those of liquorice: digested in rectified spirit they yield a strong tincture, which lofes nothing of its tafte or flavour in being inspissated to the consistence of an extract. elegant root has not come much into practice among us, though it promises, from its sensible qualities, to be a medicine of confiderable utility; it is greatly esteemed by the German physicians as an aperient, corroborant, and fudorific: and preferred by the college of Wirtemberg, Stahl, Neumann, and others, to sarsaparilla.

SARCOCOLLA [L. E.] a concrete juice, brought from Persia and Arabia, in small, whitish, yellow grains, with a few of a reddish, and sometimes of a deep red colour, mixed with them; the whitest

whitest tears are preserved, as being the freshest: its taste is bitter, accompanied with a dull kind of sweetness. This drug dissolves in watery liquors, and appears to be chiefly of the gummy kind, with a small admixture of resinous matter. It is principally celebrated for conglutinating wounds and ulcers (whence its name σαρούκολλα, sless glue) a quality, which neither this, or any other drug, has a just title to. It is an ingredient in the pulvis e cerussa compositus [L.]

SARSAPARILLA [L. E.]: a root brought from the Spanish West-Indies. It consists of a great number of long strings hanging from one head: the long roots (the only part made use of) are about the thickness of a goose quill, or thicker, flexible, composed of fibres running their whole length, fo that they may be stript into pieces from one end to the other: they have a glutinous, bitterifh, not ungrateful tafte; and no smell. This root was first brought into Europe by the Spaniards, about the year 1563, with the character of a specific for the cure of the lues venerea, which made its appearance a little before that time, and likewise of several obstinate chronic disorders. Whatever good effects it might have produced in the warmer climates, it proved unsuccessful in this; insomuch that many have denied it to have any virtue at all. It appears however, from experience, that though greatly unequal to the character which it bore at first, it is in some cases of considerable use as a fudorific, where more acrid medicines are improper. The best preparations are a decoction and extract made with water: a decoction of half an ounce of the root, or a drachmof the extract, which is equivalent thereto, may be taken for a dole.

SASSAFRAS [L. E.]: the root of a large American tree (arbor ex Florida ficulneo folio C. B.) brought to us in long straight pieces, very light, and of a spongy texture, covered with a rough fungous bark; outwardly of an all colour, inwardly of the colour of rufty iron. It has a fragrant smell, and a sweetish aromatic, subacrid take: the bark taftes much stronger than any other part; and the fmall twigs stronger than the large pieces. As to the virtues of this root, it is a warm aperient and corroborant; and frequently employed with good fuccess, for purifying and sweetening the blood and juices. For these purposes, insufons made from the rasped root or bark, may be drank as tea. In some constitutions, these liquors, by their fragrance, are apt, on first taking them, togasfect the head: in fuch cases they may be advantageously freed from their flavour by boiling; a decoction of fassafras, boiled down to the confillence of an extract, proves simply bitterish and subastringent. Hoffman assures us, that he has frequently given this extract, to the quantity of a scruple at a time, with remarkable success, for strengthening the tone of the viscera in cachexies: as also in the decline of intermittent fevers, and in hypocondriacal spasms. Sassafras vields in distillation an extremely fragrant oil, of a penetrating pungent tafte, fo ponderous (notwithstanding the lightness of the drug itself) as to fink in water. Rectified fpirit extracts the whole taste and smell of fassafras: and elevates nothing in evaporation; hence the spirituous extract proves the most elegant and efficacious preparation, as containing the virtue of the root entire.

The only officinal preparation of fassaffast is the essential oil [L. E.] The salfast itself is an ingredient

ш

in the decoction of the woods [E.] and the compound lime waters [L. E.] and the oil in the clixir guaiacinum [E.]

SATUREIÆ folia: Satureiæ hortensis, siwe cunilæ satiwæ Plinii C. B. Summer savory; the leaves [E.]

This herb is raifed annually in gardens for culinary purposes. It is a very pungent warm aromatic; and affords in distillation with water a subtile essential oil, of a penetrating smell, and very hot, acrid taste. It yields little of its virtues by infusion to aqueous liquors: rectified spirit extracts the whole of its taste and smell, and elevates nothing in distillation.

SATYRII MARIS radix: Orchidis morionis maris foliis maculatis C. B. Orchis; the root [E.]

This plant is frequent in shady places and moist meadows: each plant has two oval roots, of a whitish colour, a viscid sweetish taste, and a faint unpleasant smell. They abound with a glutinous slimy juice. with regard to their virtues, like other mucilaginous vegetables, they thicken the thin ferous humours and defend the folids from their acrimony: they have also been celebrated, though on no very good foundation, for an aleptic and aphrodifiac virtues; and frequently made use of in these intentions. in the Edinburgh pharmacopæia, the root is directed to be candied.

SALEP [E.] a celebrated restorative among the Turks, is probably the prepared root of certain plants of the orchis kind. This drug, as sometimes brought to us, is in oval pieces, of a yellowish white colour, somewhat clear and pellucid, very hard, and almost horny, of little or no smell, and tasting

like gum tragacanth. Satyrion root, boiled in water, freed from the skin, and afterwards suspended in the air to dry, gains exactly the same appearance; the roots thus prepared, dissolve in boiling water into a mucilage. Geosfroy, who first communicated this preparation of orchis, recommends it in confumptions, in bilious dysenteries, and disorders of the breast proceeding from an acrimony of the juices.

SAXIFRAGÆ ALBÆ folia, radix: Suxifragæ albæ radice granulosa J. B. White flowered suxifrage; the leaves, and the roots [E.] which last are improperly called (from their consisting of little grains) seeds.

SAXIFRAGÆ VULGARIS folia femen: Sefelis pratenfis nostratis Raii. Meadow saxifrage; the leaves and seeds.

These herbs grow wild, the first in dry sandy ground, the second in fields and meadows: the first is not very common, and hence its leaves and roots have been generally supplied by the leaves and feeds of the second. Neither of them are at present in much esteem, notwithstanding the aperient, duretic and lithontriptic virtues formerly attributed to them.

SCABIOSÆ folia: Scabiosæ majoris communioris birsutæ, folio lacimiato Raii. Scabious; the leaves.

This is a rough hairy plant growing wild in pasture grounds; of a nauseous bitterish taste. It stands recommended as an aperient, sudorisic, and expectorant; but the prefent practice has little dependance on it.

SCAMMONIUM [L. E.] Scammony; a concrete juice extracted from the roots of a large climbing climbing plant growing in the Asiatic Turkey. The best comes from Aleppo, in light, spongy masses, easily friable, of a shining ash colour verging to black; when powdered, of a light grey or whitish colour: an inferior fort is brought from Smyrna, in more compact, ponderous pieces, of a darker colour, and full of fand and other impurities. This juice is chiefly of the refinous kind : rectified spirit dissolves five ounces out of fix, the remainder is a mucilaginous subflance mixed with drof: proof spirit totally dissolves it, the impurities only being left. It has a faint unpleasant smell; and a bitterish, somewhat acrimonious taste.

Scammony is an efficacious and firong purgative. Some have condemned it as unsafe, and laid sundry ill qualities to its charge; the principal of which is, that its operation is uncertain, a full dose proving sometimes ineffectual, whilst at others a much smaller one occafions dangerous hypercatharfes. This difference however is owing entirely to the different circumstances of the patient, and not to any ill quality, or irregularity of operation, of the medicine: where the intestines are lined with an excessive load of mucus, the scammony passes through, without exerting itself upon them; where the natural mucus is deficient, a small dose of this or any other resinous cathartic, irritates and inflames. Many have endeavoured to abatethe force of this drug, and correct its imaginary virulence, by expoling it to the fune of fulphur. dissolving it in acid juices, and the like: but this could do no more than destroy as it were a part of the medicine, without making any alteration in the rest. Scammony in tubitance, judiciously managed, flands not in need of any corrector:

if triturated with sugar or with almonds, as we have formerly recommended for other refinous purgatives, it becomes sufficiently safe and mild in operation. It may likewise be conveniently dissolved. by trituration, in a strong decoction of liquorice, and then poured off from the feces: the college of Wirtemberg assures us, that by this treatment it becomes middly purgative, without being attended with gripes, or other inconveniencies; and that it likewise proves inoffenfive to the palate. The common dose of scammony is from three to twelve grains.

Scammony gives name to an officinal compound powder and electary [L.] and is an ingredient in the compound powder of sena, the cathartic extract, the coloquintida pills, mercurial pills [L.] and purgative deobstruent pills [E.]

SCHŒNANTHUS, vide Juncus odoratus.

SCILLÆ radix: Scillæ radice alba C. B. wel Scillæ vulyaris radice rubra C. B. The squill, or seaonion; its root [E.]

This is a fort of onion, growing spontaneously upon dry sandy shores in Spain and the Levant, from whence the root is anually brought into Europe. It should be chosen plump, found, fresh, and full of a clammy juice; fome have preferred the red fort, others the white, though neither deserves the preference to the other; the only difference perceivable betwixt them, is that of the colour; and hence the college allow both to be used promiseuously. This root is to the taste very nauseous, intensely bitter and acrimonious: much handled, it exulcerates the skin. With regard to its medical virtues, it powerfully stimulates the solids,

and attenuates viscid juices; and by these qualities, promotes expectoration, urine, and (if the patient is kept warm) sweat: if the dose is considerable, it proves emetic, and sometimes purgative. The principal use of this medicine is where the primæ viæ abound with mucous matter, and the lungs are oppressed by tenacious phlegm. Dr. Wagner (in his clinical observations)recommends it given along with nitre, in hydropical swellings, and in the nephritis; and mentions several cures which he performed. by giving from four to ten grains of the powder for a dofe, mixed with a double quantity of nitre: he fays, that thus managed, it almost always operates as a diureretic, though fometimes it vomits or purges. The most commodious form for the taking of fquills, unless when designed as an emetic, is that of a bolus or pill: liquid forms are to most people too offenfive, though these may be rendered less disagreeable both to the palate and stomach, by the addition of aromatic distilled waters. This root yields the whole of its virtues, both to aqueous and vinous menstrua, and likewise to vegetable acids. Its officinal preparations are, baked squills [L.] and the baked squills made into troches [L.] defigned as an ingredient in theriaca [L.]: dried squills [L.] a fyrup, vinegar, oxymel [L. E.] and pills  $\{E.\}$ 

SCINCORUM ventres [L] The belly of the skink; a kind of small lizard, brought dry from Egypt. It stands recommended as a great restorative: whatever virtues it may have as used fresh by the Egyptians, it has none as it comes to us, and serves to uselessly increase the articles of the mithridate.

SCLAREA, vide Horminum.

SCOLOPENDRIUM, vide Lin-

SCORDII folia: Chamædryos palufiris canescentis Tourn. Watergermander; the leaves [L. E.]

This is a small, somewhat hairy plant, growing wild in some parts of England, though not very common; the shops are generally supplied from gardens. It has a bitter tafte, and a strong disagreeable ſmell. Scordium is of no great esteem in the present practice, notwithstanding the deobstruent, diuretic, and fudorific virtues which it was formerly celebrated for. It enters the mithridate, theriaca, and cataplasm of cummin seed [L.]: and gives name to two compound powders and an electary [L.] though not the most valuable of their ingredients.

SCORZONER Æ radix: Scorzoneræ latifoliæ finuat r C. B. Vipers grafs; the root [E.]

Scorzonera is met with only in gardens. The roots abound with a milky juice, of a bitterish sub-acrid taste; and hence may be of some service, for strengthening the tone of the viscera, and promoting the sluid secretions. They were formerly celebrated as alexipharmacs, and for throwing out the measles and small-pox; but have now almost entirely lost their character.

SCROPHULARIÆ VULGA-RIS folia, radix: Scrophulariæ nodosæ fætidæ C. B. Fig-wort; the leaves and root [E.]

This herb grows wild in woods and hedges: the roots are of a white colour, full of little knobs or protuberances on the furface; this appearance gained it formerly some repute against scrophulous disorders and the piles; and from from hence it received its name: but modern practitioners expect no fuch virtues from it. It has a faint unpleasant smell, and a somewhat bitter disagreeable taste.

SCROPHULARIÆ AQUATICÆ MAJORIS, folia: Scrophulariæ maximæ radice fibrofa J.B. Greater water figwort; the leaves

This is a large plant, met with chiefly on the sides of rivers. The leaves have a bitter taste and an ungrateful smell: they are principally celebrated, though on no very good grounds, as a corrector of sena. See the article Sena.

SEBESTEN: Mixa five Sebesten J. B. A fort of plum, brought half dried from the East-Indies: it is of a dark or blackish brown colour, with whitishor ash coloured cups; the steff sticks close to the stone, which contains sometimes one and sometimes two kernals. This fruit has a sweet, very glutinous taste; and hence has been employed for softening acrimonious humours, in some kinds of hoarseness, and in coughs from thin sharp destuxions: at present, it is not often met with in the shops.

SECALE: Secale bybernum well majus C. B. Rye; the feeds. These are little regarded as an article of the materia medica: as food, they are accounted more detergent than most other kinds of grain.

SEDI MAJORIS, feu Semperwiwi majoris folia: Sedi majoris vulgaris C. B. Greater house-leek; the leaves [E.]

This is a low fleshy-leaved plant growing on old walls, and on the teps of houses. It stands recommended as a cooler, though its sen-

fible qualities discover no great foundation for any virtue of this kind; the taste is herbaceous, with a flight degree of pungency. It is remarkable of this plant, that its juice purified by filtration (when it appears of a dilute yellowish colour) upon the admixture of an equal quantity of rectified spirit of wine, forms a beautiful white, light coagulum, like the finer kinds of pomatum: this proves extremely volatile; freed from the aqueous phlegm, and exposed to the air, it in a very little time totally exhales. From hence it is concluded (in the medicor. Silesiac satyræ) that houseleek contains a volatile alkaline falt: but there are many substances besides these falts which coagulate with spirit of wine.

SEMPERVIVUM, vide Se-

SENA [L. E] the leaves of a shrubby plant (sena Alexandrina foliis acutis C. B.) cultivated in Perfia, Syria, and Arabia; from whence they are brought, dried and picked from the stalks, to Alexandria in Egypt; and thence imported into Europe. They are of an oblong figure, sharp pointed at the ends, about a quarter of an inch broad, and not a full inch in length, of a lively yellowish green colour, afaint not very disagreeable smell, and a subacrid, bitterish, nauseous talles Some inferior forts are brought from Tripoli and other places: these may easily be distinguished by their being either narrower, longer. and sharper pointed; or larger, broader, and round pointed, with fmall prominent veins; or large and obtuse, of a fresh green colour, without any yellow cast.

Sena is a very useful cathartic, operating middly, and yet effectually; and, if judiciously dosed and managed,

rarely occasioning the ill consequences which too frequently follow the exhibition of the stronger purges. The only inconveniencies complained of in this drug are, its being apt to gripe, and its nause-The griping quality ous flavour. depends upon a refinous substance, which, like the other bodies of this class, is naturally disposed to adhere to the coats of the intestines: the more this refin is divided by fuch matters as take off its tenacity, the less adhesive, and consequently the less irritating and griping it will prove; and the less it is divided, the more griping: hence sena given by itself, or infusions made in a very fmall quantity of fluid, gripe feverely, and purge less than when diluted by a large portion of suitable menstruum, or divided by mixing the infusion with oily emulsions. The ill flavour of this drug is faid to be abated by the greater water figwort : but we cannot conceive that this plant, whose smell is manifestly fetid, and its tafte nauseous and bitter, can at all improve those of sena: others recommend bohea tea, though neither has this any confiderable ef-The fmell of fena refides in its more volatile parts, and may be discharged by lightly boiling infofions of it made in water: the liquor thus freed from the peculiar flavour of the sena, may be easily rendered grateful to the tafte, by the addition of any proper aromatic tincture or distilled water. The colleges, both of London and Edinburgh, have given several very elegant infusions of this drug (which may be seen in part III. chap. iii.) as also spirituous tinctures [L. E.] compound powders [L. E.] and a fyrup [E.] The dose of sena in fubstance is from a scruple to a drachm, in infusion from one to three or four drachms.

It has been customary to reject the pedicles of the leaves of sena as of little or no use: Geoffroy however observes, that they are not much inserior in efficacy to the leaves themselves. The pods, or seed-vessels, met with among the seed-vessels, met with among the lege of Brussels preferred to the leaves: they are less apt to gripe, but proportionably less purgative.

## SENECIO, vide ERIGERUM.

SENEKA [E.] Senecka, rattle-snake root; the root of a species of polygola, which grows spontaneously in Virginia, and bears the winters of our own climate. This root is usually about the thickness of the little singer, variously bent and contorted, and appears as if composed of joints, whence it is supposed to resemble the tail of the animal whose name it bears: a kind of membranous margin runs on each side, the whole length of the root. Its taste is at first acid, afterwards very hot and pungent.

This root is not at present much known in the shops. The Senegaro Indians are faid to prevent the fatal effects which follow from the bite of the rattle-snake, by giving it internally, and applying it externally to the wound. It has of late been strongly recommended in pleurifies, peripneumonies, and other inflammatory distempers: in these cases, Lemery, du Hamel, and Juffieu, experienced its good success (see the French memoirs for the years 1738, 1739.) Its more immediate effects are those of a diuretic, diaphoretic, and cathartic: fometimes it proves emetic: the two last operations may be occafionally prevented, by giving the root in small doses, along with aromatic simple waters, as that of cinnamon. The usual dose of the powder is thirty grains or more.

Some have likewife employed this root in hydropic cases, and not without success: Bouvart (in the memoirs above-mentioned, 1744) relates examples of its occasioning a plentiful evacuation by flool, urine, and perspiration, and by this means removing the disease, after the common diurerics and hydragogues had failed: where this medicine operates as a cathartic, it generally proves successful: if it acts by liquifying the blood and juices, without occasioning a due discharge, it should either be abstained from, or assisted by proper additions.

SERICUM et folliculi bombycis. Silk and filkworms bags These are scarce ever made use of for any medicinal purposes. In their crude state they are certainly very insignificant: though if burnt in a clote vessel, after the same manner as sponge, they would probably prove a medicine of similar, and perhaps of superior virtue. They yield a larger quantity of volatile salt, than any other animal substance I know of.

SERPENTARIA VIRGINI-ANA [L. E] Virginian fnakeroot; the root of a species of aristolochia, growing in Virginia and Carolina.

It is a small, light, bushy root, consisting of a number of strings or sibres, matted together, issuing from one common head; of a brownish colour on the outside, and paler or yellowish within. It has an aromatic smell, like that of valerian, but more agreeable; and a warm, bitterish, pungent taste. This root is a warm diaphoretic and diuretic: it has been greatly celebrated as an alexiphar-

mac, and esteemed one of the principal remedies in malignant fevers and epid mic diseases. In these intentions, it is given in subflance from ten to thirty grains, and in infusion to a drachm or two. Both watery and spirituous menstrua extract its virtues by infusion, and elevate some share of its flavour in distillation: along with the water a small portion of essential oil arises. A spirituous tincture [L. E] and compound decoction [E.] of it are directed as officinals it enters also the cephalic tincture, compound tincture of Peruvian bark, sodorific tincure, tinctura facra, ilomachic elixir. theriaca Edinensis [E] and cataplasm of cummin seed [L.]

SERPYLLI folia: Serpilli valgaris minoris C. B. Mother of thyme; the herb [E.]

This is a small creeping plant, common on heaths and dry paflure grounds. Its taste, smell, and medical virtues are similar to those of thyme, but weaker.

SFSAMI femen: Digitalis orientalis fesam dida Tourn. the seeds called oily purging grain

This plant is cultivated in the eastern countries, from whence the feeds are brought to us. They very properly deserve the name of oily, as they yield upon expression a larger quantity of oil, than almost any other known vegetable. The appellation purging, they have no title to; among the Indians, they are said to be used as food.

SESELIS VULGARIS femen: Ligustici quod sesselli officinarum C. B. Common hartwort; the seeds [L E.]

SESELIS MASSILIENSIS femen: Seselis Massiliensis ferulæ folio. C. R. C. B. Italian hartwort; the feeds [L. E.]

These plants grow spontaneously in the warmer climates; amongst us they are met with only in the gardens of the curious. The seeds and roots of both sorts have an agreeable aromatic smell and taste; and in this light might be occasionally employed, though at present they are in disuse: being scarcely otherwise regarded than as the seeds of the first fort are an ingredient in mithridate and theriaca.

SESELI PRATENSE, vide SAXIFRAGA VULGARIS.

SIGILLI SALOMONIS, seu Polygonati radix: Polygonati latifolii vulgaris C. B. Solomon's feal; the root [E]

This grows wild in woods, but is not very common: the root has feveral joints, with some flat circular depressions, supposed to refemble the stamp of a seal. It has a sweetish mucilaginous taste. As to its virtues, practitioners do not now expect any considerable ones from it, and pay very little regard to the vulnerary qualities which it was formerly celebrated for.

SILER MONTANUM, vide Seseli vulgare.

SIMAROUBA [E.]: a bark with pieces of the wood adhering to it, brought from Guiana, in long tough pieces, of a pale yellowish colour, and a pretty strong bitter taste. It has lately come into esteem in dysenteric sluxes: a decoction of halfadrachm is given for a dose, and repeated at intervals of three or four hours.

SINAPIS semen: Sinapis rapi

folio C. B. Mustard; the seeds [L. E]

This plant is sometimes found wild, but for culinary and medicinal uses is cultivated in gardens. Mustard, by its acrimony and pungency, slimulates the solids, and attenuates viscid juices; and hence stands deservedly recommended for exciting appetite, promoting digestion, increasing the fluid fecretions, and for the other purposes of the acrid plants called antiscorbutic. It imparts its taste and fmell in perfection to acqueous liquors, whilst rectified spirit extracts extremely little of either: the whole of the pungency arises with water in distillation. Committed to the press, it yields a confiderable quantity of a foft infipid oil, perfectly void of acrimony: the cake left after the expression is more pungent than the mustard was at first. The oil is directed as an officinal [L. E.] These seeds are sometimes employed externally as a stimulant; and give name to a composition for this intention in the Edinburgh dispensatory.

SISON, vide Amomum vulgare.

SMYRNIUM, vide HIPPOSE-LINUM.

SOLANI VULGARIS folia: Solani hortenfis seu vulgaris J. B. Common nightshade; the leaves.

SOLANI LETHALIS, feu Belladonnæ folia: Solani melanocerafi C. B. Deadly nightshade; the leaves.

These plants grow wild; the first in cultivated grounds, the second in shady waste ones. They have both been supposed cooling and discutient in external applications, and poisonous when taken

L2 inter-

internally. Late experience has shewn, that an infusion of half a grain or a grain of the dried leaves of either may be taken with safety, and that in many cases the dose may be increased by degrees to five or fix grains: that they generally occasion some considerable evacuation, and fometimes, especially in the larger of the above doses, alarming nervous symptoms, which however cease with the operation of the medicine. It has been expected, that a cautious use of these very active plants would afford relief in some obstinate disorders: but though in some instances they promised great benefit, the general event of these trials has not been very favourable. The Edinburgh college, who retained these plants at the preceding revisal of their Pharmacopæia in the year 1744, have, at the Jate one in 1756, rejected them

SOLANUM LIGNOSUM, vide Dulcamara.

SOLDANELLA, vide Brassica marina.

SOPHIÆ CHIRURGORUM
femen: Nasturtii sylvestris tenuissime
divis C. B. Flixweed; the seeds.
This plant had formerly a great
character as a vulnerary, and for
stopping fluxes; but its effects have
not been considerable enough to
continue it in practice.

SORBI SYLVESTRIS cortex: Sorbi sylvesiris foliis domesticæ similis C. B. Wild service, or quicken tree; its bark [E.]

The bark of this tree has a faint unpleasant smell, and a bitter taste, from whence we may presume that it is not destitute of medical virtues, though what par-

ticular uses it has been applied to I cannot learn.

SPERMA CETI [L. E.] improperly fo called; an unctuous flaky substance, of a snowy whiteneis, a fost butyraceous taste, without any remarkable smell; faid to be prepared from the fat of the brain of the whale, by boiling and purifying it with alkaline lixi-The virtues of this concrete are those of a mild emollient: it is of confiderable use in pains and erofions of the intestines, in coughs proceeding from thin sharp defluxions, and in general in all cases where the folids require to be relaxed, or acrimonious humours to be foftened. For external purposes, it readily dissolves in oils; and for internal ones, may be united with aqueous liquors into the form of an emulsion, by the mediation of almonds, gums, or yolk of an egg. Sugar does not render it perfectly miscible with water; and alkalies, which change other oils and fats into foap, have little effect upon sperma ceti. This drug ought to be kept very closely from the air, otherwise its white colour foon changes into a yellow; and its mild unctuous taste, into a rancid and offensive one. After it has suffered this disagreeable alteration, both the colour and quality may be recovered again by fleeping it in alkaline liquors, or steeping it in a sufficient quantity of spirit of wine.

SPICA VULGARIS, vide La-VENDULA ANGUSTIPOLIA.

SPICA NARDI, vide Nardus Indica.

SPINÆ ALBÆ, seu Oxyacanthæ vulgaris, stores, baccæ: Mespili apii foliis, sylvestris, syinosæ sive oxyacanthæ cantbæ C. B. White thorn, or hawthorn; its flowers and berries [E.]

The reputation which these formerly had, in nephritic and calculous complaints, continues them in most catalogues of the materia medica, though common practice has long rejected them as insignificant. The slowers have a very pleasant smell: the berries are mucilaginous, and sweetish.

SPINÆ CERVINÆ baccæ: Rbamni cathartici C. B. Buckthorn; the berries [L. E.]

This tree, or bush, is common in hedges: it flowers in June, and ripens its fruit in September, or the beginning of October. In our markets, the fruit of some other trees, as the frangula or black berry-bearing alder, and the cornus famina or dog-berry tree, have of late years been frequently mixed with, or substituted for, those of buckthorn: this abuse may be discovered by opening the berries: those of buckthorn have almost always four feeds, the berries of the frangula two, and those of the cornus famina only one. Buckthorn berries, bruised on white paper, give it a green tincure, which the others do not. Those who sell the juice to the apothecaries, are faid to mix with it a Jarge proportion of water.

Buckthorn berries have a faint disagreeable smell, and a nauseous bitter taste. They have long been in considerable esteem as cathartics; and celebrated in dropsies, rheumatisms, and even in the gout; though, in these cases, they have no advantage above other purgatives, and are more offensive, and operate more churlishly, than many which the shops are furnished with: they generally occasion gripes, sickness, dry the mouth and throat, and leave a thirst of

long duration. The dose is about twenty of the fresh berries in sub-stance, and twice or thrice this number in decoction, an ounce of the expressed juice, or a drachm of the dried berries. A syrup prepared from the juice is kept in the shops; in this preparation, the nauseous flavour of the buckthorn is somewhat alleviated by sugar, and the addition of aromatics.

SPIRITUS VINOSUS REC-TIFICATUS. Reclified spirit of wine; " a spirit distilled from wine " or other fermented liquors, pu-" rified as much as possible from " its fetid smell, and the phlegm " that arises with it in the first di-" stillation." [L.] This purification is effected, by repeating the distillation in a very gentle heat, with certain additions, to keep down the phlegm and the gross oil in which the ill flavour resides (see Part III. chap. v.) These spirits, whatever vegetable subjects they have been produced from, are, when perfectly pure, one and the same. They have a hot pungent tafte, without any particular flavour: they readily catch flame, and burn entirely away, without leaving any marks of an aqueous moisture behind: distilled by a heat less than that of boiling water, they totally arise. the last runnings proving as flavourless and inflammable as the first: they dissolve essential vegetable oils and resins into an uniform transparent fluid. These spirits are the lightest of almost all known liquors: expressed oils, which swim upon water, fink in these to the bottom: a measure which contains ten ounces by weight of water, will hold little more than eight and a quarter of pure spirit.

The uses of vinous spirits, as menstrua for the virtues of other Q 2 medicines,

medicines, we shall see hereaster, and in this place confider only their own. Pure spirit coagulates all the fluids of animal bodies, except urine, and hardens the solid parts. Applied externally, it strengthens the vessels, thickens the juices in them, and thus powerfully restrains hæmorrhages. It inflantly contracts the extremities of the nerves it touches, and deprives them of sense and motion; by this means easing them of pain, but at the same time destroying their use. Hence employing spirituous liquors in fomentations (notwithstanding the specious titles of vivifying, heating, restoring mobility, resolving, dissipating, and the like, usually attributed to them) may sometimes be attended with unhappy consequences. These liquors, received undiluted into the flomach, produce the same effects, thickening the fluid, and contracting all the solid parts which they touch, and destroying, at least for a time, their use and office: if the quantity is confiderable, a palfy or apoplexy follows, which end in death. Taken in small quantity, and duly diluted, they brace up the fibres, raise the spirits, and promote agility: if farther continued, the senses are disordered, voluntary motion destroyed, and at length the same inconveniences brought on as before. Vinous spirits, therefore, in fmall doses, and properly diluted, may be applied to useful purposes in the cure of diseases; whilst in larger ones, or if their use is long continued, they act as a poison of a particular kind.

SPIRITUS VINOSUS TE-NUIOR. Proof spirit: " the " same spirit, containing an ad-" mixture of an equal quantity of " water: the best proof spirit is " that distilled from French wine;

but for common uses may be employed the spirit drawn from melasses, or the syrupy matter " that runs from fugar in the pu-" rification, commonly called me-" lasses spirit." [L.] The spirits usually met with under the name of proof, are those distilled from different fermented liquors, freed from their phlegm and ill flavour only to a certain degree. purity, with regard to flavour, may be easily judged from the taste, especially if the spirit be first duly diluted. It were to be wished, that we had a certain standard with regard to their strength, or the quantity of water contained in them; a circumftance which greatly influences fundry medicinal preparations, particularly the tinctures: for as pure spirit dissolves the refin and volatile oil, and water only the gummy and faline parts of vegetables, it is evident that a variation in the proportions wherein these are mixed, will vary the dissolving powder of the menstruum, and confequently the virtue of the preparation. The common methods of estimating the quantity of phlegm contained in these spirits, are liable to uncertainty: it should therefore feem necessary, for the nicer purposes, and where a perfectly flavourless proof spirit is required, to make use of the pure rectified spirit, mixed with a certain determined proportion of water; equal quantities of these liquors, whether taken by weight or measure, compose a spirit somewhat weaker that what has been generally looked upon as proof; the exact proportions are, one hundred parts by weight of pure spirit, and eighty-fix of water.

SPONGIA [L. E.] Sponge; a foft, light, very porous and compressible substance, readily imbing

bibing water, and distending thereby. It is found adhering to rocks, particularly in the Mediterranean fea, about the islands of the Archipelago. It is generally supposed to be a vegetable production: nevertheless some observations, lately made by Juffieu, give room to suspect that it is of animal origin. Chemical experiments favour this supposition; analysed, it yields the fame principles with animal substances in general: the volatile salt is in larger quantity than I have obtained from any animal matter, except the bags of the filkworm. On this falt which is generated by fire, feem to depend the virtues of the officinal fpongia usta [L.] (See Part III. chap. i.) Crude sponge, from its property of imbibing and distending by moinure, is fometimes made use of as a tent for dilating wounds and ulcers.

STANNUM [L. E] Tin is the lightest and easiest of susion of all the metals. Heated, it becomes fo brittle as to fall in pieces by a blow; and by agitation (when just ready-to melt) into a powder; hence the officinal method of pulverifing this metal, to be described in its place. The proper menstroum of tin is the marine acid, or aqua regis : vegetable acids likewise dissolve it in considerable quantity, though it has long been supposed not to be at all soluble in them, unless previously well calcined.

With regard to the virtues of this metal, it was formerly accounted a specific in disorders of the uterns and lungs; a calx of tin and antimony is still retained in some dispensatories, under the name of an antihectic: but these are virtues, to which it certainly has little claim. It has of late been celebrated, on better foundation, as an anthelmintic; and said to destroy some kinds of worms which elude the force of many other medicines: possibly the cause of this effect may be very different from what may be fulpected, an admixture of a portion of arfenic.

Tin has a strong affinity with arfenic; infomuch that when once united therewith, the arfenic, notwithstanding its volatility in other circumstances, cannot be totally expelled either by flow calcination, or by a vehement fire. most all the ores of tin contain more or less of this poisonous mineral, which is not entirely feparable in the common processes by which the ores are run down, or the metal farther purified. ings of tin held in the flame of a candle, emit a thick fume, smelling of garlic; which smell is universally held, in mineral substances. to be a certain criterion of arsenic. Henckel has discovered a method of separating actual arlenic from tin: this is effected by folution in agua regis and erystallization: Mr. Margraff has (in a late volume of the Berlin memoirs) given a farther account of this process; and relates, that from the tins usually reputed pure, he has obtained one-eighth their weight of crystals of arsenic. the preparations of tin, see the fecond part of this work.

STAPHIDISAGRIÆ femen : Delphinii platoni folio Tourn. Stavesacre; the feeds [E.]

These are large rough seeds, of an irregularly triangular figure, of a blackish colour on the outside, and yellowish or whitish within: they are usually brought from Italy; the plant is not very common in this country, though it bears our severest colds. They have a disagreeable smell. and a very nauseous bitterish, burning taste. Stavesacre was employed by the ancients as a cathartic; but it operates with so much violence both upwards and downwards, that its internal use has been, among the generality of practitioners, for some time laid It is chiefly employed, in aside. external applications, for some kinds of cutaneous eruptions, and for destroying lice and other insects: insomuch, that it has from this virtue received its name, in different languages, berba pedicularis, berbe aux poux, lauskraut, lou fewort.

STERCUS anseris, canis, columbæ, equi, ovis, pavonis, porci. The dung of the goose, dog, pigeon, horse, sheep, peacock, hog. These sulforme medicines, which nothing but the most fantastic visionaries could have introduced, are now expunged from practice, and from our pharmacoposias.

## STIBIUM, vide Antimonium.

STECHAS: Stæchas purpurea C. B. Arabian stechas, or French lavender slowers [L. E.]

This is a shrubby plant, con-

fiderably smaller than the common lavender: the flowery heads are brought from Italy and the southern parts of France; they are very apt to grow mouldy in the passage, and even when they escape this inconvenience, are generally much inferior to those raised in our gardens. The best stechas which we receive from abroad, has no great smell or tasse: Pomet assirms, that

fuch as the shops of Paris are

supplied with, is entirely destitute

of both; whilst that of our own

growth, either whilst fresh or when carefully dried, has a very fragrant smell, and a warm, aromatic, bitterish, subacrid taste; distilled with water, it yields a considerable quantity of a fragrant essential oil; to rectified spirit it imparts a strong tincture, which imposses a strong tincture, which in prissed proves an elegant aromatic extract. This aromatic plant is rarely met with in prescription; the only officinal compositions which it is admitted into, are the mithridate and theriaca.

There is another plant called stechas, which, from the beauty and durability of its flowers, has of late years had a place in our gardens, and whose aromatic qualities render worthy of one in the shops; this is the elicbrysum seu stachas citrina latiore folio C. B. golden stechas, goldilocks, or yellow cassidony: its slowers stand in umbels on the tops of the branches; they are of a deep shining yellow colour, which they retain in perfection for many years; their fmell is fragrant and agreeable, fomewhat of the musky kind; their taste warm, purgent, and subastringent; they impart their flavour to water in distillation, and by infusion to rectified spirit.

STYRAX CALAMITA [L. E.] Storax; an odoriferous refinous substance, exuding, in the warmer climates, from a tree called by C. Bauhine styrax folio mali cotonei. It has been customary to distinguish three forts of storax, though only one is usually met with in the shops.

the cane, so called from its having been formerly brought inclosed in reeds from Pamphylia: it is either in small distinct tears, of a whitish or reddish colour, or in larger masses composed of such.

2. Storax

2. Storax in the lump, or red storax. This is in masses of an uniform texture, and yellowish red or brownish colour, though sometimes likewise interspersed with a few whitish grains. Of this fort there has been some lately to be met with in the shops, under the name of storax in the tear.

3. The common florax of the shops is in large masses, considerably lighter and less compact than the foregoing. It appears, upon examination, to be composed of a fine refinous juice, mixed with a quantity of faw-dust. For what purpose this addition is made, I shall not here inquire; observing only, that it can scarce be supposed to be done with any fraudulent view, fince the faw-dust appears at fight. This common storax is much less esteemed than the two first forts; though, when freed from the woody matter, it proves superior, in point of fragrancy, Rectified spito either of them. rit, the common menstruum of refins, dissolves the storax, leaving the wood behind: nor does this tincture lose considerably of its valuable parts, in being inspissated to a folid confistence; whilst aqueous liquors elevate almost all the fragrancy of the florax.

Storax is one of the most agreeable of the odoriferous resins, and may be exhibited to great advantage in languors and debilities of the nervous system; it is not, however, much used in common practice, unless as an ingredient in the traumatic balam, the compound powder and electary of scordium, the storax pill, confectio Paulina, mithridate, and theriaca [L].

STYRAX LIQUIDA [E.] Liquid storax. What is most commonly met with under this name, is a soft resinous substance, of a

grey colour, a weak smell, similar to that of the foregoing folid ftorax: it is supposed to be compounded of solid storax, refin. wine, and oil, beaten up together into a proper consistence. genuine liquid storax, according to Petiver's account (Phil. Transact. No 313.) is obtained from a tree growing in the island Cobros in the Red sea. The preparers of this commodity yearly clear off the bark of the tree, and boil it in fea-water to the confistence of birdlime: the refinous matter, which floats upon the surface, is taken off, liquefied again in boiling water, and passed through a strainer. The purer part which passes through, and the more impure which remains on the strainer, and contains a confiderable portion of the substance of the bark, are both fent to Mocca, from whence they are fometimes, though very rarely, brought to us. The first is of the confidence of honey, tenacious, of a reddish or ash brown colour, an acrid uncluous taste, approaching in smell to the folid florax, but so strong as to be difagreeable: the other is full of woody matter, and much weaker in smell.

Liquid storax is among us scarce ever made use of in medicine, and not often found in the shops: hence the London college has expunged it from the catalogue of officinals; that of Edinburgh employs it as an ingsedient in the mercurial plaster.

SUBER: cortex Suberis latifolis perpetus virentis C. B. Cork, a fort of evergreen oak, growing in the warmer parts of Europe; its bark. This has been by some accounted astringent, and recommended as such in dysenteries and other fluxes; but modern

dern practice applies to it no fuch uses, and expects from it no virtues of any kind.

It may here be proper to take notice, that fundry liquors undergo sensible alterations from cork stoppers. Neumann observes, that acids, alkalies, both fixed and volatile, the dulcified alkaline and scid spirits, some neutral saline liquors, lime water, blue vegetable juices, and syrups made from them, are changed more or less to a yellow or brown colour.

SUCCINUM [L. E.] Amber; a folid, brittle, bituminous fubfance, dug out of the earth, or found upon the sea shores: the largest quantities are met with along the coasts of Polish Prussia and Pomerania. It is of a white. yellow, or brown colour, fometimes opake, and fometimes very clear and transparent: the dark coloured and opake forts, by digestion with certain expressed oils and animal fats, become clearer, paler coloured, more pellucid, and confiderably harder. Amber boiled in water, neither foftens or undergoes any fenfible alteration: exposed to a greater heat, without addition, it melts into a black mass like some of the more common bitumens: fet on fire, its fmell resembles that which arises from the finer kinds of pitcoal: distilled in a retort, it yields an oil and a volatile acidulous falt (see Part Ill. chap. viii.)

Amber in subflance has very little smell or taste; and hence it has by some been reckoned a mere inactive earthy body. It was formerly accounted an absorbent, and as such had a place in the compound powder of crabs claws: it certainly has no title to this class of medicines, as not being acted upon by any acid. It is supposed to

be of service in the fluor albus. gleets, hysteric affections, &c. and in these intentions is sometimes given in the form of impalpable powder, to the quantity of a drachm. A tincture of amber made in rectified spirit (to which it imparts a bitterish aromatic taste and a fragrant (mell) promises to be of real service in these difor-Boerhaave extels this tineture as having incredible efficacy in all those distempers which proceed from weakness and relaxation, and in hypochondriacal, hysterical, and cold languid cases: if part of the spirit be abstracted by a gentle heat, the remainder proves a very elegant aromatic balfam, which is, perhaps, one of the most useful preparations obtainable from this concrete. Amber is levigated in the shops into an impalpable powder, which gives name to a compound powder, [L.] and is an ingredient in mithridate and theriaca [L.] A tincture of it in dulcified spirit of vitriol [E.] and the distilled oil and falt [L. E.] are likewise officinals: the oil is an ingredient in the volatile aromatic spirit, powder for promoting delivery, gum pills, cephalic balfam, and cephalic platter [E.]

SUCCISA, vide Morsus DIA-BOLI.

SULPHUR [L. E.] Sulphur, or brimstone, is a yellow substance, of the mineral kingdom, substance, of the mineral kingdom, fusible in a small degree of heat, totally volatile in a stronger, readily inflammable, burning with a blue slame, which is accompanied with a suffocating acid sume. It dissolves in alkaline liquors and in oils, not in acids, water, or vinous spirits.

Greatest

Greatest part of the sulphur met with in the shops is obtained from certain ores by a kind of distillation, or artificially composed by uniting the vitriolic acid with inflammable matters: at some of the Saxon fulphur works (from whence we are chiefly supplied) certain miherals abounding with vitriolic acid, but containing little or no fulphur, being stratified with wood, and the latter fet on fire, a large quantity of fine sulphur is produced. It is usually brought to us in large irregular masses, which are afterwards melted and cast into cylindrical rolls, with the addition of some coarse refin, flower, or the like; whence the paler colour of Sulphur is also not unthe rolls. frequently found native in the. earth, fometimes in transparent pieces of a greenish or bright yellow colour; but more commonly in opake grey ones, with only fome streaks of yellow. This last is the fort which is understood by the name Sulphur VIVUM [E.] though that met with under this name in the shops is no other than the drofs remaining after the fublimation of fulphur. All the forts of fulphur are, when perfectly pure, in no respect different from one another, notwithstanding the preference given by some to the more uncommon fossil forts; these last are, of all others, the least proper for medicinal purposes, as being the most subject to an admixture of foreign matter, both of the metallic and arfenical kind.

Pure sulphur loosens the belly, and promotes insensible perspiration: it seems to pass through the whole habit, and manifestly transpires through the pores of the skin, as appears from the sulphureous smell of persons who have taken it, and silver being stained in their pockets of a blackish co-

lour, which is the known effect of fulphureous fumes. It is a celebrated remedy against cutaneous diseases, both given internally, and externally applied. It has likewise been recommended in coughs, afthmas, and other disorders of the breast and lungs: in these cases. however, it has no very confiderable effect, unless, as Hoffman obferves, where the disease proceeds from the blood being tainted by ferophulous or scorbutic humours: where this happens, the prudent use of sulphur is said to do good fervice, throwing out a plentiful eruption upon the skin, and by degrees carrying off the peccant matter. The common dose of sulphur rarely exceeds a scruple, though Geoffroy goes as far as two drachms. The trochisci e sulphure of the dispensatory are one of the most elegant forms for the taking of it. It enters fix officinal preparations for external use, and gives name to one of them. Some have imagined that sulphur used externally is dangerous: that as it throws the morbific matter outwards, when given inwardly, it must, in like manner, drive it into the blood. when applied externally. opinion, which is supported by fome late writers, has no just foundation: fulphur has nearly the same effects, whether used internally or externally: in both cases. the eruptions become frequently more copious after the first use of it.

It is remarkable of this concrete, that though itself a medicine of considerable efficacy, it nevertheless restrains that of some others of the most powerful kind. Mercury is rendered, by the admixture of sulphur, inactive; and the virulent antimonial regulus, almost so: hence, when antimonial and mercurial medicines exceed in operation,

fulpher

fulphur has been given for abating their violence, and fometimes restrains their further action. Even the corrosive posson, arsenic, by the addition of sulphur, becomes almost innocent; and hence if a small portion of arsenic should be contained in sulphur, it possibly may not receive from thence any poisonous qualities.

SUMACH folia, femen: frutieis quæ rbus folio ulmi C. B. Common sumach; the leaves and seeds

IE ]

This tree, or shrub, is cultiwated in some places, on account of the culinary uses of its fruits, and for the purposes of the dyers, &c. among us, it is met with only in the gardens of the curious. The feeds or berries are of a red colour, in shape round and stat. Both these and the seaves are moderately astringent, and have sometimes been exhibited in this intention, but are now become strangers to the shops.

SYMPHYTUM, vide Conso-

TACAMAHACA [E]: a refin obtained from a tall tree (tacamaha a populo similis, frudu colore Proniæ simili J. B.) which grows fpontaneoully on the continent of America, and in a sheltered situation bears the winters of our own climate. Two forts of this refin are fometimes to be met with. The best, called (from its being collected in a kind of gourd shells) tacamahaca in shells, is somewhat unctuous and foftish, of a pale yellowish or greenish colour, an aromatic taste, and a fragrant delightful fmell, approaching to that of lavender and ambergris. This fort is very rare: that commonly found in the shops is in semitransparent grains or glebes, of a whit-

ish, yellowish, brownish, or greenish colour, of a less grateful smell than the foregoing. The first is faid to exude from the fruit of the tree, the other from incisions made in the trunk. This resin is said to be employed among the Indians, externally, for discussing and maturating tumours, and abating pains and aches of the limbs: it is an ingredient in the anodyne, hysteric, cephalic, and stomachie platters of the Edinburgh pharma-The fragrance of the copœia. finer fort sufficiently points out its being applicable to other purposes.

TAMARINDUS [L. E.] Tamarind; the fruit of a tree grow-The .ing in the East and West Indies. called by C. Bauhine filiqua Arabica quæ tamarindus. It is a pod resembling a bean cod, including several hard seeds, together with a dark coloured viscid pulp of a pleasant acid take: the East-India tamarinds are longer than the West-India sort; the former containing fix or feven feeds each, the latter rarely above three or four. The pulp of these fruits, taken in the quantity of 2 or 3 drachms, or an ounce or more, proves gently laxative or purgative; and at the same time, by its acidity, quenches thirst, and allays immoderate heat. It increases the action of the purgative fweets, cafia and manna, and weakens that of the refinous cathartics. Some have supposed it capable of abating the virulence of antimonial preparations; but experience shews, that it has rather a contrary effect, and that all vegetable acids augment their power. Tamarinds are an ingredient in the electary of casia [L.] the lenitive electary [E] and decoction of tamarinds with fena [E.]

TAMA-

TAMARISCI folia, cortex: Tamaricis alterius folio tenuiore, sive Gallicæ C. B The tamarisc tree; its bark and leaves.

These are moderately astringent: they are never met with in prescription, and have long been entire strangers to the shops.

TANACETI folia, flores, semen: Tanaceti wulgari, lutei C. B. Tan-fy: the leaves, [L.] flowers, and seeds [E]

Tanfy grows wild by road-fides. and the borders of fields, and is frequently also cultivated in gardens both for culinary and medicinal uses: it flowers in June and July. Considered as a medicine, it is a moderately warm bitter, accompanied with a strong, not very disagreeable flavour: some have had a great opinion of it in hysteric disorders, particularly those proceeding from a deficiency or Suppression of the uterine purgations. The leaves and feeds have been of confiderable effeem as anthelmintics: the feeds are less bitter, and more acrid and aromatic than those of rue, to which they are reckoned fimilar; or of fantonicum, to which they have been frequently substituted.

TAPSI BARBATI, seu Verbasci folia, stores: Verbasci maris latisolii lutei C. B. Mullein; the leaves and flowers.

This is met with by road-fides, and under hedges: it is clothed all over with foft downy leaves, and produces long spikes of yellow flowers in July. The taste discovers in it a glutinous quality; and hence it stands recommended as an emollient, and is in some places held in great esteem in confumptions. The slowers of mullein have an agrecable, honey-like sweetness; an extract prepared

from them by rectified spirit of wine tastes extremely pleasant.

TARAXACUM, vide Dens LEONIS.

TARTARUM [L.E.] Tartar is a faline substance, thrown off from wines, after fermentation, to the fides and bottom of the calk: it proves of a red or white colour, and more or less foul or droffy, according to the colour and quality of the wine; the white is generally looked upon as the pureft: of either fort, fuch as is clean. folid, somewhat transparent, and has its outside covered over with fmall fhining cryftals, is preferable to fuch as appears porous, droffy, opake, and less bright. This substance, tho' truly faline, is scarce acted upon by cold water: the purest fort, or such as has been purified by art, requires four-andtwenty times its weight of boiling water to dissolve in : the folutions of both the tartars pass the filter colourless, and shoot, in the coldinto small, white, semitransparent crystals. All fuch earths as are soluble in vinegar, and alkaline falts, render tartar more easily soluble in water: hence the refiners at Montpelier are said to employ a certain earth for promoting its folutions with fome particular managements for making it shoot into large crystals. This addition may occasion a considerable alteration in the falt, infomuch that the finer forts of white tartar are perhaps preferable, on many occafions, to the common crystals. The virtues of tartar are those of a mild, cooling, aperient, laxative medicine. Taken from half an ounce to an ounce, it proves a gentle, though effectual purgative. Angelus Sala relates, that he was cured of an habitual colic, by purging himself a few times with fix drachms of the crude salt, after many other medicines had been tried to no purpose. For the preparations of tartar, see Part III. chap. viii. sect. 7. This salt is likewise an ingredient in the common infusion of sena, compound powder of sena [L.] and is used for dissolving or corroding some metallic bodies, particularly antimony, from which it receives a strong emetic impregnation.

## TELEPHIUM, vide CRASSULA.

TEREBINTHINÆ. Turpentines; refinous juices extracted from certain trees. There are four kinds of turpentine diffinguished in the shops.

TEREBINTHINA CHIA, five CYPRIA [L. E.] Chian, or Cy-

prus turpentine.

This is generally about the confiftence of thick honey very tenacious, clear and almost transparent, of a white colour, with a cast of yellow, and frequently of blue: it has a warm, pungent, bitterish taste, and a fragrant smell, more agreeable than any of the other

turpentines.

This juice is the produce of the serebinthus vulgaris C. B. common terebinth, an evergreen tree or thrub, which grows spontaneously in the warmer climates, and endures the colds of our own. The turpentine brought to us, is extracted in the islands whose names at bears, by wounding the trunk and branches a little after the buds have come forth: the juice issues limpid, and clear as water, and by degrees thickens into the confiftence in which we meet with it. A like juice exuding from this tree in the eastern countries, inspiffated by a flow fire, is of frequent use, as a masticatory, among the

Persian ladies, who (as Kompfer informs us) are continually chewing it, in order to fasten and whiten the teeth, sweeten the breath, and promote appetite.

TEREBINTHINA VENETA

[E.] Venice turpentine.

This is usually thinner than any of the other forts, of a clear, whitish, or pale yellowish colour, a hot, pungent, bitterish, disagreeable taste, and a strong smell, without any thing of the fine aromatic slavour of the Chian kind.

The true Venice turpentine is obtained from the larix folio deciduo conifera J. B. larch, a large tree, growing in great abundance upon the Alps and Pyrenean mountains, and not uncommon in the English gardens What is usually met with in the shops, under the name of Venice turpentine, comes from New England; of what tree it is the produce, we have no certain account: the finer kinds of it are in appearance and quality not considerably different from the true sort above described.

TEREBINTHINA ARGEN-TORATENSIS [L.E.] Straßurgh

turpentine.

This, as we generally meet with it, is of a middle confidence betwirt the two foregoing, more transparent and less tenacious than either; its colour a yellowish brown. Its smell is very fragrant, and more agreeable than that of any of the other turpentines, except the Chian: in taste it is the bitterest, yet the least acrid.

This refin is obtained from the two forts of fir trees mentioned in page 72, which are the most plentiful, and perhaps the only ones that grow spontaneously in Europe. There is another whose refin is much superior to the common turpentine, and has some

times been brought to us from abroad under the name of BALSA-MUM CANADENSE. This species is the abies minor, peclinatis foliis, Virginiana, conis parwis subrotundis Pluk. Virginian, or Canada sirtree, which, though not a native of this climate, has been found to endure its severest colds.

TEREBINTHINA COMMU-NIS [L. E] Common turpentine is the coarsest, heaviest, in taste and smell the most disagreeable, of all the sorts: it is about the consistence of honey, of an opake brownish white colour.

This is obtained from the pinus fylvestris C B. wild pine, a low unhandsome tree, common in different parts of Europe: this tree is extremely resinous, and remarkably subject to a disease from a redundance and extravasation of its resin, infomuch that, without due evacuation, it swells and bursts. The juice, as it issues from the tree, is received in trenches made in the earth, and asterwards freed from the grosser impurities by colature through wicker baskets.

All these juices vield, in distillation with water, an highly penerating essential oil, a brittle insipid refin remaining behind With regard to their medical virtues, they promote urine, cleanse the parts concerned in the evacuation thereof, and deterge internal ulcers in general, and at the fame time, like other bitter hot substances, frengthen the tone of the vessels: they have an advantage above most other acrid diuretics, that they gently loosen the belly They are principally recommended in gleets, the fluor albus, and the like; and by fome in calculous complaints: where these last proceed from fand or gravel, formed into a mass by viscid muccus matter, the turpentines, by disfolving the mucus, pro-

mote the expulsion of the fand; but where a calculus is formed, they can do no service, and only ineffectually irritate or inflame the parts. In all cases accompanied with inflammation, these juices ought to be abitained from, as this symptom is increased, and not unfrequently occasioned by them. It is observable, that the turpentines impart, foon after taking them, a violet smell to the urine a and have this effect, though applied only externally to remote parts, particularly the Venice fort. This is accounted the most powerful as a diuretic and detergent, and the Chian and Strasburgh as corroborants: the Strasburgh is an ingredient in the mercurial pills and Locatellus's balfam, and the Chian in mithridate and theriaca [L.] The common turpentine, as being the most offensive, is rarely given internally. Its principal use is in plasters and ointments, among farriers, and for the distillation of the oil or spirit, as it is called. dose of these juices is from a scruple to a drachm and a half: they are most commodiously taken in the form of a bolus, or dissolved in watery liquors by the mediation of the yolk of an egg or mucilage. Of the distilled oil, a few drops are a sufficient dose : this is a most potent. stimulating, detergent diuretic, oftentimes greatly heats the constitution, and requires the atmost caution in its exhibition.

TERRA JAPONICA, vide Ja-

TERRA LEMNIA et SILE-SIACA, vide Bolus.

THAPSIÆ folia: Thapfiæ froz turbith garganici femine latissimo J. B. Deadly carrot; the root. This plant does not ill deserve its epithet; a small dose sperating with extreme violence both upwards and downwards. It is an entire stranger to the shops, and met with only in the gardens of the curious.

THEÆ folio [E.] Tea; the leaves of a shrub (thea frutex, folio cerafi, flore rosæ sylvesfiris, &c. Kæmpf.) cultivated in China.

The several sorts of tea met with among us, are the leaves of the same plant, collected at different times, and cured in a somewhat different manner: the small young leaves very carefully dried, are the finer green; the older afford the ordinary green and bohea. The two first have a sensible flavour of violets, the other of roses: the former is the natural odour of the plant; the latter, as Neumann observes. is probably introduced by art. Some of the dealers in this commodity in Europe are not ignorant that bohea tea is imitable by the leaves of certain common plants, artificially tinctured and impregnated with the rose slavour. The taste of both sorts is lightly bitterish, subastringent, and some-The medical virwhat aromatic. tues attributed to these leaves are fufficiently numerous, though few of them have any just foundation: little more can be expected from the common infusions, than that of a diluent, acceptable to the palate and stomach: the diuretic, diaphoretic, and other virtues which they have been celebrated for, depend more on the quantity of warm fluid, than any particular qualities which it gains from the tea. Nothing arises in distillation from either fort of tea with rectified spirit: water elevates the whole of their flavour.

THLAPSIS, femen: Treacle, or mithridate, mustard; the seeds [L.]

Two forts of thlaspi are used promiscuously, thlaspi arvense siliquis latis C. B. and the thlaspi arvense vaccariæ incano solio majus C. B. they both grow wild, the latter most plentifully. These seeds have an acrid biting taste like common mustard, with which they agree in medical qualities: their principal use is as ingredients in the compositions whose name they bear.

THUS MASCULUM, vide OLIBANUM.

THUS VULGARE [L. E.] Common frankincense: a solid, brittle refin, brought to us in little glebes or masses, of a brownish, or yellowish colour on the outside, internally whitish, or variegated with whitish specks; of a bitterish, acrid, not agreeable taste, without any confiderable imell. It is fupposed to be the produce of the pine tree which yields the terebinthina communis, and to concrete on the surface of rhe terebinthinate juice foon after it has issued from the plant.

It is an ingredient in mithridate, the gum plaster, strengthening plaster, and stomach plaster [L.]

THYMI folia: Thymi vulgaris folio tenuiore C. B. Common thyme; the leaves [E.]

This plant is frequent in our gardens, and flowers in June and July. It has an agreeable aromatic smell, and a warm pungent taste; which it imparts by infusion to rectified spirit, and sends over in distillation with water; along with the water arises an effential oil extremely hot and pungent.

THYMI CITRATI folia: Serpyli foliis citri odore C. B. Lemonthyme; the leaves [L].

This

This is found wild in dry mountainous places, but the shops are supplied from gardens. In taste and smell it is less acrid and more grateful than the common thyme; its smell, in particular, is remarkably different, approaching to that of lemons. It gives over its slavour in distillation both with water and spirit: with the former an elegant effential oil arises. The distilled spirit is an agreeable aromatic cordial liquor, not inferior to anything of this kind.

TILIÆ flores: Tiliæ fæminæ folio majore C. B. The lime or linden tree; its flowers [L. E.]

The lime tree has been much valued on account of its quick growth and pleasant shade; it flowers in July, and loses its leaves soon after. The slowers are made use of chiefly on account of their agreeable flavour, which water extracts from them by infusion, and elevates in distillation. Among the writers on the materia medica, they have the character of an antiepileptic, and a specific in all kind of spasms and pains. derick Hoffman relates, that he knew a chronical epilepsy cured by the use of an infusion of these flowers drank as tea.

THYMELÆÆ baccæ: Thymelææ foliis lini C.B. Spurge flax; its berries, called grana enidia.

TITHYMALI radix. Spurge; the root.

Several forts of spurge are mentioned in catalogues of the materia medica. The Edinburgh college, in their last edition, retained two (ESULA MAJOR, tithymalus palustris fruticosus C. B. German spurge; and ESULA MINOR, tithymalus foliis pini C. B. pine-spurge); both the Edinburgh and London

college have now rejected them

The spurges and grana cnidia are extremely acrid, irritating cathartics, and operate with so much violence as to be altogether unfit for internal use.

TINCAR, vide BORAX.

TORMENTILLÆ radix: Tormentillæ filvesfiris C. B. Tormentil, or septsoil; the root [L. E.]

Tormentil is found wild in woods and on commons: it has long flender stalks, with usually seven long narrow leaves at a joint; the root is, for the most part, crooked and knotty, of a blackish colour on the outfide, and reddish within. This root has an austere styptic taste, accompanied with a flight kind of aromatic flavour; it is one of the most agreeable and efficacious of the vegetable astringents, and is employed with good fuccess in all cases where medicines of this class are proper. It is more used, both in extemporaneous prescription and in officinal composition, than any of the other strong vegetable astringents: it is an ingredient in the two compound powders of bole [L.] the two powders and electary of fcordium [L.] the compound white decoction [E.] and japonic confection [E.] A tincture made from it with rectified spirit possesses the whole aftringency and flavour of the root, and loses nothing of either in inspissation.

TRAGACANTHA, vide Gum-

TRICHOMANIS folia: Trichomanis five polytrichi officina um C. B. English maidenhair; the leaves. [L. E.]

This is one of the herbs called, from the smallness of their stalks.

R capil-

capillary: it is found wild in different parts of England, upon old walls, and in shady places. The leaves have a mucilaginous, sweetish, subattringent taste, without any particular flavour: they are esteemed useful in disorders of the breast, proceeding from a thickness and acrimony of the juices ! and are likewise supposed to promote the expectoration of tough phlegm, and to open obstructions of the vifcera. They are usually directed in infusion or decoction, with the addition of a little liquorice. A syrup prepared from them has frequently supplied the place of that made from the adianthum verum: fome have substituted a still cheaper ingredient, and perhaps not much to the disadvantage of the medicine; both the maidenhairs yielding little more than a mucilaginous juice, greatly resembling the substitute made use of. The syrup brought from abroad has an admixture of orange flower water.

TRIFOLII PALUDOSI folia: Trifolii palufiris C. B. Marsh trefoil, or buck-beans; the leaves [L. E.]

This plant grows wild in moist marshy places: it has three oval leaves, standing together upon one pedicle which issues from the root; their taffe is very bitter, and somewhat nauseous. Marsh trefoil is an efficacious aperient and deobstruent, promotes the fluid secretions, and, if liberally taken, gently loosens the belly. It has of late gained great reputation in fcorbutic and scrophulous disorders, and its good effects in these cases have been warranted by experience; inveterate cutaneous diseases have been removed by an infusion of the leaves, drank to the quantity of a pint a day, at proper intervals, and continued fome weeks.

Boerhaave relates, that he was relieved of the gout by drinking the juice mixed with whey.

#### TRISSAGO, vide CHAMEDRYS.

TRITICI farina, amylum, furfur: Tritici vulgaris glumas triturando deponentis J. B. Wheat; the meal or flour, and starch [L E.] (prepared from the meal by maceration in fresh quantities of water)

and bran [E]

Wheat, a common article of our food, is more glutinous and nutritious than most other kinds of grain. The flour, or the starch, prepared from it, form with water a fost viscid substance, which has been taken with good success in diarreas and dysenteries. Starch is an ingredient in the compound powder of gum tragacanth and the white pectoral troches [L.] and gives name to a lohoch [E.]

Bran contains, besides the husks, or shells of the wheat, a portion of its farinaceous matter: this is less glutinous than the siner slower, and is supposed to have a detergent quality: infusions of bran are not unfrequently employed in this intention externally, and sometimes likewise taken inwardly.

Bread, carefully toasted, and infused or lightly boiled in water, imparts a deep colour, and a sufficiently agreeable restringent taste. This liquor, taken as common drink, has done good service in a weak lax state of the stomach and intestines, and in bilious vomiting and purging, or the cholera morbus: examples are related in the Edinburgh essays of several cases of this kind cured by it, without the use of any other medicine.

TUNICA, vide CARYOPHYL-LUS HORTENSIS.

TURPETHUM, five Turbith [E.] Turbith; the cortical part of the root of an Indian convolvulus, brought to us in oblong pieces, of a brown or ash colour on the outfide, and whitish within : the best is ponderous, not wrinkled, easy to break, and discovers a large quantity of refinous matter to the eye: its taste is at first sweetish; chewed for a little time, it becomes acrid, pungent, and nauseous. This root is a cathartic, not of the safest or most certain kind; the resinous matter, in which its virtue resides, appears to be very unequally distributed, infomuch that some pieces, taken from a scruple to a drachm. purge violently; while others, in larger doses, have scarce any effect An extract made from the root is more uniform in strength, though not superior or equal to purgatives more common in the fhops.

TUSSILAGINIS five farfarac folia, flores: Tusfilaginis vulgaris C. B. Coltssoot; the leaves and flowers [E].

This grows wild in watery places, producing yellow flowers in February and March; these soon fall off, and are succeeded by large roundish leaves, hairy underneath: their taste is herbaceous, somewhat glutinous and subacrid. Tussilago stands recommended in coughs and other disorders of the breast and lungs: the slowers are an ingredient in the pectoral decoction of the Edinburgh pharmacopæia.

TUTIA [L. E.] Tutty; an impure sublimate of zinc, or an argillaceous substance impregnated therewith, formed into tubulous pieces like the bark of a tree. It is moderately hard and ponderous, of a brownish colour, and full of small

protuberances on the outfide, fmooth and yellowish within: some pieces have a blueish cast, from mineral globules of zinc being thrown up by the heat in its metallic form. Tutty is celebrated as an ophthalmic, and frequently employed as fuch in unguents and collyria: it gives name to an officinal ophthalmic ointment [L. E.] See the article Zincum.

VALERIANÆ HORTENSIS MAJORIS radix: Valerianæ majoris odorata radice J. B. The greater garden valerian; its roots

This is an oblong wrinkled root, with several fibres at the bottom, of a brownish or ash colour on the outside, and whitish within; of an aromatic smell and taste, approaching to nard. It is accounted less essications as a medicine than the following.

VALERIANÆ SILVESTRIS radix: Valerianæ sylvestris majoris montanæ C. B. Valerianæ sylvestris majoris foliis angustioribus Morison. plant. umbellif. Wild valerian (the narrow-leaved fort, growing on open, dry, mountainous places) its root [L. E.]

This root confifts of a number of strings or fibres matted together, issuing from one common head; of a whitish or pale brownish colour: its smell is strong, like a mixture of aromatics with fetids; the taste unpleasantly warm, bitterish, and subacrid. There is another wild valerian, with broader leaves, of adeeper and shining green colour, met with in watery places. Both forts have hitherto been used indiscriminately; and Linnæus has joined them into one species, under the name of valeriana foliis omnibus pinnatis. Our college have restrained the shops to the first, which

R 2 is

is confiderably the strongest, and loses of its quality, if transplanted into such soils as the other naturally delights in. The roots, produced in low watery grounds, have a remarkably faint smell in comparison of the others, and sometimes scarce any at all. Wild valerian is a medicine of great use in nervous disorders, and is particularly ferviceable in epilepfies proceeding from a debility of the nervous system. It was first brought into esteem in these cases by Fabius Columna, who by taking the powdered root, in the dose of half a spoonful, was cured of an inveterate epilepsy, after many other medicines had been tried in vain. Repeated experience has fince confirmed its efficacy in this disorder, and the present practice lays considerable stress upon it. The common dose is from a scruple to a drachm; in infusion from one to two drachms. Its unpleasant flavour is most effectually concealed by a fuitable addition of mace.

A tincture of valerian in proof spirit and in volatile spirit are kept in the shops [L]. This root gives name also to a compound water [E.] and is an ingredient in the cephalic tincture [E.] epileptic powder [E.] mithridate and the-

riaca [L.]

VERATRUM, vide Helle-BORUS ALBUS.

VERBASCUM, vide Tapsus BARBATUS.

VERBENÆ folia, radix: Verbenæ communis flore cerulæo C. B. Common wild vervain; the leaves and root [E.]

This is one of the medicines which we owe to the superstition of former ages; the virtues it has been celebrated for, both as an internal medicine and externally as an amulet, are extremely numerous: and possibly it has an equal title to them all: to the taste and smell it appears almost simply herbaceous.

VERONICA FOEMINA, vide ELATINE.

VERONICÆ MARIS, seu Betonicæ Pauli solia: Veronicæ maris supinæ et vulgatissimæ C. B. Male speedwell; the leaves [E.]

This is one of the veronicæ which produce their flowers in clusters at the joints of the stalks: it is a rough procumbent plant, not unfrequently met with on dry commons, and in fandy grounds. In taste, smell, and medical virtues, it is similar to the betonica. of which in its place; though the veronica is commonly supposed to have more of an aperient and pectoral virtue, and betony to be rather nervine and cephalic. Hoffman and Joh. Francus have written express treatises on this plant. recommending infusions of it, drank in the form of tea, as very falubrious in many disorders, particularly those of the breast.

VINCETOXICI, Asclepiadis, seu Hirundinariæ radix: Asclepiadis store albo C. B. Swallow wort, or tame

poison; the root [E.]

This is a native of the warmer climates; it is fometimes met with in our gardens, but rarely perfects its feeds. It is reckoned by botanists a species of apocynum, or dogsbane; from all the poisonous forts of which it may be distinguished, by yielding a limpid juice, whilst that of the others is milky. The root has a strong smell, especially when fresh, approaching to that of valerian, or nard; the taste is at first sweetish

and aromatic; but soon becomes bitterish, subacrid, and nauseous. This root is esteemed sudorisic, diuretic, and emmenagogue, and frequently employed by the French and German physicians as an alexipharmac, sometimes as a succedaneum to contrayerva; whence it has received the name of contrayerva Germanorum. Among us, it is very rarely made use of: it appears, from its sensible qualities, to be a medicine of much the same kind with valerian, which is indisputably preserable to it.

VINUM. Wine; the fermented juice of the grape. Among the great variety of wines in common use among us, five are employed in the shops as mensirua for medicinal simples.

Vinum album [L.] vinum album Hispanicum [E.] Mountain.

Vinum album Gallicum [E.] French white wine.

Vinum Canarinum [L. E.] Canary or fack.

Vinum Rhenanum [L. E.] Rhe-

Vinum rubrum [L.] Red port.

The uses of these liquors as menstrua and vehicles of the virtues of other medicines, will be given hereafter; in this place we shall consider only their effects on the human body. These are, to chear the spirits, warm the habit. promote perspiration, render the vessels full and turgid, raise the pulse, and quicken the circulation. The effects of the full bodied wines are much more durable than those of the thinner: all sweet wines, as Canary, abound with a glutinous nutritious substance; whilst the others are not nutrimental, or only accidentally fo, by strengthening the organs employed in digestion: sweet wines in general do not pass off freely by prine, and heat the

constitution more than an equal quantity of any other, though containing full as much spirit: red port, and most of the red wines, have an astringent quality, by which they strengthen the tone of the stomach and intestines, and thus prove serviceable for restraining immoderate secretions: those which are of an acid nature, as Rhenish. pals freely by the kidneys, and gently loosen the belly: it is supposed that these last exasperate or occasion gouty and calculous disorders, and that new wines of every kind have this effect.

VIOLÆ folia, flores: Violæ Martiæ purpureæ, flore simplici odoro C. B. The single March violet; its flowers [L. E.] and leaves [E.]

This is often found wild in hedges and shady places, and flowers in March; the shops are generally supplied from gardens. our markets we meet with the flowers of a different species. named by botanists viola Martia major, hirfuta, inodora: these may be distinguished from the foregoing by their being larger, of a pale colour, and of no smell. The officinal flowers have a very pleafant smell, and a deep purplish blue colour, denominated from They impart their them violet. colour and flavour to aqueous liquors. A syrup made from this infusion has long maintained a place in the shops, and proves an agreeable and uleful laxative for children.

VIPERA [L. E] The viper, or adder, is one of the viviparous reptiles, without feet, about an inch in thickness, and twenty or thirty in length. The poison of this serpent is confined to its mouth. At the basis of the sangs, or long teeth which it wounds with, is R 3 lodged

lodged a little bag, containing the poisonous liquid; a very minute portion of which, mixed immediately with the blood, proves fatal: our viper-catchers are faid to prevent the mischiefs otherwise following from the bite, by rubbing oil olive warm on the part. flesh of the viper is perfectly innocent, and strongly recommended as a medicine of extraordinary fervice in scrophulous, leprous, and other obslinate chronical disorders: its virtues, however, in these cases, are probably too much exaggerat-The viper is, doubtless, an ed. high nutritious food; and hence in some kinds of weaknesses, and emaciated habits, is not undefervedly looked upon as a good restorative. To answer any valuable purposes, fresh vigorous vipers (not such as have been long kept alive after they are caught) should be liberally used as food. The wines and tinctures of them can scarce be supposed to receive any considerable virtue from the animal. The dry flesh brought us from abroad, is intirely infignificant.

In the shops, a broth is directed to be prepared from fresh vipers, and a vinous tincture from dried ones [L.]: the dried slesh is also an ingredient in theriaca, and the fat in the ointment of tutty [L.] this fat being supposed peculiarly useful in disorders of the eyes, for which that ointment is designed.

VIRGÆ AUREÆ folia: Virgæ aureæ angustifoliæ, minus serratæ C. B. Golden rod; the leaves [E.]

This is found wild on heaths and in woods, producing spikes of yellow flowers in August. The leaves have a moderately astringent bitter taste, and hence prove serviceable in debility and laxity of the viscera, and disorders proceeding from that cause.

VISCI QUERNI lignum, folia : Vifci baccis albis C. B. Misseltoe; the wood and leaves [E.]

This is a bushy plant, growing on the trunk and branches of different trees: that met with on the oak is generally preferred, perhaps on account of its being the most rare. It may, however, be propagated by art on any trees, by rubbing the berries against the bark : this office has hitherto been performed by the thrush (who feeds on the berries in the winter) in clearing his bill from the feeds that slick about it. This plant was held in veneration by the fuperstition of former ages: it was hung about the neck to prevent witchcraft, and, taken internally, to expel poisons. Of late times it has been celebrated as a specific in epilepsies, palsies, &c. virtues, which it were greatly to be wished that experience gave any countenance to.

#### VITEX, vide Agnus castus.

VITIS VINIFERA. The vine tree. The leaves of this tree were formerly celebrated as aftringents, but have, for a long time, been intirely difregarded: their tafte is herbaceous, with only a flight roughness .- The trunk of the tree, wounded in the spring, yields a clear, limpid, watery juice: this tear of the vine has been accounted excellent for fore eves; and by some recommended likewise in ardent and malignant fevers, and as a diuretic.-The flowers have a pleasant smell, which water elevates from them in distillation; along with the water, a fmall portion of an elegant effential oil is said to arise, possessing in great persection the fragrance of the flowers.—The unripe fruit is of a very harsh, rough, sour tafte:

taste: its expressed juice, called verjuice, omphacium agressa [E] was of great esteem among the ancients, and still continues so in some places, as a cooling aftringent medicine: a rob and syrup were formerly prepared from it.—The ripe fruit or grapes, of which there are several kinds, properly cured and dried, are the raisins and currants of the shops: the juice by sermentation affords wine, vinegar, and tartar; of all which in their places.

VITRIOLUM. Vitriol is a faline crystalline concrete, composed of metal, and an acid similar to those of sulphur and alum. There are but three metallic bodies, which this acid is capable of perfectly disfolving or being united with into a crystalline appearance, zinc, copper, and iron: with the first it forms a white, with the second a blue, and with the third a green salt.

 ${f VITRIOLUM}$  ALBUM [L. F.] White vitriol, or vitriol of zinc; found in the mines of Goslar, sometimes in transparent pieces, but more commonly in form of white efflorescences, which are dissolved in water, and afterwards reduced by evaporation and crystallization into large masses. We rarely meet with this fort of vitriol pure : after the zinc, which is its proper basis, has been revived by inflammable fluxes, there remains a substance which is attracted by the magnet, and discovers itself, on other trials also, to be iron: a solution of the vitriol deposites on standing an ochery fediment, which generally gives a blue tincture to volatile al kalies, and hence appears to contain copper. White vitriol is sometimes given from five or fix grains to half a drachm, as an emetic: it operates very quickly, and if pure, without violence. Externally, it is employed as an opththalmic, and often made the basis of collyria, both in extemporaneous prescription, and in dispensatories: a solution of it is directed in this intention both by the Edinburgh and London colleges.

VITRIOLUM CŒRULEUM [L. E.] Blue vitriol, or vitriol of copper, falfely called Roman vitriol. Greatest part of the blue vitriol at present met with in the shops, is faid to be artificially prepared, by uniting copper with the vitriolic acid. This salt has a highly acrid, auftere, and very naufeous taste: it is a strong emetic, too violent to be exhibited with any tolerable degree of fafety. Its principal use is externally as an escharotic, and for flopping hæmorrhages, which it effects by coagulating the blood, and contracting the mouths of the vessels; it gives name to an officinal water for this intention.

VITRIOLUM VIRIDE [L. E.] Green vitriol, or vitriol of iron, commonly called copperas, the Roman vitriol of the Italian and other foreign writers. This is prepared in large quantity at Deptford, by diffolving iron in the acid liquor which runs from certain fulphureous pyritæ exposed for a length of time to the air. When pure, it is similar in quality to the officinal fal martis.

The green and blue vitriols (as well as the white) are in many places found native in the earth; though usually, in this state. cither fort is free from an admixture of the other; hence vitriol are met with of all the intermediate colours betwixt the grass green of the one and the sapphire blue of the

other

The acid of these salts has the greatest affinity with zinc, next to this with iron, and with copper the least of all. Hence, solutions of white vitriol deposite, on standing, greated part of the irony and cupreous matter, which they contain, and, if some fresh zinc be added, the whole: in like manner, upon adding bright polished iron to folutions of green vitriol, if it holds any cupreous matter, this will be thrown down. By this means the white and green vitriols may be purified from other metallic bodies.

ULMARIÆ, seu Reginæ prati folia, stôres: Ulmariæ barbæ cupri storibus compactis C. B. Meadowsweet, or queen of the meadows; the leaves and flowers [E.]

This herb is frequent in moist meadows, and about the sides of rivers; it flowers in the beginning of June, and continues in flower a considerable time. The slowers have a very pleasant flavour, which water extracts from them by infusion, and elevates in distillation. The leaves are herbaceous.

URTICÆ MAJORIS VUL-GARIS folia, semen: Urticæ racemiseræ majoris perennis Raii. Stinging nettle; the leaves and seeds [E.]

URTICÆ ROMANÆ folia femen: Urticæ urentis pilulas ferentis femen lini C. B. Roman nettle; the leaves and feeds.

These have had sundry virtues attributed to them, which the present practice pays no regard to. The young leaves of the first sort are by some used in the spring as a wholesome pot herb.

UVÆ PASSÆ [L.] majores [E.] Raisins of the sun; the dried grapes of the wisis Damastena.

UVÆ PASSÆ minores [E.] Currants; the dried grapes of the vitis Corinthiaca.

The principal use of these is as an agreeable sweet; they impart a very pleasant slavour both to aqueous and spiritous menstrua. The seeds or stones are supposed to give a disagreeable relish, and hence are generally directed to be taken out: nevertheles I have not found that they give any taste at all. The raisins of the sun are an ingredient in the pectoral decoction, tincture of sena, and stomachic tincture [L.]

WINTERANUS CORTEX[E.] Winter's bark; the produce of a tree growing in Jamaica, Barbadoes, &c. called by Sir Hans Sloane periclymenum redum, foliis laurinis, cortice acri uromatico. It was first discovered on the coast of Magellan by captain Winter, in the year 1567: the failors then employed the bark as a spice, and afterwards found it serviceable in the scurvy; for which purpose it is, at present also, sometimes made use of in dietdrinks. The true Winter's batk is not often met with in the shops, canella alba being generally subtlituted to it, and by many reckoned to be the same: there is nevertheless a considerable difference betwixt them in appearance, and a greater in quality: the Winter's bark is in larger pieces, of a more cinnamon colour than the canella, and tastes much warmer and more pungent.

ZEDOARIA [L. E.] Zedoary; the root of an Indian plant brought over in oblong picces about the thickness of the finger, or in roundish ones about an inch in diameter. Both forts have an agreeable fragrant smell, and a warm, bitterish aromatic tasse.

In distillation with water it yields an essential oil, possessing the smell and flavour of the zedoary in an eminent degree; the remaining decoction is almost simply bitter. Spirit likewise brings over some small share of its flavour; nevertheless the spirituous extract is considerably more grateful than the zedoary itself. An extract made from it with proof spirit (which is inferior to that prepared with reclified spirit) is an ingredient in the confectio cardiaca [L.]: the root in substance enters the confectio Paulina, mi-Thridate and theriaca [L.]

ZIBETHUM [E.] Civet; a foft unctuous substance, of a white, brown, or blackish colour, brought from the Brazils, the coast of Guinea, and the East-Indies: it is met with in certain bags, situated in the lower part of the belly of an animal said to be of the cat kind. The chief use of this drug is in persumes; it is rarely, if ever, employed for any medicinal purposes.

ZINCUM. Zinc; a metal, differing from all the other bodies of that class, in being inflammable per se, sublimable into flowers, which afterwards remain fixed in the strongest fire, soluble in every acid, not miscible in susion with sulphur, changing copper into a yellow metal, brass. Several pro-


ductions of this metal, though not generally known to be such, are kept in the shops; as its rich ore calamine, the white vitriol, the pure white flowers of zinc, called pompholyx, and the more impure compound tutty. The preparations of zinc are employed principally in external applications as opthalmics. The flowers levigated into an impalpable powder, form with oily substances an useful unguent. and with rose water, and the like. elegant collyria, for defluxions of thin sharp humours upon the eyes: they are moderately altringent, and act, if the levigation has been duly performed, without acrimony or irritation. Taken internally, they prove emetic.

ZINGIBER [L. E.] Ginger; a root brought from China and the East and West Indies; of a fragrant smell, and a hot, biting, aromatic talle. Reclified spirit extracts its virtues, by infusion, in much greater perfection than aqueous liquors; the latter elevates its whole flavour in distillation, the former little or nothing. Ginger is a very useful spice, in cold flatulent colics, and in laxity and debility of the intestines: it does not heat so much as those of the pepper kind, but its effects are more durable. It gives name to an officinal fyrup [L. E.] and enters a great number of the compositions.

General titles including several simples.

The five opening roots:

Smallage,
Afparagus,
Fennel,
Parfley,
Butchers-broom.



The simples of each of the above classes have been often employed together, under the respective general appellations. This practice has entirely ceased among us; and accordingly these denominations are now expunged both from the London and Edinburgh pharmacopæias, though still retained in foreign ones.

#### General rules for the collection and preservation of simples.

#### ROOTS.

Annual roots are to be taken up before they shoot out stalks or flowers: biennial ones, chiefly in the autumn of the same year in which the feeds were fown: the perennial, when the leaves fall off, and therefore generally in the autumn. Being washed clean from dirt, and freed from the rotten and decayed fibres, they are to be hung up in a [warm] shady, airy place, till sufficiently dried. The thicker roots require to be flit longitudinally, or cut transversely into Such roots as lose thin flices. their virtues by exficcation for are defired to be preserved in a fresh state, for the greater conveniency of their use in certain forms] are to be kept buried in dry fand [E.]

THERE are two seasons, in which the biennial and perennial roots are reckoned the most vigorous, the autumn and spring; or rather the time when the stalks or leaves have fallen off, and that in which the vegetation is just going to begin again, or soon after it has begun; which times are sound to differ considerably in different plants.

The college of Edinburgh, in the two first editions of their pharmacopæia, directed them to be dug in the spring, after the leaves were formed; in the third edition, the autumn is preserved, and this rule is continued in the succeeding ones. The generality of roots appear, indeed, to be most efficacious in the spring; but as at this time they

are also the most juicy, and consequently shrivel much in drying, and are rather more difficultly preserved, it is commonly thought most advisable to take them up in autumn. No rule, however, can be given, that shall obtain universally: arum root is taken even in the middle of summer, without suspicion of its being less active than at other seafons; while angelica root is inert during the summer, in comparison of what it was in the autumn, spring, or winter.

#### HERBS and LEAVES.

HERBS are to be gathered when the leaves have come to their full growth, before the flowers unfold; but of fome plants the flowery tops are preferred. They are to be dried in the same manner as roots [E.]

For the gathering of leaves there cannot, perhaps, be any universal rule, any more than for roots; for though most herbs appear to be in their greatest vigour about the time of their flowering, or a little before, there are some in which the medicinal parts are more abundant at an earlier period.

Thus mallow and marshmallow leaves are most mucilaginous when young, and by the time of flowering approach more to a woody nature. A difference of the same kind is more remarkable in the leaves of certain trees and shrubs: the young buds, or rudiments of the leaves of the black poplar tree, have a strong fragrant smell, approaching to that of storax; but by the time that the leaves have come

to their full growth, their fragrance is exhausted.

Herbs are directed by most of the pharmaceutic writers to be dried in the shade; a rule which appears to be very just, though it has sometimes been misunderstood. They are not to be excluded from the fun's heat, but from the strong action of the folar light, by which last their colours are very liable to be altered or destroyed, much more fo than those of roots. Slow drying of them in a cool place is far from being of any advantage: both their colours and virtues are preserved in greatest perfection, when they are dried hallily, by a heat of common fire as great as that which the fun can impart: the juicy ones in particular require to be dried by heat, being otherwise subject to turn black. Odoriferous herbs, dried by fire till they become friable, discoverindeed, in this arid flate, very little finell; not that the odorous matter is dislipated; but on account of its not being communicated, from the perfectly dry subject, to dry air; for as foon as a watery vehicle is supplied, whether by infuling the plant in water, or by exposing it, for a little time, to a moist air, the odorous parts begin to be extracted by virtue of the aqueous moisture, and discover themselves in their full force.

Of the use of heat in the drying of plants, we have an instance in the curation of tea among the Chinese: according to the accounts of travellers, the leaves, as soon as gathered, are brought into an apartment furnished with a number of little furnaces or stoves, each of which is covered with a clean, smooth iron plate: the leaves are spread upon the plates, and kept relling with the hands till they begin to curl up about the edges;

they are then immediately swept off on tables, on which one person continues to roll them, while another fans them that they may cool hastily. This process is repeated two or three times, or oftener, according as the leaves are disposed to unbend on standing.

#### FLOWERS.

Flowers are to be gathered when moderately expanded, on a clear dry day, before noon. Red rofes are taken before they open, and the white heels clipped off and thrown away [E.]

THE quick drying, above recommended for the leaves of plants, is more particularly proper for flowers; in most of which both the colour and smell are more perishable than in leaves, and more subject to be impaired by flow exficcation. Of the flowers which come fresh into the apothecaries hands, the only ones employed dry in the London pharmacopœia, are red roses; and these, in all the compolitions in which they are used in a dry state, are expressly ordered to be dried hastily (cclereter arefalle.) One of the most valuable aromatics of European growth, faffron, is a part of a flower, dried on paper on a kind of kiln, with a heat sufficient to make it sweat, with care only not to endanger the scorching of it.

It may here be observed, that the virtues of flowers are confined to different parts of the flower in different plants. Saffron is a singular production, growing at the end of the stile or pittil: the active part of camomile flowers is the yellow disk, or button in the middle; that of lilies, roses, clove-july-flowers, violets, and many others, the petals or flower-leaves; while rosemary has

little

little virtue in any of these parts, the fragrance admired in the slowers of this plant residing chiefly in the cups.

SEEDS and FRUITS.

SEEDS should be collected when ripe and beginning to grow dry, before they fall off spontaneously. Fruits also are to be gathered when ripe, unless they are ordered to be otherwise [E.]

Or the fruits whose collection comes under the notice of the apothecary, there are few which are used in an unripe state: the principal is the floe, whose virtue, as a mild astringent, is greatly diminished by maturation. The fruit of the orange-tree, raised in our gardens or green-houses, is sometimes gathered in a state of much greater immaturaty, foon after it is formed on the tree, before it has acquired its acid juice: at this time it proves an elegant aromatic bitter, greatly refembling what are called Curassao oranges, which appear to be no other than the same fruit gathered at the same period, in a warmer climate.

The rule for collecting feeds is more general than any of the others, all the officinal seeds being in their greatest persection at the time of their maturity. secds contain little watery moisture, they require no other warmth for drying them than that of the temperate air in autumn: fuch as abound with a gross expressible oil, as those commonly called the cold feeds, should never be exposed to any confiderable heat; for this would hasten the rancidity, which, however carefully kept, they are very liable to contract. Seeds are best preserved in their natural hulks, or coverings, which should be separated only at the time of using; the husk, or cortical part, serving to defend the seed from being injured by the air.

Woods and BARKS.

THE most proper season for the felling of woods, or shaving off their barks, is generally the winter [E.]

THE only woods of our own growth, retained in the catalogues of simples of our pharmacopæias, are the juniper and box; the first of which is rarely or flever kept in the shops, or employed in practice; the other is procured from the turner, and it is indifferent at what season it has been cut down, being at all times sufficiently sit for the only use it is applied to, the yielding an empyreumatic oil by distillation in a strong fire.

Of the barks of our own growth. the London college has not retained one. In the Edinburgh pharmacopæia there are several, viz. those of the ash tree, birch tree, oak, elm, sloe, wild service, black alder, and elder, which, however, have been so rarely used in medicine, that the feafons of their greatest perfection cannot be ascertained from experience. It may be doubted, whether barks are not generally more replete with medicinal matter in the fummer and fpring than in winter. The barks of many trees are, in fummer, fo much loaded with refin and gum, as to burst spontaneously, and discharge the redundant quantity. It is faid that the bark of the oak answers best for the tanners, at the time of the rifing of the fap in fpring; and as its use in tanning depends on the same aftringent quality for which it is used in madicine, it should seem to be fitted

for medicinal purposes also in the fpring. It may be observed, likewise, that it is in this last season that barks in general are most conveniently peeled off.

and the viper, to which the Edinburgh add snails, earth-worms, and bees. Whatever virtues these bodies may have, they are supposed to be best when they have attained

#### ANIMALS and MINERALS.

Animals and minerals are to be chosen in their most perfect state, unless they are ordered otherwise [E.]

THE animals of the London pharmacoposia are only millepedes

and the viper, to which the Edinburgh add snails, earth-worms, and bees. Whatever virtues these bodies may have, they are supposed to be best when they have attained to their common full growth. As there are no distinctions of maturity or immaturity in the mineral kingdom, the only rule for directing our choice here must be, the purity of the subjects from any mixture of other bodies: none of them are ever to be used in an impure state.

#### PART III.

### Pharmaceutical Preparations.

#### CHAPTER I

#### THE MORE SIMPLE PREPARATIONS.

TERREORUM, aliorumque quæ aqua non dissolventur corporum præparatio. The preparation of EARTHY and such other pulverable bodies as will not dissolve in water.

THESE substances are first to be pulverised in a mortar, and then levigated with a little water, upon a hard and smooth marble, into an impalpable powder: this is to be dried upon a chalk stone, and asterwards set by for a few days, in a warm, or, at least, very dry place. L.

After this manner are to be prepared,

Ærugo, werdigris. L.

Antimonium, antimony, L. E.

Chelæ cancrorum, crabs clasvs, L. E.

Corallum, coral. L. E.

Creta, chalk. L. E.

Lapis bezoar, bezoar flone; which is
to be moistened, in the levigation,
with spirit of wine instead of wa-

ter. L.

Lapis calaminaris, calamine flone, previously calcined for the use of those who make brass. L. Where this is not to be had, the mineral may be calcined, by heating it three times red hot, and quenching it as often in water. E.

Lapis hæmatites, blood stone. L. E. Lapis lazuli. E. Margaritæ, pearls. L. E.

Oculi cancrorum, crabs eyes, fo called. L. E.

Ostreorum testæ, ogster shells, washed clean from dirt. L. may also be prepared by exposing them, for some days, to the sun, and then rubbing them in a marble mortar till they come into a kind of . paste; this is to be again dried in the sun, and afterwards rubbed into an impalpable powder: the hollow shells are preferred [E.] on account of their containing more of the fine white earth. in proportion to the outward rough coat, than the thinner flat ones: the rough matter appears to be largely impregnant with marine falt.

Ovorum

Ovorum testa, egg shells, freed by boiling from the skin that adheres to them. L.

Succinum, amber. L. E.

Tutia, tutty. L. E.

In preparing antimony, calamine, and tutty, particular care ought to be taken to reduce them into the most subtile powder possible. L.

Where large quantities of the foregoing powders are to be prepared, it is customary, instead of the stone and muller, to employ hand-mills made for this use, consisting of two stones, the uppermost of which turns horizontally upon the lower, and has an aperture in the middle, for the conveniency of supplying fresh matter, or of returning that which it has already passed, till it is reduced to a proper degree of sineness.

For the levigation of hard bodies, particular care should be taken, whatever kind of instruments is made use of, that they be of sufficient hardness, otherwise they will be abraded by the powders. The hæmatites, a hard iron ore, is most conveniently levigated betwixt two iron planes: for is the common levigating stones are made use of, the preparation, when sinished, will contain almost as much of foreign matter from the instrument as of the hæmatites.

It has been customary to moisten several powders, in levigation, with rose, balm, and other distilled waters: these, nevertheles, have no advantage above common water; since in the subsequent exsication they must necessarily exhale, leaving the medicine possessed of no other virtue than what might be equally expected from it when prepared with the cheaper element.

Some few substances, indeed, are more advantageously levigated with spirit of wine than with water. Thus bezoar has the green colour, usually expected in this costly preparation, confiderably improved thereby. A little spirit may be added to the other animal substances, if the weather is very hot, and large quantities of them are prepared at once, to prevent their running into putrefaction; an accident which. in those circumstances, sometimes happens when they are levigated with water only. Crabs eyes, which abound with animal gelatioous matter, are particularly liable to this inconvenience.

The caution given above for reducing antimony, calamine, and tutty, to the greatest subtility possible, demands particular attention. The tenderness of the parts to which the two last are usually applied, requires them to be perfectly free from any admixture of gross irritating particles. The first, when not thoroughly comminuted, might not only by its sharp needle-like spicula wound the stomach, but likewise answers little valuable purpose as a medicine, proving either an useless load upon the viscera, or at best raffing off without any other fenfible effect than an increase of the grosser evacuations; whilst, if reduced to a great degree of fineness, it turns out a medicine of confiderable efficacy.

The most successful method of obtaining these powders of the requisite tenuity, is, to wash off the finer parts by means of water (see page 54.) and continue levigating the remainder, till the whole becomes fine enough to remain, for some time, suspended in the fluid; a process received in the Edinburgh pharmacopaia, and there

directed as follows:

Edinb.

Edinb.

A quantity of water is to be poured upon the levigated powder, in a large vessel, and the vessel repeatedly shaken, that the siner parts of the powder may be dissuited through the water: the liquor is then to be poured off, and set by till the powder settles. The gross part, which the water would not take up, is to be further levigated, and treated in the same manner.

After this method are prepared antimony; calamine, tutty, bloodstone, chalk, and lapis

lazuli.

By this method, which is that commonly practifed in the preparation of colours for the painter, powders may be obtained of any required degree of tenuity; and without the least mixture of the gross parts, which are always found to remain in them after long continued levigation: all the coarfer matter settles at first, and the finer powder continues suspended in the water, longer and longer, in proportion to the degree of its The same process may finencis. likewise be advantageously applied to other hard pulverable bodies of the mineral kingdom. or artificial preparations of them; provided they are not foluble in, or specifically lighter than water. The animal and absorbent powders, crabs claws, crabs eyes, oyfter shells, egg shells, chalk, pearl, coral and bezoar, are not well adapted to this treatment; nor Thefe indeed do they require it. fubstances are readily foluble in acid juices without much comminution: if no acid is contained in the first passages, they are apt to concrete, with the mucous matter usually lodged there, into hard indissoluble masses:

greater degree of fineness they are reduced to, the more are they disposed to form such concretions, and enabled to obstruct the orifices of the small vessels. See page 62.

AXUNGIÆ PORCINÆ, SEVI-QUE OVILLI curatio.

The purification or trying of bogs lard and mutton fuet. Lond.

Chop them into small pieces, and melt them by a gentle heat, with the addition of a little water; then strain them from the membranes.

THE use of the water is to prevent the fat from burning and turning black; which it does very effectually, though it fomewhat prolongs the process, and is likewife apt to be in part imbibed by the fat. The Edinburgh dispenfatory directs the fat to be first freed from the skins, blood vessels and fibres, then washed in fresh quantities of water till it no longer gives the liquor any bloody tinge, afterwards melted, strained. and kept close from the injuries of the air. The shops are usually supplied with these fats ready prepared.

AXUNGIÆ VIPERINÆ curatio.

The purification of vipers fat.

Lond.

Let the fat, separated from the intestines, be melted by a gentle fire, and then pressed through a thin linen cloth.

THE quantity of this fat usually purified at a time is so small, that the heat may be easily regulated so as to prevent burning, without the addition of any water.

It is not necessary, as Dr. Pemberton observes, to be very curious in picking out the fat; it is sufficient if the heart, liver, and other bloody parts, are taken away; for the rest of the membranes crisp up while the fat melts, so as to be easily separated by straining.

MELLIS DESPUMATIO.

The despumation or clarifying of boney.

Let the honey be liquefied in a water-bath, (that is, by fetting the vessel containing the honey in a vessel of hot water) and the four which arises taken off.

THE intention of this process is to purify the honey from wax or other drossy matters that have been united with it by the violence of the press in its separation from the comb; and from meal and such like substances, which are sometimes fraudently mingled with it. When the honey is rendered liquid and thin by the heat, these lighter matters rise freely to the surface.

### SCILLÆ COCTIO. The baking of fquills.

Lond.

Let the squill (freed from the outer skin, and the hard part to which the little fibres adhere) be inclosed in a paste made of wheat slour and water, and baked in an oven, till the passe becomes dry, and the squill soft and tender throughout.

This preparation is as old as the theriaca, and is continued in our dispensatory, for no other use than making the troches of squills, which are one of its principal ingredients. The Edinburgh dispensatory having now dropt the theriaca, has dropt also the baked squills and the troches, and admitted them formerly only in compliance with custom, giving expressly the preference to squills moderately dried. The intention of baking the root is to abate its acrimony.

### SCILLÆ EXSICCATIO. The drying of Squills.

Lond.

Let the squill, cleared from its outer skin, be cut transversely into thin slices, and dried with a very gentle heat.

By this method, the squill dries much fooner than when only its several coats are separated, 28 has been usually directed; the internal part being here laid bare, which, in each of the entire coats, is covered with a thin skin, which impedes the exhalation of the moisture. The root loses, in this process, four-fifths of its original weight; the parts which exhale appear to be merely watery: hence fix grains of the dry root are equivalent to half a drachm of it when fresh; a circumstance to be particularly regarded in the exhibition of this medicine. In the preceding editions of our dispensatory, a particular caution was given not to use an iron knife for cutting squills, but one of wood, ivory, or other bone: the foundation of this caution is said to be, not so much that the squill would receive any ill qualities from the iron: as that its acrid juice, adhering to the knife, might render a wound received by it extremely painful, or even dangerous.

RHABARBARI et NUCIS MOSCHATÆ torrefactio.

The roasting of rhubarh and nutmeg.

Lond.

Roast them with a gentle heat, until they become easily friable.

Nurmegs, in their natural state. are so soft and unctuous, as scarce to be reducible into powder, a form in which they are occasionally wanted; and rhubarb is very difficultly fo, unless it be thorough-The torrefaction renders ly dry. them easily pulverable, and as soon as this point is obtained, should be immediately discontinued, otherwife the drugs will be confiderably injured. This treatment is supposed by some to increase the astringency of the subjects, perhaps on no very good foundation: it undoubtedly renders the rhubarb less purgative, and the nutmegs less aromatic. Both drugs may be reduced into sufficiently fine parts for most purposes, by means of a grater, without any alteration being made in their native quality.

## SPONGIÆ USTIO. The burning of sponge. Lond.

Burn the sponge in a close earthen vessel, until it becomes black, and easily friable: then powder it in a glass or marble mortar.

This medicine, now first received in the dispensatory, has been in use for a considerable time; and employed against scrophulous disorders and cutaneous soulnesses, in doses of a scruple and upwards. Its virtues seem to depend upon a volatile salt, just formed, and combined with its own oil: if the sponge be distilled with a stronger heat, it yields a large proportion of that salt in its proper form. The salt is in this

preparation so far extricated, that if the burnt sponge be ground in a brass mortar, it corrodes the metal, so as to contract a disagreeable taint, and sometimes an emetic quality.

Bees, earthworms, and other animal substances, have by some been prepared in the same manner, and recommended in different diseases; but as these substances fall greatly short of sponge in the quantity of volatile salt producible from them by sire, they are probably inferior also in medicinal efficacy. Of all the animal maters that have been tried, raw silk is the only one which exceeds or equals sponge, in the produce of salt.

A good deal of address is requifite for managing this process in perfection. The sponge should be cut small, and beaten for some time in a mortar, that all the stony matters may be got out, which, compared with the weight of the sponge when prepared, will sometimes amount to a confiderable quantity. The burning should be discontinued as soon as ever the matter is become thoroughly black. If the quantity put into the vessel at once is large, the outside will be fufficiently burnt before the infide is affected; and the volatile falt of the former will in part escape, before that in the latter is begun to be formed. 'The best method of avoiding this inconvenience feems to be, to keep the sponge continually stirring, in such a machine as is used for the roasting of coffee.

### CORNU CERVICALCINATIO.

The calcination of hartshorn.

Lond.

Burn pieces of hartshorn in a potter's furnace, till they become S 2 persectly perfectly white; then powder and levigate them after the same manner as the other earthly bodies.

THE intention here is, totally to burn out and expel the oil, falt, and other volatile parts; so as to leave only a white infipid animal earth. For this purpose, a strong fire, and the free admission of air, are necessary. The potter's furnace is directed merely for the take of convenience: where this is not to be had, any common furnace or flove may be made to ferve: on the bottom of the grate spread fome lighted charcoal, and above this lay the horns. The whole will burn vehemently: the vegetable matter is reduced to ashes: and the horns are burnt to whiteness, still retaining their original form, by which they are easily diftinguished from the other: they ought to be separated as soon as grown cold, to prevent their imbibing any fixt falt from the vegetable ashes moistened by the air. The horns left after the distillation of the volatile falt and oil of hartshorn are as proper for this use as any other; that process only collecting such parts as are here dissipated in the air.

Calcined harshotrn is the purest of the animal absorbent powders; as being perfectly free from any glutinous or oily matter, which most of the others abound with. It appears nevertheless to be one of the weakest in absorbent power, or the most difficult of solution in

acids.

## PULPARUM EXTRACTIO. The extraction of pulps. Lond.

Unripe pulpy fruits, and ripe ones, if they are dry, are to be boiled in a fmall quantity of water un-

til they become soft: then press out the pulp through a strong hair sieve, and afterwards boil it down to a due confissence, in an earthen vessel, over a gentle sire; taking care to keep the matter continually stirring, to prevent its burning. The pulp of casia sistularis is in

The pulp of casia sistularis is in like manner to be boiled out from the bruised pod, and reduced afterwards to a proper consistence, by evaporating the

water.

The pulps of fruits that are both ripe and fresh, are to be pressed out through the sieve, without any previous boiling.

### STYRACIS COLATIO. The straining of storax. Lond.

Soften florax calamita in hot water; then press it out betwixt warm iron plates; and separate the florax, now purished, from the water.

THE storax commonly met with, stands greatly in need of purifica-It contains a large quantity of woody matter, which this process effectually frees it from, though in other respects liable to fome inconveniences. The woody substance in some measure defends the storax from the action of the press, and retains part of it behind: at the same time that the florax is apt to fuffer a confiderable dissipation of its volatile parts, in which its fragrance and principal virtue confift. To prevent as much as possible this last inconvenience, the operator ought carefully to avoid using a greater heat than is absolutely necessary; and as foon as the storax is sufficiently fostened, to be expeditious in the straining of it. It has been queried whether this refin does not

communicate somewhat to the water it is boiled in; as benzoine, with which it agrees in its other pharmaceutical characters, imparts to water a faline matter similar to the sublimed flowers. On trial it could not be observed that any saline matter was thus separated from storax, though it impregnated the water considerably with its fragrance.

Storax may be excellently purified by means of spirit of wine, which this refin totally dissolves in, so as to pass through a filtre, the impurities alone being left. If the storax is afterwards wanted in a solid form, it may be recovered from this solution by gently distilling off the spirit, which will elevate very little of its slavour, or by pouring to it a quantity of water. See chap. vi. sect. 3.

OPIUM COLATUM, vel EXTRACTUM THBBAICUM.

Strained opium, or the shebaic extract.

Lond.

Take of opium, cut into slices, one pound: dissolve it into the consistence of a pulp, in a pint of boiling water, with care to prevent its burning: and whilst it remains quite hot, strongly press it from the secess through a linen cloth: the strained opium is then to be reduced, by a water bath or other gentle heat, to its original consistence.

Opium thus foftened by a small quantity of water, passes the strainer entire, the seces only being lest behind. If it was dissolved in a large quantity of water, its resinous and gummy parts would be separated from one another.

WHERE large quantities of opium are purified at once, the inspissation is most commodiously performed in a water-bath; but fmall quantities may be very fafely inspissated, by placing the vessel immediately over a gentle fire, the matter being kept ftirring, and the vessel occasionally removed from the fire whenever there is any fofpicion of its becoming too hot. The groffer impurities of the opium are by this process effectually separated; but some of its heterogeneous admixtures, confifting chiefly of dust and farinaceous matters, are so fine, as partly to pass along with it through the pores of the strainer when dilated by the press; this manifestly appears upon boiling the strained opium in water, and afterwards in spirit; when a confiderable quantity of earthy matter will be left, which is not foliable in either of those menstrua.

THE OTHER GUMS, as ammoniacum, galbanum, afafætida, and the like, are purified after the fame manner, only here a larger quantity of water may be made use of without injury lf the resinous part happens to subfide, take it out, and reserve it to be added again towards the end of the inspissation, that it may unite with the rest into one uniform mass.

Any gum that melts easily, as galbanum, may likewise be purished by including it in a bladder, and keeping it in boiling water, until the gum becomes soft enough to be pressed from its impurities through a canvas strainer [L.]

In the straining of all the gums, care should be taken, that the heat be neither too great, nor too long continued; otherwise a considerable portion of their more active S 2 volatile

volatile matter will be lost: an inconvenience which cannot, by any care, be wholly avoided: Lence, as the faculty of Paris observes, the purer tears, unstrained, are preserable, for internal use, to the strained gums. The last of the above methods, that of foftening the gum in a bladder by external heat, without the addition of water, appears to be the most eligible for all those that will admit of being thus liquefied fufficiently; both as exhalation is prevented during the liquefaction; and as the strained gum returns in cooling to its original confisence, without the further heat which is requifite in the other method for evaporating the water. Opium is perhaps less injured by heat than the rest of the gums, the virtues of this drug feeming to reside more in its fixed than in the volatile parts: it is never-theless expedient, that the smell of the opium, which affords an useful mark of its genuineness, be as much as possible preserved; this, if the quantity of water was large, would be destroyed by the long evaporation which would then become necessary.

In the Edinburgh dispensatory, opium, and the fouler kinds

of aloes, are directed to be purified, by diffolying them in a fufficient quantity of water with a gentle heat, straining the folutions, and evaporating them to the consistence of honey. The other gums are not required to be purified.

It were to be wished that the confishence, to which the strained solutions are to be reduced, was determined with more precision, particularly in regard to opium, that there might be as little uncertainty as possible in its dose.

#### MILLEPEDARUM PRÆPA-RATIO.

Preparation of millepedes. Lond.

The millepedes are to be inclosed in a thin canvas cloth, and suspended over hot spirit of wine, in a close vessel, till they are killed by the steam, and rendered friable.

Edinb.

Let them be included in a proper vessel, and dried with a very gentle heat.

Both these are convenient ways of rendering millepedes pulverable, without endangering any lost of such virtues as they may be possessed of.

#### CHAPTER IL

Substances extracted from vegetables by expression.

#### SECT. L

#### Juices.

UICES are obtained from the fucculent parts of plants, by including them, after being properly cut, bruifed, &c. in a hair bag, and preffing them, betwirt wooden cheeks, in the common forew press, so long as any liquor

drops from them.

The harder fruits require to be previously well beaten or ground; but herbs are to be only moderately bruised; for if these are over bruised, a large quantity of the herbaceous matter will be forced out along with the juice. Hempen or woollen bags are apt to communicate a diagreeable flavour; the threads of these likewise swell in proportion as they imbibe mositure, so as in great measure to prevent the free percolation of the juice.

The fluids thus extracted from fucculent fruits, both of the acid and sweet kind, from most of the acrid herbs, as scorvy-grass and water-creffes, from the acid herbs. as forrel and wood-forrel, from the aperient lactescent plants, as dandelion and hawkweed, and from fundry other vegetables, contain great part of the peculiar tafte and virtues of the respective subjects. The juices, on the other hand, extracted from most of the aromatic herbs, as those of mint and the fragrant Turkey balm, commonly called balm of Gilead, have scarcely any thing of the flavour of the plants, and feem to differ little from decoctions of them, made in water, boiled till the volatile odorous parts have been diffipated. Many of the odoriferous flowers, as the lily, violet, hyacinth, not only impart nothing of their fragrance to their juice, but have it totally destroyed by the previous bruifing. want of sufficient attention to these particulars, practitioners have been frequently deceived in the effects of preparations of this class: juice of mint has been often prescribed as a stomachic, though it wants those qualities, by which mint itself, and its other preparations, operate in that intention.

The juices, thus forcibly preffed out from plants, differ from those which flow spontaneously or from incisions; these last consisting chiefly of fuch fluids as are not diffused through the whole substance of the vegetable subject, but elaborated in distinct vessels. or secreted into particular receptacles. From poppy heads, flightly wounded, there issues a thick milky liquor, which dries, by a moderate warmth, into opium; whilst the juice obtained from them by pressure is of a dark green colour, and far weaker virtue.

Juices, newly expressed, are generally thick, viscid, and very impure: impure: by colature, a quantity of gross matter is separated, the juice becomes thinner, limpid, and better fitted for medicinal purposes, though as yet not entirely pure: on standing, it becomes again turbid, and apt to run into a fermentative or putresactive state. Clarification with whites of eggs renders the juices more perfectly sine; but there are few that will bear this treatment without a manifest injury to their stavour, taste, and virtue.

The most effectual method of purifying and preferving these liquors, is, to let the strained juices stand in a cool place, till they have deposited their groffer feces, and then gently pass them several times through a fine strainer till perfectly clear; when about one-fortieth part their weight of good spirit of wine may be added, and the whole fuffered to fland as before: a fresh fediment will now be deposited, from which the liquor is to be poured off, strained again, and put into small bottles that have been washed with spirit and dried. little oil is to be poured on the furface, so as very nearly to fill the bottles, and the mouths closed with leather, paper, or stopt with firaw, as the flasks in which Florence wine is brought to us; this ferves to keep out dust, and suffers the air, which in process of time arises from all vegetable liquors, to escape; which air would otherwise endanger the burfting of the glasses, or, being imbibed afresh, render their contents vapid The bottles are to be and foul. kept on the bottom of a good cellar or vault, placed up to the necks By this method, juices in fand. may be preferved for a year or two; and fome for a much longer time.

It has already been observed,

that there are great differences in juices, in regard to their being accompanied, in the expression, with the virtues of the subjects: there are equal differences in regard to their preserving those virtues, and this independently of the volatility of the active matter, or its disposition to exhale, Even the volatile virtue of scurvygrass may, by the above method, be preserved almost intire in its juice for a considerable time; while the active parts of the juice of the wild cucumber quickly separate and settle to the bottom, leaving the fluid inert. Juices of arum root, iris root, bryony root, and fundry other vegetables, throw off ia like manner their medicinal parts to the bottom.

### SUCCI SCORBUTICI. The fcorbutic juices. Lond.

Take the juice of
Garden four ygrafs, two pints;
Brooklime,
Water creffes, each one pint;

Seville oranges, a pint and quarter.

Mix them together, let them stand till the seces have subsided, and then either pour the liquor off clear, or pass it thro' a strainer.

#### Edinb.

Take of
Juice of garden scurvygrass,
oranges, each one pint
and a half;

Water cresses,

Brooklime, each one pint; White fugar, ten ounces; Compound horferadish water, half a pint.

Mix the juices with the fugar, and depurate them according to art; then add the horferadift water. BOTH these compositions are of considerable use for the purposes expressed in the title; the orange juice is an excellent assistant to the scurvy grass and other acrid and antiscorbutics, which, when thus mixed, have been found from experience to produce much better effects than when employed by themselves. These juices may be taken, from

an ounce or two to a quarter of a pint, two or three times a day: they generally increase the urinary secretion, and sometimes introduce a laxative habit. Preserved with the cautions above mentioned, they will keep good for a considerable time; though, whatever care be taken, they are found to answer better when fresh.

## S E C T. II. Expressed oils.

XPRESSED oils are obtained , chiefly from certain feeds and kernels of fruits, by thoroughly pounding them in a stone mortar, or, where the quantities are large, grinding them in mills, and then including them in a canvas bag, which is wrapt in a hair cloth, and strongly prested betwixt iron plates. The canvas, if employed alone, would be squeezed so close to the plates of the press, as to prevent the oil from running down: by the interpofition of the hair-cloth, a free passage is allowed it.

SUNDRY machines have been contrived, both for grinding the subject, and pressing out the oil, in the way of business To facilitate the expression, it is customary to warm either the plates of the press, or the subject itself after the grinding, by keeping itstirring, in a proper vessel over the fire: the oil, liquefied by the heat, separates more freely and more plentifully. When the oil is defigned for medicinal purposes, this practice is not to be allowed; for heat, especially if its degree is sufficient to be of any confiderable advantage for promoting the separation, renders the oil less soft and palatable, impresses

adifagreeable flavour, and increases its disposition to grow rancid: hence the colleges both of London and Edinburgh expressly require the operation to be performed without heat.

Nor are the oils to be kept in a warm place after their expression. Exposed but for a few days to a heat no greater than that of the human body, they lose their emollient quality, and become highly rancid and acrimonious. Too much care cannot be taken for preventing any tendency to this acrid irritating, state, in medicines so often used for abating immoderate irritation.

So much are these oils disposed to this injurious alteration, that they frequently contract an acrimony and rancidity while contained in the original subjects. Hence great care is requisite in the choice of the unctuous seeds and kernels, which are often met with very rancid: almonds are particularly liable to inconveniences of this kind.

Expressed oils are prepared for mechanic uses from sundry different subjects, as nuts, poppy seed, hemp-seed, rape-seed, and others. Those directed for medicinal purposes in the London and Edinburgh pharmacopoias, are,

**OLEUM** 

OLEUM AMYGDALINUM.

Oil of almonds.

OLEUM SEMINUM LINI. Oil of linfeed.

OLEUM SEMINUM SINAPI.

Oil of muftard-feed.

THE oil of almonds is prepared from the sweet and bitter almonds indifferently; the oils obtained from both forts being altogether Nor are the differences of the other oils very confiderable. the discriminating qualities of the fubjects not refiding in the oils that are thus obtained by expression: the oil of mustard-seed is as foft, infipid, and void of pungency, as that of sweet almonds, the pungency of the mustard remaining entire in the cake left after the expression. The several oils differ in some of their properties from one another; but in medicinal qualities they appear to be all nearly alike, and agree in one common emollient virtue. They foften and relax the folids, and obtund acrimonious humours: and thus become serviceable, internally, in pains, inflammations, heat of urine, hoarseness, tickling coughs, &c. in glysters, for lubricating the intestines, and promoting the ejection of indurated feces; and in external applications, for tension and rigidity of particular parts. Their common dose is half an ounce: in some cases, they are given to the quantity of three or four ounces. The most commodious forms for their exhibition, we shall see hereafter, in the chapter of Emulfions.

THE oils expressed from aromatic substances, differ from the foregoing, in retaining, for the most part, an admixture of the aro-

matic matter of the subject. Thus nutmegs and mace yield, upon expression, an oil impregnated with the slavour of the spices; and an oil expressed from aniseeds, has a great share of the peculiar smell of the seeds. A purgative oil also is extracted in America from the purgative seeds of the ricinus. It does not appear that other qualities of vegetables are communicated to their expressed oils.

THE rinds of the several varieties of oranges, lemons and citrons, yield by a kind of expresfion their effential oils almost pure, and nearly fimilar to those which are obtained from them by distillation. The essential oils, in which the fragrance and aromatic warmth of these fruits reside, are contained in numerous little vesicles, which may be diffinguished by the naked eye, spread all over the surface of the peel. If the rind is cut in flices, and the flices separately doubled or bent in different parts, and squeezed between the fingers, the vesicles burst at the bending, and discharge the oil in a number of fine flender jets. A glass plate being set upright in a glass or por-celane vessel, and the slices squeezed against the plate, the little jets unite into drops upon the plate, and trickle down into the vessel beneath. But though this process affords the true native oil, in the same state wherein it existed in the subject, unaltered by fire or other agents, it is not practicable to advantage, unless where the fruit is very plentiful; as only a small part of the oil it contains can thus be extracted or collected.

The oil is more perfectly feparated by rubbing the rind upon a lump of fugar. The fugar, by the inequality of its furface, produces the effect of a rasp, in tearing open the oily vesicles: and in proportion as the vesicles are opened, the sugarimbibes theoil. When the outward part of the lump is sufficiently moistened, it is scraped off, and the operation continued on the fresh surface. The oil thus combined with the fugar, is fit for most of the uses, to which it is applied in a fluid state. Indeed the pure essentials oils, obtained by distillation, are often purposely mixed with sugar, to render their use the more commodious.

CHAP.

#### CHAPTER III.

#### Infusions in different menstrua.

#### SECT. I.

#### Infusions and decoctions in water.

WATER, the direct memstruum of gums and falts, extracts readily the gummy and faline parts of vegetables. Its action, however, is not limited to these; the resinous and oily principles being, in most vegetables, so intimately blended with the gummy and faline, as to be in great part taken up along with them: some of the refinous cathartics, and most of the aromatic herbs, as well as bitters and aftringents, yield to water greatest part of their smell, taste, and medicinal virtue. Even of the pure essential oils, and odorous refins of vegetables, separated from the other principles, water imbibes a part of the flavour; and by the artificial admixture of gummy or faline matter, the whole substance of the oil or resin is made dissoluble in water.

Of pure falts, water dissolves only certain determinate quantities (see page 39): by applying heat, it is generally enabled to take up more than it can do in the cold, and this in proportion to the degree of heat; but as the liquor cools, this additional quantity separates, and the water retains no more than it would have dissolved without heat. With gummy substances, on the other hand, it unites unlimitedly, dissolving more and more of them till it loses its fludity: heat expedites the action of the

water, but cannot enable it to take up more than it would do, by allowing it longer time, in the cold. The active parts extracted from most vegetables by water, and oils and refins made soluble in water by the artificial admixture of gum, partake of this property of pure gums, being dissoluble without saturation.

It has been imagined that vegetables in a fresh state, while their oily, refinous, and other active parts, are already blended with a watery fluid, would yield their virtues to water more freely and more plentifully, than when their native moisture has been dissipated by dry-Experience however shews. that dry vegetables, in general, give out more than fresh ones, water feeming to have little action upon them in their recent state. If, of two equal quantities of mint, one be infused fresh in water, and the other dried, and then infused in the like quantity of water for the same length of time, the infusion of the dry herb will be remarkably the strongest: and the case appears to be the same in all the vegetables that have been tried.

In all the preparations described in this chapter, it is to be understood that the subjects must be moderately and newly dried; unless when they are expressly ordered to be taken fresh; in which case it is to be judged that their virtues are defroyed or impaired by drying.

The native colours of many vegetables are communicated to water along with their medicinal matter; many impart a colour different from their own; and others, though of a beautiful and deep colour themselves, give scarcely any to the menstruum. Of the first kind are the yellow and red slowers; of the second, the leaves of most plants; of the third, some of the blue flowers, as those of cyanus and larkspur. Acid liquors change the insusions of most flowers.

ers, the yellow ones excepted, to a red; and alkalies, both fixt and volatile, to a green.

From animal fubstances, water extracts the gelatinous and nutritious parts, whence glues, gellies, broths, &c. and along with these, it takes up principles of more activity, as the acrid matter of cantharides. It dissolves also some portion of calcined calcareous earths, both of the animal and of the mineral kingdom, but has no action of any other kind of earthy matter.

#### ARTICLE I. Infusions in cold water.

Infusum CARDUI.

Infusion of carduus.

Take an ounce of the dried leaves of carduus benedictus, and a pint of common water. Let them steep for fix hours, without heat, and then filter the liquor through paper.

By this management, only the finer parts of the carduus are extracted, and the infusion proves an agrecable light bitter; it fits eafier on the stomach than any other medicine I know of the bitter kind; whereas, by long continued maceration, or by the application of heat, the groffer and more ungrateful parts are taken up, and the liquor becomes nauseous, so as to provoke vomiting. I have often given the light infusion, with great benefit, in weaknesses of the stomach, where the common bitters did not agree. It may be flavoured at pleasure with aromatic materials: instead of pure water, a mixture thereof, with some grateful distilled spirituous water, as twelve ounces of common water, and four of the spirituous water of orange-peel, may be

used for the menstruum. The little quantity of spirit contained in this compound will not considerably vary the dissolving power of the water.

MANY other vegetables may be advantageously treated in the same manner. From those which are weak in virtue, rich infusions may be obtained, by returning the liquor upon fresh quantities of the subject; the water loading itself more and more with the active parts. These loaded infusions are doubtless applicable to valuable purposes in medicine, as they contain, in a small compass, the siner, more subtile, and active principles of vegetables, in a form readily miscible with the sluids of the human body.

### TINCTURA MENTHÆ. Tincure of mint. Edinb.

Take half an ounce of the dry leaves of spearmint, and a pint of simple mint water. Steep them in a close vessel, in a warm place, for sour hours, and then strain out the tincture.

Тнв

THE distilled water of mint is impregnated with as much of the volatile parts of the herb, as water can be made to retain by distillation. By infusion, however, it still takes up more, being equally effectual, as a menstruum, with fresh water, hence the tincture proves very rich in the virtue of the mint. This is another useful method of obtaining strong infusions from vegetables, and it may be varjed at discretion: the distilled water of one plant may be employed as a menstruum for another.

Infusion of Peruvian bark.

Take an ounce of Peruvian bark reduced into fine powder, and twelve ounces of water. Macerate without heat for twenty-four hours, occasionly shaking the vessel; then pour off the clear liquor, and pass it through a fine

ftrainer.

THE extraction of the virtues of Peruvian bark, with aqueous liquors, has hitherto been attempted by ftrong coction: but this drug, contrary to most other vegetables, has lately been observed to give out more to cold than to boiling water. In boiling, a refinous matter, containg the astringency of the bark, is hastily melted out by the hear, but not truly dissolved by the water, and hence, in cooling, it begins to separate, renders the liquor turbid, and at length fettles to the bottom; whereas, by maceration in cold water, the affringent and bitter parts are gradually extracted together, and the former as well as the latter, are retained by the water in a state of perfect folution. The infusion appears to be one of the best preparations of the bark for weak flomachs, and may be given in doses of two or three ounces, in

intermitting fevers, and in other diforders where the corroborating virtues of bark are required.

AQUA PICEA.

Tar water.

Take of
Tar, two pounds;
Water, one gallon.
Stir them strongly together with a wooden rod; and after standing to settle for two days, pour off

the water for use.

TAR water has lately been recommended to the world as a certain and fafe medicine in almost all diseases; a slow yet effectual alterative in cachexies, scurvies, chlorotic, hysterical, hypochondriacal, and other chronical complaints; and a sudden remedy in acute distempers which demand immediate relief, as pleurifies, peripneumonies, the small pox, and all kinds of fevers in general. The medicine, though certainly far inferior to the character that has been given of it, is doubtless in many cases of confiderable utility: it fenfibly raises the pulse; and occasions some considerable evacuation, generally by perspiration or urine, though sometimes by stool or vomit: hence it is supposed to act by increasing the vis vitæ, and enabling nature to expel the morbific humours.

I shall here insert, from the first public recommender of this liquor (bishop Berkeley) some observations on the manner of using it. "Tar water, when right, is "not paler than French, nor deeper colour than Spanish white wine, and full as clear; if there be not a spirit very sensibly perceived in drinking, you may conclude the tar water is not good. It may be drank either cold or warm: in colics, I

" take

" take it to be best warm. As to " the quantity, in common chroni-" cal indispositions, a pint a day " may suffice, taken on an empty " ftomach, at two or four times, to " wit, night and morning, and " about two hours after dinner and " breakfast: more may be taken " by strong stomachs. But those " who labour under great and in-" veterate maladies, must drink a " greater quantity, at least a quart " every twenty-four hours: all of " this class must have much pa-" tience and perseverance in the " use of this, as well as of all " other medicines, which, though " fure, must yet in the nature of " things be flow in the cure of in-" veterate chronical disorders. In " acute cases, fevers of all kinds, " it must be drank in bed, warm, " and in great quantity (the fever " still enabling the patient to " drink) perhaps a pintevery hour, " which I have known to work fur-" prizing cures. But it works fo " quick, and gives such spirits, " that the patients often think " themselves cured before the fe-" ver hath quite left them."

## AQUA CALCIS SIMPLEX. Simple lime water. Lond.

Take a pound of quicklime, and a gallon and a half of water. Pour the water gradually upon the lime, and when the ebullition is over, let the whole stand to fettle; then filter the liquor through paper.

Edinb.

Takea pound of fresh-burnt quicklime, and two gallons of water. Pour the water by little and little upon the lime, and when the ebullition is over, strongly shake the vessel: then let the whole stand at rest, that the lime may settle, and after two days, filter the liquor, which is to be kept in vessels closely stopt.

A lime water may be prepared in the same manner from calcined oyster shells.

THE reason of adding the water by degree to the lime is, that when poured on at once, it reduces the external part to a kind of muddy substance, or soft paste, which in some measure defends the internal part from being acted upon by the water. It does not appear that the different proportions of water, in the two above prescriptions, occasion any fensible difference in the strength of the product; the quicklime is far from yielding all its soluble parts to either proportion; the remainder giving a strong impregnation to many fresh quantities of water, though not fo strong as to the first. The caution of keeping the water in close-stopt vessels, ought to bestrictly attended to; for in open ones, the calcare. ous matter, dissolved in the liquor, foon begins to separate, and forms a white crust upon the surface. This crust is not of a saline nature, as some have imagined; but an infipid earth, no longer mifcible with watery liquors.

Lime water has been found of great fervice in scrophulous and fcorbutic complaints, in some kinds of alvine fluxes, female weaknesses. and other diforders, proceeding from a laxity and debility of the folids: particularly in corpulent and phlegmatic habits. It appears likewife to be possessed of a lithontriptic power, and in fundry calculous cafes has procured confiderable relief: the lime water prepared from calcined oyster shells is found to be, in this intention, more efficacious than that of the common stone or chalk lime. It is given internally, in the

dof

dose of a quarter of a pint, three or four times a day; and likewise used externally for washing foul ulcers.

#### A Q U A C A L C I S. COMPOSITA.

Compound lime water.

Edinb.

Take of

Sasiafras, root and bark, shaved, two ounces:

Nutriegs, well bruised, three drachms;

Liquorice, fliced, one ounce; Lime water, fresh prepared, four pints.

Digest them together for two days, in a very close vessel; and then strain the liquor.

### AQUA CALCIS MINUS COMPOSITA.

Lime water less compounded.

London.

Take of

Liquorice, one ounce; Saffafras bark, half an ounce; Simple lime water, fix pints.

Macerate without heat for two days, and then strain off the liquor.

#### AQUA CALCIS MAGIS COMPOSITA.

Lime water more compounded.

Lond.

Take of

Guaiacum wood, shaved, half a pound:

Liquorice, one ounce; Saffafras bark, half an ounce; Coriander feeds, three drachms; Simple lime water, fix pints.

Macerate without heat for two days, and then strain off the liquor.

THIS last water has been used .

for some time in our hospitals, under the title of AQUA LIBERANS. As the guaiacum wood difficultly communicates its virtues to the cold liquor, some have proposed boiling it in the lime water before the other ingredients are added; but though this treatment more perfeetly extracts the virtues of the wood, it very much injures those of the lime water, greatest part of the matter it had taken up from the lime being separated and thrown off in the boiling. Nor indeed is there any occasion to have recourse to expedients of this kind; the quantity of the wood in the above prescription being so large, that the liquor receives a sufficient impregnation from it by maceration in the cold. If however, on this or other occasions, it should be though expedient to increase the dissolving power of lime water by boiling, we may do it without any injury to the lime water, by the method directed by the London college for obtaining a folution of fulphur in this menstruum, viz. by adding some quicklime in substance, which will continue to give a fresh impregnation to the water, affer the lime at first dissolved in it has been separated by the boiling.

In all these compositions, the additional articles take off the ill stavour of the lime water, render it more grateful both to the palate and stomach, and at the same time considerably promote its medicinal essimatory, especially when intended against cutaneous disorders, and soulness of the blood and juices. They may be taken in the same quantities as the simple lime water, and continued for some time; the patient keeping moderately warm

during their use.

ARTICLE

#### ARTICLE II. Infusions in boiling water.

## INFUSUM AMARUM. Bitter infusion. Edinb.

Take of
Gentian root, two drachms;
Leffer centaury tops, half an
ounce;

Boiling water, a pint.

Infuse them for four hours, then filter the liquor, and add to it an ounce of aqua aromatica, or of spirituous cinnamon water.

#### INFUSUM AMARUM SIMPLEX.

Simple bitter infusion. Land.

Take of

Gentian root,

Fresh yellow rind of lemon peel, carefully freed from the inner white part, each half an ounce;

Dry yellow rind of Seville orange peel, freed in like manner from the white, one drachm and a half;

Boiling water, three quarters of

Macerate for an hour or two, then filter the liquor thro' paper, or pass it through a strainer, without pressure.

Both these liquors are very elegant and useful bitters; the latter, in particular, is as agreeable a one as can well becontrived, the peels communicating a fine flavour, which is the only addition that the gentian stands in need of. The committee of the London college observe, that most of the ingredients, which usually enter the composition of bitter infusions, being prepared by them separately, amongst all the strong bitters, gentian gave the most unexceptionable colour,

w but it wants the affifiance of " fome ingredient to furnish an ac-" ceptable flavour; scarce any of " the bitters accompanied with fla-" vour, fuch as zedoary, calamus " aromaticus, and the like, ap-" peared to be truly grateful, ex-" cept orange peel and cardomom " seeds: but cardomom seeds are e mucilaginous, and render the li-" quor cloudy, and orange peel is " accompanied with a hot oil that " requires it to be but sparingly " used: lemon peel, in its outer " rind, to which all its flavour is " confined, is not a bitter, but sup-" plies the gentian most success-" fully with what is wanted; tho' " the composition, by a moderate " addition of orange peel, becomes " yet more perfect."

# INFUSUM AMARUM PURGANS. Purging bitter infusion. Lond.

Take of

oena ; Yellow rind of lemon peel, fresh,

each three drachms;

Gentian root;

Yellow rind of Seville orange peel, dry;

Leffer cardamom feeds, freed from the hufks, each half a drachm:

Boiling water, five ounces by measure.

Macerate them together, and when cold, ftrain off the liquor.

### INFUSUM AMARUM cum SENA.

Bitter infusion with sena. Edinb.

Take of Sena, one drachm;

Gen-

Gentian root, Sweet fennel feeds, each half a drachm; Boiling water, a quarter of a

pint,

Infuse them for four hours, and then strain the liquor.

This infusion may likewise be prepared with two, three, or more times the quantity of sena.

BOTH these are useful purging bitters. The quantities here prescribed are intended for one dose: the first is the largest, and the other the smallest dose that sens is usually given in.

#### INFUSUM SENÆ COMMUNE. Common infusion of sena.

Lond.

Take of

Sena, an ounce and a half; Crystalsof tartar, three drachms; Lesser cardamom seeds, freed from the huses, two drachms;

Water, one pint.

Boil the cryftals of tartar in the water, until they are diffolved; then pour the water, whilst it continues boiling, upon the other ingredients; and when cold, strain off the liquor for use.

In ourformer pharmacopœia, an alkaling falt was used in the infufion of fens, inflead of the acid one here directed. The first was supposed to promote the operation of the medicine, by fuperadding a degree of purgative virtue of its own, and by enabling the water to extract formewhat more from the capital ingredient, than it would be capable of doing by itself; whilst acids have rather a contrary effect. Experience, however, has sufficiently shewn (as the committee as-Inre us) " that this infusion and the

" following one with lemon jaice, " do not fail in their intention: " and in a medicine, very nauseous " to many, it is of principal con-" fequence to prepare it so, that " the lightest and least disgussful " parts may be extracted." Alkaline salts increase the offensiveness of the fena; whilst crystals of tartar confiderably improve the colour of the infusion, and likewise render the taste, to some persons, less disagrecable. Soluble tartar should feem a good ingredient in these kinds of compositions, as it not only improves the tafte, but promotes the purgative virtue of the medicine: this addition also renders the infusion less apt to gripe, or occasion flatulencies.

#### INFUSUM SENÆ, LIMONIATUM. Infusion of sena with lemon.

Take of

Sena, an ounce and a half; Yellow rind of lemon peel, fresh, one ounce;

Lond.

Lemon juice, one ounce, by measure;

Boiling water, one pint.

Macerate them together, and when
cold firain off the infusion.

This is a very pleasant and fusiciently efficacious purge: the committee observe, that it is the most agreeable form they have been able to contrive for the exhibition of fena to such as are more than ordinarily offended with its flavour. The dose is from two ounces to four.

INFUSI SENÆ UNCIÆ QUATUOR.

A four ounce infusion of sena-

Edinb.

Take of

Sena, three drachms; Ginger, Ginger, one scruple;
Boiling water, four ounces.
Infuse for four hours, and then
strain off the liquor.

This infution is tolerably grateful, the ill flavour of the fena being in good measure covered by the ginger, the quantity of which is here increased to double of that in former editions of the pharmacoposia. Formerly two drachms of the greater water-figwort were The water-figwort has added. been discovered to be the Brazilian. herb iquetaia, celebrated as a specific corrector of the flavour of fena: that plant, however, has not been found from experience to anfwer this purpose so effectually as it was supposed to do before it was commonly known.

## INFUSUM RHABARBARI. Infusion of rhubarb. Edinb.

Take of rhubarb, fliced, one ounce:

Cochineal, one scruple;
Boiling water, one pint.
Insuse them for a night, and to
the strained liquor add one
ounce of spirituous cinnamon
water.

This appears to be one of the best preparations of rhubarb when designed as a purgative; water extracting its virtue more effectually than either vinous or spirituous menstrua: in this respect rhubarb differs from most of the other vegetable cathartics.

## TINCTURA ROSARUM. Tindure of roses. Lond.

Take of

Red rose buds, freed from the white heels, half an ounce;

Strong spirit (called oil) of vitriol, one scruple;

Boiling water, two pints and a half:

Double-refined fugar, one ounce and a half.

First mingle the spirit of vitriol with the water, in a glass, or glazed earthen vessel, and in this mixture macerate the roses; when the liquor is grown cold, strain it, and add the sugar.

#### Edinb.

Take of

Red rofes, cleared from the heels, and dried, one ounce; Spirit of vitriol, one drachm; Boiling water, two pints;

White fugar, two ounces.

Mix the acid spirit with the water,
and insuse the roses therein for
four hours; then silter the tincture, and add to it the sugar.

Some have directed the oil of vitriol to be dropped upon the roses before the water is put to them: but this method is certainly faulty: for fuch of the roses as this canftic liquor falls upon undiluted, will be burned up by it, and have their texture destroyed. Others have made an infusion of the roses in water first, and then added the acid, from an apprehension, that if this acid is added to the water, it would weaken its power as a menfroum; but, as the committee obferve, whatever the acid spirit will hinder the water from extracting, it must precipitate, if added afterwards; though in this preparation the oil of vitriol bears so small a proportion to the water, that its effect, in this respect, will be very little. The infusion should be made in a glass, or stone-ware vessel, rather than a glazed earthen one; for the acid will be apt to corrode the glazing of the latter.

T 2 This

This tincture is of an elegant red colour, and makes a very grateful addition to juleps in hæmorrhages, and all cases that require mild coolers and subaftringents: it is sometimes taken with boluses or electaries of the bark, and likewise makes a good gargle.

INFUSUM LINI.
Infusion of linseed.

Take of

Linfeed, whole, two spoonfuls; Liquorice, sliced, half an ounce; Boiling water, four pints.

Let them fland in infusion by the fire for some hours, and then flrain off the liquor.

An ounce of coltsfoot leaves is fometimes added to these ingredients; which addition procures this medicine the title of INFU-SUM PECTORALE, pelloral infusion. Both insusions are soft, emollient, mucilaginous liquors, and as such they are directed in defluxions of thin acrid rheums, and erosions of the vessels. They are given to the quantity of a pint a day.

INFUSUM ANTISCORBUTICUM.

Antiferbutic infusion.

Take of

Buckbean leaves, two ounces; Curaffao oranges, half an ounce; Compound horferadish water, four ounces;

Common water, four pints.

Let the common water, boiling, be poured on the buckbean and orange, and fuffered to stand in a close vessel for a night; then strain out the liquor, and add to it the horseradish water.

This infusion is a very useful, and not inelegant antiscorbutic: buckbean appears from experience to be a very efficacious herb in this intention: the aromatic material, here joined to it, alleviates its ill flavour, and at the fame time promotes its virtue. A quarter of a pint of the liquor may be taken three or four times a day.

INFUSUM CEPHALICUM.

Cephalic infusion.

Take of

Wild valerian root, two ounces; Rofemary, or fage, half an ounce:

Aromatic water, four ounces; Common water, four pints.

Let the common water be poured, boiling, on the herb and root, and suffered to stand for a night in a close vessel; then strain out the infusion, and add to it the aromatic water.

This infusion is calculated against epileptic disorders, and other like affections of the nervous system. The dose is a quarter of a pint, to be taken twice a day.

INPUBUM ALCALINUM.

Alkaline infusion.

Take of

Salt of tartar, half an ounce; Saffron, half a drachm; Liquorice root, two ounces; Boiling water, three pints.

Let them fland together in a warm place for eight or ten hours, and then strain out the liquor for use.

This infusion is of service in a lentor, or viscidity of the blood and juices, the consequence of an obstructed perspiration, and oftentimes the origin of inflammatory distempers: it attenuates thick humours, and promotes the natural secretions. It is to be taken warm, in little quantities at a time, but frequently repeated.

INFUSUM

1

INPUSUM DIURBTICUM. Diuretic infusion.

Take of

Wormwood leaves, dried, half on ounce ; Salt of tartar, two scruples;

Compound juniper water, two ounces;

Common water, twelve ounces. Pour the common water, boiling, on the wormwood and falt of tartar, and, when grown cold, firain off the liquor, and mix with it the juniper water.

This infusion is much of the fame nature with the foregoing. It is directed in the obstructions of the viscera which frequently succeed a long continuance of bilious fevers, or frequent relapses into them, and which generally end in a dropfy, jaundice, or irregular intermittent. The quantity here prescribed is to be taken every day, at three doses, and a purgative occasionally interposed. If intermittent fevers return after the care of the other disorders. they are then successfully treated by the bark.

Preparations of this kind are likewise of considerable use in maniacal disorders; in which, as Dr. Mead observes, evacuations by the kidneys are of greater consequence than is generally supposed; especially if the mania is of the furious kind, and accompanied with febrile heat. Alkaline salts, given in large doses, are here the most effectual dinretics.

INFUSUM PARALYTICUM. Paralytic infufion. Take of

Horseradish root, shaved, Mustard-seed, bruised, each four ounces;

Boiling water, four pints.

Let them steep together, in a close veffel, for 24 hours.

This infusion is strongly impregnated with the pungency of the mustard-seed and horse-radish, which, by this simple process, give out the whole of their virtues. Though the medicine is defigned chiefly (as its title expresses) for a stimulant in paralytic complaints, there are several other disorders in which it may be employed to good advantage; in scorbutic cases, in particular, it promises to be a remedy of great utility: it generally promotes the urinary discharge, and, if the patient is kept warm, perspiration. It is taken sometimes to half a pint, twice a day,

THEA ANTIPHTHISICA. Antiphthisical tea.

Take of

Avens root, two ounces; Male speedwell.

Ground-ivy, each one ounce and a half:

Liquorice, one ounce; Sweet fennel feeds, 3 drachms.

THESE ingredients are to be cut, bruised, and well mixed together, and half an ounce of the composition insused, for a few minutes, in five or fix tea-cups full of boiling water. In consumptive cases, and disorders of the breaft, one cup of the infusion, with a tea-spoonful of honey, may be drank every hour or two. After the same manner, medicated teas may be prepared from other vegetable substances, as chamomile flowers, linfeed, orange-peel, fumitory, &c.

INFUSUM CINAMOMI. Infusion of cinnamon. Take two ounces of powdered cinnamon, and two pints of boiling boiling water. Infuse them in a close vessel, in a moderate heat, for half an hour; and then filter the liquor.

This infusion is agreeably im-

pregnated with the flavour and warmth of the spice, and may, on many occasions, supply the place of the simple cinnamon water.

#### ARTICLE III. Decoctions.

THE effect of boiling differs from that of infusion in some material One of the most particulars. obvious differences is, that as the effential oils of vegetables, in which their specific odours reside, are volatile in the heat of boiling water, they exhale in the boiling along with the watery stream, and thus are loft to the remaining decoction; whereas both in cold and hot infusions they are preserved. Odorous substances. and those in general whose virtues depend on their volatile parts, are, therefore, unfit for this treatment. The foluble parts of these may, nevertheless, be united in this form with those bodies of a more fixed nature; by boiling the latter till their virtues are fufficiently extracted, and then infusing the former in this decoction.

The extraction of the virtue of the subject is usually promoted or accelerated by a boiling heat; but this rule is less general than it is commonly supposed to be. have already observed, that Peruvian bark gives out its virtue more perfectly by cold infusion than by coction. In some cases, boiling occasions a manifest disunion of the principles of the subject: thus, when almonds are triturated with cold water, their oil, blended with the mucilaginous or other soluble matter of the almond, unites with the water into a milky liquor called an emulfion: but

on boiling them in water, the oil feparates and rifes to the furface; and if the most perfect emulsion be made to boil, a like separation happens.

### DECOCTUM ALBUM. The white decoction. Land.

Take of
Calcined hartshorn, prepared,
two ounces;
Gum Arabic, two drachms;
Water, three pints.
Boil them till only two pints remain, and then strain off the liquor.

Edinb.

Take of
Calcined hartshorn, prepared,
one ounce;
Gum Arabic, two drachms;
Common water, three pints;
Cinnamon bruised, one drachm;
White sugar, two drachms.
Boil the calcined hartshorn and
gum in the water till only two
pints remain, adding the cinnamon towards the end: in this
decoction, unstrained, dissolve
the sugar.

THESE decoctions are used as common drink in acute diseases, attended with a looseness, and where acrimonious humours abound in the primæ viæ. The gum is added in order to render the liquor lightly glutinous, and thus enable it to sustain more of the calx, which is the ingredient

that

that the colour, but probably not the virtue, of the medicine depends upon. Calcined hartshorn has no quality from which it feems capable either of confiringing and frengthening the veffels, giving a greater degree of confiftency to thin fluids, or odtending acrimosous humours. It blunts and abforbs acid juices; but acrimony and acidity are very different: there are few (perhaps none of the acute) diforders of adults attended with the latter; and few of infants are unaccompanied therewith. Some have proposed starch as an ingredient in these kinds of decoctions: a fmall quantity of this foft, gelatinous, farinaceous substance should seem to be greatly preferable to the earthly calx. It may be observed, that the water is not enabled by the boiling to diffolve any part of the calx; and that in the decoction, the earth is only diffused in substance through the water, as it would be by agitation.

#### DECOCTUM ALBUM COM-POSITUM.

Compound white decoction.

Edinb.

Take of
Comfry roots,
Tormentil roots,
Calcined harthorn,
Chalk,
White fugar, of each half an
ounce;
Cinnamon bruifed, one drachm;
Common water, three pints.
Boil the roots in the common wa-

Boil the roots in the common water, till such time as the liquor, when strained, will amount only to a quart, adding the cinnamon towards the end: strain the decoction, add to it the calcined hartshorn, chalk, and sugar, and mix them well together.

Ł.

This is a very well contrived composition for the purposes of a mild, lightly incrassating restringent. A quarter of a pint, more or less, may be taken occasionally, according to the argency of the symptoms. The calcined hartshorn and chalk appear to be the least useful of its ingredients.

# DECOCTUM CRETACEUM. Chalk Decoction. Edinb.

Take of

White chalk, prepared, one ounce;

Nutmeg, bruifed, one drachm; Gum Arabic, two drachms; White fugar, half an ounce; Common water, three pints.

Boil the water with the chalk and gum, till it is reduced to a quart, adding the nutmeg towards the end; and in the turbid decoction diffolve the fugar.

HERE, as in the white decoction, the absorbent earth is only mixed in substance with the water, and the use of the gum is to prevent its subsiding. As a medicine, chalk is more effectual than calcined hartshorn, in all the intentions for which these kinds of earths are given.

# DECOCTUM JAPONICUM. Japonic decoction. Edinb.

Take of

The confectio japonica (describhereaster among the electaries) one ounce;

Common water, a pint and a half;

Spirituous cinnamon water, Syrap of meconium, each one

Boil the confection in the common water, till the liquor, after firstning, firaining, will amount to a pint, to which, while turbid, add the cinnamon water and the fyrup.

This decoction is used both in draughts and in glysters, as an anomodyne and reftringent in fluxes. The quantity here prescribed contains two grains and a half of opium, exclusive of the syrup.

## DECOCTUM ad ICTERICOS. Decoction for the jaundice. Edinb.

Take of

Celandine, roots and leaves, Turmeric.

Madder, each one ounce; Millipedes, two hundred;

Water, three pints.

Boil the celandine, turmeric, and madder in the water, till only a quart of liquor remains after fraining; then, having pressed out the juice of the millepedes, add this to the decoction when grown cold.

THE ingredients of which this decoction is composed, have been long held by many as specifics for the cure of the disease expressed in its title. The medicine; though not a little unpleasant, is well calculated to answer many useful phriposes, if well managed and properly assisted. A quarter of a pint may be taken twice a day, or oftener,

# DECOCTUM LIGNORUM. Decoction of the woods. Edinb.

Take of

Guaicum shavings, three ounces; Raisins of the sun, stoned, two ounces;

Sasiafras wood, shaved, one ounce;

Liquorice, fliced, half an ounce; Water, one gallon.

Boil the guaicum and raisins with the water over a gentle fire, to the consumption of one half; adding, towards the end, the fassafras and liquorice. Strain out the liquor, and having suffered it to rest for some time, pour off the clear from the feces.

This decoction is very well contrived, and, if its use is duly continued, will do great service in some cutaneous diseases, soulness of the blood and juices, and some disorders of the breast, particularly in cold phlegmatic habits. It may be taken by itself, in the quantity of a quarter of a pint, two or three times a day, or used as an assistant in a course of mercurial or antimonial alteratives; the patient, in either case, keeping warm, in order to promote the operation of the medicine.

#### DECOCTUM ad NEPHRITICOS.

Nephritic decoction. Edinb.

Take of

Marshmallow roots, one ounce and a half;

Liquorice,

Linfeed, each half an onnce; Pellitory of the wall, one ounce; Raifins of the fun, floned, two ounces;

Water, fix pints.

Boil the water, with the markmallow roots and raifins, to four pints, adding the other ingredients towards the end. Strain out the liquor, and let it fettle till fine.

This decoction is intended chiefly as an emollient, to be liberally

berally drank of in nephritic paroxysms; in which cases, by softening and relaxing the parts, it frequently relieves the pain, and procures an easy passage for the fabulous matter. The medicine is now made more simple than before, without any diminution of its virtue, by the rejection of wild carrot feed, restharrow root, and figs, the place of which is abundantly supplied by an increase of the marshmallow root, linseed, and liquorice. The carrot feeds were indeed unfit for this form, as they give out little of their virtue to watery liquors.

## DECOCTUM NITROSUM. Nitrous decodion. Edinb.

Take of
Pure nitre, half an ounce;
White fugar, two ounces;
Cochineal, one scruple;
Water, two pints and a half.
Boil to two pints, then suffer the whole to rest for some time, and pour off the clear decoction.

This is an elegant way of difguifing nitre, and rendering it agreeable to the patient, both which intentions are fully answered by the cochineal and sugar. There does not feem to be any occasion for so long boiling; for the water will dissolve a much larger quantity of the nitre and sugar than is directed above, without any heat, and it easily extracts a sine colour from cochineal.

The virtues of nitre have been already mentioned in the preceding part. This or other similar forms are the most commodious for the exhibition of it; for when given in a solid form, it often occasions great uneasiness about the somach. Two or three ounces

of this decoction may be taken for a dose.

## DECOCTUM PECTORALE. Pettoral decoction. Lond.

Take of
Common barley,
Stoned raifins,
Figs, each two ounces;
Liquorice, half an ounce;
Water, four pints.

First boil the water with the barley, then add the raisins, and, lastly, (just before the end of the process) the sigs and liquorice. The boiling is to be continued so long, that the liquor, when strained, may be no more than two pints.

Edinb.

#### Take of

ske or
Stoned raifins of the fun,
Barley, each one ounce;
Fat figs, in number four;
Florentine orris root,
Liquorice,
Coltsfoot flowers, each half an
ounce;
Water, fix pints.

Boil the water with the raifins, barley, and figs, until only four pints remain; adding, towards the end, the other ingredients; then ftrain out the liquor for use.

Both these decoctions are useful soft pectorals, and very agreeable to the palate, particularly the sirft. They are good auxiliaries in sharp defluxions on the breast and lungs, and have sometimes done service by themselves. They may be drank at pleasure.

### DECOCTUM SERPENTARIÆ COMPOSITUM.

Compound decoction of fnakeroot.

Edinb.

#### Take of

Virginian fnakeroot, 6 drachms; Edinburgh theriaca (described hereaster hereafter among the electaries) half an ounce; Cochineal, one scruple; Water, two pints.

Boil the water with the fnakeroot

to one half, adding the theriaca and cochineal towards the end: then strain out the liquor for use.

This preparation is an useful sudorific and alexipharmac, containing nearly all the virtue of the snakeroot, and great part of that of the theriaca. The quantity of theriaca here prescribed holds nearly three grains and a half of opium; so that about a sist of a grain of opium, or somewhat more, goes to an ounce measure of the decoction.

# DECOCTUM TAMARINDORUM cum SENA.

Decoction of tamarinds with sena.

Edinb.

Take of

Tamarinds, fix drachms;
Cryftals of tartar, two drachms;
Sena, one drachm;
Syrup of violets, one ounce;
Simple cinnamon water, half an ounce;

Common water, a pint and a half.

Boil the common water with the tamerinds and crystals of tartar, fo long that there may be a pint of strained liquor; in which, whilst hot, infuse the sens for four hours; afterwards strain off the liquor, and add to it the syrup of violets and cinnamon water.

This decoction may likewife be prepared with two, three, or more times the quantity of sena.

THIS is a fufficiently efficaciens, and not disagreeable cooling. purge. The quantity here preferibed, is intended for a dose, which may be divided into three or four parts, to be taken at thort intervals, as the stomach will bear it.

## AQUA HORDEATA. Barley evater. Lond.

'Take of Pearl barley, two ounces; Water, four pints.

First wash the barley from the mealy matter that adheres to it, with some cold water: then boil it a little with about half a pint of fresh water, which will acquire a considerable tinge from it. Throw away this tinged water; put the barley into the water prescribed, made arft to boil; and continue the boiling till half the water is wasted.

This liquor is to be drank freely, as a diluter, in fevers and other disorders; hence it is of consequence that it should be prepared fo as to be as elegant and agreeable as possible: for this reason, it was inferted in the pharmacopœia, and the several circumstances which contribute to its elegancy fet down: if any one of them is omitted, the beverage will be less gratefui. However trivial medicines of this class may appear to be, they are of greater importance, in the cure of acute difeases, than many more laborious preparations.

#### MUCILAGO SEMINUM CYDONIORUM. Mucilage of quince seeds.

Lond.

Take of

Quince feeds, one drachm; Water, fix ounces by measure.

Boil

Boil them over a foft fire, till the water grows slimy, almost like the white of an egg; then pass it through a linen cloth.

This is a pleafant foft mucilage, of a fomewhat fweetish taste, and a light agreeable smell: in these respects, and its easy solubility in water, it differs from the mucilage of gum tragacanth, which some have supposed it similar to: it has another difference, to its disadvantage, being apt to grow mouldy in keeping.

# GELATANI CORNU CERVI. Jelly of harifborn. Edinb.

Take of

Hartshorn shavings, half a pound; Water, three quarts;

White fugar, fix ounces;
Mountain wine, a quarter of a

Orange (or lemon) juice, one ounce.

Boil the hartshorn with the water, by a gentle heat, in a glazed earthen vessel, till two parts are wasted; strain out the remaining liquor, add to it the other ingredients, and boil the whole over a gentle fire, to the consistence of a soft jelly.

# JUS VIPERINUM. Viper broth. Lond.

Take a middle fized viper, freed from the head, skin, and inteftines, and two pints of water. Boil them to a pint and a half; then remove the vessel from the fire; and when the liquor is grown cold, let the fat, which congeals upon the surface, if the viper was fresh, be taken off. Into this broth, whilst warm, put a pullet of a moderate size, drawn and freed from the skin,

and all the fat, but with the flesh intire. Set the vessel on the sire again, that the liquor may boil; then remove it from the fire, take out the chicken, and immediately chop its slesh into little pieces: put these into the liquor again, set it over the sire, and, as soon as it boils up, pour out the broth, sirst carefully taking off the scum.

HERE all the circumstances subfervient to the persection of the broth are carefully set down: and even plain chicken broth, for the use of the sick, ought to be made in a similar manner.

This seems to be one of the best preparations of the viper; all the benefit that can be expected from that animal being by this means obtained. It is a very nutritious and reflorative food; continued for. a length of time, it has sometimes done good fervice in leprous and other obstinate cutaneous diseases. The dried flesh of the vipers. brought from abroad, is not at all superior to the fresh vipers of our own country: the wines and tincture of the animal, probably, have little virtue: the volatile falt, however strongly recommended by fome, does not appear to differ from that producible from every animal substance. See chap. viii. fect 2.

### DECOCTUM ANTIHECTICUM. Antihedic decodion.

Take of

Comfry root,

Eryngo root, each half an ounce; Conferve of rofes, two ounces; Dulcified spirit of vitriol, forty

drops; Water, three pints.

Boil the water with the roots and the conferve, till one pint is wasted; then strain off the remaining maining liquor, and add to it the dulcified spirit.

This decoction is usefully given in hectic cases, where thin acrimonious humours abound, and in beginning consumptions. The dose is a quarter of a pint, to be taken two or three times a day.

DECOCTUM VULNERARIUM.
Vulnerary decoction.

Take of

The herb ground-ivy, Plantane leaves.

White sugar, each half an ounce; Water, three pints.

Boil the herbs in the water so long that there may be only two pints of strained liquor, in which dissolve the sugar.

The herbs which give virtue to this decoction, have long been cebrated as specifics for the cure of internal contusions and ulcerations, of coughs and pulmonary phthises, proceeding either from bruises, or an erosion of the viscera from a spontaneous acrimony of the hamours Though the real virtues of these plants fall short of the character which has been usually given of them, yet experience has shewn that they are superior to numerous others which have been very strongly recommended.

DECOCTUM ANTIFEBRILE.

Antifebrile decoction.

Take of

Virginian snakeroot, bruised, Peruvian bark, in powder, each three drachms;

Water, one pint.

Water, one pint.

Boil them to half a pint, and, having strained off the liquor, mix with it, of Spirituous cinnamon water, an ounce and a half;

Syrup of clove july-flowers, two drachms.

In the putrid malignant fever, arising from foul air in crowded bospitals and jails, this medicine has been given with remarkable fuccess. In the low state of this dangerous disease, when the pulse, before quick, begins to fink, the stupor to increase, and petechize to appear, it promises to be a very useful remedy for supporting the vis vitæ, promoting a critical diaphoresis, and correcting the putrid humours. Four spoonfuls of the decoction are to be taken every four or fix hours; and moderate quantities of wine or cordial boluses, with volatile salts, interposed at proper intervals.

DECOCTUM PEBRIPUGUM.

A febrifuge decedion.

Take of

Chamomile flowers, dried, two

Salt of tartar, two drachms; Water, three pints.

Boil the water with the chamomile flowers, till one pint of the liquor is wasted; then strain out the remaining decoction, and diffolve it in the alkaline salt.

In a thick viscidstate of the blood and juices, and obstructions of the abdominal viscera, a quarter of a pint of this decoction, taken three or four times a day, has sometimes removed intermittent severs, after the Peruvian bark had been tried in vain. It is nearly similar to the alkaline and diuretic insusions described above.

Apozema aperiens.

Aperient apozem.

Take of
Rhubarb,
Madder, each three drachms;
Salt of tartar, two drachms;
Water, three pints.

Boil them together for an hour, and having strained out the decoction, add to it three ounces of fyrup of ginger.

This promifes to be a very powerful aperient and attenuating medicine, of great service in isterical and hydropic cases. The dose is three ounces, which may be repeated thrice a day.

DECOCTUM ASTRINGENS.

Afteringent decoction.

Take of

Tormentil root, one ounce; Pomegranate peel,

Plantane leaves, each half an ounce;

Syrup of dry roses, one ounce; Water, three pints.

Boil the water with the tormentil, granate peel, and plantane, till one pint is wasted, adding the cinnamon towards the end: then strain off the decoction, and mix with it the syrup.

THE title of this preparation sufficiently expresses its virtues. The dose, in sluxes where the morbid matter has been evacuated, and astringency is the only indication, is from one to sour ounces, three or sour times a day.

DECOCTUM BARDANE.

Decoction of burdock.

Take of

Burdock roots, two ounces; Vitriolated tartar, one drachm; Water, three pints:

Boil the water with the roots, fo long, that the liquor, when firained, may amount only to a quart; to which add the vitriolated tartar.

This decoction is drank to the quantity of a pint a day, as a mild aperient, diuretic, and fweetener,

in scorbutic and rheamatic complaints.

Decoctum Campechense.

Decocion of logwood.

Take of

Shavings of logwood, three ounces;

Cinnamon, two drachms; Water, four pints.

Boil the water with the logwood till half the liquor is waited, adding the cinnamon towards the end of the boiling; then strain out the decoction for use.

This is an agreeable mild reftringent, in diarrhœas and other fluxes, where stronger astringents would be improper or unsafe. It is given in the hospitals in doses of a quarter of a pint three or four times a day. It generally tinges the stools red, which has occasioned some to be alarmed, as if the colour proceeded from a discharge of blood: the patient, therefore, is to be cautioned against any surprize on that account.

Decoctum diureticum.

Diuretic decoction.

Take of.

Parsley, or fennel roots, one ounce;

Wild carrot feeds, 3 drachms; Pellitory of the wall, half an ounce;

Raisins, two ounces; Nitre, one drachm;

Water, three pints.

Boil the water with the roots, feeds, pellitory, and raifins, fo long, that there may be only two pints of liquor after straining; in which dissolve the nitre.

Take of 2.
Grafs roots, two ounces;
Sorrel, or wood forrel leaves,
one handful;

Tamarinds,

half t

Nitre, two drachms: Barley water, three pints.

Boil the roots in the barley water, till one pint of the liquor is wasted, adding towards the end the forrel, tamarinds, and nitre: then strain out the apozem for use.

Take of Marshmallow roots, fresh, one pound; Fennel roots, half a pound; Nitre, half an ounce;

Water, one gallon. Boil the water with the roots, till one-fourth of the liquor is wasted; then strain off the remaining decoction, and dissolve in it the nitre.

THESE cooling aperient liquors are nsed like the nephritic decoction already described as common drink for promoting urine in nephritic diseases. They may be taken with safety, and often with good effect, in inflammatory cases. where the hot stimulating diureties would be manifestly prejudicial.

DECOCTUM PERUVIANUM. Peruwian decoction.

Take of

Peruvian bark, in powder, two ounces;

Water, three pints.

Boil them together, till one pint of the liquor is wasted, and then strain off the remaining decoction for use.

This decoction should be passed only through a coarfe strainer, and drank whilst turbid: if suffered to fland till clear, the more efficacious parts of the bark will subside. We have formerly observed, that the

Tamarinds, an ounce and a virtues of this drug confift chiefly in its refinous substance, which, though it may be totally melted out by the heat of boiling water, remain only partially suspended in that menstruum. See p. 270.

> DECOCTUM SENERAL. Decoction of seneka. - Take of

Seneka, rattle-soake root, one ounce;

Water, a pint and a half. Boil to one pint, and ftrain.

THE virtues of this decoction will be eafily understood from those of the root which it is prepared from. See page 225. The dose, in hydropic cases, and rheumatic, or arthritic complaints, is two ounces, to be repeated three or four times a day, according to its effect.

DECOCTUM TERRES TABONICE. Decoction of Japan earth. Take of

Japan earth, two drachms; Spirituous cinnamon water. Syrup of quinces, each two ounces :

Common water, one pint. Boil the common water with the Japan earth, till about one fourth of the liquor is wasted; then fuffer the decoction to fettle, and having poured off the clear part, add to it the spirituous water and the fyrup.

This decoction is a very agreeable and useful medicine in fluxes, that are not critical or fymptomatic, and in a weak, lax flate of the intestines. A spoonful or two may be taken every hour or oftener: thus managed, it produces much better effects than if larger dofes are given at once.

FOTUS COMMUNIS.

The common fomentation.

Lond.

Take of

Abrotanum leaves, dried, Sea wormwood tops, dried, Chamomile flowers, dried, each one ounce;

Bay leaves, dried, half an ounce: Water, fix pints.

Lightly boil them, and firain out the decoction for use.

IT is left to the choice of the apothecary to take either the male or female abrotamum, that is, fouth-' ernwood, or lavender cotton: which, though differing from one another in some respects, may be looked upon as fimilar with regard to the purposes for which this composition is intended: nor, indeed, can either of them give much affistance to chamomile flowers and wormwood. The use of this decoction is expressed in its title: spirit of wine, which is commonly added in fomentations, is left to be directed by the prescriber, in fuch quantity as particular cases may require.

DECOCTUM COMMUNE pro CLYSTERE.

The common decoction for glysters.

Lond.

Take of

Mallow leaves, dried, one ounce; Chamomile flowers, dried, Sweet fennel feeds, each half am ounce;

Water, one pint.

Boil them together, and strain out the decoction for use.

THE title of this decoction sufficiently expresses its use, as the basis of glysters. The ingredients should be very lightly boiled, or at least the chamomile slowers and fennel seeds not put in till towards the end, a part of the virtue of these being soon lost by boiling.

DECOCTUM COMMUNE.

The common decoction. Edinb.

Take of

Chamomile flowers; one ounce; Elder flowers, Sweet fennel feeds, each half an ounce;

Water, two quarts.

Make them just boil, and then strain out the liquor. The virtues of the ingredients may be sufficiently extracted also, by infusing them, for some hours, in the boiling water.

This decoction is intended to answer the purposes of both the foregoing. It is less loaded with the ingredients than either; but not, perhaps, for that reason the less useful.

Fotus Anodynus.

Anodyne fomentation.

Take of

Garden poppy heads, one ounce; Elder flowers, half an ounce; Water, three pints.

Boil them till one pint is wasted, and then strain out the liquor for use.

This fomentation is prescribed for tumested and inflamed parts, to abate the inflammation and pain. Whether the opiate matter in the poppy heads contributes much to this intention, may be questioned; as the effects of the composition may be attributed, perhaps, more to the warm fluid softening and relaxing the skin, than to the particular qualities of the matters which it is impregnated with.

Fotus Aromaticus,
Aromatic fomentation.

Take of
Cloves,
Mace, each one drachm;
Red wine, one pint.

Boil them a little, and strain off the liquor.

This preparation is intended not only as a mere topical application for external complaints, but likewise for relieving the internal parts. The pains of the bowels which accompany dysenteries and diarrhoeas, flatulent colics, uneasiness at the stomach, and reachings to vomit, are frequently abated by fomenting the abdomen and region of the stomach with the warm liquor.

FOTUS ROBORANS.

Strengthening fomentation.

Take of
Oak bark, one ounce;

Granate peel, half an ounce; Alum, two drachms;

Smith's forge water (that is, water in which red-hot iron has been feveral times quenched) three pints.

Boil the water with the oak bark and granate peel, to the confumption of one-third; then firain the remaining decoction, and diffolve it in the alum.

This is a strong aftringent liquor, in which intention it is directed, both as a fomentation for strengthening relaxed parts, and as an injection in the sluor albus.

#### SECT. II.

Wheys.

#### SERUM SOLUTIVUM.

Laxative whey.

Take of

Damask rose buds, fresh, one ounce;

Whey, two pints.

Steep them together for a night, and then firain out the whey for ule.

WHEY, thus impregnated with the virtues of the damask rose, operates very gently by stool, and for this purpose is held by some in great esteem. Its action may be quickened, and its taste rendered more agreeable, by the addition of a suitable proportion of crystals of tartar.

SERUM SINAPINUM.
Muftard whey.

Take of

Mustard seed, bruised, three spoonfuls;

Cows milk, two pints. Set the milk over the fire to boil, and add to the mustard feed, that a curd may be formed, from which the whey is to be carefully separated.

This is not an inelegant form for the exhibition of mustard seed; its pungency and medicinal virtues, depending thereon, being, in great measure, communicated to the whey.

### SERUM ALUMINOSUM. Alum whey.

Lond.

Take of

Cows milk, one pint;
Alum, in powder, two drachms.
Boil them till the milk is curdled,
then carefully separate the whey.

THIS medicine is a firong, tho' not very grateful, astringent. It is given in immoderate uterine fluxes, and sometimes in the diabetes, in which last intention it is recommended by Dr. Mead. The dose

is a quarter of a pint three or four times a day. It has been recommended also in intermittent fevers, the quantity above prescribed to be taken before the approach of a sit, divided into different doses: but in this disorder, great caution is requisite in the use of so strong an attringent.

SERUM SCORBUTICUM.

Scorbutic whey.

Lond.

Take of

Cows milk, one pint; Scorbutic juices, a quarter of a pint.

Boil them till the milk is curdled, , and then carefully separate the whev.

This whey may be used as common dripk in scorbutic cases: the quantity above directed, at least, ought to be taken every day, if any considerable effect is expected from it.

#### SECT. III.

#### Vinegars.

JINEGAR extracts the virtues of several medicinal substances in tolerable perfection; but at the same time its acidity makes a notable alteration in them, or superadds a virtue of a different kind; and hence it is more rarely employed in this intention, than purely aqueous, or spirituous menstrua. Some drugs however, vinegar, for particular purposes, excellently assists, or coincides with, as squills, garlick, ammoniacum, and others: and in many cases where this acid is it felf principally depended on, it may be advantageously impregnated with the flavour of certain vegetables; most of the odoriferous flowers impart to it their fragrance, together with a fine purplish, or red colour: violets, for instance, if fresh parcels of them are infused in vinegar in the cold for a little time, communicate to the liquor a pleasant flavour, and deep purplish red colour. Vinegar, like other acids, added to watery infations or decoctions, generally precipitates a part of what the waser had diffolved.

## ACETUM ROSACEUM. Vinegar of roses. Edinb.

Take of

Red roses dried, one pound; Strong vinegar, one gallon. Expose them to the sun in a close vessel, for forty days, and then strain off the liquor.

This is scarce otherwise made use of than for embrocating the head and temples, in some kinds of head ach, &c. in which it has now and then been of service.

# ACETUM SCILLITICUM. Vinegar of fquills. Lord.

Take of Dried squills, one pound:

Vinegar, fix pints.

Macerate the squills in the vinegar with a gentle heat; then press out the liquor, and set it by till the seces have subsided; the vinegar being afterwards poured off, add to it about one-twelfth its quantity of proof spirit, that it may keep the longer from growing mothery.

It should seem most convenient to add the spirit before the vinegar is decanted; for by this means, the purification is accelerated and rendered more persect; and the liquor, prevented from growing soul a second time, which it is apt to do upon the assussment of the spirit, however carefully it may have been depurated before.

#### Edinb.

Take of
Squills, cut into thin flices, one
pound;
Strong vinegar, fix pints.
Expose them to the sun after the
mannerdirected for making vinegar of roses; and afterwards
itrain out the liquor.

The root in this last prescription seems intended to be used fresh, in which case the vinegar proves much weaker of the squills than the first: a pound of the fresh squills being searcely equivalent to three ounces when dry. Vinegar has been supposed by some to diminish the virtue of squills: but this does not appear from experience, the acidity of the liquor only rendering the pungent bitterness of the root somewhat less perceivable.

Vinegar of iquills is a medicine of great antiquity: we find, in a treatile attributed to Galen, an account of its preparation, and of many particular virtues then afcribed to it. It is a very powerful fimulant, aperient, and attenuant of tenacious juices: and hence is frequently used, with good succels, in disorders of the breast occasioned by a load of thick viscid phlegm, for promoting urine in hydropic cases, &c. See the section of acrids, page 69. The dose of this medicine is from a drachm to half an ounce; where crydities abound in the first passages, it may

be given at first in a larger dose, to evacuate them by vomit. It is most conveniently exhibited along with cinnamon, or other agreeable aromatic waters, which prevent the nausea it would otherwise, even in small doses, be apt to occasion.

ACETUM PROPHYLACTICUM.

Prophyladic vinegar.

Paris.

Take of
Fresh tops of common wormwood,
Roman wormwood,
Ro emary,
Sage,
Mint,

Rue, each one ounce and a half; Lavender-flowers, dried, two ounces;

Ounces;
Garl ck,
Calamus aromaticus,
Cinnamon,
Cloves,

Nutmegs, each two drachms; Strong vinegar, eight pints. Digest them, by the heat of the fun or a sand bath, in a matrass closely stopt, for twelve days; then strongly pressout and strain the siquor, and having afterwards siltered it, add half an ounce of camphor dissolved in

spirit of wine.

This composition is designed, as its title expresses, for an antipestilential. It is said, that during the plague at Markeilles, four persons, by the use of this preservative, attended, unburt, multitudes of those who were insected; that under colour of these services, they robbed both the sick and the dead; and that one of them being afterwards apprehended, saved himself from the gallows by discovering the remedy. The preparation is hence called Vinaigre des quatre voleurs,

leurs, the vinegar of the four thieves. It is not to be doubted, that vinegar impregnated with antiseptic vegetables, will contribute greatly to prevent the effects of contagious air.

### ACETUM THERIACALE.

Treacle vinezar. Edinb.

Take of

Edinburgh theriaca, described hereafter among the electaries, one pound;

Strong vinegar, four pints. Digest them together, in a very gentle hear, for three days; and then strain out the vinegar for ufe.

This medicine has been greatly celebrated in acute and contagious diseases, as a sudorific and alexipharmac. Some have chosen to employ the vinegar as a vehicle, rather than as a menitruum, for the theriaca; in either case, it is indisputably, for sundry purposes, an useful addition. To half an ounce by measure of the compofition here prescribed, there goes fimewhat more than half a grain of opium; though it does not appear, that the medicine has all the effect which might be expected from that article.

#### ACETUM LITHARGYRITES. Vinegar of litharge.

Take of

Litharge, four ounces: Strong vinegar, one pint. Digell in a fand heat for three days, frequently thaking them; then filter the liquor for use.

This liquor is of the same nature with folutions of faccharum faturni, of which hereafter. It is only used externally, as a cosmeric. against cutaneous eruptions, rednels, inflammations, &c. But even here, it is not void of danger; there are examples of its continued use having occasioned fundry ill consequences.

### SEÇT. IV.

HE original intention of medicated wines was, that medicines, which were to be continued for a length of time, might be taken in the most familiar and agreeable form; by this means, a course of remedies was complied with, notwithstanding the repugnance and aversion which the sick often manifest to those directly furnished from the shops; and hence the inferior fort of people had their medicated Nevertheless, as vinous liquors excellently extract the virtues of several simples, and are not ill fitted for keeping, they have been

employed as officinal mensirua also; and fubitances of the greatest efficacy are trusted in this form. compounds of water and inflammable spirit, they take up such parts of vegetables and animals as are foluble in those liquors; though most of them abound at the fame time with a mucilaginous or viscous substance, which renders them less effectual menstroa than purer mixtures of water and spirit. They contain likewise a subtile acid, which somewhat further obstructs their action on certain vegetable and animal matters, but en-

U 2 ables ables them, in proportion to its quantity, to dissolve some bodies of the metallic kind, and thus impregnate themselves with the corroborating virtues of steel, the alterative and emetic powers of antimony, and the noxious qualities of lead.

NOTE.

To all the medicated wines, after they have been strained, you may add about one-twentieth their quantity of proof spirit, to preserve them from fermentation. They may be conveniently kept in the same kind of glass bottles that wines generally are for common uses, which should likewise be corked with the same care [L.]

# VINUM ALOETICUM ALKALINUM. Alkaline aloetic wine.

Alkaline aloetic wine.
Lond.

Take of

Any fixed alkaline falt, eight ounces;
Socotorine aloes,
Saffron,
Myrrh, each one ounce;
Sal ammoniac, purified, fix drachms;

Mountain wine, two pints.

Macerate without heat for a week
or longer; then filter the wine
through paper.

This is the Elixir Proprigations which affect the compounder rather than the composition. It is observable, that the sal ammoniac is decompounded in this process, after the same manner as in the distillation of the spiritus falis ammoniaci (see chap. viii. sect. 2) its acid being absorbed by, and neutralizing a part of the fixed alkali, and its volatile alkaline salt being set at liberty; so that the result is the same as if as much pure volatile

falt was added as the fal ammoniac is capable of affording, viz. near half an ounce, with about fix drachms of marine falt.

Helmont's elixir, in our preceding pharmacopæia, isthus directed:

Take of

Red tartar,
Nitre, each twelve ounces;
White wine, two pints;
Aloes.

Saffron, each one ounce and a half.

Let the nitre and tartar be reduced into powder, and the mixture thrown by degrees into an hot crucible: when sufficiently calcined, pour the matter into a glass mortar, and add the wine, so as to make a ley thereof; with which ley, a tincture is to be drawn from the aloes and faffron.

Take also of

Sal ammoniac, eight ounces; Spring water, twenty ounces; White wine, one pint;

Myrrh, an ounce and a half.
Dissolve the sal ammoniac in the water, strain the solution, and evaporate it to dryness. One ounce of this dry salt is to be dissolved in the wine; and with this solution, draw a tineture from the myrrh.

Mix both tinctures together, in a close vessel, so as to make them into an elixir.

THE preparation made after this troublefome method is not different from the foregoing. The nitre and tartar, when calcined together, form an alkaline falt fimilar to those which the shops are supplied with at a cheaper rate.

Helmont and others have entertained a very high opinion of this medicine, and looked upon it as "a vivifying and preferving bal-"fam, capable of continuing health

" and

" and prolonging life to the utmost " possible limits." The medicine is doubtless a very efficacious and useful one for many purposes: it may be so managed as to attenuate. viscid juices and open obstructions in the remoter parts, and promote evacuation by almost all the emunctories. In doses of one, two, or three drachms, it increases the urinary fecretion; and if the patient is kept moderately warm, generally proves diaphoretic or fudorific; in larger doles, it gently loofens the belly.

#### VINUM AMARUM.

Bitter wine. Lond.

Take of
Gentian root,
Yellow rind of lemon peel, fresh,
each one ounce;
Long pepper, two drachms;
Mountain wine two pints.
Macerate without heat, and strain
out the wine for use.

This is a very elegant bitter, which the addition of the long pepper renders confiderably warmer than the watery infusion. Gentian and lemon peel, as we have already seen, make a bitter of a very grateful flavour: "the spice here added was selected after the trial of many other materials."

## VINUM ANTIMONIALE. Antimonial wine. Lond.

Take of

Crocus of antimony, washed,

Mountain wine, a pint and a half.

Digest without heat, and filter the wine through paper.

#### VINUM EMETICUM.

Emetic wine. Edinb.

Take of

Crocus metallorum, one ounce; Mountain wine, one pint. Stir them well together; then let the mixture stand till it has perfectly settled, and carefully pour off the wine.

However carefully the fettling and decantation are performed, the filtration of the wine through paper appears to be necessary, lest some of the finer parts of the crocus should chance to remain sufpended in substance. It is not here. as in most other wines and tinctures. where the matter left undissolved by the menstruum is of little consequence: the antimonial crocus. after the action of the wine, continues as virulent as ever, and capable of impregnating fresh parcels of the liquor as strongly as the first, and this, in appearance, inexhaustibly; yet after thirty repeated infusions, it has been found scarce sensibly diminished weight.

The antimonial wine possesses the whole virtues of that mineral, and may be so dosed and managed. as to perform all that can be effected by any antimonial preparation; with this advantage, that as the active part of the antimony is here already diffolved and rendered miscible with the animal fluids, its operation is more certain. Given from ten to fifty or fixty drops, it acts generally as an alterative and diaphoretic; in larger doses, as a diuretic and cathartic; whilft three or four drachms prove for the most part violently emetic. It has been chiefly used, in this last intention, in fome maniacal and apoplectic cafes; and hence gained the name of emetic wine.

U 3

VINUM

V.NUM CHALYBEATUM.
Steel wine.

Lond.

Take of
Iron filings, four ounces;
Cinnamon,
Mace, each half an ounce;
Rhenish wine, four pints.
Macerate without heat for a month,
frequently shaking the vessel,
and strain off the wine for use.

Edinb.

Take of
Iron filings, three ounces;
Cochincal, half a drachm;
Rhenish wine, two pints.
Digest them together for twenty
days, frequently shaking the veffel; and then pass the wine
through a filter.

BOTH these wines are sufficiently elegant ones: Rhenish is an excellent menstruum for steel, and diffolves a confiderable quantity of it: the cochineal, in the second, imparts a fine colour; and the spices, ·in the first, give the liquor an agrecable flavour, make it fit easier on the stomach, and likewise promote its medicinal efficacy. In the preceding edition of the Edinburgh pharmacopoia, the digestion was ordered to be performed in a fand heat, continued for ten.days. Some have objected to the use of heat, that it impregnated the wine more strongly with the metal, and thus relidered it more unpleafant? to the talle: but if this was the only inconvenience, the remedy would be easy, diluting it with more wine. Heat has another effect, much less desirable, and which art cannot remedy; making a disagreeable alteration in the quality of the wine itself.

Steel wine is a very useful preparation of this metal, and frequentity exhibited in chlorotic and other indispositions where chalvbeates are proper. Boerhaave recommends it as one of the noblest medicines he was acquainted with, for promoting that power in the body by which blood is made, when weakened by a bare debility of the over-relaxed folids, and an indolent, cold, aqueous indisposition of the juices: for in this case, says he, no virtue of any vegetable or animal substance, no diet, nor regimen can effect that, which is effected by iron: but it proves hurtful, where the vital powers are already too strong, whether this proceeds from the fluids or the folids. The dose is from a drachm to half an ounce; which may be repeated two or three times a day.

Some direct folutions of iron, made in wine or other vegetable acids to be evaporated to the confidence of an extract, under the title of Extractum martis. These preparations have no advantage, in point of virtue, above the common chalybeates; though in some forms, that of pills in particular, they may be rather more commodicully exhibited, than most of the officinal chalybeates of equal efficacy. They may be made into pills by themselves, and are tenacious enough to reduce other sub-

stances into that form.

#### VINUM CROCEUM.

Saffron wine. Lond.

Take of
Saffron, one ounce;
Canary, one pint.
Macerate without heat, and firain
off the wine.

CANARY has been objected to by fome, as an improper menstruum for medicinal simples, since it contains a large quantity of unctuous matter, which impedes its disolving

power;

power; a pint of this fort of wine left, upon evaporation, two ounces of a mellaginous substance, not unlike honey boiled hard. It is nevertheless, for saffron, a very well adapted mentiruum, as not only fufficiently loading itself with its virtues, but likewise coinciding in the general intention of the medicine, that of a cordial. preparation made with Canary is also better fitted for keeping, than when wines that have any tendency to acidity are employed; for tinctures of faffron drawn with these last, soon lose their fine colour; whilst those made with the first, retain it for a much longer time. The dose of this tincture is from one drachm to three or more.

# VINUM IPECACOANHÆ. Wine of ipecacounha. Lond.

Take of

Ipecacoanha, two ounces; Yellow rind of Seville orange peel, dried, half an ounce; Canary, two pints.

Macerate without heat, and strain out the wine.

#### TINCTURA IPECACU-ANHÆ.

Tindure of ipecacuanha. Edinb.

Take of

lpecacuanha, in powder, one ounce;

Mountain wine, one pint.

After two days digestion, let the tincture be filtered for use.

BOTH these wines are very mild and safe emetics, and equally serviceable, in dysenteries also, with the ipecacoanha in substance; this soot yielding nearly all its virtues both to the mountain and Canary wines here ordered, as it does a good share of them even to aqueous liquors. The common dose is an ounce, more or less, according to the age and strength of the parient. The college of Edinburgh added formerly a scruple of cochineal, which imparts a fine red colour to the liquor: this article is now omitted, on a complaint, that the red colour of the matters evacuated sometimes alarmed the patient, as if it proceeded from a discharge of blood.

### VINUM VIPERINUM. Viper wine.

Lond.

Take of
Dry vipers, two ounces;
Mountain, three pints.
Macerate with a gentle heat for a
week, and then strain off the
wine.

It has been disputed, whether live or dry vipers are preferable for making this medicine: fuch as are moderately and newly dried, are perhaps the most eligible, fince by exficcation they feem to lose only their phlegmatic or aqueous parts. Whether they communicate to the wine, either when used fresh or dry, To much virtue as they are supposed to do, is greatly to be doubted. Some compositions under this name have been highly celebrated, as restoratives, in debilities and decays of conftitution : but what virtues of this kind they possessed, were supplied chiefly from other ingredients.

# VINUM MILLEPEDARUM. Wine of millepedes. Edinb.

Take of

Live millepedes, bruised, two ounces;

Rhenish wine, one pint, Insuse them together for a night, and afterwards press the liquor through a strainer.

THIS

Ters wine has been commended as an admirable cleanfer of all the viscera, yielding to nothing in the jaundice, and obstructions of the kidneys or urinary passages, of excellent service in almost all chronical distempers, even in scrophulous and firumous swellings, and in defluxions of rheum upon the eyes. But those who expected these extraordinary virtues from it, have often been deceived; and at present, there are few who have any great dependance on it. It is directed to be given from half an ounce to two ounces.

#### TINCTURA CEPHALICA. Cephalic tinclure. Edinb.

Take of

Wild valerian root, four ounces; Virginian fnakeroot, one ounce: Rosemary tops, half an ounce; French white wine, fix pints. Digest them together for three days, and then filter the tincture.

This preparation promifes to be a medicine of confiderable utility as a cephalic, that is, in diforders of the nervous Estem, wherein the membranes of the brain are often principally affected, as in / TINCTURA RHABARBARI vertiginous, epileptic, and paralytic complaints. The composition is improved from former editions, by the rejection of some ingredients, of which the best were fuperfluous; viz. cafumunar, white dittany roots, peony roots, misletoe of the oak, and peacocks dung. Casumunar is doubtless an article of importance, but much inferior, in the present intention, to the ingredients now retained.

Here it may be proper to obferve, that though some of the distilled waters, to be treated of hereafter, receive many supernu-

merary ingredients, without any confiderable injury to the produce; yet in medicines prepared by infusion, it is far otherwise. there, ingredients, which give nothing over, do little harm: but as all those commonly employed in infusions communicate some thing to the mentruum; fo, if superfluous ones are admitted, they load the liquor with an useless matter, and occupy in it the place that ought to be possessed by the more efficacious.

#### TINCTURA CEPHALICA PURGANS.

Purging cephalic tinaure. Edinb.

This is made by adding to the foregoing, of Sena, two ounces; . Black heliebore roots, ounce: French white wine, two pints.

PURGATIVES are often very necessary additions to medicines of Those here the foregoing class. made choice of are well adapted to the purpose, and in such quantity as to make the wine gently lexative in doses of two ounces.

### VINOSA.

Vinous tindure of rhubarb. Lond.

Take of Rhubarb, two ounces; Lesser cardamom seeds, freed from the husks, half an ounce; Saffron, two drachms; Mountain wine, two pints. Macerate without heat, and then Arain off the tincture.

This is a warm, cordial, laxative medicine. It is used chiefly in weakness of the stomach and bowels, and fome kinds of loofeneffes, nesses, for evacuating the offending matter, and strengthening the tone of the viscera. It may be given from half a spoonful to three or four spoonfuls or more, according to the circumstances of the disorder, and the purposes it is intended to answer.

### TINCTURA SACRA.

Take of

Socotorine aloes, eight ounces; Canella alba, two ounces; Mountain wine, ten pints.

Reduce the aloes and canella separately into powder, then mix, and pour on them the wine; afterwards macerate without heat for a week or longer, occasionally shaking the vessel; lastly, strain off the wine. It will be convenient to mix

with the powders fome white fand, well washed from dirt, to prevent the aloes from concreting, which it is apt to do upon being moistened.

Edinb.

Take of

Socotorine aloes in powder one ounce;

Ginger,

Virginian snakeroot, each one drachm:

Cochineal, one scruple;

Mountain wine, a pint and a half.

Digest for three days, and then strain off the tincture.

This medicine has long been in great esteem, not only as a cathartic, but likewise as a stimulus; the wine dissolving all that part of the aloes in which these qualities reside, a portion only of the less active resinous matter being lest. The aromatic ingredients are added, to warm the medicine, and somewhat alleviate the ill sta-

vour of the aloes: canella alba, or cloves, are faid, among numerous materials that have been made trial of, to answer this end the most successfully. The snakeroot, in the second of the above prefcriptions, feems designed for promoting the stimulating virtue of the aloes, and thus extending. its action to farther purposes than it is by itself capable of. Probably in the same intention, asarum was made an ingredient in our former pharmacopæias; in the edition preceding the present, the tincture is directed as follows: Take of

Aloes, eight ounces;
Afarum,
Cinnamon,
Zedoary,
Cardamom feeds,
Saffron, each four drachms;
Cochineal, a fcruple;
Mountain wine, ten pints.

Pour the wine on the other ingredients reduced into powder, digest them together, and afterwards strain off the tincture for use.

The tindura facra appears, from long experience, to be a medicine of excellent fervice in languid, phlegmatic habits, not only for cleanfing the primæ viæ, but likewife for attenuating and diffolving viscid juices in the remoter parts. for stimulating the solids, warming the habit, promoting or exciting the uterine purgations, and the hamorroidal flux. The dose, as a purgative, is from one to two ounces, or more: it may be introduced into the habit, so as to be productive of excellent effects, as an alterant, by giving it in fmall doses, at proper intervals; thus managed, it does not for a confiderable time operate remarkably by flool; but at length proves purgative, and occasions a lax habit of much longer continuance, than that produced by the other common cathartics.

### TINCTURA AD STOMACHICOS.

Stomachic tinaure.

Edinb.

Take of Calamus aromaticus. Gentian root, each one ounce and a half: Peruvian bark, in powder, two Ounces : Curaffoa oranges. Leffer centaury tops, Cardous benedictus feeds, each one ounce; Iron filings (to be tied up in a bag) three ounces; French white wine, one gallon. Digest for the space of three days, and then filter the tincture This tincture may likewise be made without the iron.

This tincture is a very efficacious medicine in weakness of the flomach and chylopoietic organs, and in a lax flaccid flate of the viscera in general. It is here rendered much more elegant and grateful than as it flood in former editions, by the rejection of fome exceptionable ingredients, as galangal, zedoary, chamomile, and wormwood: the carduus feeds and centaury tops might still perhaps be spared without injury. as they do not feem to have any virtues which gentian root does not possessin a greater degree. The Seville orange peel of former editions is here exchanged for the varipe young fruit of the orange tree, called Curaffao oranges, an article well adapted to compositions of this kind, being an aromatic bitter of a very agreeable flavour.

TINCTURA THEBAICA.

Thebaic tindure.

Lond.

Take of
Strained opium, two ounces;
Cinnamon,
Cloves, each one drachm;
Mountain wine, one pint.
Macerate without heat for a week,
and then filter the tincture
through paper.

THIS is the LIQUID LAUDA-NUM of SYDENHAM, with the exchange of Canary wine for mountain, and the omission of an ounce of faffron. The aromatics in the form above are in so small quantity, that the prescriber can scarce expect any confiderable effect from them, the proportion of each that goes to a grain of opium, amounting to no more than the fixteenth part of a grain: even these minute proportions, however, are in good measure sufficient to take off the ill odour of the opium, which seems to be all that is intended by them.

The principal advantages of exhibiting opium in this form are, that by being already dissolved, it exerts itself the sooner in the body; and that, by some persons, liquids are more commodiously taken, than a bolus or pill. common doses of the tincture are from ten drops to forty, fifty or more, according to the exigencies of the case. It were to be wished, that the dose could be more exactly ascertained, by weight or measure; as the drops may, according to different circumstances, vary in quantity, though in number the same; and as an error therein may, in some cases, be of mischievous consequence. Twenty drops contain, at a medium, about one grain of opium, or rather, to much as that quantity of

wine will extract from one grain; for the liquor does not dissolve . the whole substance of the opium, nor is the solution equivalent, in its effect, to the full quantity of opium employed in it.

A liquid opiate, free from the inconveniences here complained of, will be described under the head of spirituous tinctures.

VINUM AROMATICUM.

Aromatic wine.

Take of
Cloves,
Ginger, each half an ownce;
Cinnamon,
Nutmegs, each one ounce;
Canary wine, fix pints.
Beat the fpices into a coarse powder, and steep them in the wine for some days; then pass the liquor through a strainer.

This wine is a very high cordial, and greatly commended for warming the habit and strengthening the nervous system. It is so hot of the spices as to require being diluted for use, and to be taken only in small quantities at a time. Mixed with a little lemon juice, and a large proportion of milk, it forms a pleasant and useful whey in low severs.

VINUM ANTISCORBUTICUM.

Antiscorbutic avine.

Paris.

Take of

Leaves of Buckbean,
Water creffes,
Brooklime,
Dittander,
Scurvy-grafs,
Jack by the hedge,

Roots of horseradish, each one ounce;

Florence orris, two drachms; White wine, half a gallon. The herbs and roots, all fresh gathered and cut small, are to be steeped in the wine, in a vessel very closely stopt, for twenty-four hours; after which the wine is to be filtered for use.

This composition is not ill contrived for answering the purpose expressed by its title; though some of the ingredients are not onexceptionable. An ounce of the herbaceous brooklime is altogether infiguificant in half a gallon of an infusion of such powerful materials; and it may be doubted whether the fresh arris root communicates any of its virtues to the liquor. The roots of the Florentine, as well as of the common orris, raised in our gardens, are, while fresh, strong purgatives; but their purgative matter is for little disposed to solution in watery menstrua, that it separates from the expressed juices and settles to the bottom. In drying they change their nature; and the Florentine species, in a dry state, might be an useful addition, for giving an agreeable flavour to the wine. The flavour which this root communicates to vinous liquors greatly resembles that of raspberries.

VINUM SCORBUTICUM.

Scorbutic wine.

Take of

Garden scurvy grass, one handful;

Horseradilh root, scraped, half an ounce;

Winter's bark, two drachms; Mountain wine, two pints. Let them steep together in the cold for three days.

This wine is so far impregnated with the virtues of the ingredients, as to do considerable service in soci-

scorbutic habits. It is used chiefly in the spring, in the quantity of a common wine glass two or three times a day. Though far more simple than the preceding, it is not perhaps less efficacious.

VINUM SCORBUTICUM MUNTINGII.

Muntingius's scorbutic wine.

Take of

The roots of the greater waterdock, fix ounces;

Gentian root,

Liquorice, Cinnamon,

Ì

Black pepper,

Mace. each three ounces:

Saffron, two ounces;

Mountain wine, fixteen pints;

Strong vinegar, four pints; Yolks of three fresh eggs.

Reduce the roots and spices into a gross powder, and pour on them the wine, vinegar, and yolks of the eggs: digest the whole in a close vessel, with a gentle warmth, for three days; and then strain out the liquor for use.

THE author of this composition recommends it as a medicine of infallible efficacy against inveterate scurvies, and all kinds of scorbutic complaints, particularly fuch as are not accompanied with a fever or inflammation: even palfies, and the venereal lues, he fays, have yielded to it. The dose is from three to fix ounces, to be taken in the morning on an empty flomach, and continued for fourteen or twenty days, or longer: some quantity of it is likewise to be mixed with the patient's common drink, which he directs to be either good Rhenish wine, or found malt liquors not too new. If the patient complains of heat, dryness, a violent cough, or where there are any symptoms of a confumption, the black pepper is ordered to be omitted, and the liquorice increased in its room to fix ounces.

A composition differing from the above only in the omission of vinegar, and employing spirit of wine for the menstruum, is said to have come lately into esteem at Paris; against the gout.

VINUM PEBRIPUGUM.
Febrifuge wine.
Paris.

Take of

Peruvian bark, in powder, two ounces:

Rough red wine, two pints.

Digest them together in a circulatory vessel, with a moderate heat, for forty-eight hours, occasionally shaking the vessel; then suffer the whole to cool, and pass the wine through a strainer.

This is the preparation of bank made use of by fir Robert Tabor or Talbot (an English gentleman refiding in France) who was one of the first that retrieved the character of the medicine itself, at the time that some ill consequences following its imprudent use had brought it into disesteem. He kept this preparation a fecret, till Lewis XIV. purchased it for a confiderable fum, and communicated it to the public. It was not however the preparation, but a proper method of managing the medicine. upon which the fuccess of his practice depended. See page 196. It appears from experience, that this wine is less certain in the cure of agues, than the bark given in substance; nor is it equal, in this intention, for general use, to the watery infusion described in page 270; the wine preventing its being taken so freely as is in many cases requisite. It nevertheless has its uses, in those intermittent severs where a large quantity of the bark is not necessary; and is particularly serviceable in a laxity and debility of the stomach and intestines.

VINUM GUAIACINUM.
Guaiacum wine.

Take of

Guaiacum wood, Yellow faunders, each two ounces;

Orange peel, dried,

Leffer cardamom feeds, each one ounce;

Mountain wine, one gallon. Let them steep together for a week, and then strain out the wine for use.

This is a moderately warm and corroborating wine. It is used in nervous weaknesses, in decays of constitution from cold pituitous humours; and proves an useful preservative against rheumatic and arthritic complaints. Two ounces, or an ordinary wine glass, may be

taken two or three times a day, and continued for a month or two.

VINUM GUATACINUM COM HELLEBORO.

Guaiacum wine with bellebore.
Take of

Guajacum wood,

Black hellebore root, each two ounces:

Lesser cardamom seeds,

Orange peel, dried, each one ounce;

Mountain wine, four pints.

Let these ingredients steep together
for a week or longer, and thea

strain out the wine for use.

From the warm stimulating, deobstruent qualities of this wine, it may be used to good advantage in cold phlegmatic habits, where the humours stagnate in the remote vessels, and where there is a disposition to gouty, rheumatic, or hydropic disorders. It is to be taken chiefly over night, in such small doses as not to run off by shool.

#### SECT.. V.

Ales.

MEDICATED ales are intended as diet drinks in chronical indispositions. There are two ways of impregnating malt liquors with the virtues of medicinal substances; macerating the subject in the liquor after the fermentation is completely finished; and fermenting it along with the liquor, or at least adding it towards the end of the fermentation, that, by the resolutive power of that process, its texture may be opened, and its

medicinal parts more fully extracted. Neumann observes, that the active powers of many vegetables are not only effectually extracted, but extended, as it were, by fermentation: that so much pounded nutmeg, as will lie on the point of a knife, gives a flavour to a large vat of fermenting ale; whereas when the fermentation is sinished, the quantity of liquor to which it gives a like impregnation is comparatively inconsiderable.

CERE-

CEREVISIA AMARA.

Bitter ale.

Take of
Gentian root,
Lemon peel, fresh, each four
ounces;
Long pepper, one ounce;
Ale, one gallon.

Let them steep together without heat.

This is an agreeable bitter from chic ale, much superior to the common purls, or any of the compositions of this kind to be met with in the extemporaneous recipe writers.

CEREVISIA APERIENS.

Aperient ale.

Take of
Mustard seed, unbruised, ten
ounces;
Long birthwort root, six ounces;
Lest. centaury tops, two ounces;
Savin tops, one ounce;

New small ale, ten gallons.

THIS is an useful aperient dietdrink in eachestic and chlorotic indispositions, and in all cases where obstructions begin to form in the viscera. It is to be taken, to the quantity of half a pint at a time, twice a day.

CEPEVISIA BUTLERI.

Dr. Butler's ale.

Dr. Butler's ale.'

Take of
Betony,
Sage.
Agrimony,
Garden scurvy graf,
Roman wormwood, each three
handfuls;
E'ecampane roots,
Horseadish roots, each sour
onnces;
New ale, sour gallors.

The herbs and roots are to be put

in a bag, and hung in the ale while it works.

THIS liquor has fo far obtained among the common people, as to have been frequently made and fold in public houses. It is used in the spring, for purifying the blood, and preventing scorbutic disorders.

CRREVISIA CEPHALICA.
. Cephalic ale.
Take of

Wild valerian root, ten ounces; Mustard feed, whole, fix ounces; Virginian stakeroct, two ounces; Rosemary, or sage, three ounces; New small ale, ten gallons.

THE ingredients of this composition are all of the warm and stimulating kind; and consequently tend to invigorate the nervous system, and promote the circulation of the sluids. In passes, epilepsies, and vertigoes, some beness may be expected from thia liquor used as common drink,

CEREVISIA DIURETICA.
Diuretic ale.

Take of
Mustard seed, whole,
juniper berries, each eight
ounces;
Wild carrot seeds, three ounces;
Common wormwood, two ounces;
New small ale, ten gallons.

Take of 2.

Broom tops,

Muftard feed, each fixteen ounces;

Flower-de-luce roots,

Sharp-pointed dock roots, each twelve ounces;

Winter's bark,

Elder bark,

Wild carrot feeds,

Juniper

Juniper berries, each two pounds; New ale, twelve gallons.

In hydropic cases, and corpulent scorbuic habits, these aperient and diuretic liquors are very uteful diet drinks. Half a pint of either may be taken two or three times a day.

CEREVISIA AD SCORBUTICOS.

Scorbutic ale.

Take of
Horseradish root, fresh, one
pound;
Sharp-pointed dock roots, half
a pound;

Canella alba, two ounces;
Buckbean leaves, fresh, eigle ounces; or dried, three ounces;
New small ale, ten gallons.

In scorbutic disorders, and impurities of the blood and juices, this liquor, used as common drink, generally does good service. All the ingredients are very effectual for the intention, and well suited to the form. If the sharp pointed dock roots were exchanged for those of the great water dock, the composition would be still more powerful.

#### SECT. VI

#### Spirituous Tinetures.

RECTIFIED spirit of wine is the direct menstruum of the resins and essential oils of vegetables; and totally extracts these active principles from sundry vegetable matters, which yield them to water either not at all or only in part. It dissolves likewise the sweet saccharine matter of vegetables; and generally, those parts of animal bodies, in which their peculiar smells and tastes reside.

The virtues of many vegetables are extracted almost equally by water and rectified spirit; but in the watery and spirituous tinclures of them there is this difference, that the active parts, in the watery extractions, are blended with a large proportion of inert gummy matter, on which their folubility in this menstruum in great measure depends, while rectified fpirit extracts them almost pure from gum. Hence, when the spisituous tinctures are mixed with watery liquors, a part of what the spirit had taken up from the subject generally separates and subsides, on account of its having been freed from that matter which, being blended with it in the original vegetable, made it soluble in water. This however is not universal; for the active parts of some vegetables, when extracted by rectified spirit, are not precipitated by water, being almost equally dissoluble in both menstrua.

Rectified spirit may be tinged by vegetables of all colours, except blue: the leaves of plants in general, which give out but little of their natural colour to watery liquors, communicate to spirit the whole of their green tincture, which for the most part proves elegant, though not very durable.

Fixed alkaline falts deepen the colour of spirituous tinctures; and hence have been supposed to promote the dissolving power of the menstruum, though this does not appear from experience: in

the trials that have been made to determine this affair, no more was found to be taken up in the deep coloured tinctures, than in the paler ones, and often not so much; if the alkali be added after the extraction of the tincture, it will heighten the colour as much as when mixed with the ingredienta at first. Nor is the addition of these falts in making tinctures, useles only, but likewise preju-

dicial, as they, in general, injure the flavour of aromatics and superadd a quality, fometimes contrary to the intention of the medicine.—Volatile alkaline falts, in many cases, promote the action of the spirit.—Acids generally weaken it; unless when the acid has been previously combined with the vinous spirit into a compound of new qualities, called dulcified spirit.

#### General rules for extracting tinetures; from the Edinburgh Pharmacopæia.

Ŧ

The vegetable substances ought to be moderately and newly dried, unless they are expressly ordered otherwise. They should likewise be cut and bruised, before the menstruum is poured on them.

11

If the digettion is performed in balneo, the whole success depends upon a proper management of the fire: it ought to be all along gentle, unless the hard texture of the subject should require it to be augmented; in which case, the heat may be increased so as to make the menstruum boil a little, towards the end of the process.

III.

Very large circulatory vessels ought to be employed for this purpose, which should be heated before they are luted together.

Circulatory veffels are those which are so contrived, and of such a height, that the vapour, which arises during the digestion, may be cooled and condensed in the upper part, and fall down again into the liquor below: by this means the dissipation, both of the spirit and of the volatile

parts of the ingredients, is prevented. They are generally composed of two long-necked matraffes or boltheads, the mouth of one of which is to be inferted into that of the other, and the juncture fecured by a piece of wet bladder, The use of heating the vessels is, to expel a part of the air; which otherwise, rarefying in the process, would endanger burfting them, or blowing off the uppermost matrass. A fingle matrass with a long neck, or with a glass pipe inserted in its mouth, is more commodious than the double vestel. See page 44.

The vessel is to be frequently shaked during the digestion.

All tinctures should be suffered to fettle before they are committed either to the filter or strainer.

VI.

In the tinctures (and distilled spirits likewise) designed for internal use, no other spirit (drawn from malt, melasses, or other fermented matter) is to be used; than that expressly preferibed.

VII. Refin

VII.

Refins and refinous gums yield tinctures more fuccessfully, if, after being ground into powder, they be mixed with some white fand, well washed and dried, which will prevent their running into lumps by the heat. If the powders prescribed are sufficient for this purpose, such an addition is unnecessary.

## TINCTURA AMARA. Bitter tinsure. Lond.

Take of
Gentian root, two ounces;
Yellow rind of Seville orange
peel, dried, one ounce;
Leffer cardamom seeds, freed
from the husks, half an ounce;
Proof spirit, two pints.
Digest without heat, and strain off

the tincture.

This is a very elegant spirituous bitter. As the preparation is defigued for keeping, lemon peel, an excellent ingredient in the watery bitter infusions, has, on account of the perishableness of its flavour, no place in this. The cardamom feeds are here a wery commodious ingredient, as in this spirituous menstruum they are free from the inconvenience which they are attended with in other liquors, of rendering them untransparent. The Edinburgh pharmacopœia has a composition similar in intention to this, under the gitle of

### ELIXIR STOMACHICUM. Stomachic elixir.

Take of
Gentian root, two ounces;
Curassao oranges, one ounce;
Virginian snakeroot, half an
ounce;
Coghineal, half a drachm:

French brandy, two pints. Let them fleep for three days, and then filter the elixir.

This clixis differs from that of former editions, in the substitution of Curassao granges to fresh orange peel, and in the addition of half an ounce of Virginian snakeroot. The sirst is a grateful aromatic bitter, and the latter superadds a degree of pungency coinciding with the intention. Both this and the preceding composition are very useful stomachic bitters.

# TINCTURA AROMATICA. Aromatic tincure. Lond.

Take of
Cinnamon, fix drachms;
Lesser cardamom seeds, freed
from the huses, three drachms;
Long pepper,
Ginger, each two drachms;
Proof spirit, two pints.
Digest without heat, and then strain
off the tincture.

This is a very warm aromatic, too much so to be given without dilution. A tea-spoonful or two may be taken in wine, or any other convenient vehicle, in languors, weakness of the stomach, statulencies, and other like complaints. The stomachic tincture, described hereaster, is similar in intention to this, but contrived less hot of the spices, that it may be taken by itself.

The Edinburgh pharmacoposia directs a composition similar to the foregoing, under the title of

TINCTURA AROMATICA, vulgo SALUTIFERA;

which is as follows;

Take of

Calamus asomaticus root,
X Cinnamon.

Cinnamon, each half an ounce;
Leff-r cardamom feeds,
Angelica feeds, each three
drachms;
Long repper, one drachm;
French brandy, two pints.
Macerate for three days, and filter
the tincure.

This preparation is improved from the preceiting editions by the omiffion of fome articles either fuperfluous or foreign to the intention; galangal, gentian, zedoary, and bayberries. As now reformed, it is a fufficiently elegant warm aromatic.

# TINCTURA BALSAMICA. Balsamic tinæure. Edinb.

Take of

Balfam of Copaiba, one ounce and a half:

Balfam of Peru, half an ounce; English faffron, one drachm; Rectified spirit of wine, one

Digest these ingredients together,
in a sand-heat, for three days;
and then pass the tincture thro
a strainer.

This tincture is an excellent balfamic, both for internal and external purposes. It is usually given in doses of ten, twenty, or thirty drops, in the fluor albus, gleets, cachexies, some kinds of afthmas and nephritic complaints, for threngthening the tone of the viscera, and corroborating the nervous fystem in general. Some caution is requisite in the use of these refinous warm medicines: in cold, languid, phlegmatic habits, they have, for the most part, good effects; but in bilious and plethoric constitutions, where there is any tendency to inflammation, or immoderate heat, they are ma-

nifestly prejudicial, and raise or continue febrile symptoms.

TINCTURA CANTHARIDUM. TinBure of Cantharides. Lond

Take of

C: ntharides, bruised, 2 drachms; Cochineal, half a drachm; Proof spirit, a pint and a half. Digest them together, and afterwards filter the tincture through paper.

Edinb.

Take of Cantharides, two drachms; Balfam of Copaiba, one ounce and a half;

Cochineal, one drachm; Rectified spirit of wine, a pint and a half.

Digest the cantharides in the spirit for two days; then strain off the liquor, and add to it the balsam and cochineal. Digest again, in a sand heat, for sour days, and then strain off the tincture.

These tinctures possess the whole virtues of the fly, and are the only preparations of it defigned for internal use; tinctures being by far the most commodious and safe form for the exhibition of this active drug. The two tinctures are scarcely different in virtue from one another; the rectified and proof spirit equally extracting the medicinal parts of the The cochineal is cantharides. used only as a colouring ingredient: the balfam in the second prefeription, and the gum guaiacum, camphor, and effential oil of juniper berries, which were formerly added, however well adapted to the intentions of cure, can be of little consequence in a medicine limited to so small a dose.

If any additional substances should be thought requisite for promoting the effect of the cantharides, whe ther as a diuretic, as a detergent in ulcerations of the urinary pasfages, or as a specific restringent of feminal gleets and the fluor albus, they are more advantageously joined extemporaneously to the tineture, or interposed by themselves at proper intervals. The vival dose of these tindures is from ten to twenty drops, which may be taken in a glass of water, or any other more agreeable liquor, twice a day; and increased by two or three drops at a time, according to the effect.

## TINCTURA CARDAMOMI. Tindure of cardamoms. Lond.

Take of
Lesser cardamom seeds, husked,
half a pound;
Proof spirit, two pints.
Digest without heat, and strain
the tincture.

TINCTURE of cardamoms has been in use for a considerable time, though now first received into the Dispensatory. It is a pleasant, warm cordial, and may be taken, along with any proper wehicle, from a drachm to a spoonful or two.

# TINCTURA CASTOREI. Tincture of castor. Lond.

Take of
Russia castor, powdered, two
ounces;
Proof spirit, two pints.
Digest for ten days without heat,

#### Edinb.

and firain off the tincture.

Take of Ruffia castor, an ounce and a half;

Rectified (pirit of wine, one pint. Digest them with a gentle heat for three days, and asterwards strain out the liquor.

An alkaline falt was formerly added in this last prescription, which is here judiciously rejected, as being at least an useless, if not prejudicial ingredient. It has been disputed, whether a weak or rectified spirit, and cold or warm digestion, are preferable for making this tincture. To determine this point, the following experiment has been brought. "Some fine Si-" beria castor having been in-" fused in good French brandy, " without heat, for twenty days, " the tincture proved very weak: " on the same individual castor (the magma or refiduum of the " former tincture) the same quan-" tity of reclified spirit was pour-" ed, as before of brandy; and " after a few hours warm digef-" tion. a tincture was extracted " much stronger than the other." But this experiment is not fatiffactory; the effects of the two menstrua, and of heat, having been respectively compared in very different circumstances. From the trials which I have made, it appears, that castor, macerated without heat, gives out its finer and most grateful parts to either spirit. most perfectly to the rectified : that heat enables both menstrua to extract greatest part of its groffer and more naulcous matter; and that proof spirit extracts this last more readily than redified.

The tincture of castor is recommended in most kinds of nervous complaints, and hysteric disorders; in the latter, it sometimes does fervice, tho' many have complained of its proving inestectual. The dose is from twenty drops to forty, fifty, or more,

2 TINCTURA

### TINCTURA CASTOREI COMPOSITA.

Compound tindure of castor. Edinb.

Take of

Russia castor, one ounce;
Asasocida, half an ounce;
Volatile oily spirit, (commonly called spirit of sal volatile)
one pint.

Digest for fix days in a close-stopped phial, frequently shaking the vessel, and then strain the

tincture.

Thre composition is a medicine of real efficacy, particularly in hysterical disorders, and the several symptoms which accompany them. The wolatile oily spirit, here intended, is the second of those hereaster described under that title: it is an excellent mensitruum both for the castor and the assaction, and greatly adds to their virtues.

### TINCTURA CINNAMOMI. Tindure of Cinnamon. Lond.

Take of

Cinnamon, an ounce and a half; Proof spirit, a pint. Digest without heat, and strain off

the tincture.

THE tincture of cinnamon posfesses the restringent virtues of the cinnamon, as well as its aromatic cordial ones; and in this respect it differs from the distilled waters of the spice.

TINCTURA CORTICIS
PERUVIANI SIMPLEX.
Simple tindure of Peruvian bark.
Lond.

Take of

Peruvian bark, four ounces; Proof spirit, two pints.

Digest and strain.

A medicine of thiskind has been. for a long time pretty much in esteem, and usually kept in the shops, though now first received into the dispensatory Some have employed highly-reclified spirit of wine as a menstruum, which they have taken care fully to faturate, by digettion on a large quantity of the bark. Others have thought to affift the action of the spirit, by the addition of a little fixt alkaline falt, which does not, however, appear to be of any advantage; and others have given the preference to the vitriolic acid, which was supposed, by giving a greater confiftence to the spirit, to enable it to sustain more than it would be capable of doing by itself; at the same time that the acid improves the medicine, by increasing the roughness of the bark. This last tincture, and that made with rectified spirit, have their advantages; though, for general use, that above directed is the most convenient of any; .the proof spirit extracting nearly all the virtues of the bark. It may be given from a tea spoonful to half an ounce or an ounce, according to the different purposes it is intended to answer. Sec PERU-VIANUS CORTEX, p. 195.

TINCTURA CORTICIS
PERUVIANI VOLATILIS.
Volatile tinuure of Peruvian bark.
Lond.

Take of

Peruvian bark, four ounces; Spirit of falammoniac, two pints. Digestwithoutheat, in a vessel close stopped; and afterwards strain the tincture.

This tincture is but lightly impregnated with the virtues of the bark, and is so acrimonious, that the largest dose, which can with safety be given of it, can contain only a very small quantity of the subject. The medicine, neverthetheless, has its uses, and may be fer-

iceabl

viceable in fome cases where the ftronger are improper, as in difficulty of breathing, obstructions, and oppressions of the breast. Stronger sinctures of this kind may be obtained by means of dulcified spirit of fal ammoniac, or the spirit prepared with quick-lime. All the three may be employed where a large quantity of bark is not required, as at the close of the cure of intermittents, in weakness of digestion, attended with a cold fenfation at the fromach, and some fluxes, particularly those from the uterus, where the circulation is languid, the fibres relaxed, and where there is a periodical return of flight feverish complaints theie cases, I have often experienced salutary effects from a tincture in dulcified spirit of sal ammoniac, given to the quantity of a tea-spoonful, five or fix times a day, in any appropriated vehicles,

TINCTURA CORTICIS PERUVIANI.

Compound tindure of Peruvian bark. Edinb.

Take of
Peruvian bark, in powder three
ounces;
Virginian inakeroot,
Gentian, each two drachms;
Prench brandy, two pints.
Let them fleep together for three

Let them steep together for three days, and afterwards filter the tincture.

THE substances here joined to the bark, in many cases, promote its efficacy in the cure of intermittents; and, not unfrequently, are absolutely necessary. In some ill habits, particularly where the juices are sluggish and tenacious, the viscera and abdominal glands obstructed, the bark, by itself, proves unsuccessful, if not injuriaous; whilst given in conjunction with corrobarant stomachies and deobstruents, it rarely fails of the due effect. Gentian and Virginian snakeroot are among the best additions for this purpose, to which it is often necessary to join chalybeat medicines also.

TINCTURA CROCI.

Tindure of faffron.

Edinb.

Take of
English saffron, one ounce;
French brandy, one pint.
After digesting them for three days, let the tincture be strained out for use.

This tincture is similar in virtue to the saffron wine. A spirituous menstruum is here preserred to the wine; as a tincture drawn with the former retains its elegant colour longer, and is not apt to deposite in keeping any part of what it had taken up from the faf-The shops have been accustomed to employ treacle water as a menstruum for saffron, with a view to the promoting its efficacy in the intention of an alexipharmac; but the acid in that compound water foon destroys the colour of the tincture.

# TINCTURA FŒTIDA. Fætid tināure. Lond.

Take of
Asafætida, four ounces;
Rechified spirit of wine, 2 pints.
Digest and strain.

This tincture, now first received into the pharmacopoia, has been in use for a considerable time: it possesses the virtues of the asasocida itself, and may be given from 10 drops to 50 or 60. It was first X 3 proposed

proposed to the college to be made with proof spirit: this dissolves more of the asasectida than a rectissed one, but the tineture proves turbid; and therefore rectified spirit, which extracts a transparent one, is very justly preserred.

# TINCTURA FULIGINIS. Tincture of foot. Lond.

Take of
Wood foot, two ounces;
Afafætida, one ounce;
Proof spirit, two pints.
Digest and strain.

Edinb. .

Take of

Shining wood foot, one ounce; Afafœtida, half an ounce; French brandy, a pint. Digest for three days, and strain.

THE proof spirit is not liable to the same objection here as in the foregoing tincture: for when foot is added, whatever spirit be employed, the tinclure will not prove transparent. Fuller, in his Pharmacopœia Domestica, has a medicine under the title of Hys-TERIC TINCTURE, fimilar to thefe, only with a little myrrh, which is no very material addition to afafætida and foot. These medicines are found serviceable, not only in hysterie cases, but likewise in other hervous disorders. They may be given from a tea-spoonful to a common spoonful twice a day.

### TINCTURA GUAIACINA VOLATILIS.

Volatile tincture of guaiacum. Lond.

Take of

Gum guaiacum, four ounces; Volatile aromatic spirit, a pint and a half.

Digest without heat, in a vessel

close stopped; and afterwards let the tineture be passed through a strainer.

I HIS is a very elegant and efficacious tineture; the volatile spirit excellently dissolving the gum, and at the same time promoting its medicinal virtue. In rheumatic cases, a tea spoonful, taken every morning and evening in any convenient vehicle, has proved of sinegular service.

# TINCTURA JALAPII. Tincure of jalap. Lond.

Take of

Jalap root, eight ounces;

Proof spirit, two pints.

After proper digestion, strain off
the tincture.

This tincture is an useful and mild purgative; the menstruum, here employed, taking up so much of the gummy parts, as corrects the griping quality which the resso is attended with. It may be taken by itself, from a drachm to half an ounce, or mixed in smaller quantity with cathartic insusions, or the like.

# TINCTURA JALAPPÆ. Tinaure of jalap. Edinb.

Take of
Jalap, in coarfe powder, three
ounces;
French brandy, one pint.
Digest them for three days, and
then strain the tincture.

RECTIFIED spirit of wine was formerly ordered for the preparation of this tincture; but rectified spirit, dissolving little more than the pure resinous parts of the jalap, rendered the use of the medicine somewhat less commodious

than

than that of the tincture prepared with proof spirit. Most of the tincures made in reclified spirit. diluted with water fo as to be fit for taking, form a turbid white mixture: many of hem are fafely taken in this form, without any further addition: but the cathartic ones are never to be ventured on without an admixture of fyrup or mucilage to keep the refin united with the liquor; for if it feparates, in its pure undivided state. it never fails to produce violent gripes.

Some have preferred to the tinctures of jalap, a folution in spirit of wine of a known quantity of the refin extracted from the root; and observe, that this solution is more certain in strength than any tincture that can be drawn from the root directly: for as the purgative virtue of jalap refides in its refin, and as all jalap appears, from experiment, not to be equally refinous, (some forts yieldin five, and others not three ounces of refin from fixteen, it follows, that although the root be always taken in the same proportion to the menstruum, and the menstruum always exactly of the same strength, it may, nevertheless, according to the degree of goodness of the jalap, be impregnated with different quantities of refin, and confequently prove different in degree of efficacy. Though this objection against the tincture does not reach so far as some seem to sup pose, it certainly behaves the apothecary to be careful in the choice l he inferior forts of the root. may be employed for making the refina jalapii, which they yield in

as great perfection, though not in

fo large quantity, as the best. Neu-

mann thinks even the worm eaten

jalap as good, for that purpose, as

any other.

TINC YURA JALAPPÆ COMPOSI 1 A. Compound to fure of jalap. Edinb.

Take of
Jalap fix drachms;
Black heliebore roots, three
drachms;
Juniper berries,

Guaiacum shavings, each half an ounce; French brandy, a pint and a

half.
Digeft for three days, and afterwards firain the tineture.

This tincture requires to be taken in larger quantity than either of the foregoing, if intended to act fully as a cathartic. It may, in some cases, be employed to advantage, in small doses, as an alterant. The quantity of the purgative materials, that goes to an ounce of the tincture, is fifteen grains of jalap, and seven and a half of the black hellebore root.

TINCTURA JAPONICA.

Japonic tineture.

Lond.

Take of
Japan earth, three ounces;
Cinnamen, two ounces;
Proof spirit, two pints.
After proper digestion, let the tincture be passed through a strainer.

A tincture of this kind, with the addition of Peruvian bark, ambergris, and musk, to the ingredients above directed, was formerly kept in the shops. The tincture here received is preserable for general use; where any other ingredients are required, tinctures of them may be occasionally mixed with this in extemporaneous prescription. The cinnamon is a very useful addition to the Japan earth, not only as it warms the stomach, &cc.

&c. but likewise as it improves the roughness and astringency of the other.

This tincture is of good service in all kinds of defluxions, catarrhs, loofenesses, uterine sluors, and other like disorders, where mild astringent medicines are indicated. Two or three tea-spoonfuls may be taken, every now and then, in red wine, or any other proper vehicle.

# TINCTURA LACCÆ. Tincture of gum lac. Edinb.

Take of

Gum lac, powdered, an ounce; Myrrh, powdered, half an ounce; Spirit of scurvy-grass, a pint and and a half.

Digest in a sand-heat for fix days: after which, strain off the tincture for use.

· This tincture is principally employed for strengthening the gums, and in bleedings and scorbutic exulcerations of them: it may be fitted for use in these intentions, by mixing it with honey of roses, or the like. Some recommend it internally against scorbutic complaints, and as a corroborant in gleets, female weaknesses, &c. Its warmth, pungency, and manifestly astringent bitterish taste, point out its virtues, in these cases, tobe confiderable, though common practice, among us, has not yet received it.

### TINCTURA FLORUM MARTIALIUM.

Tinsture of the martial flowers.

Lond.

Take of
Martial flowers, four ounces;
Proof spirit, one pint.
Digest and strain.

## TINCTURA MARTIS. Tincture of iron. Edinb.

Take of

Iron filings, three ounces;
Dulcified spirit of salt, two
pounds;

Digest them together in a gentle heat of fand, and then filter the tincture.

### TINCTURA MARTIS in SPIRITU SALIS.

Tincture of iron in spirit of salt.

Lond.

Take of

Iron filings, half a pound; Glauber's fpirit of falt, three pounds;

Rectified spirit of wine, three

Digest the iron filings in the spirit of falt, without heat, as long as the spirit acts upon the iron: after the seces have subsided, evaporate the liquor to one pound, and add thereto the vinous spirit.

ALL the tinctures of steel are no other than real folutions of the metal made in acids, and combined with vinous spirits. The three tinctures, here directed, differ from one another only in strength, the acid being the same in all: the first is the weakest, and the last the strongest. our former pharmacopæia there was a tincture from the matter which remains after the sublimation of the martial flowers, which, though it appears to be a good one, is now expunged as superfluous. Some have recommended dulcified spirit of nitre as a menstruum; but though this readily dissolves the metal, it does not keep it suspended. The marine is the only acid that can be employed for this use..

All these tinctures are greatly preserable to the calces or croci of

iron,

iron, as being not only more speedy, but likewise more certain in their operation: the latter, in fome cases, pass off through the intestinal tube with little effect; whilst the tinctures scarce ever fail. From ten to twenty drops of either of the tinctures, may be taken, two or three times a day, in any proper vehicle; though it is feldom advisable to extend the do e fo far as the last of these quantities, especially in regard to the tincture in spirit of salt, which is exceeding strong of the iron.

TINCTURA MELAMPODII.

Tincture of melampodium, or black hellebore. Lond. and Edinb.

Take of Black hellebore roots, four ounces; Cochineal, two scruples; Proof spirit, two pints. Digest them together, and afterwards filter the tincture through paper.

This is perhaps the best preparation of hellebore, when defigned for an alterative, the menstruum here employed extracting the whole of its virtues. It has been found, from experience, particularly ferviceable in uterine obstructions; in fanguine constitutions, where chalybeates are hurtful, it seldom fails of exciting the menstrual evacuations, and removing the ill consequences of their sup-So great is the power preffion. of this medicine, that wherever, from an ill conformation of the parts, or other causes, the expected. discharge does not succeed upon the use of it, the blood, as Dr. Mead' has observed, is so forcibly propelled, as to make its way through other passages. A tea-spoonful of the tincture may be taken twice in a

day, in warm water, or any other convenient vehicle.

The college of Edinburgh had formerly a tincture of this root with wine. Proof spirit is undoubtedly preferable, both as a menstroum, and as being better fitted for keeping.

TINCTURA MYRRHÆ. Tindure of myrrh. Lond.

Take of Myrrh, three ounces; Proof spirit, two pints. After due digestion, strain off the tincture.

Edinb.

Take of Myrrh, in powder, an ounce and a half;

Rectified spirit of wine, a pint. Digest them together in a sandheat for fix days; then strain off the tincture for we.

THE pharmaceutical writers in general have been of opinion, that no good tincture can be drawn from myrrh by spirit of wine alone. without the affistance of fixed alkaline salts. But it appears, from proper experiments, that these salts only heighten the colour of the tincture, without enabling the menstruum to dissolve any more than it would by itself. fied spirit extracts, without any addition, all that part of the myrrh, in which its peculiar smell and tafte refide, viz the refin; and proof spirit dissolves almost the whole of the drug, except its imparities.

Tincture of myrrh is recommended internally, for warming the habit, attenuating viscid juices, strengthening the folids, opening obstructions, particularly those of the uterine vessels, and residing putrefaction. Boerhaave greatly esteems

esteems it in all languid cases, proceeding from fimple inactivity; in those female disorders which are occasioned by an aqueous, mucous, fluggish indisposition of the humours, and a relaxation of the vessels; in the fluor albus, and all diseases arising from a like cause. The dose is from fifteen drops to forty or more. The medicine may, doubtless, be given in these cases to advantage; though with us it is more commonly used externally, for cleaning foul ulcers, and promoting the exfoliation of carious bones.

### TINCTURA MYRRHÆ et ALOES.

Tincture of myrrh and alocs.

Edinb.

Take of

Myrrh, in powder, one ounce and a half.

Hepatic aloes, in powder, one ounce;

Rectified spirit of wine, two pints.

Digett in a fand heat for fix days, and then let the tincture be ftrained off.

This tincture is employed only in chirurgical dreffings, for cleanfing foul ulcers, reftraining the progress of gangrenes, &c. in which intention the aloes is an useful addition to the myrrh. The hepatic aloes is reckoned more effectual for these purposes than the finer socotorine.

#### TINCTURA OPII, vulgo LAUDANUM LIQUIDUM. Tincture of opium, commonly called liquid laudanum.

Edinb.

Take of

Crude opium, two ounces; English saffron, one ounce; Aqua aromatica, or spirituous cinnamon water, twenty

Digest in a sand-bath, with a gentle heat, and afterwards strain off the tincture.

It is made also without saffron.

This is a very elegant liquid opiate, the menstruum dissolving nearly the whole substance of the opium, and effectually covering its ill flavour. The proportion of menstruum is somewhat larger than in the vinous tincture formerly described (see page 298): one grain of opium goes to about twenty drops of that tincture, and twenty-five of this: nevertheless, as there appears to be more of the opium dissolved here than in the other, this tincture may posfibly be the strongest of the two. It were to be wished that the shops were furnished with a liquid opiate, in which the proportion of merstruum was still much larger, so as to admit of the dole being determined by weight or measure; the method, by drops, feeming too precarious for a medicine of fo powerful a kind. The following preparation is contrived with this view.

Take of

Thebaic extract, half a drachm; Highly-rectified spirit of wine, called alcohol, ten ounces; Simple cinnamon water, twenty ounces.

Digest them together until the opium is dissolved, and then filter the solution through paper.

This preparation I apprehend to be free from all the inconveniences attending the common opiate tinctures. The menstruum difsolves the whole of the opium, except the impurities, and confequently the tincture is not liable to any uncertainty in point of firength. The dose may be ascertained to the greatest exactness: one grain of opium is contained in one ounce by measure, which is equal nearly to seven drachms by weight. Neither the tinctures in wine or proof spirit are so well adapted for keeping as could be wished; in long standing, a part of the opium is gradually thrown off from both, and, consequently, the tinctures become gradually weaker: the part, which thus feparates, amounts fometimes, as I have been informed, to near onefourth of the quantity of opium at first dissolved: it floats on the surface of the vinous tincture, and in the spirituous finks to the bottom. In the preparation here recommended, it has not been observed that any feparation happens.

Instead of the cinnamon water, pure water may be employed in the mixture; and where aromatic additions are wanted, either in a medicinal intention, or for covering the ill smell of the opium, any proper tincture or distilled water may be extemporaneously joined. Saffron, an addition employed by the Edinburgh college, has been looked upon as a corrector of opium; but the qualities it was supposed to correct are merely imaginary: nor indeed can that article be of much importance in any intention, in the small quantity that enters a dose of the tincture; a grain of opium being accompanied with only half a grain

TINCTURA RHABARBARI

of faffron.

SPIRITUOSA.

Spirituous tincture of rhubarb.

Lond.

Take of Rhubarb, two ounces; Lesser cardamom seeds, husked,

half an ounce;
Saffron, two drachms;
Proof spirit, two pints.
Digest without heat, and strain off
the tincture for pse.

TINCTURA RHE! AMARA.

Bitter tincture of rhubarb.

Edinb.

Take of
Rhubarb, one ounce;
Gentian root, a drachm and a
half;
Virginian finakeroot, 1 drachm;
Cochineal, one fcruple;
French brandy, one pint.
Digest for two days, and then
strain the tincture.
This tincture may likewise be made
with mountain wine.

TINCTURA RHEI DULCIS.

Sweet tindure of rhubarb.

Edinb.

Take of
Rhubarb, two ounces;
Leffer cardamoms, half an ounce;
French brandy, two pints.
Digest for two days; and then,
having strained out the tincture,
add to it four ounces of white
sugar-candy in powder, and digest again until the sugar is dissolved.

THE last of these preparations is somewhat improved from the former edition. Two ounces of liquorice and one of raisins are supplied by an increase of the sugar; and two drachms of canella alba, by increasing the cardamom seeds from two to four drachms.

All the foregoing tinctures of rhubarb are defigned as stomachics and corroborants, as well as purgatives. Spirituous liquors excellently extract those parts of the rhubarb in which the two first qualities reside, and the additional ingredients considerably promote their

their efficacy. In weakness of the flomach, indigetion, laxity of the intestines, diarrhocas, colicky and other like complaints, these medicines are frequently of good fervice: the second is also, in many cases, an useful addition to the Poruvian back, in the cure of intermittents, particularly in cachectic habits, where the viscera are obstructed. In these intentions, a spoonful or two may be taken for a dose, and occasionally repeated.

#### TINCTURA SATURNINA. Saturnine tindure.

Lond.

Take of

out heat.

Sugar of lead. Green vitriol, each two ounces; Rectified spirit of wine, two pints.

Reduce the falts feparately into a powder; then add the spirit, and digest them together without heat; afterwards filter the tincture through paper.

> TINCTURA ANTIPHTHISICA. Antiphtbifical tindure. Edinb.

Take of Sugar of lead, an ounce and a Vitriol of iron, an ounce; Rectified spirit of wine, a pint. Let a tincture be extracted with-

THE reducing of the falts feparately into powder, and performing the digestion without beat, are very necessary circumstances; for if the ingredients ate attempted to be pulverized together, they will grow foft and almost liquid; and if heat is made use of, scarce any tincture will be obtained.

These tinctures are sometimes given, from twenty to thirty drops, for reftraining immoderate fecretions, particularly the colliquative sweats attending hectic fevers and phthisical disorders; whence the name antiphthifical tincture. They are undoubtedly medicines of great efficacy in these cases, but too dangerous ones to be rashly ventured on. Some have supposed, that they do not contain any of the fugar of lead; but experiments, made for that purpose, have shewn that they do: and therefore the London college has very judiciously changed the title of their tincture into one expressing its being a preparation of lead.

#### TINCTURA SENÆ. Tineture of Jena. Lond.

Take of Raifins, floned, 16 ounces; Sena, one pound: Caraway feeds, one ounce and a half; Lesser cardamoms, husked, half an ounce; Proof spirit, one gallon. Digest without heat, and then Arain the tincture.

#### TINCTURA SENÆ COMPOSITA, vulgo ELIXIR SALUTIŠ.

Compound tindure of sena, commonly called elixir of bealth. Edinb.

Take of

Sena, two ounces; Rhubarb, one ounce: Sweet fennel feeds, Juniper berries, Guaiacum shavings, each half an ounce;

French brandy, three pints. Digett for the space of three days; then strain off the tincture, and add to it three ounces of powdered fugar-candy.

Both

Born these tinaures are useful carminatives and cathartics, especially to those who have accustomed themselves to the use of spiritous liquors; they oftentimes relieve flatulent and colicky complaints, where the common cordials have little effect: the dose is from one to two ounces. Several preparations of this kind have been offered to the public, under the name of Daffy's elixir: the two above are equal to any, and fuperior to most of them. guaiscum, in the last of the above formulæ, is a very useful ingredient, as it is found to have very good effects, when joined with purgatives: 2 drachms of fena, infufed in half a pint of decoction of guaiacum, work as brifkly as 3 drachms infused in plain water, and with greater ease to the patient.

#### TINCTURA SERPENTARIÆ. Tindure of Inakeroot. Lond.

Take of

Virginian inakeroot, three ounces: Proof spirit, two pints. Digest without heat, and strain off the tincture.

THE tincture of inakeroot was, in our last pharmacopæia, directed with the tindura salis tartari, which being now expunged, it was proposed to the college to employ rectified spirit; but as the heat of this spirit prevents the medicine from being taken in so large a dose as it might otherwise be, a weaker spirit was made choice of. The tincture made in this men-Aruum, which extracts the whole virtues of the root, may be taken, to the quantity of a spoonful or more, every five or fix hours.

The college of Edinburgh directs this tincture to be drawn with plague water, a distilled spirituous water, impregnated with masterwort, angelica, and elder flowers, and mixed with distilled vine-

Take of

Virginian snakeroot, 2 ounces: Cochineal, one drachm; Plague water, two pints. Digest in a gentle heat for three days, and then frain the tinc-

THE plague water is equally efficacious, as a menttruum, with the simple proof spirit or rectified spirit, and likewife coincides with the general intention of the medicine, as an alexipharmac in tevers.

#### TINCTURA STOMACHICA. Stomachic tinaure. Lond.

Take of

Raifins, floned, four ounces; Cinnamon, half an ounce: Caraway seeds, Lesser cardamoms, husked, Cochineal, each two drachms: Proof spirit, two pints. Digest without heat, and strain off the tincture.

This is a moderately warm stomachic tincture, much more pleafant than the usqueBaugh of our former pharmacopœias. may be taken, without any vehicle, to half an ounce or an ounce, tho' oftener used in mixtures.

#### TINCTURA STYPTICA. Styptic tinaure. Lond.

Take of

Green vitriol, calcined, one drachm;

French brandy (fuch as has acquired a yellow tinge from . the cask) two pints.

Mix

Mix them together, that the spirit may grow black; then pass it through a strainer.

Some have supposed, that no other spirit than French brandy would succeed in striking the black colour, for which this tincture is valued. But any spirit, that has gained an impregnation from the oak casks, which these kinds of liquors are generally kept in, or from other vegetable aftringents, will equally exhibit this phænomenon; and French brandy will not do it without such assistance. The title of this tincture expresses its medicinal intention. lebrated STYPTIC OF HELVETIUS (which is faid to be the same with that of EATON) differs from it no otherwise, than in being more operose in composition. They are recommended both for internal use, and for restraining external hæmorrhages: their virtues do not feem to depend so much on the iron, as on the menstruum, the quantity of metal dissolved being extremely small. In keeping, the iron is apt to separate, and the liquor to lose its black colour.

### TINCTURA SUCCINI. Tincture of amber.

Take of

Yellow amber, two ounces; Reclified spirit of wine, twenty ounces.

Digest in a sand heat for eight days, and asterwards silter the tincture.

This is a very elegant preparation of amber, of a grateful balfamic tafte, and fragrant smell. Boerhaave, Hoffman, and others, strongly recommend it in disorders proceeding from a lax state of the solids and debility of the nervous system; in suppressions of the menstrual discharges, the stuor albus, seminal gleets, rheumatic complaints, and some kinds of epilepsies. It is directed to be taken from 10 to 100 drops, in Canary or other rich wine.

The medicine is, doubtless, an efficacious one; though it would be much more so, if a part of the spirit was drawn off, so as to leave what it had extracted from the amber, concentrated into the confistence of a balfam: a tea-spoonful of this may be taken three or four times a day, with fugar, or in any convenient vehicle. The spirit distilled off, which is richly impregnated with the amber smell, may be referred for extracting a fresh tincture from another parcel of amber. A tineture of amber, made in this spirit, possesses the whole virtue of the concrete, and appears to be one of the most valuable preparations of it.

Fixed alkaline falts have been commonly employed in the preparation of this tincture; but with no good effect; for they not only do not promote the dissolution of the amber, but likewise injure the medical virtue of the preparation. Scarcely any of the substances that have been made trial of, give any considerable assistance to spirit of wine in dissolving this concrete, except the vitriolic acid, which, when intimately combined with it into a dulcified spirit, forms a menstruum, said to be much more efficacious for amber than the fimple vinous spirit. The college of Edinburgh have, accordingly, in the late reformation of their pharmacopæia, made choice of this menstruum, and directed the tincture as follows:

fare as rottoms

Take of

Yellow amber, two ounces;
Dulcified spirit of vitriol, one
pint.
Digest

Digest them in a fand-bath, with a gentle heat for four days, and then filter the tincture.

TINCTURA SUDORIFICA.

Sudorific Tindure.

Edinb.

Take of

Virginian snakeroot, 6 drachms; Cochineal,

English saffron, each 2 drachms;

Opium, one scruple; Spirit of Mindererus, one pint.

Digest them together in a gentle heat for three days, and then pass the tincture through a strainer.

This composition is an efficacious sudorisic; the ingredients be ing of the mo" powerful kind, and the mentiruum not only extracting those parts of them in which their virtues confift, but cooperating strongly in the same intention. Russia castor, a supernumerary ingredient in former editions, is now omitted: and cochi neal, which, from the quantity of it formerly employed, seemed to have been defigned with a medicinal view, is now reduced to one half, and nothing more is expected from it than to furnish an agreeable colour to the tincture. an ounce of the tincture, by meafure, contains five-eighths of a grain of opium

TINCTURA SULPHURIS.

Tindure of Sulphur.

Take of

Rectified spirit of wine, one pint.

Hepar fulphuris (that is, a mixture of fulphur and fixed alkaline falt melted together) four ounces.

Grind the hepar into powder whilft hot from the fire; add to it the fpirit, and digest in a moderate heat for twenty-four hours; then pour off the tincture from the feces.

THE digestion may be commodiously performed in a glass receiver: put the spirit first into the vessel, and pour the hot powder upon it: then shake them together, and, to prevent the exhalation of any part of the spirit during the digestion, insert a glass tube into the mouth of the re-

ceiver.

This tincture is of a rich gold colour, a hot aromatic taste, and a particular. not ungrateful smell. Its virtues are those of a warm attenuating, aperient, and anti-acid medicine. Some have recommended it as a last resource in phthisics and ulcerations of the lungs; but in these cases it promises little service, and has been sometimes found prejudicial. The dose is from 10 to 60 drops: it is most commodiously taken in Canary or other rich wines.

## TINCTURA ANTIMONII. Tindure of antimony. Lond.

Take of

Any fixed alkaline falt, 1 pound; Antimony, half a pound; Rectified spirit of wine, two

pints.

Reduce the antimony into powder, mix it with the falt, and melt them together with a strong fire, for an hour. Then pour out the matter, pulverize it, add the spirit, and digest them for three or four days: after which strain off the tincture for use.

Edinb.

Take of

Antimony, in powder, four ounces;
Salt of tartar, fix ounces;
Rectified spirit of wine, 2 pints.

Mix

Mix the antimony with the falt of tartar, and inject them by little and little into a crucible placed in a strong fire. The mixture melts thin, and is to be continued in this state for half an hour: after which it is to be poured out into a hot and dry Powder the mais iron mortar. while hot, put it into a heated matrais, and pour thereon the Digest them together. spirit. for three days, in a gentle heat of fand; and then decant the tinclurc.

In these processes the alkaline salt unites with the sulphur of the antimony into a hepar, which communicates to the spirit a tracture similar to the foregoing. This antimonial tincture is supposed to contain likewise some of the reguline parts of the mineral, and is said to have sometimes provoked a puke when taken on an empty stomach, even in a small dose. It stands recommended, in doses of from 10 to 60 drops or more, as a deobstruent, promoter of urine, and purisher of the blood.

TINCTURA ANTIMONII DIA-PHORETICA.

Tindure of diaphoretic antimony. Take of

Diaphoretic antimony, fixteen ounces;

Nitre, four pounds;

Tartarized spirit of wine, three pints.

Let the antimony and nitre be finely powdered, mixed, injected by a fpoonful at a time into a red-hot crucible, and kept in a strong melting heat for half an hour. Then pour the matter into a warm iron mortar, powder it whilst hot, and immediately add the vinous spirit. Digest for

three days, and filter the tincure for use.

This tincture is recommended for the same purposes as the foregoing, and in the same dose. It is very fragrant in smell, and agreeable to the take.

TINCTURA SALIS TARTARI.

Tinclure of falt of tartar.

Take of

Salt of tartar, fix ounces.

Melt it in a crucible till it acquires a greenish colour; pulverize it whilst hot, and immediately pour upon it, in a strong long necked matrass, as much rectified spirit of wine as will stand three or four inches above it: digest, for several days, in a pretty strong sand-heat, that a tincture may be obtained.

This preparation is taken from the preceding edition of our pharmacopœia. It has been usually expected to be of a red hue; but. (as the committee observe) if neither the falt nor the spirit have any oily tincture, the spirit, though it acquires from the alkali a hot pungent tafte, will scarce receive any degree of colour, unless by some spark of coal, which may accidentally fall into the crucible, while the falt is calcining. For this rea-Son, this tincture has been usually prepared in a counterfeit manner, by adding some portion of antimony to the falt, whereby it refembled too much the tincture of antimony, for both to be retained at the late revisal

TINCTURA TOLUTANA.

Tindure of balfam of Tolu.

Edinb.

Take of
Balfam of Tolu, an ounce and a
half;

Reclified

tincture.

Rectified fpirit of wine, a pint. Digest in a fand-heat, until the balfam is dissolved: and then strain the tincture.

This folution of balfam of Tolu possesses all the virtues of the balfam itself. It may be taken internally, in the several intentions for which this valuable balfam is proper, to the quantity of a tea-spoonful or two, in any convenient vehicle. Mixed with the plain syrup of sugar; it forms an elegant balfamic syrup.

### TINCTURA VALERIANÆ SIMPLEX.

Simple tindure of valerian.

Lond.

Take of
Wild valerian root, four ounces;
Proof spirit, two pints.
After due digestion, strain off the

The valerian root ought to be reduced into fine powder, otherwise the spirit will not sufficiently extract its virtues. The tincture proves of a deep colour, and confiderably strong of the valerian; though it has not been found to answer so well in the cure of epileptic disforders, as the root in substance exhibited in the form of powder or bolus. The dose of the tincture is, from half a spoonful to a spooful or more two or three times a day.

#### · TINCTURA VALERIANÆ VOLATILIS.

Volatile tindure of valerian. Lond.

Take of

Wild valerian root, four ounces;

Volatile aromatic spirit, two pints.

Digest without heat, in a vessel closely stopt, and asterwards strain off the tincture.

The volatile spirit is here an excellent menstruum, and at the same time considerably promotes the virtues of the valerian, which in some cases wants an assistance of this kind. The dose may be a tea-spoonful or two.

#### TINCTURA VERATRI.

Tincture of veratrum, or white hellebore. Lond.

Take of
White bellebore root, eight
ounces;
Proof spirit, two pints.

Digest them together, and filter the tincture through paper.

This tincture is fometimes used for acuating cathartics, &c. and as an emetic in apoplectic and maniacal disorders. It may likewise be so managed, as to prove a powerful alterative and deobstruent, in cases where milder remedies have little effect. But a great deal of caution is requisite in its use: the dose, at first, ought to be only a sew drops; if considerable, it proves violently emetic or cathartic.

## BALSAMUM GUAIACINUM. Balfam of guaiacum. Lond.

Take of
Gum guaiacum, one pound;
Balfam of Peru, three drachms;
Recufied spirit of wine, two pints
and a halt.

Digest till the gum is dissolved, and then strain off the balfam.

## ELIXIR GUAIACINUM. Elixir of guaiacum. Edinb.

Take of.
Gum guaiacum, in powder, two ounces;
Balfam of Peru, two drachms;
Y

>

Essential oil of Sassafras, one drachm;

Volatile oily spirit, one pint.

Digest the gum guaiacum and balsam in the volatile spirit for six
days in a close stopt phial; which
is now and then to be shaken;
afterwards strain the tincure and

add to it the effential oil.

BOT H these compositions are medicines of great efficacy, and capable of answering many useful purpoles. They warm and strengthen the habit, and promote insensible perspiration. Twenty or thirty drops may be taken two or three times a day, or oftener, in any proper vehicle, in rheumatic complaints, cutaneous defedations &c. particularly where the patient is of a cold phlegmatic temperament. and the folids weak and relaxed. In hot bilious constitutions, and tensity or rigidity of the vessels, like other stimulating medicines, they are evidently improper.

### BALSAMUM COMMENDATORIS. Boume de commandeur.

Take of

Dry Peruvian balfam, one ounce; Storax in the tear, two ounces; Benjamin three ounces; Socotorine aloes, Myrrh, Olibanum, Angelica Roots, St. John's worth fibwers, each half an ounce; Spirit of wine, two pounds eight

ounces by weight.

Let them fland together in the fun during the dog days, in a glafs veffel, closely stopt; and atterwards strain out the balfam thro' a linen cloth.

THIS balfam has been inferted, with little variation, in some soreign pharmacopæias, and likewise

kept a secret in private hands, under the titles of Balfamum Perficum, Balsam of Berne, Wade's balsam, Friars balsam, Jesuits drops, The form above is taken from the original receipt published. by Pomet (Histoire des drogues, edit. 2. tom. ii. p. 56) It stands greatly recommended, externally for cleansing and healing wounds and ulcers. for discussing cold tumors, allaying gouty, rheumatic, and other old pains and aches; and likewife internally for warming, and strengthening the stomach and intestines, expelling flatulencies, and relieving colicky complaints. Outwardly it is applied cold on the part with a feather; inwardly a few drops are taken at a time, in wine or any other convenient vehicle.

#### BALSAMUM. TRAUMA FICUM. Traumatic or vulnerary balfam. Lond.

Take of
Benzoine, three ounces,
Storax, firained, two ounces,
Ballam of Tolu, one ounce,
Secotorine aloes, half an ounce,
Rectified spirit of wine, twopints.

Digeft, that the gums may as much as possible be diffolved; and then strain off the balfam for use.

This is an elegant reformation of the preceding composition, confiderably more simple, yet not inferior in esseasy. The balsam of Tolu supplies, with advantage, the dry Peruvian balsam, a drug very rare to be met with in this country: the olibanum, myrrh, and angelica roots here omitted, were certainly superfluous in a medicine containing so much more powerful materials; and the St. John's wort slowers are as deservedly thrown out, as

bav-

baving little else to recommend them than prejudice or superstition.

Edinb.

Take of

Benzoine, powdered, three ounces;

Balsam of Peru, two ounces; Hepatic aloes, in powder, half an ounce;

Rectified Spirit of wine, two pints.

Digest them in a sand heat, for the space of three days; and then strain the balsam.

THIS is a further contraction of the baume de commandeur, without any injury to it as a medicine, at least with regard to the purposes for which the title shews it designed. Socotorine aloes is here judiciously exchanged for the hepatic, which appears from experience to be the most serviceable in external applications.

## ELIXIR ALOES. Elixir of aloes. Lond.

Take of
Tincture of myrrh two pints;
Socotorine aloes,
Saffron each, three ounces,
Digest them together, and strain
off the elixir.

### ELIXIR PROPRIETATIS. Edinb.

Take of.

Myrrh in powder; two ounces; Socotorine aloes, an ounce and an half;

English Saffron one ounce; Rectified spirit of wine, two pints.

Digest the myrrh with the spirit, in a sand bath, for the space of four days: then add the aloes in powder, and the Saffron; continue the digestion for two days

longer, suffer the seces to subside, and pour off the clear elixir.

This is the elixir proprietatis of Paracelfus, improved with regard to the manner of preparation. myrrh faffron and aloes, have been usually directed to be digested in the spirit together; by this method the menitruum foon loads itself with the latter: so as scarce to take up any of the myrrh; whilst a tincture, extracted first from the myrrh, readily dissolves a large quantity of the others. The alka-·line falt, commonly ordered in these preparations, with a view to promote the diffolution of the myrrh, we have already obscrved to be useless; and accordingly it is now omitted.

This medicine is greatly recommended, and not undefervedly, as a warm stimulant and aperient. It strengthens the stomach and other viscera, cleanses the first passages from tenacious phlegm, and promotes the natural fecretions in ge-Its continued use has frequently done good fervice in cachectic and icteric cases, uterine obstructions, and other like disorders; particularly in cold, pale phlegmatic habits; where the patient is of a hot, bilious constitution and florid complexion, this warm stimulating medicine is less proper, and fometimes prejudicial. The dose may be, from twenty drops to a tea-spoonful or more, two or three times a day, accordto the purpoles which it is intended to answer.

#### ELIXIR PROPRIETATIS. VITRIOLICUM. Edibb.

Take of
Myrrh, in powder, two ounces;
Socotorine aloes, in powder one

English saffron, half an ounce; Y2 Dulcised Dulcified spirit of vitriol, one pint and a half.

Digest them in a sand-heat for the space of six days; and having then suffered the seces to subside, pour off the clear elixir.

HERE the dulcified spirit of vitriol is very judiciously substituted to the fpirit of fulphur, ordered in other books of pharmacy to be added to the foregoing preparation; for that strong acid precipitates from the liquor great part of what it had before taken up from the other ingredients; whereas, when the acid is previously combined with the vinous spirit, and thereby dulcified, as it is called, it does not impede its diffolving power. This elixir possesses the general virtues of the preceding, and is, in virtue of the menstruum, preferred to it in hot constitutions, and weaknesses of the stomach. See Elixir vitrioli in the following page.

### ELIXIR PAREGORICUM. Paregorick elixir.

Lond.

Take of
Flowers of benzoine,
Opium strained, each one drachm;
Camphor, two scruples;
Essential oil of aniseeds, half a drachm;
Rectified spirit of wine, two pints.
Digett and strain.

This elixir is taken from Le Mort with the omission of three unnecessary ingredients, honey, liquorice, and alkaline salt. It was originally prescribed under the title of ELIXIR ASTHMATICUM, which it does not ill deserve. It contributes to allay the tickling, which provokes frequent coughing; and at the same time, is supposed to

open the breast, and give greater liberty of breathing: the opium procures (as it does by itself) a temporary relief from the symptoms; whilst the other ingredients tend to remove the cause, and prevent their return. It is given to children, against the chin-cough, &c. from five drops to twenty; to adults, from twenty to an hundred. Half an ounce by measure contains about a grain of opium.

## ELIXIR PECTORALE. Pedoral elixir. Edinb.

Take of
Balfam of Tolu, two ounces;
Balfam of Peru, one ounce;
Flowers of benzoine,
English saffron, each half an ounce;
Rectified spirit of wine, two pints.
Digest them in a sand-heat for three days, and then strain off the

elixir.

This ballamic elixir is given to the quantity of a tea-spoonful, two or three times a-day, as an expectorant and detergent, in coughs and ulcerations of the breast. The balsam of Peru is a new ingredient, introduced in the present edition; and the slowers of benzoine are substituted to benzoine in substance.

#### ELIXIR VITRIOLI ACIDUM-Acid elixir of witriol. Lond.

Take of the
Aromatic tincture, one pint;
Strong spirit (called oil) of vitriol, four ounces.
Mix them together; and after the feces have subsided, filter the elixir through paper.

THIS

l

1

This preparation was originally taken from Mynsicht, and has been usually distinguished by his name. It is here prepared in a somewhat different manner from that directed by the author, and in other books of pharmacy; the oil of vitriol and spirit of wine being there first mixed together, and then digested upon aromatics.

Mynficht's elixir of vitriol is directed in our preceding pharmacopeia as follows:

Take of

Cinnamon,

Ginger.

Cloves, each three drachms; Calamus aromaticus, one ounce; Galangal, an ounce and a half; Sage,

Mint, each half an ounce; Cubebs,

Nutmegs, each two drams; Aloes wood,

Citron peel, each one dram. Reduce these ingredients into a

powder, to which add, of
Sugar-candy, three ounces;
Spirit of wine, a pint and a
half;

Oil of vitriol, one pint.

Digest them together for ewenty days, and then filter the tincture for use.

In the new edition of the Edinburgh pharmacopæia, the elixir vitrioli is thus prepared.

Take of

Cinnamon, one ounce and a half; Ginger, one ounce; Pepper-mint leaves, dried, half an ounce;

Oil of vitriol, six ounces; Rectified spirit of wine, two

pints.

Drop the oil of vitriol by little and little into the spirit of wine, and digest them together in a sandbath, with a very gentle heat, for three days; then add the

other ingredients; continue the digestion, in the same gentle heat, for three days longer; and afterwards filter the tincture in a glass funnel.

THE intention in these processes is, to obtain a tincture of aromatic vegetables, in spirit of wine, combined with a confiderable proportion of vitriolic acid. When the tincture is first drawn with vinous spirits, and the acid added afterwards, as in the first of the above prescriptions, the acid precipitates great part of what the spirit had before taken up: and on the other hand, when the acid is mixed with the spirit immediately before the extraction, as in the fecond process, it prevents the diffolution of all that it would have precipitated by the former way of treatment: by previously uniting the acid and the vinous spirit together by digestion, as in the last process, the inconvenience is somewhat lessened.

All these compositions are valuable medicines in weakness and relaxations of the stomach, and decays of constitution, particularly in those which proceed from irregularities, which are accompanied with flow febrile symptoms, or which follow the suppression of intermit-They have frequently taken tents. place after bitters and aromatics, by themselves, had availed nothing; and, indeed, great part of their virtue depends on the vitriolic acid; which, barely diluted with water, has, in these cases, where the stomach could bear the acidity, produced happy effects.

Fuller relates (in his Medicina gymnastica) that he was recovered, by Mynsicht's elixir, from an extreme decay of constitution, and continual reachings to vomit. They may all be given from ten to thirty or forty drops or more, according

Mark Street

Y 3

to the quantity of acid, twice or thrice a day, at such times as the stomach is most empty.

## ELIXIR VITRIOLI DULCE. Sweet elixir of vitriol. Lond.

Take of the

Aromatic tincture, one pint;
Dulcified spirit of vitriol, eight
ounces by weight.

Mix them together.

THIS is defigned for persons whose stomach is too weak to bear the foregoing acid elixir; to the taste, it is gratefully aromatic, without any perceptible acidity. The dulcified spirit of vitriol, here directed, occasions little or no precipitation upon adding it to the sincture.

THE college of Edinburgh, in the edition of their pharmacopæia preceding the prefent, employed dulcified spirit of vitriol as the menstruum. The composition was as follows; Take of

Dulcified spirit of vitriol, two pounds:

Effential oil of mint, half an ounce;

of lemon peel,

of nutmegs, each two drams.
Gradually drop the oils into the fpirit, and mix the whole well together.

This elixir; if the effential oils are good, and the dulcified spirit made as it ought to be, (if it is not, it will not dissolve the oils) proves a very elegant and grateful stomachic, similar to the foregoing sweet elixir: a tea spoonful of either may be taken two or three times a day

A medicine of this kind was formerly in great effeem under the title of Vigani's volatile elixir OF VITRIOL; the composition of which was first communicated to the public in the Pharmacopæia reformata. It is prepared by digesting some volatile spirit of vitriol upon a fmall quantity of mint leaves curiously dried, till the liquor has acquired a fine green colour. the spirit, as it frequently does, partakes too much of the acid, this colour will not succeed: in such case it should be rectified from a little fixt alkaline salt, as hereaster directed in chap. viii. fect. 5. The mint is most commodiously suspended in the spirit in a fine linen cloth: this prevents the necessity of filtration, during which the more volatile parts would exhale.

# ELIXIR MYRRHÆ. COMPOSITUM. Compound elixir of myrrb. Lond.

Take of

Extract of savin, one ounce,
Tincture of castor, one pint;
Tincture of myrrh, half a pint.
Digest them together, and then
strain the elixir.

This preparation is improved from one described in some former dispensatories under the name of ELIXIR UTERINUM. It is a medicine of great importance in uterine obstructions, and in hypocondrical cases; though, possibly, means might be contrived of superadding more effectually the virtues of savin to a tinclure of myrrh and castor. It may be given from sive drops to twenty or thirty, or more, in pennyroyal water, or any other suitable vehicle.

### ELIXIR SACRUM. Edinb.

Take of Rhubarb, cut finall, ten drams; SocotoSocotorine aloes, in powder, fix drams;

Leffer Cardamon feeds, half an ounce;

French brandy, two pints.

Digeft for two days, and then strain
the elixir.

SPIRITUS VINOSUS.
CAMPHORATUS.
Campberated spirit of wine.
Lond. and Edinb.

Take of

Camphor, two ounces;
Rectified spirit of wine, two pints
Mix them together, that the camphor may be dissolved.

This folution of camphor is employed chiefly for external uses, against rheumatic pains, paralytic numbnesses, inflammations, for discussing tumors, preventing gangrenes, or restraining their progress. It is too pungent to be exhibited internally, even when diluted, nor does the dilution succeed well; for on the admixture of aqueous liquors, the camphor gradually separates and runs together into little masses.

Hoffman, Rothen, and others, mention a camphorated spirit not subject to this inconvenience. It is prepared by grinding the camphor with somewhat more than an equal weight of fixt alkaline falt, then adding a proper quantity of proof spirit, and drawing off one half of it by distillation. This spirit was proposed to the college to be received into the pharmacopæia, at the late revifal, under the title of SPIRITUS CAMPHOR ETARTA-REZATUS. But upon trial, it did not answer expediation; some of the camphor, as the committee observe, rises with the spirit in distillation, though but a finall quantity; whence, mixed with a large portion of water, it does not sensibly render it turbid; but in a proper quantity, it exhibits the same appearance as the more common camphorated spirit; it did not appear, that spirit distilled from camphor, with or without the alkaline salt, differed at all in this respect.

The most convenient method of uniting camphor with aqueous liquors, for internal use, seems to be by the mediation of almonds, or of mucilages; triturated with these, it readily mingles with water into the form of an emulsion, at the same time that its pungency is considerably abated. It may also be commodiously exhibited in the form of an oily draught, expressed oils totally dissolving it.

### Tenctura Benzoini. Tindure of benzoine.

Take of

Benzoine, four ounces: Rectified spirit of wine, one pint.

Digest them together in a sand-heat for three or four days, and then decant off the tincture.

THIS tincure hands recommended in asthmas, and other disorders of the lungs, in doles of from twenty to fixty or feventy drops. It has, however, been principally made use of externally, as a cosmetic, for clearing and imoothing the ikin : for these purposes it is mixed with a large proportion of water, when it forms a white liquor called LAC VIRGINIS.. If this be fuffered to rest for some time, the benzoine precipitates, in form of a white magistery (of a very pleafant smell, and not disagreeable taste), which in the Brandenburgh pharmacopœia is preferred to the flowers of benzoine, as being free from the empyreumatic flavour which thefe are generally attended with: it is, however, of a different Y 4

ent nature from the flowers, being no other than the benzoine in its whole substance; whereas the flowers are a distinct part of it; not resinous, like the rest of the mass, but rather, as we shall see hereaster, of the saline kind. The precipitation is directed to be made with rose water.

GUTTE VITE.

Drops of Life.

Take of
Opium, four ounces;
Saffron, one ounce;
Virginian inakeroot,
Cochineal, each half an ounce;
Nutmegs,
Zedoary, each two ounces;
Camphor, one ounce;
Tincture of diaphoretic antimony; one pint;
Western two ounces

Water, two pints.
Digest the opium with the water in a scalding heat, till as much as possible of it is dissolved, and pass the solution through a strainer. Digest the other ingredients in the antimonial tincture for three or four days. Mix both liquors together, let them stand in digestion for two days longer, and after the seces have subsided, pour off the clear for use.

This medicione has been recommended as preferable to the common opiates, and less apt to leave a nausea on the stomach: the dose is from ten drops to forty or sifty.

TINCTUREA feu Essentia au-BRB.

Tincture or effence of ambergris.

Parif.

Take of
Ambergris, one drachm;
Tartarized spirit of wine,
Spirit of roles, that is, highly

reclified spirit of wine drawn off from dried damask roses, each one ounce and an half.
Digest in the heat of a water-bath.

THE ambergris, if pure, is here totally dissolved into a reddish liquor, provided the heat be sufficient to make the spirit boil or simmer: with a weaker heat, or if the spirit is not highly recisied, the solution does not succeed. This tincture is a high cordial: eight or ten drops may be taken on sugar.

TINCTURA feu Essentia REGIA. The royal tincture, or effence. Parif.

Take of
Ambergris, two scruples;
Musk, one scruple;
Civet, ten grains;
Oil of cinnamon, six drops;
Oil of rhodium, four drops;
Salt of tartar, half a drachm;
Rectified spirit of wine,
Spirit of roses,
Spirit of orange slowers, each

one ounce and a half.

Grind the falt of tartar with the ambergris, musk, civet, and essential oils, till they are thoroughly mixed; then add the spirits, and digest in a warm place for some days, frequently shaking the veffel; afterwards let the liquor settle, and pour offthe clear from the dregs.

THIS tincture is a very high perfume; and by those who can bear substances of that class, may be taken like the preceding; as a cordial. A few drops give a fine flavour to a large quantity of other liquors. The ambergris diffolves here with less heat than in the foregoing preparation; the effential oils promoting its solution. :

Ē

t

TINCTURA ODO NTALGIAMYN-SICHTI.

Mynficht's tindure for the toothach. Argentoraten .

Take of Guaiacum wood, two ounces; Saffafras, Sarfaparilla, each one ounce; Pellitory of Spain, Alum, Sal prunellæ, each half an ounce; Stavefacre feeds, Henbane feeds, eachtwo drachms; Opium, Cloves, each one drachm and a half: Serpyllum, Origanum, Saffron, each one drachm; Rectified spirit of wine; Vinegar, each one pint and a

half. Reduce the dry ingredients into powder, and extract a tincture from them with the spirit and vinegar mixed.

"ALITTLE of this tincture is to " be taken warm into the mouth. " and repeated if there should be " occasion. It effectually relieves " the most violent toothachs; pre-" venting the afflux of humours, " and furprifingly extracting those " already fettled upon the parts: " the pain feems often on the " first application of it to increase, " but foon after abates and goes " off." The above composition, and this account of its virtues, is from the pharmacopæia of the college of Straiburg.

Essentia alexipharmaca STARLII.

Stabl's alexipharmac essence. Argentoratens.

Take of the roots of Masterwort, Carline thistle, Angelica,

Pimpinella alba, each half an ounce; Swallow wort, Elecampane, White dittany, Contrayerva, Wild valerian, each one ounce. Extract an effence or tincture from these ingredients, with a sufficient quantity of highly reclified spirit of wine.

This tincture, kept a secret by its celebrated author, and first published by Juncker, is greatly efteemed by many of the German phyficians as a diaphoretic and elexipharmac, for attenuating viscid humours, and gently promoting urine; both in low fevers, particularly in exanthematous ones, where the eruptions have been repelled, and in chronical disorders; the dose is twenty or thirty drops or more. It is doubtless a medicine of efficacy, though some of its ingredients might be retrenched without injury to its virtue.

> ESSENTIA LIGNORUM. Essence of the woods. Argentoratenf.

Take of Sassafras, two ounces; Guaiacum, three ounces; China root, Sarfaparilla. Red saunders. Yellow faunders, each one ounce: Spirit of wine, as much as will cover the above ingredients to the heat of four inches.

Digest for eight days, and then fil-

ter the effence.

THIS essence, or tincture, is given in venereal and catarrhous disorders, and impurities of the humours in general, from a scruple to a drachm or more. By gently drawing off half of the spirit, the remainder ·

mainder becomes proportionably fironger, and is then called effentia lignorum concentrata.

BALSAMUM VITE.

Balfam of life.

Brandenburg.

Take of

Effential oils of Lavender,
Nutmegs,
Cloves,
Rhodium,
Serpyllum, each
half a drachm;
Clonamon,
Lemon peel,
Bergamotte,

fcruples;
Balfam of Peru, one drachm;
Highly rectified spirit of lawender, fifteen ounces.

each

two

First dissolve the balkam in the spirit, then add the oils, and digest till the whole is dissolved.

Trees fragrant ballam is an improvement on one described by Hossman, in his notes on Poterius, and is probably the fame, or nearly the same, with the balsam so much celebrated afterwards in that author's practice, internally in languors, faintings, debilities of the nervous system, cholics, &c. from ten to twenty or thirty drops, and externally, applied to the nostrils, temples, &c. in vertiginous, le-thargic, and other like complaints. Thus much is certain, from Hoffman's own writings, that his balfam was composed of fragrant oils diffolved in reclined spirit of wine.

#### SECT. VII

#### Oils by infusion and decoction.

XPRESSED oils extract the refinous and oily parts of vegetables, but do not act upon, or unite with the gummy and mucilaginous: hence the oleum e mucilaginibus of the shops contains nothing of the mucilage which its ingredients abound with. These oils
may be tinged, by vegetable matters, of almost all colours; the
leaves of most plants communicate
a green; yellow slowers, a dilute
gold colour; some red flowers, a
light red; alkanet root, a beautiful and deep red.

In making the officinal oils from the leaves of plants, a good deal of care is necessary, to give them the fine green colour expected in them. It the boiling of the herb in the oil is not continued till all the aqueous moisture has exhaled (the mark of which is, the herb's being crisp) the oil will have a dingy yellowish hue; if continued longer, it turns black, and contracts an empyreumatic smell. The most convenient method of managing the process seems to be, to strain off the oil when sufficiently impregnated with the virtue of the plant, and afterwards to let it stand in a clean vessel, over a gentle sire, until, by frequent trials on a white tile, it appears to have gained the deep green colour required.

## OLEUM CHAMÆMELI. Oil of camomile. Edinb.

Take of

Camomile, with the flowers, fresh gathered and bruised, one pound;

Oil olive, three pints.

Boil them gently till the herb is almost crisp; then strain and press out the oil.

The

sediment.

The oils of other herbs are prepared in the fame manner.

> OLEUM HYPERICI. Oil of St. John's wort. Lond.

Take of

The flowers of St. John's wort, full blown, fresh gathered, and carefully freed from the cups, four ounces; Oil olive, two pints.

Pour the oil upon the flowers, and let them fland together, till the oil is fufficiently coloured.

#### OLEUM e MUCILAGINIBUS. Oil of mucilages. Lond.

Take of

Marshmallow root, fresh, half a pound:

Linfeed,

three Fenugreek feed, each ounces:

Water, two pints; Oil olive, four pints.

Bruise the roots and seeds, and gently boil them in the water for half an hour; then add the oil, and continue the boiling till all the water is wasted: afterwards let the oil be carefully poured off for use.

#### OLEUM SAMBUCINUM. Oil of elder. Lond.

Take of

Elder flowers, one pound,

Oil olive, two pints.

Boil the flowers in the oil, till they are almost crisp; then press out the oil, and fet it by till the feces have subsided.

OLEUM VIRIDE. Green oil.

Lond:

. Take of Bay leaves,

Rue leaves. Marjoram leaves, Sea wormwood leaves. Camomile leaves, each, fresh, gathered, three ounces: Oil olive, two pints. Bruife the herbs, and gently boil them in the oil till they are almost crisp; then press out the oil, let it stand to settle, and afterwards pour it off from the

ALL the foregoing oils are defigned for external applications They are supposed, besides only. the general emollient quality of the oil itself, to receive particular virtues from the ingredients: that of camomile flowers, to be a warm discutient and resolvent; that of St. John's wort flowers to be peculiarly grateful to the nerves, to give great relief in all kinds of pains and weariness, to resolve tumours, and heal wounds and ulcers; and the oil of mucilages to be fofter and more emollient than common An oil prepared in the same manner from wormwood, rubbed on the stomach and umbilical region, is faid to excite appetite, strengthen the viscera, and kill worms; and one from rue, to be of fingular efficacy against worms and colicky pains, and swellings.

It is presumed, however, that at present there are few who expect much more from these preparations than from common oil itself, which has the advantage of being less offenfive. The mucilaginous ingredients, marthmallow root and linfeed, in the oleum e mucilaginibus, make no addition to the virtue of the oil, for mucilages, as already observed, are not soluble in oils. Experience has not discovered any fuch lingular qualities in flowers of St. John's wort, that four ounces of them should communicate any

remark-

remarkable virtue to a quart of oil. Of the other herbs, the more va-Inable parts are diffipated by the boiling heat: and although the remaining matter, if it was taken internally either by itself, or dissolved in watery or spirituous liquors, might not be destitute of activity, yet it can scarcely be supposed, when combined with a large quantity of oil, to have any material effect in external applications. The number of these oils has, therefore, been judiciously retrenched, at the late reformation : the five, above retained, are not one-tenth part of those which were formerly ordered to be kept in the shops. The most certain way of answering the purposes intended by these preparations, appears to be, by mixing with the expressed oil a suitable quantity either of the native refins of vegetables, or of the effential oils and refinous extracts artificially prepared from them.

## OLEUM CAMPHORATUM. Campborated oil. Edinb.

Take of
Fresh drawn oil of almonds, or
linfeed, two ounces;
Camphor, one ounce.
Dissolve the camphor in the oil.

This oil is designed, like the foregoing ones, for external purposes. It has been in use for some time, in the infirmary of Edinburgh, against burns, rheumatic pains, &c. and is thence received into the pharmacopæia of the Edinburgh college.

OLEUM ODORIFERUM.

Odoriferous oil.

Let some fine carded cotton be dipt in oil olive, or oil of ben nuts, that it may be thoroughly imbibed with the oil, without retaining fo much as to drip fpontaneously. Lay a bed of this cotton in the bottom of a tin or porcelane vessel, and lightly spread upon it a pretty thick layer of any odoriferous flowers fresh gathered, as jasmine flowers, violets, lillies of the valley, &c. Above these, spread more of the cotton, and then more flowers, alternately, till the vessel is full: then cover it close, and let it stand for twenty-four hours in a gentle warmth. Great part of the fragtance of the flowers will be communicated to the oil in the cotton, which is to be stratified in the same manner with two or three fresh quantities of the flowers, till it is sufficiently impregnated therewith, after which the oil is to be squeezed out from the cotton in a press.

This appears to be the most effectual method of transferring into expressed oils the odoriferous matter of those tender slowers which yield little or no effential oil; the persumed oils and essences of those slowers, brought from Italy, are prepared in this manner. The odorous parts may be again separated from the oil, and transferred into water or spirit, by distillation with those liquors.

#### CHAPTER IV.

Conservation of recent vegetables and their infusions, &c. by

sugar and honey.

#### SECT. I.

#### Conserves.

ONSERVES are compositions of recent vegetable matters and sugar, beaten together into an uniform mass.

This management is introduced for preserving certain simples, undried, in an agreeable form, with as little alteration as possible in their native virtues; and to some subjects it is very advantageously ap-Vegetables, whose virtues are loft or destroyed in drying, may in this form be kept uninjured for a length of time: for, by carefully fecuring the mouth of the containing veffel, the alteration, as well as diffipation, of their active principles, is generally prevented; and the sugar preserves them from the corruption which juicy vegetables would otherwise undergo.

There are, however, fundry vegetables, whose virtues are impaired by this treatment. Mucilaginous substances, by long lying with sugar, become less gluttinous; and astringents, sensibly softer upon the palate. Many of the fragrant flowers are of so tender and delicate a texture, as almost entirely to lose their peculiar qualities on being beaten or bruised.

In general, it is obvious, that in this form, on account of the large admixture of fugar, only substances of considerable activity can be taken to advantage as medicines. And indeed, conferves are at present considered chiefly as auxiliaries to medicines of greater efficacy; or as intermediums for joining them together. They are very convenient for reducing into boluses or pills, the more ponderous powders, as mercurius dulcis, the calces of iron, and other mineral preparations; which with liquid or less consistent matters, as syrrups, will not cohere.

The shops were formerly incumbered with many conserves altogether insignificant; the sew now retained have in general either an agreeable slavour to recommend them, or are capable of answering some useful purposes as medicines. Their common dose is the bulk of a nutmeg, or as much as can be taken up at once or twice upon the point of a knife. There is in general no great danger of exceeding in this particular.

### General method of preparing conserves.

Leaves are picked from the stalks, and slowers from their cups. They are then beaten in a marble mortar, with a wooden pessle, into a smooth mass; after which, thrice their weight of double refined sugar is added by degrees. grees, and the beating continued till they are uniformly mixed.

The sugar should be pulverized by itself, and passed through a fieve, before it is mixed with the vegetable mass; otherwise it cannot easily be reduced to sufficient finenels, fo as to be duly incorporated. Some vegetables are scarce reducible to the requilite finenels by beating in a mortal : fuch is orange This is most conveniently rasped or grated off from the fruit, then well mixed with the fugar, and the compound fet by in a close vessel for some weeks; after which, it may be beaten fmooth with considerably less labour than at first. This peel and red rose buds, are commonly ground in a wooden mill made for that purpose.

CONSERVA foliorum COCHLEA-RIÆ hortensis.

Conserve of the leaves of garden ∫curvy-gra∫s. L. E.

This is the only form that fourwy grassin substance can be kept in. without the total loss of its virtues. The conserve retains the full taste and virtue of the herb for a very considerable length of time; as a year or two; provided the veffel be made perfectly close, and fet in a cool place, It may be given in fcorbutic habits, three or four times a day, or oftner; though it is more frequently used as an assistant to other medicines of fimilar intention, than depended on by itself. It is an excellent addition to arum root in rheumatic cases; and in this form, even the fresh root of arum may be taken freely, without any complaint of the excellive pungency which of itself it impresses on the tongue. An ounce of fresh arum root, beaten into a pulp, and four ounces or less of conserve of

fcurvy grafs, well mixed together form a compound, in which the pungency of the arum is hardly perceived, and which I have given, with good effect, to the quantity of a nutineg twice or thrice a day. To further sheath the acrimony of the arum, it may be beaten with equal its weight of powdered gum arabic, before the admixture of the conferve.

CONSERVA foliorum LUJULÆ. Conserve of the leaves of woodforrel. L . E

THIS is a very elegant and grateful conserve; in taste it is lightly acidulous, with a pecaliar flavour, which some resemble to that of green tea. It is taken occasionly, for quenching thirst; and cooling the mouth and fauces in hot diftempers. It may be usefully joined to the foregoing preparation, whose virtue it fomewhat promotes, at the same time that it improves the tafte.

CONSERVA foliorum MENTHÆ vulgaris.

Conserve of the leaves of spearmint. Lond.

THE conserve of mint retains the tafte and virtues of the herb. It is given in weakness of the stomach and reaching to vomit; and not unfrequently does fervice in fome cases of this kind, where the warmer and more active preparations of mint would be less proper.

CONSERVA foliorum RUTÆ Conserve of the Leaves of Rue. Lond.

This conserve is given from a drachm to half an ounce, in crudities of the prime viæ, for promoting digestion, and in hysteric

diforders: it gently stimulates the solids, attenuates viscid juices, and excites the natural secretions. Some have had a great opinion of it, taken in a morning, as a preservative against the effects of contagious air or exhalation.

CONSERVA fummitatum AB-SINTHH maritimi.

Conserve of the tops of sea wormwood.

THE conserve of wormwood has been celebrated in dropses: Matthiolus relates, that several persons were cured by it of that distemper, without the assistance of any other medicine. Where the disorder indeed proceeds from a simple laxity or slaccidity of the folids the continued use of this medicine may be of some service; as it appears to be a not inelegant mild corroborant. It is directed to be given in the dose of half an ounce, about three hours before meals.

CONSERVA florum LAVEN-DULÆ.

Conferve of lawender flowers.

Lond.

This conferve is not near so fragrant as the flowers themselves. It is nevertheless a sufficiently agreeable one; and is sometimes used as a mild cordial, and in debilities of the nervous system.

CONSERVA florum MALVÆ.

Conserve of the floruers of mollows.

Lond.

This is looked upon as an emollient, and fometimes made use of as such in disorders of the breast and urinary passages. It is the most unimportant of the conserves: nor do the flowers themselves appear to have much virtue. CONSERVA florum ROSARUM rubrarum immaturarum.

Conserve of the buds of red roses.

L. E.

THIS is a very agreeable and useful conferve. A drachm or two diffolved in warm milk, are frequently given as a light restringent, in weakness of the stomach, and likewise in coughs and phthisical complaints. In the German ephemerides, examples are related of very dangerous phthifes cured by the continued use of this medicine: in one of these cases, twenty pounds of the conserve were taken in the space of a month; and in another. upwards of thirty. Riverius mentions feveral other instances of this kind.

CONSERVA florum RORIS-MARINI. Conferve of rolemary flowers. L E.

ROSEMARY flowers in great meafure lose their peculiar fragrance by beating, and hence the conserve has very little of their flavour. Some are therefore accustomed to make this preparation from the leaves of the plant (which retain their virtues under the petile) or at least to add a portion of their to the flowers. The conserve of rosemary is directed in weakness of the nerves, and as a light cordial.

CONSERVA flavedmis CORTI-CUM AURANTIORUM Hispalensium.

Conferve of the yellow rind of Sevile orange peel, I. E.

This conserve is a very elegant one, containing all the virtues of the peel in a form sufficiently agreeable both with regard to the dose and the conveniency of taking. It is a pleasant warm stomachic,

and in this intention is frequently made use of.

#### CONSERVA FRUCTUS CY-NOSBATI. Conferve of bips.

L. E.

Hips require less sugar for reducing them into a conserve, than the substances above enumerated. Twelve ounces of the pulp of the ripe fruit, are to be mixed with only twenty ounces of sugar.

THE conserve of hips is of some esteem as a soft, cooling restringent; three or four drachms or more are given at a time, in bilious sluxes, sharpness of urine, and hot indispositions of the stomach. A good deal of care is requisite on the part of the apothecary in making this conserve: the pulp is apt to carry with it some of the prickly sibres, which the inside of the fruit is lined

with; if these are retained in the conserve, they will irritate the stomach, so as to occasion vomiting.

### CONSERVA PRUNORUM SILVESTRIUM.

Conferves of floes.

Lond.

Let the floes be put into water, and fet over the fire till they grow foft, with care that they do not burst. Then take the floes out of the water, press out their pulp, and mix with it thrice its weight of double-refined sugar.

THIS preparation is a gentle aftringent, and may be given as such in the dose of two or three drachms. The degree of its astringency will vary according to the maturity of the sloes, and the length of time that the conserve has been kept.

#### SECT. II.

#### Preserves.

RESERVES are made, by fteeping, or boiling recent fimples, first in water, and then in fyrup, or folution of sugar. The subject is afterwards either kept moist in the syrup, or taken out and dried, that the sugar may candy upon it; this last is the most usual method.

In this process, some of the valuable parts of the subject are extracted by the liquor, and consequently lost to the preparation; greater regard being here had to palatableness than medicinal efficacy. And indeed most of the preparations of this kind are considered rather as sweetmeats than as medicines; as the business of the confectioner rather than of the apo-

thecary. It would be needless therefore to mention the doses of the several articles, or give particular remarks on the manner of preparing them.

## RADIX ERYNGII CONDITA. Candied eryngo roots. Lond.

Boil them in water, till the rind will easily peel off; when peeled, slit them through the middle, take out the pith, and wash them three or four times in cold water. For every pound of the roots, so prepared, take two pounds of double-refined sugar, which is to be dissolved in a proper quantity of water, and set over the fire; as soon as the

azo

quor begins to boil, put in the roots, and continue the boiling till they are foft.

After this manner are candied ANGELICÆ CAULES.

Angelica flalks.

## CORTEX AURANTIORUM CONDUITUS. Candied orange peel.

Steep the fresh peels of Seville oranges in water, which is to be frequently renewed, until they lose their bitterness. Then, having dissolved in water a suitable quantity of double-resined sugar, boil the peels in this liquor till they become soft and transparent. After the same manner are candied.

LIMONUM CORTICES.

Lemon peels. [L.]

In the same, or a similar manner, may likewise be candied RADICES ANGELLICÆ. Angelica roots. [E.]

### RADICES HELENII. Elecampane roots. [E.]

All forts of fruits, flowers, and feeds may also be preserved, either by keeping them in syrup, or crusting them over with sugar; but these kinds of preparations scarce belong to the art of pharmacy.

Nutmegs and ginger are brought

to us ready candied from the East-Indies [E.]

## MARS SACCHARATUS. Candied feel. Edinb.

Put any quantity of clean filings of iron into a brass kettle, suspended over a very gentle fire. Add to them, by little and little, twice their weight of white sugar, boiled to the consistence of candy, with which powdered starch has been previously mixed in the proportion of a drachm to every pound; agitating the kettle continually, that the filings may be crusted over with the sugar, and taking great care to prevent their running into lumps.

This is a very agreeable preparation of steel; but has hitherto been made only by the confectioners. The college of Edinburgh received it in the former editions: but, as there described, it was almost impossible to binder the matter from concrating into lumps. They have now discovered the intermedium which prevents that inconvenience, and which the confectioners have kept a secret; the addition of a little starch to the fugar. The preparation may be given to the quantity of half a drachm, in those cases wherein cha! · ly beat medicines are proper. See page 141.

### S E C T. III.

#### Gellies.

PEGETABLE gellies are composed of the juices of fruits and sugar, boiled to a thick confistence. Independently of the admixture of sugar, the boiling appears to occasion some alteration in the quality of the juices themselves.

The recent juices of the summer fruits are prone to fermentation: after they have been boiled, they are less disposed to ferment, and at the same time they are much less liable to produce, in the human body, flatulencies, gripes, or fluxes;

though they still retain, in no fmall degree, their original antiseptic, anti-inflammatory, and aperient or restringent virtues.

#### GELATINA, seu miva CYDONIORUM.

Gelly, or marmalade, of quinces. Edinb.

Take three pints of depurated quince juice, and a pound of white fugar. Simmer them together, to a proper thickness.

This is an useful, cooling, restringent medicine: it is given in weakness of the stomach, reachings to vomit, diarrhoeas and dyfenteries, proceeding from an hot indisposition, or sharp bilious humours. It is best taken in little quantities, as a tea-spoonful or two now and then, either by itfelf, or diluted with any fuitable liquors.

#### GELATINA BERBERORUM. Gelly of barberries. Edinb.

Take a pound of barberries picked clean from the stalks, and the Boil them with a gentle heat, to a due consistence; then pass the gelly through a flannel cloth.

#### GELATINA RIBESIORUM. Gelly of currants. Édinb.

is prepared after the fame manner.

Here the trouble of expression is faved, thefe foft fruits freely giving out their juice, which incorporates with the fugar, in the process. Both these preparations are gratefully sub-acid and cooling, and in this intention are occalionally made use of, for moistening the mouth and fauces in febrile or inflammatory distempers. Diffolved in water, they afford an ufeful diluent drink, of a saponaceous nature, which mingles with the blood or its ferum when thickened (as in fome kinds of fevers) where pure water runs off by the kidneys almost unchanged. By the same qualities, they prove ferviceable likewise in chronical diforders proceeding from obstructions of the viscera, or accompanied with immoderate heat: in bilious fluxes and putrid scurvies, their liberal and continued use has fometimes had good ef-Boerhaave greatly comfects. fame quantity of white fugar. mends these kinds of preparations in the scorbutic disorders to which feafaring people are particularly subject.

#### EECT. IV.

#### Syrups.

CYRUPS are saturated solutions of fugar, made in water, or watery or vinous infusions, or in juices. They were formerly confidered as medicines of much greater importance, than they are thought to be at prefent. Syrups and distilled waters were for some ages made use of as the great As multiples of distilled waters

evacuation of any peccant humour was never attempted till, by a due course of these, it had first been regularly prepared for expullion. Hence arose the exuberant collection of both, which we meet with in pharmacopeias: and like errors have prevailed in each. alteratives: infomuch that the have been compounded from maŢ

ġ

Z

.

i

1

ø

ı

í

₫

erials unfit to give any virtue over the helm: fo numbers of fyrups have been prepared from ingredients, which in this form cannot be taken in sufficient doses to exert their virtues; for twothirds of a fyrup consist of sugar, and greatest part of the remaining third is an aqueous fluid.

Syrups are at present chiefly re-

garded as convenient vehicles for medicines of greater efficacy; and made use of for sweetening draughts and juleps, for reducing the lighter powders into boluses, pills, or electaries, and other like purposes. Some likuse may not improperly be considered as medicines themselves; as those of saffron, and buckthorn berries.

#### General rules for preparing Syrups.

All the rules laid down for making decoctions are likewise to be observed in the decoctions for syrups. Vegetables, both for decoctions and insusions, ought to be dry, unless they are expressly ordered otherwise [E.]

11

In the London pharmacopæia, only the purest or double-refined fugar is allowed.

In the Edinburgh, the directions are less explicit. For such syrups as are prepared without boiling, it is left to the choice of the operator to employ either the double-refined, or the common white fugar; which last he is directed to purify, for those syrups (not for the others) by previously dissolving it in water, clarifying the folution with whites of eggs, and boiling it down to a thick consistence. with care to take off the fcum which rifes during the boiling

In the fyrups prepared by boiling, it has been customary to perform the clarification with whites of eggs after the sugar had been dissolved in the decoction of the vegetable. This method is apparently injurious to the preparation; since not only the impurities of the sugar are thus discharged;

but a considerable part likewise of the medicinal matter, which the water had before taken up from the ingredients, is separated Nor indeed along with them. is the clarification and despumation of the fugar, by itself very adviseable, for its purification by this process is not so perfect as might be expected; after it has undergone this process, the refiners still separate from it a quantity of oily matter, which is difagreeable to weak stomachs. See page 212. It appears therefore most eligible to employ fine sugar for all the fyrups; even the purgative ones (which have been usually made with coarse sugar, as somewhat coinciding with their intention) not excepted; for as purgative medicines are in general ungrateful to the stomach, it is certainly improper to employ an addition which increases their offentiveness.

III.

Where the weight of the fugar is not expressed, twenty-nine ounces thereof are to be taken to every pint of liquor. The fugar is to be reduced into powder, and dissolved in the liquor by the heat of a waterbath, unless ordered otherwise. [L.]

Z 2 Although

Although in the formulæ of feveral of the fyrups, a double weight of fugar to that of the liquor is directed, yet less will generally be sufficient. First therefore dissolve in the liquor an equal weight of sugar, then gradually add some more in powder, till a little remains undissolved at the bottom, which is to be afterwards incorporated by setting the syrup in a waterbath [E.]

The quantity of fugar should be so much, as the liquor is capable of keeping dissolved in the cold: if there is more, a part of it will feparate, and concrete into crystals, or candy; if less, the syrup willbe subject to ferment, especially in warm weather, and change into a vinous or four liquor. If, in crystallizing, only the superstuous fugar separated, it would be of no inconvenience; but when part of the fugar has candied, the remaining fyrup, is found to have an under proportion, and is as subject to fermentation as if it had wanted fugar at first.

IV.

Copper vessels, unless they are well timed, should not be employed in the making of acid syrups, or such as are composed of the

juices of fruits [E.]

The confectioners, who are the most dextrous people at these kinds of preparations, to avoid the expence of frequently new tinning their vessels, rarely make use of any other than copper ones untinned, in the preparation even of the most acid syrups, as of and oranges lemons. theless, by taking due care. that their coppers be well fcoured and perfectly clean, and that fyrup the remain no longer in them than is absolutely necessary, they avoid giving it

any ill taste or quality from the metal. This practice, however, is by no means to be recommended to the apothecary.

The fyrup, when made, is to be fet by till next day; if any faccharine crust appears upon the furface, take it off [L.]

SYRUPUS ex ALLIO. Syrup of Garlick. Lond.

Take of

Garlick, fliced, one pound; Boiling water, two pints.

Macerate them in a close vessel for twelve hours, then strain off the liquor, and dissolve in it a proper quantity of sugar, so as to make a syrup.

This fyrup is occasionally made use of for attenuating viscid phlegm, and promoting expectoration in humoural assumas, and oppressions of the breast; in these cases, it proves a medicine of considerable efficacy, though a very unpleasant one: it tastes and smells strongly of the garlick. The college have received it as an alternative to the oxymel ex allio, for the use of those with whom honey disagrees.

SYRUPUS ex ALTHÆA.

Syrup of mar/bmallows.

Lond.

Take of

Marshmallow roots, fresh, one pound;
Double-refined fugar, four pounds;

Water, one gallon.

Nevercare,
fcourid that
louger
ely neving it

NeverBoil the water with the roots to
one half: when grown thoroughly cold, pour off and prefs out
the decoction, and fet it by for
a night to fettle: next morning,
pour off the clear liquor, and
adding to it the fugar, boil the

whole to the weight of fix pounds.

Edinb.

Take of
Marshmallow roots, three ounces;
Liquorice, one ounce;

Liquorice, one ounce; English maidenhair, one ounce and a half;

White fugar, fix pounds;

Water, fix pints.
Boil the water with the marshmallow roots to the consumption of one third, adding, towards the end, the liquorice
and maidenhair; then strain
out the remaining decoction,
and suffer it to rest for some
time. Pour off the clear liquor from the sediment, and
boil it with the sugar over a
gentle fire, keeping the matter
continually stirring, till it becomes a syrup. This syrup supplies likewise the place of the
pectoral syrup.

THE syrup of marshmallows feems to have been a fort of favourite among the dispensatorywriters, who have taken great pains to alter and amend it, but have been wonderfully tender in retrenching any of its articles. In the above prescriptions, it is lopt of its superfluities, without any injury to its virtues. It is used chiefly in nephritic cases, for sweetening emollient decoctions. and the like; of itself, it can do little service, notwithstanding the high opinion, which some have en-tertained of it; for what can be expected from two or three spoonfuls of the fyrup, when the decoction, from which two or three pounds are made, may be taken at a draught or two? The college of Edinburgh has very properly

united this and the pectoral fyrup into one: for the fyrup of marshmallows has always, till the late reformation, contained the principal ingredients of the pectoral fyrup, and its own capital ingredient coincides in the same intention.

SYRUPUS e CORTICIBUS
AURANTIORUM.
Syrup of orange peel.
Lond

Take of

Yellow rind of Seville orange peel, fresh, eight ounces; Boiling water, five pints.

Macerate them for a slight in a close vessel; next morning, strain out the liquor, and disfolve in it the proper quantity of sugar for making it into a syrup.

Edinb.

Take of the Yellow rind of orange . peel,

fresh, six ounces; Boiling water, three pints.

Infuse them for a night, in a close vessel, then strain the liquor, let it stand to settle, and having poured it off clear from the sediment, dissolve therein twice its weight of white sugar, so as to make it into a syrup without boiling.

In making this fyrup, it is particularly necessary, that the sugar be previously powdered, and dissolved in the insusion with as gentle a heat as possible, to prevent the exhalation of the volatile parts of the peel. With these cautions, the syrup proves a very elegant and agreeable one, possessing great share of the fine slaveur of the orange peel.

SYRUPUS BALSAMICUS.

Balfamic fyrup.

Lond.

Take of Balsam of Tolu, eight ounces; Water, three pints;

Boil them for two or three hours in a circulatory vessel, or at least in a long necked matrass, having its mouth lightly covered. When grown cold, strain out the liquor and mix therewith a proper quantity of sugar to make it into a syrup.

THE coction may be conveniently performed in a retort with a receiver adapted to it, the liquor which comes over being occasionally poured back; or the water may be entirely drawn off, and the sugar dissolved in the distilled liquor.

#### Edinb.

Take of the

Syrup of fugar, just made, and warm from the fire, two pounds;

Tincture of balfam of Tolu, one ounce.

When the fyrup has grown almost cold, stir into it the tincture, by little at a time, agitating them well together, till persectly united. The mixture is then to be kept in the heat of a warm-bath until the spirit has exhaled.

This method of making the balfamic fyrup was dropt in one of the preceding editions of the Edinburgh pharmacopæia, on a complaint that the spirit spoiled the taste of the syrup; which it did in a great degree when the tincture was drawn with malt spirits; the nauseous oil, which all the common malt spirits are accompanied with, being left in the syrup after the evaporation of

the pure spirituous part. Particular care therefore should be taken, that the spirit, employed for making the tincure, be persectly clean, and well rectified from all ill slavour.

The intention of the contrivers of the two foregoing processes seems to have been somewhat different. In the first, the more subtile and fragrant parts of the balsam, are extracted from the grosser resinous matter, and alone retained in the syrup: the other syrup contains the whole substance of the balsam, in larger quantity. They are both moderately impregnated with the agreeable slavour of the balsam.

In some pharmacopeias, an elegant syrup of this kind is prepared from a tincture of balsam of Peru, with rose water and a proper quantity of sugar.

SYRUPUS CARYOPHYLLO-RUM RUBRORUM. Syrup of clove-july flowers. Lond.

Take of

Clove-july flowers, fresh gathered, and freed from the beels, three pounds;

Boiling water, five pints.

Macerate them for a night, in a glass or glazed earthen veffel; then strain off the liquor, and dissolve therein its due proportion of sugar to make it into a syrup.

Edinb.

One pound of the flowers is to be infused in three pints of water, and the syrup made as above, without boiling.

This fyrup is of an agreeable flavour, and a fine red colour; and for these, it is chiefly valued. Some have substituted to it one easily parable at seasons when the

Howers are not to be procured; an ounce of clove spice is infused for some days in twelve ounces of white wine, the liquor strained, and, with the addition of twenty ounces of fugar, boiled to a proper consistence: a little cochineal renders the colour of this syrup exactly fimilar to that prepared from the clove-july flower; and its flavour is of the fame kind, though not so pleasant. The abuse may be readily detected by adding to a little of the fyrup some alkaline salt or ley; which will change the genuine fyrup to a green colour, but in the counterfeit, it will make no fuch alteration, only varying the shade of the red.

SYRUPUS CROCI.
Syrup of Saffron.
Lond.

Take of
Saffron wine, one pint;
Double refined fugar, twentyfive ounces.

Dissolve the sugar in the wine, so as to make a syrup thereof.

SAFFRON is very well fitted for making a fyrup, as in this form a fufficient dose of it is contained in a reasonable compass. This fyrup is at present more frequently prescribed than the wine from which it is made: it is a pleasant cordial, and gives a fine colour to juleps.

## SYRUPUS CYDONIORUM. Syrup of quinces. Lond.

Take of
Quince juice, depurated, three
pints;
Cinnamon, one drachm;
Cloves,
Ginger, each half a drachm;
Red Port wine, one pint;

Double refined fugar, nine

Digest the juice with the spices, in the heat of ashes, for six hours; then adding the wine, pass the liquor through a strainer; and asterwards dissolve in it the sugar, so as to make a syrup.

Ir the quinces are kept for fome time, in an airy place, before the juice is pressed out, the fyrup proves father more elegant, and richer of the fruit, than when they are taken fresh from the tree. In either case, the preparation is a very agreeable, mild, cordial restringent; and in some kinds of loosenesses and disorders of the flomach, may be either taken by itself, in the quantity of a spoonful or two at a time, or employed for reconciling to the palate and stomach, medicines of the more ungrateful kind.

### SYRUPUS KERMESINUS.

Syrup of kermes. Edinb.

This fyrup is brought to us ready made, from the fonthern parts of France.

THE syrup of kermes is of an agreeable taste, and a fine red colour. It is accounted cordial and corroborant, and supposed to be particularly serviceable in weaknesses and other disorders of pregnant women.

#### SYRUPUS e SUCCO LIMONUM. Syrup of lemon juice. Lond.

Take of
Juice of lemons, suffered to
stand till the seces have subsided, and afterwards strained,
two pints;
Z 4 Double

Double refined fugar, fifty ounces.

Diffolve the fugar in the juice, fo as to make a fyrup thereof.

#### Edinb.

Take of
Lemon juice, depurated, two
pounds;
White fugar, four pounds.
Make them into a fyrup according
to art, without boiling.

After the same manner are prepared

SYRUPUS e SUCCO MORORUM. Syrup of mulberries [L.]

SYRUPUS e SUCCO FRUCTUS
RUBI IDÆI.
Syrup of raspberries [L.]

ALL these are very pleasant, cooling syrups, and in this intention are occasionally made use of, in draughts and juleps, for quenching thirst, abating heat, &c. in billious or inflammatory distempers. They are sometimes likewise employed in gargarisms for inflammations of the mouth and tonsils.

### SYRUPUS e MECONIO, five DIACODION.

Syrup of meconium, or diacodium.

Lond.

Take of

White poppy heads, dried and cleared from the feeds, three pounds and a half:

Water, fix gallons.

Cut the heads, and boil them in the water, stirring them now and then, to prevent their burning, till only about one third part of the liquor remains, which will be almost entirely soaked up by the poppies. Then remove the vessel from the fire, strongly press out the decoction, and boil it down to about four pints: strain it whilst hot, first through a fieve, and afterwards through a fine woollen cloth, and set it by for a night, that the feces may subfide. Next morning, pour the liquor off clear, and boil it with six pounds of double-refined sugar, until the weight of the whole is nine pounds, or a little more, that it may become a syrup of a proper consistence.

This fyrup, impregnated with the opiate matter of the poppy heads, is given to children in doses of two or three drachms; to adults, from half an ounce to an ounce and upwards, for obtunding and incrassating aerimonious humours, easing pain, procuring reft, and answering the other intentions of mild opiates. Partioular care is requisite, in its preparation, that it may be always made, as nearly as possible, of the fame strength; and accordingly the college have been very minute in their description of the process.

SYRUPUS PAPAVERIS ALBI, feu de MECONIO, vulgo DIACODION.

Syrup of white poppies, or of meconium, commonly called diacodium.

Edinb.

Take of
White poppy heads, just ripe,
and moderately dried, fourteen ounces;

Boiling water, one gallon.

Let these be steeped together for a night, and then boiled until half the liquor is wasted: strain, and strongly press out the remainder, and boil it, with the addition

addition of four pounds of white sugar, to the consistence of a syrup.

This process is considerably different from the preceding. The poppy heads are not boiled so long; and their quantity, in proportion to the produce of syrup, is much less. How far these differences may affect the strength of the preparation, I shall not take upon me to determine.

#### SYRUPUS PAPAVERIS ERRATICI. Syrup of wild poppies. Lond.

Take of
Wild poppy flowers, fresh, four
pounds;

Boiling water, four pints and a half.

Pour the water on the poppies, fet them over the fire, and frequently stir them, until the flowers are thoroughly moistened: as foon as they have sunk under the water, let the whole be set by to steep for a night; next day pour off, and press out the liquor, and set it by for a night longer to settle: afterwards add the proper quantity of double-refined sugar to make it into a syrup.

THE defign of fetting the flowers over the fire is (as Dr. Pemberton observes) that they may be a little scalded, so as to shrink enough to be all immerged in the water; without this artifice, they can scarce be all got in: but they are no longer to be continued over the fire, than till this effect is produced, lest the liquor become too thick, and the syrup be rendered ropy.

This fyrup has been recommended in diforders of the breaft, coughs, spitting of blood, pleurifies, and other diseases, both as an emollient, and as an opiate. It is one of the lightest of the opiate medicines, and in this respect so weak, that some have doubted of its having any anodyne quality.

## SYRUPUS PECTORALIS Pedoral jyrup. Lond.

Take of,
English maidenhair, dried, five ounces;
Liquorice, four ounces;
Boiling water, five pints.
Macerate them for some hours, then strain out the liquor, and, with a proper quantity of double-refined sugar, make it into a syrup.

THE title of this composition expresses its medical intention: it is fupposed to soften acrimonious humours, allay tickling coughs, and promote the expectoration of tough phlegm. true maidenhair is the only fort that has been usually directed in these kinds of compositions: the use of the English is here very judiciously allowed; not only as being more easily procurable, and having been substituted to the other in the shops, but likewise as there does not seem to be any medicinal difference betwixt them. Fuller finds great fault with both these ingredients, on a supposition that all their virtues fly away in drying: but in this he was certainly mistaken; for the virtues of both these maidenhairs consist in a mucilaginous substance, which suffers no injury by being dried. There is one species indeed, the Canada maidenbair, which has a confiderable share of a pleasant smell and flavour joined to its mucilage; but this is as yet as little known in

the shops, though not uncommon in some of our gardens.

SYRUPUS e FLORIBUS
PARALYSIS.
Syrup of cowflips.

This is made from cowslip flowers, after the same manner as the syrup of clove-july-flowers.

IT has been supposed serviceable in nervous disorders; its agreeable flavour recommends it to the patient, though at present there are sew who suppose it to possess any singular virtues.

### SYRUPUS ROSARUM SOLUTIVUS.

Solutive syrup of roses.

Lond.

Take the liquor that remains after the distillation of fix pounds of damask roses;

Of double-refined fugar, five pounds.

Having pressed out the liquor from the roses, boil it down to three pints, and set it by for a night to settle; next morning, pour it off clear from the sediment, and adding the sugar, boil the mixture to the weight of seven pounds and a half.

#### SYRUPUS ROSARUM PALLIDARUM. Syrup of pale roses. Edinb.

Take of Pale roses, fresh gathered, one

Boiling water, three pints; White fugar, two pounds.

pound:

Macerate the roles in the water for a night; then strain the liquor, and adding to it the sugar, boil them into a syrup.

This fyrup may likewise be made from the liquor remaining after the distillation of rose water, depurated from its seces.

THE liquor remaining after the distillation of roses (provided the still has been perfectly clean) is as proper for making this fyrup as a fresh infusion: for the diftillation only collects those volatile parts, which are distipated in the air, whilst the infusion is boiling to its consistence. This syrup is an agreeable and mild purgative for children, in the dose of half a fpoonful, or a spoonful. It likewife proves gently laxative to adults, and in this intention may be of service in costive habits. Its principal use is in solutive glys-

## SYRUPUS de ROSIS SICCIS. Syrup of dry roses. Edinb.

Take of
Red rofes dried, half a pound;
White fugar, fix pounds;
Boiling water, four pints.
Infuse the roses in the water for a night, then boil them a little, strain out the liquor, and adding to it the fugar, boil them to the consistence of a syrup.

THIS fyrup is supposed to be mildly astringent: but is principally valued on account of its red colour. The London college have omitted it, having retained others at least equal to it in that respect.

## SYRUPUS SCILLITICUS Syrup of fquills. Lond.

Take of
Vinegar of squills, a pint and
a half;
Cinnamon,
Ginger, each one ounce;
Double-refined sugar, three
pounds and a half.
Steep

I

E

Steep the spices in the vinegar for three days; then strain out the liquor, and add the sugar, so as to make a syrup thereof.

Edinb.

Take of
Vinegar of squills, two pints.
White sugar, four pounds.
Make them into a syrup, without boiling.

THE spices, in the first of these compositions, somewhat alleviate the offensiveness of the squills, though not so much as to prevent the medicine from being disagreea-able. It is used chiefly, in doses of a spoonful or two, for attenuating viscid phlegm, and promoting expectoration, which it does very powerfully.

SYRUPUS de SENNA et RHEO-Syrup of fena and Rhubarb. Edinb.

Take of
Sena, two ounces;
Rhubarb, sliced, one ounce;
Ginger, bruised, two drachms,
White sugar, three pounds and
a half.

Current-railins, two ounces;

Water, four pints.

Boil the water with the currants to the confumption of one-fourth; and in the hot decoction infuse for a night the sena, rhubarb, and ginger. The liquor being then strained out, suffered to settle, and poured off clear from the sediment, boil it with the sugar, over a gentle sire, to the considence of a syrup.

This fyrup is defigned chiefly as a purgative for children; but is not a very agreeable one, nor among us often made use of. The former London pharmacopæia had a medicine of this kind, with some superfluous articles, which the committee, in their revisal of it, retrenched: they likewise omitted the sena, as being at best unnecessary, and retained only rhubarb for the purgative ingredient: the composition was, nevertheless, at length intirely expunged, and very justly; for, as they observe, rhubarb is easily given to young children in powder or infusion, and the taste of it cannot be rendered agreeable to them by any sweetening.

SYRUPUS SIMPLEX.

The fimple fyrup.

Lond.

Diffolve in water so much doublerefined sugar, as will make it into a syrup.

SYRUPUS SACCHARI.

Syrup of Sugar..

Edinb.

Take of White Sugar,

Water, each equal quantities. Boil them into a lyrup.

THESE preparations are plain liquid fweets, void of flavour or colour. They are convenient for fundry purpoles where these qualities are not wanted, or would be exceptionable.

SYRUPUS e SPINA CERVINA. Syrup of bucktborn. Lond.

Take of the
Juice of ripe and fresh buckthorn berries, one gallon;
Cinnamon,

Ginger,

Nutmegs, each one ounce; Double-refined fugar, feven pounds.

Set the juice by for fome days, to fettle; then pass it through a strainer.

strainer, and in some part thereof macerate the spices. the rest of the juice, adding towards the end that part in which the spices were macerated, first passed through a strainer: this part of the process must be so managed, that the whole liquor may be reduced to four pints. Lastly, put in the fugar, and make the mixture into a fyrup.

Edinb. Take of the Tuice of ripe buckthorn berries. depurated fix pounds. White fugar, four pounds. Boil them to the confiftence of a fyrup.

BOTH these preparations, in doles of three or four spoonfuls, operate as brilk cathartics. principal inconveniencies attending them are, their being very unpleasant, and their occasioning a thirst and dryness of the mouth and fauces, and sometimes violent gripes: both these may be pre-vented, by drinking liberally of water-gruel, or other warm liquids, during the operation. The un-gratefulness of the buckthorn is endeavoured to be remedied in the first of the above prescriptions, the addition of aromatics which however are scarcely sufficient for that purpose. The fecond also had formerly an aromatic material for the same intention, a drachm of the effential oil of cloves; which being found ineffectual, is now rejected.

#### SYRUPUS VIOLARUM. Syrup. of wiolet. Lond.

Take of Violets, fresh, and well colour-

ed, two pounds:

Boiling water, five pints. Macerate them for a whole day,

in a glass, or at least a glazed earthen vessel; then pour out the liquor, and pass it through a thin linen cloth, carefully avoiding even the lightest preffure: afterwards, adding the due proportion of fugar, make it into a fyrup.

#### Edinb.

Take of

March violets, fresh, one pound: Boiling water, three pints.

Steep them together for a night, in a glazed earthen vessel close covered: then strain out the liquor, and diffolve in it twice its weight of white fugar, fo as to make a syrup without boil-

This fyrup is of a very agreeable flavour, and in the quantity of a spoonful or two, proves to children gently laxative. It is apt to lose, in keeping, the elegant blue colour, for which it is chiefly valued; and hence fome have been induced to counterfeit it with materials whose colour is more permanent. This abuse may be readily discovered, by adding to a little of the suspected syrup any acid or alkaline liquor. If the fyrup is genuine, the acid will change its blue colour to a red, and the alkali will change it to a green; but if counterfeit, these changes will not happen, It is obvious from this mutability of the colour of the violet, that the prescriber would be deceived if he should expect to give any blue tinge to acidulated or alkalized juleps or mixtures, by the addition of the blue fyrup.

SYRUPUS

SYRUPUS ZINGIBERIS. Syrup of ginger. Lond.

Take of

Ginger, cut into thin flices, four

Boiling water, three pints. Macerate them for fome hours, then strain out the liquor, and make it into a syrup with a proper quantity of double-refined iugar.

Edinb.

Take of

Ginger, fliced and bruifed, three ounces:

White fugar, four pounds; Boiling water, three pints.

Steep the ginger in the water, in a close vessel, for a night; then boil them a little, and having strained out the decoction, set it by to fettle. Pour off the clear liquor, add to it the fugar, and make them into a fyrup.

THESE are agreeable and moderately aromatic fyrups, lightly impregnated with the flavour and virtues of the ginger.

CONFECTIO ALKERMES. Confection of kermes.

Take of

Juice of kermes grains, warmed and strained, three pounds; Damask rose water, six ounces by measure: Oil of cinnamon, half a scruple; Double refined fugar, one pound.

Dissolve the sugar in the rose water, by the heat of a water bath, into a syrup; then mix in the juice of kermes, and after it has grown cold, the oil of cinnamon.

Edinb.

Take of

Syrup of kermes, three pounds: Yellow Saunders, Cinnamon each fix drachnis: Cochineal, three drachms: Saffron, one drachm and an half. Evaporate the fyrup, with a gentle heat, to the confidence of honey; then mix with it the other ingredients reduced to a very fine powder:

Both these compositions are elegant and agreeable cordials: the dose, when taken by themfelves, is from a scruple to a drachm or more. The first has an advantage of mixing uniformly in juleps. without spoiling their transparency, which the powders in the fecond always do. Particular care ought to be had in the choice of the effential oil, which for the most part is grievously adulterated; it would be convenient to grind the oil with a little of the fugar, before it is added to the other ingredients: for by this means, it will mix more perfectly, and not be apt to separate in keeping.

#### SEC T. IV.

#### Honeys and Oxymels.

HE more fixt parts of vegetables, diffolved in watery liquors, may be thence transferred into honey, by mixing the honey with the watery decoction or juice

of the plant, and boiling them together till the aqueous part has exhaled and the honey remains of its original confistence.

MEL ELATINES.

Honey of fluellin.

Lond.

Take of
Depurated juice of fluellin, four

Clarified honey, four pounds; Boil them to a due consistence.

This preparation made its first appearance in the preceding edition of our pharmacopæia. It is very rareley made use of, and not often kept in the saops.

## MEL HELLEBORATUM. Honey of hellebore. Lond.

Take of

White hellebore roots, dried and cut in flices, one pound; Clarified honey, three pounds; Water four pints.

Let the roots be macerated in the water for three days, and then boiled a little; press out the liquor, and having passed it again through a strainer, boil it with the honey to a proper thickness.

PARTICULAR care ought to be had, to reduce its preparation as nearly as possible to the honey confistence, that its strength may not be too uncertain. It acts, as a drastic purgative or emetic, too violent and precarious for common use. It has been sometimes given in maniacal cases, in doses of one or two drachms or upwards; though more frequently employed in glisters. The present practice very rarely makes use of it at all.

MEL MERCURIALE.

Honey of mercury.

Take of

Juice of French-herb-mercury, Honey, each three pounds. Boil them together to the confistence of honey, taking off the foum which arises to the top.

THIS is defigned chiefly for glyfters: it is very rarely made use of, and hence is now dropt both by the London and Edingburgh colleges.

MEL ROSACEUM.

Honey of Roses.

Lond.

Take of

Red rose buds, freed from the heels, and hastily dried, four ounces:

Boiling water three pints; Clarified honey, five pounds.

Steep the roles in the water for fome hours, then strain off the liquor, mix with it the honey, and boil them to a due consistence.

Edinb.

Take of Red roses, dried, half a pound; Boiling water, four pints; Clarified honey, six pounds.

Steep the roses in the water for a night, then strain out the liquor, add to it the honey, and boil the mixture to the consistence of honey.

This preparation is not unfrequently made use of, as a mild cooling detergent, particularly in gargatisms for ulcerations and inflammation of the mouth and tonsils. The design of hastily drying the roses, as directed in the first of the above prescriptions, is, that they may the better preserve their astringency. See page 252.

## MEL SOLUTIVUM. Solutive boney. Lond.

Take of

The liquor remaining after the diffillation of fix pounds of damask roses;

. ... ...

Cum-

í

3

ſ

1

Cummin seeds, bruised a little, one ounce;

Brown sugar, four pounds; Honey, two pounds.

Having pressed out the liquor, boil it to three pints; adding, to-ward the end, the seeds tied up in a linen cloth. Then put in the sugar and honey, and boil down the mixture to the consistence of thin honey.

This composition is very well
contrived for the purpose expressed
in its title. It is principally employed in laxative glysters; and
hence brown sugar is here allowed;
whilst, for all other uses, the
double-refined is directed.

## OXYMEL ex ALLIO. Oxymel of garlick. Lond.

Take of

Garlick, cut in slices, an ounce and a half;

Caraway feeds,

Sweet fennel feeds, each two

Clarified honey, ten ounces by weight;

Vinegar half a pint.

Boil the vinegar, for a little time with the feeds bruised, in a glazed earthen vessel; then add the garlick, and cover the vessel close; when grown cold, pressout the liquor, and dissolve in it the honey, by the heat of a water-bath.

This oxymel is recommended for attenuating viscid juices, promoting expectoration, and the fluid fecretions in general. It is doubtless a medicine of considerable efficacy, though very unpleafant, the flavour of the garlick prevailing, notwithstanding the adelition of the aromatic seeds. OXYMEL PECTORALE.

Pedoral oxymel,

Edinb.

Take of

Elecampane roots, one ounce; Florence orris roots, half an ounce;

Gum ammoniacum, one ounce; Vinegar, half a pint;

Clarified honey, one pound; Water three pints.

Let the roots, cut and bruised, be boiled in the water till one third is iwasted; then strain off the liquor, let it stand to settle, and having poured it off clear from the seces, add to it the honey, and the ammoniacum, previously dissolved in the vinegar. Mix them together by boiling them a little.

THE title of this composition expresses its medical virtues. It is designed for those disorders of the breast that proceed from a load of viscid phlegm (which this medicine attenuates and promotes the expectoration of) and obstructions of the pulmonary vessels. Two or three spoonfuls may be taken every night and morning, and continued for some time.

## OXYMEL SCILLITICUM. Oxymel of fquille. Lond.

Take of

Clarified honey, three pounds; Vinegar of fquills, two pints. Boil them in a glazed earthen veffel, over a gentle fire, to the confiftence of a fyrup.

#### Edinb.

Take of

Clarified honey, four pounds; Vinegar of squills, two pints. Boil them to the consistence of a fyrup.

THE

THE honey was formerly employed for this preparation unclarified; and the scum, which in fuch cases arises in the boiling. taken off: by this means, the impurities of the honey were dif-charged; but some of the medicinal parts of the squills, which the vinegar was impregnated with, were also separated. For this reason the colleges both of London and Edinburgh have now judiciously ordered the honey, for all these kinds of preparations, to be previously clarified by it

Oxymel of fquills is an useful aperient, detergent, and expectorant, and of great fervice in humoural afthmas, coughs, and other disorders, where thick phlegm abounds. It is given in doses of from danger. Their glazing is two or three drachms, along with procured by a vitrification of two or three drachms, along with fome aromatic water, as that of lead; and vinegar by a boiling cinnamon, to prevent the great heat, may corrode so much of vinausea which it would otherwise trified lead, as to receive from it be apt to excite. In large doses, noxious qualities. See page 35. it proves emetic.

OXYMEL SIMPLEX. Simple oxymel.

[L. E]

Take of Clarified honey, two pounds; Vinegar, one pint. Boil them to a due confidence.

THIS simple preparation is not inferior in efficacy to many more elaborate compositions. It is an agreeable, mild, cooling, faponaceous, detergent, and attenuating medicine. It is often used in cooling, detergent gargarisms, and not unfrequently as an expecto-

The boiling of oxymels in glazed earthen vessels, is not free

#### CHAPTER V.

Separation and collection of those parts of vegetable and animal substances, which are volatile in the heat of boiling water.

THERE are many vegetable, and some animal substances, whose virtues reside, wholly or in part, in a matter which is capable of totally exhaling in the heat of boiling water. In most of the processes hitherto described, it is endeavoured, as much as possible to preserve this volatile matter along with the more fixt parts; whether those fixt parts were themselves medicinal, or only subservient to the union of the volatile

matter with the fluids employed. The aim, in the present chapter, is, to completely separate this volatile subtile principle, and collect it pure from the grosser fixt parts, either in a concentrated state, or diluted with water or spirit of wine. In its concentrated state, it appears commonly an oil; which, from its containing always the specific odour, and frequently the other medicinal powers of the subject, is called essential oil.

#### S E C T. I.

#### Essential Oils.

distillation in an alembic, with a large refrigeratory. A quantity of water is added to the subject, sufficient to prevent its burning; and in this water, it is likewise macerated a little time before the distillation. The oil comes over along with the water; and either swims on its surface, or sinks to the bottom, according as it is lighter or heavier than that sluid [L.]

In the Edinburgh pharmacopoeia, fome fea falt is ordered to be added to the water, sufficient to give it a slight brackish taste. The length of the maceration is to be varied according to the texture and compactness of the subject: the most tender subjects

scarce require any: those of a foft and loofe texture are to be steeped for two or three days; and the more viscous ones, for a longer time: the further the maceration is intended to be protracted, the greater quantity of sea falt must be added. From viscous substances the oil may be obtained in a shorter time, by submitting them to a slight and not too long continued, fermentation; in which case, the addition of salt is improper. Seeds and spices are to be bruised, and woods to be rasped, previously to the maceration or fermentation [E.]

ESSENTIAL oils are obtained only from odoriferous substances; A a but

but not equally from all of this class, nor in quantity proportionable to their degree of odour; some, which, if we were to reason from analogy, should seem very well fitted for this process, yielding extremely little oil, and others none at all. Roses and camomile flowers, whose strong and lasting fmell promifes abundance, found upon experiment to contain but a small quantity; the violet and jasmine flower, which persume the air with their odour, lose their fmell upon the gentlest coction, and do not afford the least perceptible mark of oil upon being distilled, unless immense quantities are submitted to the operation at once; whilst favin, whose disagreeable fcent extends to no great distance, gives out the most oil of almost any vegetable known.

Nor are the same plants equally fit for this operation, when produced in different foils or feasons; or at different times of their growth. Some yield more oil if gathered when the flowers begin to fall off than at any other time; lavender and rue for inftance. Others, as fage, afford the largest quantity when young, before they have fent forth any flowers: and others, as thyme, when the flowers have just appeared. All fragrant herbs yield a larger proportion of oil when produced in dry foils and warm fummers, than in the oppofite circumstances. On the other hand, fome of the difagreeable strong-scented ones, as wormwood, are faid to contain most, in rainy feasons and moist rich grounds.

SEVERAL of the chemists have been of opinion, that herbs and flowers, moderately dried, yield a greater quantity of essential oil, than if they were distilled when fresh. It is supposed, that the oil being already blended, in fresh plants, with a watery fluid, great part of it remains distused through the water after the distillation, divided into particles too minute to unite and be collected; whereas in drying, the oily parts, on the exhalation of the moisture which kept them divided and dispersed, run together into globules, which have little disposition to mingle with watery sluids, and easily separate from the water employed in the distillation.

This theory, however, does not appear to be altogether fatisfactory; for though the oil be collected in the subject into distinct globules, it does not rife in that form, but refolved into vapour, and blended and coagitated by the heat with the vapour of the water; and if the oil in a dry plant was less disposed to unite with aqueous fluids than in a fresh one, the dry ought to yield a weaker infusion than the fresh, the contrary of which is generally found to obtain. the oil of the dry plant is most perfectly extracted, and kept diffolved by the water before the difillation, I can fee no reason why it should have a greater tendency to separate from the water after-

The opinion of dry plants yielding most oil seems to have arisen from an observation of Hossman, which has, I think, been missinderstood: A pound, he says, of dry spike showers yields an ounce of oil, but if they were distilled fresh, they would scarce- ly yield above half an ounce; and the case is the same in balm, sage, &c. The reason is, that in drying, the watery humidity exhales; and as from two pounds of a fresh plant we do

ŀ

not obtain above one pound of dry, and little of the subtile oil evaporates in the drying, it follows, that more oil ought to be afforded by the dry than by the fresh." The meaning of which I apprehend to be no more than this, that if two pounds of a fresh plant are by drying reduced to one, without any loss of the oil, then the one pound dry ought to be equivalent to the two A late writer quotes an experiment of Neumann, which appears to be misunderstood in the same manner; for Neumann, in the place referred to, fays only, that dry wormwood is found to yield much more oil than an equal weight of the fresh plant. I do not recollect any instance, in which fresh and dry plants have been brought to a fair comparison, by dividing a quantity of the subject into two equal weights, and diftilling one while fresh, and the other after it has been carefully and moderately dried.

But whatever may be the effect of moderate exticcation, it is certain, that if the drying be long continued, the produce of oil will be diminished, its colour altered, and its smell impaired.

WITH regard to the proportion of water, if whole plants, moderately dried, are used, or the shavings of woods; as much of either may be put into the vessel, as, lightly pressed, will occupy half its cavity; and as much water may be added, as will rife up to two thirds its height. The water and ingredients, altogether, thould never take up more than three-fourths of the still; there should be liquor enough to prevent any danger of an empyreuma, but not so much as to be too apt to boil over into the receiver.

THE maceration should be continued fo long, as that the water may fully penetrate the pores of the subject. To promote this effect, woods should be thinly shaved across the grain, roots cut transversely into thin slices, barks reduced into coarse powder, and feeds lightly bruifed. Very compact and tenacious substances require the maceration to be continued a week or two, or longer; for those of a softer and looser texture, two or three days are sufficient: whilst some tender herbs and flowers not only fland not in need of any at all, but are even injured by it.

Whether the addition of fea falt is of any real fervice, is greatly to be doubted. The uses generally. alligned to it are, to penetrate and unlock the texture of the subject more effectually than simple water could do: and to prevent the fermentation or putrefaction, which the matter is apt to run into. during the length of time that the maceration is often continued. But sea salt seems rather to harden and condense, than to soften and resolve, both vegetable and animal subjects; and if it prevents putrefaction, it must, on that very account, be rather injurious than of service. The resolution, here aimed at, approaches near to a beginning putrefaction; and faline fubitances, by retarding this, prolong the maceration far beyond the time that would otherwise be necessary. It is in the power of the operator, when he perceives the process coming near this pitch, to put a stop to it at pleasure, by proceeding immediately to distillation: by this means, the whole affair will be finished in a very little time, with at least equal advantage in every other respect; provided the manual operations of A a 2

pounding, rasping, and the like which are equally necessary in either case, be scientifically complied with.

Bodies of a very viscous and compact texture are directed, in the Edinburgh pharmacopæia, to be fermented for fome days with a little yeast; half their quantity of water is sufficient for performing the fermentation; fo much more as is necessary, is to be added afterwards, before the distillation. This process undoubtedly motes the resolution of the subject, and the extrication of the oil: it happens, however, that affistances of this kind are needful. Particular care must be had not to continue the fermentation too long; or to give a bad flavour to the oil by an ill chosen ferment, or using too large a quantity of any.

Some chemists pretend, that by the addition of faits and acid spirits, they have been enabled to gain more oil from certain vegetable matters, than can possibly be got from them without such affiftance. Experiments made on purpose to settle this point seem to prove the contrary; this at least is constantly found to be true, that where there is any reason to think the yield to be greater than usual, the quality of the oil is propor-The quantity tionably injured. of true effential oil in vegetables, can by no means be increased: and what is really contained in them may be easily separated without any addition of this kind. All that faline matters can do in this respect, is, to make the water susceptible of a greater degree of heat than it can sustain by itfelf, and thus enable it to carry up a gross uncluous matter, not volatile enough to arise with pure water: this gross matter, mingling with the pure oil, increases

the quantity, but at the fame time must necessarily debase its quality. And indeed when water alone is made use of, the oil which comes over about the end of the operation is remarkably less fragrant, and of a thicker consistence, than that which arises at the beginning; distilled a second time, with a gentle heat, it leaves a large quantity of gross almost insipid resinous matter behind.

THE choice of proper instruments is of great confequence to the performance of this process to advantage. There are some oils, which pass freely over the swan neck of the head of the common still; others less volatile, cannot easily be made to rife so high. For obtaining these last we would recommend a large low head, having a rim or hollow canal round it: in this canal the oil is detained on its first ascent (and thence conveyed at once into the receiver) the advantages of which are sufficiently obvious.

With regard to the fire, the operator ought to be expeditious in raising it at first, and to keep it up, during the whole process, of such a degree, that the oil may freely distil, otherwise, the oil will be exposed to an unnecessary heat, a circumstance which ought as much as possible to be avoided. Fire communicates to all these oils a disagreeable impression, as is evident from their being much less grateful when newly distilled, than after they have stood for some time in a cool place: the longer the heat is continued, the more alteration it must produce in them.

The greater number of oils require for their distillation the heat of water strongly boiling; but there are many also which rife

with a considerably less heat: fuch as those of lemon peel, citron peel, of the flowers of lavender and rolemary, and of almost all the more odoriferous kinds of flowers. We have already observed, that these flowers have their fragrance greatly injured, or even destroyed, by beating or bruising them: it is impaired also by the immersion in water, in the present process, and the more so in proportion to the continuance of the immersion, and the heat; hence these oils, distilled in the common manner, prove much less agreeable in smell than the subjects themselves. For the distillation of fubstances of this class, I have contrived another method; instead of being immersed in water, they are exposed only to its vapour, A proper quantity of water being put into the bottom of the still, the odoriferous herbs or flowers are laid lightly in a basket, of fuch a fize, that it may enter into the still, and rest against its sides, just above the water. The head being then fitted on, and the water made to boil, the Ream, percolating through the subject, imbibes the oil, without impairing its fragrance, and carries it over into the receiver. Oils thus obtained possess the odour of the subject in an exquisite degree, and have nothing of the disagreeable scent perceivable in those distilled by boiling them in water in the common manner.

It may be proper to observe, that those oils, which rise with a less heat than that of boiling water, are generally called, by the chemical and pharmaceutical writers, light oils; and those which require the heat of water strongly boiling, are called ponderous. I have avoided these expressions, as they might be

thought to relate to the comparative gravities of the oils; with which the volatility or fixedness have no connection. Oil olive is lighter than most of the effential oils; but the heat requisite to make it distil exceeds that in which the heaviest essential oil distils, considerably more than the heat of boiling water exceeds that of ice.

THE water employed in the difillation of effential oils, always imbibes some portion of the oil; as is evident from the smell, taste, and colour which it acquires. It cannot however retain above a certain quantity; and therefore, such as has been already used and almost faturated itself, may be advantageously employed, instead of common water, in a second, third, or any suture distillation of the same subject.

Some late chemical writers recommend, not the water which comes over, but that which remains in the still, to be used a second time. This can be of no service; as containing only such parts of the vegetable as are not capable of arising in distillation, and which serve only to impede the action of the water as a menstruum, and to endanger an em-

pyreuma.

After the distillation of one oil, particular care should be had to duly cleanse the worm before it is employed in the distillation of a different plant. Some oils, those of wormwood and aniseeds for instance, adhere to it so tenaciously, as not to be melted out by heat, or washed off by water: the best way of cleaning the worm from these, is to run a little spirit of wine through it.

Essential oils, after they are distilled, should be suffered to stand A a 3 for

which it may be reduced into by mixing it with crumb of bread.

# OLEUM SEMINUM ANETHI ESSENTIALE. Effential oil of dill feeds. Lond.

This is a very warm oil; of a flavour not very agreeable, less so than that of the seeds. It is sometimes given as a carminative, in flatulencies, cholicky pains, hiccups, and the like, from one to three or four drops.

### OLEUM SEMINUM ANISI ESSENTIALE.

Essential oil of aniseeds. L. E.

This oil possesses the taste and fmell of the anifeeds in perfection. It is one of the mildest of the diflilled oils: fifteen or twenty drops may be taken at a time without danger, though common practice rarely goes to far as half this number. Its fmell is extremely durable and diffusive: milk drawn from the breast after taking it, is found impregnated with its odour: and possibly this may be, in part, the foundation of the pectoral virtues usus virtues usus virtues usus virtues usus virtues usus virtues usus virtues virtues usus virtues virtues usus virtues flatulencies and cholics, it is said by some to be less effectual than the feeds themselves.

It is remarkable of this oil, that it congeals, even when the air is not fenfibly cold, into a butyraceous confiftence; and hence, in the distillation of it, the operator ought not to be over-solicitous in keeping the water in the refrigeratory too cool; it behoves him rather to let it grow fomewhat hot, particularly towards the end of the process; .otherwise the oil, congealing, may fo stop up the worm, as to endanger blowing off the head of the still, at least a confiderable quantity of oil will remain in it.

#### OLEUM SEMINUM CARUI ESSENTIALE.

Essential oil of caraway seeds.

The flavour of this exactly refembles that of the caraway. It is a very hot and pungent oil; a fingle drop is a moderate dofe, and five or fix a very large one. It is not unfrequently made use of as a carminative; and supposed by some to be peculiarly serviceable for promoting urine, to which it communicates some degree of its smell.

#### OLEUM CARYOPHYLLORUM AROMATICORUM ESSENTIALE.

Essential oil of cloves. L. E.

This oil is so ponderous as to fink in water, and is not easily elevated in distillation; if the water which comes over be returned on the remaining cloves, and the distillation repeated, some more oil will generally be obtained, though much inferior in quality to the The oil of cloves is usually described as being " in taste exces-" fively hot and fiery, and of a gold yellow colour." (Boerb. process. 27.) Such indeed is the composition which we receive under this name from Holland; but the genuine oil of cloves is one of the milder oils : it may be taken with great safety (duly diluted) to the quantity of ten or twelve drops or more. Nor is its colour at all yellow, unless it has been long and carelessly kept, or distilled by too violent a fire: when in perfection, it is limpid and colourless, of a pleasant, moderately warm and pungent tafte, and a very agreeable fmell, much refembling that of the spice itself. The Dutch oil of cloves contains a large quantity of expressed oil, as evidently appears

İ

i

ı

appears upon examining it by distillation. This however, cannot be the addition to which it owes its acrimony. A small proportion of a resinous extract of cloves communicates to a large one of oil a deep colour, and a great degree of acrimony.

#### OLEUM FLORUM CHAMÆ... MELI ESSENTIALE

Essential oil of camomile slowers.

Lond.

This is a very pungent oil, of a firong not ungrateful finell, refembling that of the flowers: its colour is yellow, with a cast of greenish or brown. It is sometimes given in the dose of a few drops, as a carminative, in hysteric disorders, and likewise as a vermisuge: it may be conveniently made into pills with crumb of bread.

The oil above described is that obtained from the common garden camomile, which is the only fort directed in our dispensatories (see the foregoing part, page 124.) There is another species, more frequent in corn-fields than in our gardens, (chamemælum vulgare Ger. Raii spnops. ed. 3. 288.) which yields a beautiful blue oil: this colour, if the oil is carefully kept, remains for many years; but if the air is not perfectly excluded, soon degenerates into a yellow like that of the foregoing.

## OLEUM CINNAMOMI. Oil of cinnamon. L. E.

This valuable oil is extremely hot and pungent, of a most agreeable stavour, like that of the cinnamon itself. In cold languid cases, and debilities of the nervous system, it is one of the most immediate cordials and restoratives. The dose is one, two or three drops: which must always be carefully di-

luted by the mediation of fugar. &c. for so great is the pungency of this oil, that a fingle drop let fall upon the tongue, undiluted, produces, as Boerrhaave observes, a gangrenous eschar. In the distillation of this oil, a smart fire is required; and the low head, with a channel round it, above recommended for the distillation of the less volatile oils (page 356.) is particularly necessary for this, which is one of the least volatile, and which is afforded by the spice in exceeding small quantity. The distilled water retains no small portion of the oil: but this oil being very ponderous, great part of it subsides, from the water, on standing for two or three weeks in a cool place.

### OLEUM SEMINUM CYMINI ESSENTIALE.

Effential oil of cummin feeds.

Lond.

This is one of the warmer and less pleasant oils. It is employed chiefly in cold, flatulent, hysteric complaints, in doses of two or three drops. It gives its smell strong to the urine, and is supposed peculiarly serviceable for promoting its discharge.

### OLEUM SEMINUM FOENICULI ESSENTIALE.

Essential oil of fennel seeds. Edinb.

The oil obtained from sweet sennel seeds is much more elegant and agreeable than that of the common sennel (see page 142.) It is one of the mildest of these preparations: it is nearly of the same degree of warmth with that of aniseeds, to which it is likewise similar in slavour, though far more grateful. It is given from two or three drops to ten or twelve, as a carminative, in cold indispositions of the sto-mach:

much: and in some kinds of coughs, for promoting expectoration.

OLEUM foliorum HYSSOPI. ESSENTIALE.

Essential oil of by sop leaves. The oil of hyssop is moderately acrid, of a strong not very agreeable fmell, exactly refembling the original herb: its colour is yellowish, with a flight cast of green; which in keeping changes to a brown-It is commended in humoural asthmas, for promoting expectoration, &c. from one to two or three drops: but it is rarely made use of, and not often kept in the shops; for which reason, it is now omitted both by the London and Edinburgh colleges.

OLEUM baccarum JUNIPERI ES-SENTIALE.

Essential oil of juniper berries. Ľ. E.

This oil is a very warm and pungent one, of a strong flavour, not unlike that of the berries. In the dose of a drop or two, it proves a ferviceable carminative and stomachic: in one of fix, eight, or more, a stimulating, detergent, diuretic and emmenagogue: it feems to have somewhat of the nature of the turpentines, or their distilled oil; like which it communicates a violet fmell to the urine.

The oil of these berries resides partly in velicles spread through the fubstance of the fruit, and partly in little cells contained in the feeds: when the berry is dry, and the oil hardened into a refinous substance, it becomes visible, upon breaking the feeds, in form of little transparent drops. In order therefore to obtain this oil to advantage, we ought, previous to the distillation, to bruise the berry thoroughly; so as to break the feeds, and entirely lay open the oily receptacles.

OLEUM florum LAVENDULÆ ESSENTIALE.

Essential oil of lawender flowers. L. E.

This oil, when in perfection, is very limpid, of a pleasant yellowish colour, extremely fragrant, posfessing in an eminent degree the peculiar finell generally admired in the flowers. It is a medicine of great use, both externally and internally, in paralytic and lethargic complaints, rheumatic pains, and debilities of the nervous system. The dose is from one drop to five

Lavender flowers yield the most fragrant oil, and in considerably the largest quantity, when they are ready to fall off spontaneously, and the feeds begin to shew themselves: the leaves give out extremely lit-The flowers may be separated from the rest of the plant, by drying it a little, and then gently beating it: they should be immediately committed to distillation, and the process conducted with a well regulated gentle heat: too great heat would not only change the colour of the oil, but likewife make a disagreeable alteration in its fmell.

OLEUW baccarum LAURI ESSEN-TIALE.

Essential oil of bayberries.

The oil of bayberries is thin and limpid, moderately pungent, of a strong and tolerably grateful smell. It is given in flatulent colics, hyfteric complaints, and for allaying the pains consequent upon delivery, the dole, from two drops to five or fix. It is likewife made an ingredient in carminative glysters; and in some hysteric cases, is applied externally.

ESSEN-

ŀ

ESSENTIA LIMONUM [L.]
OLEUM corticum LIMONUM
[E.]

Essence of lemons, or the essential oil of lemon peel.

This is a pleafant oil, of a fine smell, very near as agreeable as that of the fresh peel: it is one of the lightest and most volatile effential oils we have, perfectly limpid, and almost colourless. It is taken in doses of two or three drops, as a cordial, in weakness of the stomach, &c. though more frequently used as a persume. It gives a sine slavour to the officinal spiritus volatilus aromaticus, and occasions the soap pills to sit easy on the stomach.

OLEUM MACIS ESSENTIALE.

Essential oil of mace.

The effential oil of mace is moderately pungent, very subtile and volatile, of a strong aromatic smell, like that of the spice itself: it is thin and limpid, of a pale yellowish colour, with a portion of thicker and darker coloured oil at the bottom. This oil is celebrated in vomiting, hiccups, colicky pains, &c. both given internally from one to four drops, and applied externally to the stomach and umbilical region. It is however but rarely made use of, and not often met with in the shops.

#### OLEUM MAJORANÆ ESSENTIALE.

Essential oil of marjoram leaves.

This oil is very hot and penetrating, in flavour not near so agreeable as the marjoram itself: when in persection, it is of a pale yellow colour; by long keeping, it turns reddish: if distilled with too great a heat, it arises of this colour at

first. It is supposed to be peculiarly serviceable in relaxations, obstructions, and mucous discharges of the uterus: the dose is one or two drops.

#### OLEUM MENTHÆ ESSENTIALE.

Essential oil of the leaves of common mint.

 $\{L, E, \}$ This oil smells and tastes strongly of the mint, but is in both respects somewhat less agreeable than the herb itself. It is an useful stomachic medicine; and not unfrequently exhibited in want of appetite, weakness of the stomach, reachings to vomit, and other like disorders, when not accompanied with heat or inflammation: two or three drops, or more, are given for a dose. It is likewise employed externally for the same purposes: and is an excellent ingredient in the stomachic plasters of the shops.

OLEUM MENTHÆ PIPERITI-DIS ESSENTIALE.

Essential oil of the leaves of pepper mint.

Lond.

This possesses the smell, taste, and virtues of the pepper mint in persection; the colour is a pale greenish yellow. It is a medicine of great pungency and subtility; and dissusses, almost as soon as taken, a glowing warmth through the whole system. In colics, accompanied with great coldness, and in some hysteric complaints, it is of excellent service. A drop or two are in general a sufficient dose.

#### OLEUM NUCIS MOSCHATÆ ESSENTIALE.

Essential oil of nutmegs.

L. E.

The effential oil of nutmegs poffesses the slavour and aromatic virtues of the spice in an eminent degree. It is similar in quality to the oil of mace, but somewhat less grateful.

#### OLEUM ORIGANI ESSENTIALE.

Essential oil of the leaves of origanum.

[L. E.]

This oil has a very pungent acrimonious tafte, and a penetrating smell. It has been chiefly employed externally as an errhine, and for easing pains of the teeth.

### OLEUM ESSENTIALE PIPERIS JAMAICENSIS.

Essential oil of Jamaica pepper. Edinb.

This is a very elegant oil, and may be used as a succedaneum to those of some of the dearer spices. It is of a sine pale colour, in slavour more agreeable than the oil of cloves, and not far short of that of nutmegs. It sinks in water, like the oils of some of the eastern spices.

#### OLEUM PULEGII ESSENTIALE.

Essential oil of the leaves of pennyroyal.

[L. E.]

This oil, in smell and taste, refembles the original plant; the virtues of which it likewise possesses, It is given in hysteric cases, from one to sour or sive drops.

#### OLEUM RORISMARINI ES-SENTIALE.

Essential oil of rosemary.
[L. E.]

The oil of rosemary is drawn from the plant in flower. When in perfection, it is very light and thin, pale, and almost colourless; of great fragrancy, though not quite so agreeable as the rosemary itself.

It is recommended, in the dose of a few drops, in nervous and hysteric complaints. Boerhaave holds it in great esteem against epilepsies, and suppressions of the uterine purgations, occasioned by weakness and inactivity.

#### OLEUM LIGNI RHODII ESSENTIALE.

Effential oil of rhodium. L. E.

This oil is extremely odoriferous, and principally employed as a perfume in scenting pomatums, and the like. Custom has not as yet received any preparation of this elegant aromatic wood into internal use.

### OLEUM RUTÆ ESSENTIALE. Essential oil of rue leaves.

L. E.

The oil of rue has a very acrid tafte, and a penetrating smell, refembling that of the herb, but rather more unpleasant. It is sometimes made use of in hysteric disorders, and as an anthelmintic; as also in epilepsies proceeding from a relaxed state of the nerves.

Rue yields its oil very sparingly. The largest quantity is obtained from it when the slowers are ready to fall off, and the seeds begin to shew themselves: suitable maceration, previous to the distillation, is here extremely necessary.

#### OLEUM SABINÆ ESSENTIALE. Tential oil of favin leave:

Essential oil of savin leaves
L. E.

Savin is one of the plants which in former editions of the Edinburgh pharmacopæia were directed to be lightly fermented before the distillation; this, however, is not very necessary; for savin yields, without any fermentation, and even without much maceration, a very large

1

¥

è

1,

Į

large quantity of oil: the foregoing herb stands more in need of a treatment of this kind. The oil of favin is a celebrated utrine and emmenagogue: in cold phlegmatic habits, it is undoubtedly a medicine of good service, though not capable of performing what it has been usually represented to do. The dose is two or three drops or more.

#### OLEUM SASSAFRAS. ESSENTIALE. Essential oil of fassafras. L E.

This is the most ponderous of all the known effential oils, but rises in distillation with sufficient ease: it appears limpid as water, has a moderately pungent taste, a very fragrant smeel, exactly resembling that of the saffasras. It stands greatly commended as a sudoristic, and for purifying the blood and juices: it is likewise supposed to be of service in humoural asthmas and coughs. The dose is from one drop to eight or ten; though Geoffrey goes as far as twenty.

The decoction remaining after the diffillation of the oil, affords by inspifiation (see chap. vi.) an useful extract, of a mild bitterish, subastringent taste. Hossman says, he has given it with great benefit, in doses of a scruple, as a corroborant in cachecitic cases, in the decline of intermitting severs, and for abating hypochondriacal spasses.

## OLEUM TEREBINTHINÆ. Oil of turpentine. L. E.

This is distilled in the same manner as the foregoing oils: and is strictly an effential one, though not usually ranked in this class: it is commonly, but improperly as the college observe, called spirit of turpentine. It is employed in large quantities for some mechanic purposes and hence the distillation of it is become a particular business.

This oil is a very hot stimulating medicine (see page 239.) It is sometimes given as a sudorific and diuretic, in the dose of two or three drops: in larger doses, it is apt to greatly heat the body, occasion pain of the head, and effufion of the femen and liquor of the prostrate glands. It has nevertheless been of late taken in considerable doses (along with honey or other convenient vehicles) against the sciatica; and, as is said, with good fuccess. Some have recommended it against venereal runnings; but here it has produced misehievous consequences, inflaming the parts and aggravating the disorder. Externally it is not unfrequently employed against rheumatic pains, aches, sprains, for discusting cold tumours, and restraining hemorrhages.

After the distillation of the turpentine, there remains in the still a brittle resinous substance, of a yellow colour, called refina flava yellow resin [L]

The only use of this is in external applications, for giving confistence to plasters, and the like purposes.

Most of the foregoing oils are drawn by our chemits, and easily procurable in a tolerable degree of perfection; those of cinnamon, cloves, nutmegs and mace, excepted. These are usually imported from abroad; and are for the most part so much adulterated, that it is difficult to meet with such as are at all fit for use.

Nor are the adulteration of these kinds of preparations easily discoverable. The grosser abuses indeed may be readily detected;

thus

thus if the oil is mixed with spirit of wine, it will turn milky on the addition of water; if with expressed oils, rectified spirit will dissolve the effential, and leave the other behind, if with oil of turpentine, on dipping a piece of paper in the mixture, and drying it with a gentle heat, the turpentine will be betrayed by its sinell. But the more subtile arrists have contrived other methods of sophistication, which elude all trials of this kind.

Some have looked upon the specific gravity of oils as a certain criterion of their genuineness; and accordingly we have given a table of the gravity of several in page 38. This however is not to be absolutely depended on; for the genume oils, obtained from the same fubject, oftentimes differ in gravity as much as those drawn from different ones. Cinnamon and cloves, whose oils nsually fink in water, yield, if flowly and warily diftilled, an oil of great fragrancy, which is nevertheless specifically lighter than the aqueous fluid employed in the distillation of it; whilst on the other hand, the last runnings of fome of the lighter oils prove fometimes fo ponderous as to fink in water.

As all effential oils agree in the general properties of folubility in spirit of wine, indissolubility in water, missibility with water by the intervention of certain intermedia. volatility in the heat of boiling water, &c. it is plain that they may be variously mixed with one another, or the dearer sophisticated with the cheaper, without any poifibility of discovering the abuse by any trials of this kind. And indeed, it would not be of much advantage to the purchaser, if he had infallible criteria of the genuineness of every individual oil. It is of as much importance, that they be good, as that they be genuine; for I have often feen genuine oils, from incurious diffillation, and long and careless keeping, weaker both in smell and tafte than the common

fophliticated ones.

The smell and taste seem to be the only certain tests that the nature of the thing will admit of. a bark should have in every respect the appearance of good cinnamon, and should be proved indisputably to be the genuine bark of the cinnamon tree; yet, if it wants the cinnamon flavour, or has it but in a low degree, we reject it: and the case is the same with the oil. is only from use and habit, or comparison with specimens of knowa quality, that we can judge of the goodness, either of the drugs themfelves, or of their oils.

Most of the effential oils, indeed, are too hot and pungent to be tasted with safety; and the smell of the subject is so much concentrated in them, that a small variation in this respect is not easily distinguished. But we can readily dilute them to any affignable degree. A drop of the oil may be diffolved in spirit of wine; or received on a bit of sugar, and dissolved by that intermedium in water. The quantity of liquor, which it thus impregnates with its flavour, or the degree of flavour which it communicates to a certain determinate quantity, will be the measure of the degree of goodness of the oil.

I shall here subjoin some experiments, of the quantity of effential oil obtained from different vegetables, reduced into the form of a table. The first column contains the names of the respective vegetable substances, the second the quantity of each which was submitted to the distillation, and the third the quantity of oil obtained. In every other part of this book,

w bere

3

ŀ

•

÷

.

1

f

1

5

ø

where pound weights are mentioned, the troy pound of twelve ounces is meant; but these experiments having been all made by a pound of fixteen ounces, it was thought expedient to fet down the matter of fact in the original weights; especially as the several materials, in the large quantity commonly required for the distillation of oils, are purchased by weights of the same kind. But to remove any ambiguity which might arise from hence, and enable the reader to judge more readily of the yield, a reduction of the weights is given in the next column; which shews the number of parts of each of the subjects, from which one part of oil was obtained. To each article is affixed the author's name from whom the experiment is taken; those to which no name is added, are experiments of my own. different distillations of one subject, feveral of which are inferted in the table, shew how variable the yield of oil is, and that the exotic spices, as well as our indigenous plants, do not always contain the fame proportion of this active principle. though it must be observed, also that part of the differences may probably arise from the operation itself having been more or less carefully performed.

Table of the quantity of effential oil obtained from different vegetables.

			3		•			
Agallochum v	700d	-	10 lb.]	) .	6 4 dra. 7	1 !	320	Hoff.
Angelica root		-	ılb.		t dra.		128	Carth.
Anifeed -		-	ı lb.		4 dra.		32	Neum.
Anifeed -		-	3 lb.		j oun.		48	
Anifeed		-	4 lb.		ı oun.		64	
Asafetida		-	4 oun.	٠.	ı dra.		32	Neum.
Calamus aron	naticus -	<b>-</b>	50 lb.		2 OUB.	E	185	Hoff.
Calamus aron	naticus -	-	ı lb.		2 scrup.	from	192	Neum.
Caraway feed	s -	-	4 lb.		2 oun.		. 32	
Caraway feed	<b>s</b> -	•	2 lb.	ŀ	9 dra.	2.	28½	
Caraway seed	ş <b>-</b>	•	I Cwt.	oil	83 oun	obtained	211	
Carline thiftle		-	ı lb.	=	2 fcrup.		153	Neum.
Cardamom se	eds -	-	ı oun.	1	ı scrup.	2.8	24	Neum.
Carrot feeds	- •	-	2 lb.	effential	ı i dra.	₿	171	
Cascarilla		-	ılb.		'ı dra.	- <del>-</del> 5	128	
Camomile flow	vers -	-	л lb.	jo	30 gra.	Jo	256	Carth.
Common came		-	6 lb.	yielded	5 dra.	ر د د	153	v
Wild camomil		-	ılb.	Ŗ	20 gra.	part	384	Carth.
Wild camomi		-	6 lb.	Ž.	2½ dra.	1	307	
Chervil leaves	, fresh -	-	9 lb.		30 gra.	one	2304	Neum.
Cedar wood		-	ı lb.		2 dra.	=	64	Margg.
Cinnamon	<b>-</b> · <b>-</b>	-	ı lb		ı dra.	that	128	Sala.
Cinnamon		-	ı lb.		2½ scrup.	o l	153	Neum.
Cinnamon		-	4 lb.		6 dra.		853	Lemery.
Cinnamon	• •	-	ı lb.		2 dra.		64	Carth.
Cinnamon		-	ı lb.		8 fcrup.		451	
Clary feeds		•	4 lb.		2 dra.		256	
Clary in flowe	r, fresh	-	130 lb.		3 <del>½</del> oun.		594	-
Cloves -	• •	-	ı lb. j		i doun.	. 1		Teichm.
					_		•	Cloves

368 Pharma	ceutical	Pref	parations.	•	F	art II
Cloves	-  1 lb."	) 1	2 t oun.	. 1	7#	Carth
Cloves -	2 lb.	1 1	5 oun.		63	Hoff.
Copaiba balfam	ı lb.	1 1	боuв.		$2\frac{2}{3}$	Hoff.
Copaiba balfam	ı lb.	r 1	8 oun.		2	]
Cummin feed	. I bush.	1 (	21 oun.	1		1
Dictamnus Creticus .	ı lb.	1 1	30 gra.		256	Ì
Dill feed	4 lb	1 1	2 oun.		32	
Elecampane root	2 lb.	1 1	3½ scrup.		245	Neum
Elemi -	ı lb.	1 1	I oun.		iś	Neum
Fennel feed, common .	2 oun	1 1	ı scrup.		48	Neum
Fennel feed, fweet .	l bush.	ll	18 oun.		'	
Galangal root -	ı lb.	1 1	ı dra.		128	Carth
Garlick root, fresh	2 lb.	1 1	30 gra.		256	
Ginger -	. ⊿ lb.	1 1	I dra.		128	
Horseradish root, fresh	Soun.	lI	15 gra.			Neum
Hyffop leaves	2 lb.	1 1	1 dra.		237	Neum
Hyssop leaves -	ı lb.	1 1	1 dra.		85	Carth
Hystop leaves	ı lb.	1 1	2 dra.	1	64	Carth
Hystop leaves, fresh	2 cwt	1 l	6 oun.	2	597	
Hyssop leaves, fresh	10 lb.	1 1	3 dra.	from	427	1
Hyssop leaves, fresh	30 lb	1 1	9 dra.	9 [	427	1
Juniper berries -	8 lb.	<del> </del>	3 oun.	obtained	427	Hoff.
Juniper berries -	lb.		3 dra.	<u>.</u>		Carth
Lavender in flower, fresh	48 lb.	2.	12 oun.	ڄ	64	
Lavender in flower, fresh	30 lb.	1.5	64 oun.	S	72	
Lavender in flower, fresh	134cwt	effential	60 oun.	was	403	
Lavender flowers, fresh	2 lb.	ام	4 dra.	lio j	64	Hoff.
Lavender flowers, dried	4 lb.		2 oun.	ojo	32	٠ ريوت
Lavender flowers, dried	2 lb.	yielded	1 oun.	S I	32	Hoff.
Lavender flowers, dried	4 lb.	I 후 I	' 3 oun.	part	217	Hoff.
Broad-leaved lavender ?	4 lb.	^	ı oun.	-	64	Hoff.
flowers, dry	ı lb	1 1	2 dra.	one	64	
Lovage root -	lb.	1 (	ı dra.	=	128	Carth
Mace -	ı lb.	1 1	5 dra.	that	253	
Mace -	. Ib.	i I	6 dra.	ی	211	Certh
Marjoram in flower, fresh		1 1	3‡ oun.	_	347	
Marjoram in flower, fresh	134 lb.	1 1	3 <sup>1</sup> / <sub>2</sub> dra.		493	•
:	34 lb.	1 1	ı i oun.		362	
Marjoram leaves, fresh	. 18¥ lb.	1 1	4 dra.	1	592	
Marjoram leaves, dried	4 lb.	] !	ı oun.			Hoff.
Masterwort root	. ib.	1	30 gra.		256	Neum
1 · 1	14 lb.	1 1	4 dra.	[	448	
	6 lb.	1 1	4 dra.	l	177	l
Mint leaves, dried	4 lb.	1 (	ı don.	l		Hoff.
Peppermint, fresh	4 lb.		3 dra.		1703	
Myrrh -	. ib.	i l	2 dra.		64	Hoff.
- · ·	T	1	3 dra.		427	Neun
Nutmegs -	i lb.	1 1	i oun,		1 4-1	Hoff.
Nutmegs -	1 lb. 1 lb.	! !	ı oun.			Geoff.
Nutmegs -	ı lb.	1 1	4 dra.	i	22	Neum
Nutmegs :	ı lb.	l l	6 dra.		31	Sala.
- · · · · · · ·	,	, (	,	, ,	L ~ · 5	Nutme

i

1

.

Chap. 5		369		
Nutmegs	-  i lb.	) j 5 dra. 1	<b>1</b> [25	3 Carth.
Parsley seeds -	- 2 lb.	ı dra.	25	6
Parsley leaves, fresh	-238 lb.	2 oun.	1 10	04
Parnfnep feeds -	- 8 lb.	2 dra.	51	
Penny-royal in flower, fresh	- 13 lb.	6 dra.	27	
Black pepper -	- 2 lb.	6 dra.	الما	23
Black pepper -	-l 1 lb.	24 dra.	1 1 '	82 Neum
Black pepper -	- I lb.	4 scrup.		of Carth.
Black pepper -	- 1 lb.	i dra.	1	28 Heister
Black pepper -	- 6 lb.	3 dra.	2	56 Geoff.
Pimento -	Loun.	30 gra.		16 Neum.
Rhodium wood -	- 1 lb.	3 dra.	4	23 Neum.
Rhodium wood -	- 1 lb.	2 dra.	6	4 Sala.
. Rhodium wood -	- 1 lb.	3 dra.	4	22 \ala.
Rhodium wood -	- i jp.	3 cra.	4	23 Carth.
Rhodium wood	- ilb.	4 dra.	_	2 Carth.
Rofemary in flower	- I CWt.	8 oun.	2	24
Rosemary leaves -	- I b.	2 dra.	6	4 Sala.
Rosemary leaves -	- I lb.	3 dra.	<b>5</b> 4	23 Sala.
Rosemary leaves -	- 3 lb.	36 dra.	LE 12	- i - ·
Rosemary leaves -	i lb.	ı dra.	-p 12	
Rosemary leaves -	- I lb.	ਰ ਾਡੇ dra.		3 Cartb.
Rosemary leaves, fresh	o lb.	I I Soun. I	22	
Rofes -	-100 lb.	ੂੰ 4 dra.	0 32	00 Tachen.
Rofes -	- 100 lb.	4 dra. 1 oun. 30 gra.		Oo Homb.
Rofes -	10 lb.	30 gra.	} ₹ 76	
Rue -	10 lb.	5 2 dra.	ā 64	
Rue -	4 lb.	9 4 dra. 1 dra. 2 dra.	5 3 <sup>2</sup>	o Hoff.
Rue in flower -	60 lb.	I dra.	21 20 20 20 20 20 20 20 20 20 20 20 20 20	
Rue in flower -	72 lb.	2 oun.	E 50	
Rue with the seeds Saffron -	i lb.	1 dra.	10 that one 28 24 24 24 24 24 24 24 24 24 24 24 24 24	
	lb.	5 fcrup.	85	- 10
Sage leaves - Sage in flower, fresh	- 34 lb.	2½ oun	E 7	′ I
Sage of virtue in flower	27 lb.	6 dra.	□   54 ©   57	
Sage of virtue in flower	. 8 lb.	2 dra.	68	, I
Saffafras -	- 6 lb.	ı∄ oun.	5	177 ~
Sassafras -	- 6 lb.	2 oun.		8 Neum.
Savin -	2 lb.	5 oun.		6º Hoff.
Saunders, yellow	_ Ilb.	2 dra.		4 Carib.
Smallage feeds	_ I lb.	21 scrup.	15	
Stechas in flower, fresh	- 53 lb.	2 dra.	36	т (
Thyme in flower, fresh	- 2 CW .	ς≟ oun.	65	
Thyme in flower, dry	- 3 tb.	i dra.	29	
Lemon thyme in flower, fresh	-  <1 lb.	ı¼ oun.	65	
Lemon thyme in flower, tresh	- 98 lb.	zi oun.	62	
Lemon thyme dried a little	-1104 lb.	3 oun.	55	
Wormwood leaves, dry	- 4 lb.	i oun.		4
Wormwood leaves, dry	- 18 lb.	ı½ oun.	119	2
Wormwood leaves, dry	- 25 ib.	3½ oun.	11	4
Zedoary _	- i lb.	i i dra.	į 12	8 Neum.
_	Вь	-	•	SECT.

日本時間 在日本的時 即以日本日本日

THE PROPERTY OF THE PARTY OF TH

9

-

quor that runs from the nose of the worm, the distillation is to be

ftont.

This is the method of diftillation commonly practifed for the officinal waters. It is accompanied with one imperfection, affecting chiefly those waters, whose principal value consists in the delicacy of their flavour; this being not a little injured by the boiling heat usually employed, and by the coagitation of the odorous particles of the subject with the water. Sometimes also a part of the plant sticks to the sides of the still, and is so far scorched as to give an ungrateful taint to the liquor.

THERE is another method of managing this operation, which I have already recommended for the distillation of the more volatile esfential oils, and which is equally applicable to that of the waters. In this method, the advantages of the foregoing ones are united, and their inconveniencies obviated. A quantity of water being poured into the still, and the herbs or flowers placed in a basket over it, there can be no possibility of burning: the water may be made to boil. but so as not to rise up into the basket, which would defeat the intention of this contrivance. hot vapour of the water passing lightly through all the interftices of the subject matter, imbibes and carries over the volatile parts unaltered in their native flavour. this means the distilled waters of all those substances, whose oils are of the more volatile kind, are obtained in the utmost perfection, and with sufficient dispatch; for which last intention the still may be filled quite up to the head.

In the distillation of effential oils, the water, as observed in the

foregoing section, imbibes always a part of the oil. The distilled liquors, here treated of, are no other than water thus impregnated with the effential oil of the fubject; whatever fmell, taste, or virtue, is here communicated to water, or obtained in the form of a watery liquor, being found in a concentrated The effential oil, state in the oil. or some part of it, more attenuated and subtilized than the rest, is the direct principle, on which the title of spiritus rector, or prefiding spirit, has been bestowed.

All those vegetables therefore which contain an effential oil, will give over some virtue to water by distillation : but the degree of the impregnation of the water, or the quantity of water which a plant is capable of fatiating with its virtue, are by no means in proportion to the quantity of its oil. The oil fatiates only the water that comes over at the same time with it: if there is more oil than is sufficient for this fatiation, the furplus feparates, and concretes in its proper form, not miscible with the water that arifes afterwards. Some odoriferous flowers, whose oil is in fo little quantity, that scarcely any visible mark of it appears, unless fifty or an hundred pounds or more are distilled at once, give nevertheless as strong an impregnation to water, as those plants which abound most with oil.

MANY have been of opinion, that distilled waters may be more and more impregnated with the virtues of the subject, and their strength increased to any assigned degree, by cobobation, that is, by redstilling them a number of times from fresh parcels of the plant. Experience, however, shews the contrary; a water skilfully drawn in the first distillation, proves, on every repeated

Ľ

ģ

è

g;

Ė

;¢

ŗ

ſ.

ø

\$

1

1

į.

peated one, not stronger, but more Aqueous liquors are disagreeable. not capable of imbibing above a certain quantity of the volatile oil of vegetables, and this they may be made to take up by one, as well as by any number of distillations, the oftener the process is repeated, the ungrateful impression which they generally receive from the fire, even at the first time, becomes greater and greater. plants which do not yield at first waters fufficiently strong, are not proper subjects for this process, fince their virtue may be obtained much more advantageously by Others.

General rules for the distillation of the officinal simple Waters.

Ī.

Plants and their parts ought to be

fresh gathered [E.]

Where they are directed fresh, such only must be employed; but some are allowed to be used dry, as being easily procurable in this state at all times of the year, though rather more elegant waters might be obtained from them whilst green [L.]

Having bruised the subject a little, pour thereon thrice its quantity of spring water: this quantity is to be diminished or increased, according as the plants are more or less juicy than ordinary [E.]

When fresh and juicy herbs are to be distilled, thrice their weight of water will be fully sufficient: but dry ones require a much larger quantity. In general there should be so much water, that after all intended to be distilled has come over, there may be liquor enough left to prevent the matter from burning to the still.

m.

The distillation may be performed in an alembic with a refrigeratory, the junctures beingluted [E.]

IV.

The diffillation is to be continued as long as the water which comes over is perceived to have any fmell or tafte of the plant [E.]

Plants differ so much according to the soil and season of which they are the produce, and likewise according to their own age, that it is impossible to fix the quantity of water to be drawn from a certain weight of them, to any invariable standard. The distillation may always be continued as long as the liquor runs well slavoured of the subject, and no longer.

If the herbs are of prime goodness, they must be taken in the weights prescribed. But when fresh ones are substituted to dry, or when the plants themselves are the produce of unsavourable seasons, and weaker than ordinary; the quantities are to be varied according to the discretion of the

artist [L.]

After the odorous water, alone intended for use, has come over, an acidulous liquor arises, which has sometimes extracted so much from the copper head of the still, as to prove emetic. To this are owing the anthelmintic virtues attributed to certain distilled waters.

In the preceding edition of the Edinburgh pharmacopæia, some vegetables were ordered to be slightly fermented with the addition of yeast, previously to the distillation.

The principle on which this management is founded, is cer-Bb 3 tainly tainly just; for the fermentation somewhat opens and unlocks their texture, so as to make them part with more in the subsequent distillation than could be drawn over from them without some assistance of this kind. Those plants, however, which require this treatment, are not proper subjects for simple waters to be drawn from; their virtues being obtainable to better advantage by other processes.

#### VI.

If any drops of oil swim on the surface of the water, they are to be carefully taken off [E.]

#### VII.

That the waters may keep the better, about one-twentieth part their weight of proof spirit may be added to each, after they are distilled [L.]

A great number of distilled waters were formerly kept in the shops, and are still retained in foreign pharmacopoias. The faculty of Paris direct, in the last edition of their codex medicamentarius, no less than one hundred and twenty-five different waters, and one hundred and thirty different ingredients in one single water. Near one half of these preparations have scarcely any virtue or slavour from the subject, and many of the others are insignificant.

The colleges of London and Edinburgh have rejected these oftentatious superfluities; and given an elegant and compendious set of waters, sufficient for answering such purposes as these kinds of preparations are applied to in practice. Distilled waters are employed chiefly as grateful diluents, as suitable vehicles for medicines of greater efficacy, or for rendering disgussful ones more acceptable to the palate

and flomach; few are depended on in any intentions of consequence, by themselves.

#### AQUA ALEXETERIA SIMPLEX.

Simple alexeterial water. Lond.

Take of

Spearmint leaves, fresh, a pound and a half;

Sea wormwood tops, fresh; Angelica leaves, fresh, each one pound;

Water, as much as is fufficient to prevent an empyreuma. Draw off by diffillation three gallons.

#### Edinb.

Take of

Elder flowers, moderately dried, two pounds;

Angelica leaves, fresh gathered, one pound;

Water, a sufficient quantity. Distil off three gallons.

THESE waters are sufficiently elegant with regard to taste and smell; though sew expect from them such virtues as their title seems to imply. They are used occasionally for vehicles of alexipharmac medicines, or in juleps to be drank after them, as coinciding with the intention; but in general are not supposed to be themselves of any considerable efficacy.

## AQUA SEMINUM ANETHI. Dill feed water Lond.

Take of

Dill feeds, a pound and a half; Water, as much as is sufficient to prevent an empyreums. Draw off by distillation one gallon.

This water, which turns out pretty strong of the dill feeds, is fometimes ۲.

ż

\*

11

£

Ł

1

Ė

f

fometimes employed as the basis of carminative juleps. It is similar in flavour to a water drawn from caraway seds; but less agreeable.

### AQUA ANGELICE. Angelica water.

Take of

Angelica leaves, fresh, any quan-

Water, three times as much.

Diftil as long as the liquor runs
well flavoured of the plant.

THIS water is among us very rerely made use of. It smells and tastes considerably of the angelica, but does not prove so agreeable, as might be expected.

### AQUA ARTEMISIA. Mugwort Water.

Take of

Mugwort leaves, fresh, as much as you please;

Water, a fufficient quantity; Yeaft, a little.

Let them fland together in a warm place, till they begin to ferment, and then diftil according to art.

MWGWORT water has been held by many in great efteem as an uterine, but the herb itfelf has little title to that class, and the distilled water none at all It is at present scarce ever called for, or kept in the shops.

# AQUA CORTICUM AURANTIORUM SIMPLEX. Simple orange peel water. Lond.

Take of

Yellow peel of Seville oranges, dried, four oances; Water, as much as is sufficient to prevent burning.

Distil off one gallon.

THIS water proves very weak of the orange peel. It is defigned for a diluter, in fevers, and other diforders where the flomach and palate are subject to receive quick difgust; in which cases (as the committee observe) cordial waters, especially if their use is to be long continued ought to be but lightly impregnated with any flavour, however agreeable.

### Aqua cardui Bentdicti. Carduis water.

This is prepared from the leaves of cardnus benedictus, after the fame manner as the aqua artemifia.

This water has been looked upon as a fudorific and alexipharmac; and in this intention is fill frequently preferibed by foreign physicians, in juleps and draughts. Among us, it has long been difused, and held intirely infignificant; this plant, however opened by fermentation, giving nothing valuable over the helm. The decoction, which remains after the diftillation, duly depurated and infpiffated, proves a medicine of some use; it is a moderately strong bitter, similar to the extract of cardaus; see chap. vi. in keeping, a considerable quantity of effential salt sometimes shoots in it.

## AQUA CASTOREI. Cafter water. Lond.

Take of

Russia castor, one ounce; Water, as much as will prevent burning.

Draw off two pints.

CASTOR yields almost all its slavour in distillation to water; but treated in the same manner with spirit of wine, gives over nothing.

B b 4

The spirit of castor formerly kept in. the shops, had none of the smell or virtues of the drug; whilst the water, here directed proves, when fresh drawn, very strong of it.

It is remarkable, that the virtues of this animal substance reside in a volatile oil, analogous to the effential oils of vegetables: some are reported to have obtained, in distilling large quantities of the drug, a small portion of oil, which smelt extremely strong of the castor, and diffused its ungrateful scent to a great distance.

This water is made use of in hysteric cases, and some nervous complaints, though it has not been found to answer what many people expect from it: it loses greatly of its flavour in keeping.

AQUACERASORUM NIGRORUM Black cherry water.

Let any quantity of black cherries be bruised, so as that the stones may be broken; and then diftilled according to art, with only a small portion of water.

This is a very grateful water, and has long maintained a place in the shops. It has frequently been employed by physicians as a vehicle, in preference to the other diftilled waters: and among nurses. and others who have the care of young children, has been the first remedy against the convulsive disorders to which children are so often subject.

This water has nevertheless of late been brought into disrepute, and by some 'looked upon as poifonous. They observe that it receives its flavour principally from the cherry stones; and that these kernels, like many others, bear a resemblance in taste to the leaves of the lauro-cerasus, which have some time past been discovered to

yield, by infusion or distillation. the most sudden poison known: some physicians of Worcester have lately found, by trial purpofely made, that a diffilled water very strongly impregnated with the flayour of the cherry kernels (no more than two pints being distilled from fourteen pounds of the cherry stones) proved in like manner poifonous to brutes: the committee of the London college repeated the fame experiment, and found the effects agreeable to those gentle-

men's report.

It by no means follows from these trials, nor after such long experience can it be imagined, that black cherry water, when no ftronger than the shops have been accustomed to prepare it, is unsafe. These kernels, as the committee observe, plainly resemble opium, and some other things, which poifon only when taken in too great a quantity; the water from the very laurel leaves is harmless when duly diluted; and even spirit of wine proves a poison of a kind not greatly different, if drank to a certain degree of excess. Nor can it be concluded, from the trials with the strong black cherry water on dogs, &c. that even this will have the same effects in the human body; the kernels of many forts of fruits being in fubitance poisonous to brutes, though innocent to man.

It is possible, however that this water in any degree of strength may not be altogether fafe to the tender age of infants, where the principles of life are but just beginning as it were to move: it is possible, that it may there have had pernicious effects, without being sufpected; the symptoms it would produce, if it should prove hurtful, being fuch as children are often thrown into from the difeafe

which

:

Ē

ø

ľ.

3

ď

5

3

which it is imagined to relieve. On these considerations, both the London and Edinburgh colleges have chosen to lay it aside; more especially as it has been too often counterseited with a water distilled from bitter almonds, which are known to communicate a poisonous quality.

### AQUA CINNAMOMI SIMPLEX.

Simple cinnamon water, Lond.

Take of
Cinnamon, one pound;
Water, as much as will prevent
burning.
Diftil off one gallon.

### AQUA CINNAMOMI SINE VINO.

Cinnamon water without wine.

Edinb.

Take of

Cinnamon, one pound;
Water, a gallon and a half.
Steep them together for two days;
and then diffil off the water, till
it ceases to run milky.

This is a very grateful and useful water, possessing in an emineat degree the fragrance and aromatic cordial virtues of the fpice. Great care should be had, in the choice of the cinnamon, to avoid the too common impolition of casia being substituted in its room: this latter yields a water much less agreeable than that of cinnamon, and whose flavour is manifeftly empyreumatic. two drugs may be easily diffinguished from one another by the marks laid down under the respective articles in the second part of this work: see page 121 and 127.

The virtues of all these waters depend upon their containing a

portion of the oil of the subject. The oil of cinnamon is very ponderous, and arises more difficultly than that of any of the other vegetable matters from which fimple waters are ordered to be This observation directs drawn. us, in the distillation of this water, to make use of a quick fire, and a low veffel. For the fame reason, the water does not keep so well as might be wishthe ponderous oil parting from it in time, and falling to the bottom, when the liquor loses its milky hue, its fragrant fmell, and aromatic tafte. Some recommend a finall proportion of fugar to be added, in order to keep the oil united with the water.

## AQUA CHAMÆMELI. Chamomile water Edinb.

Take any quantity of chamomile flowers, and so much water as will prevent burning. Distil off the water so long as it proves sufficiently strong of the flavour of the flowers.

Chamomile flowers were ordered in the former editions to be fermented previously to the distillation, a treatment which they stand little in need of: for they give over, without any fermentation, as much as that process is capable of enabling them to do. In either case, the smell and peculiar flavour of the flowers arise, without any thing of the bitterness; this remaining behind in the decoction; which, if duly depurated and inspissated, yields an extract fimilar to that prepared from the flowers in the common manner. . The distilled water has been used in flatulent colics, and the like,

٠.

but is at present held in no great effeem.

## AQUA FOENICULI. Fennel water Land.

Take of

Sweet fennel feeds, one pound; Water, as much as is sufficient to prevent an empyreuma. Distilloss one gallon.

Edinb.

Take of

Feanel leaves, fresh, any quantity;

Water, three times as much.

Diftil as long as the water runs well flavoured.

THE first of these waters is a fufficiently grateful one, and the other is not unpleasant; the leaves should be taken before the plant has run into flower; for after this time, they are much weaker and less agreeable. Some have observed, that the upper leaves and tops, before the flowers appear, yield a more elegant water, and a remarkably finer effential oil, than the lower ones; and that the oil obtained from the one swims on water, whilst that of the other finks. No part of the herb, however, is equal in flavour to the feeds.

## AQUA HYSSOPI. Hyffop water. Edinb.

This is distilled from the fresh leaves of hysfop, after the same manner as the water of fennel leaves.

Hyssor waterhas been held by fome in confiderable efteem as an uterine and a pectoral medicine. It was directed in the last edition of the Edinburgh Pharmacopæia,

for making up the black pectoral troches, but is now exchanged for common water. Few at prefent expect any fingular virtues from it, nor is it often made use of, or met with in the shops.

## AQUA MELISSÆ. Balm water. Edinb.

This is prepared by distilling the green leaves of balm, as in the foregoing process.

In the former editions of the Edinburgh pharmacopæia, this water was ordered to be cohobated, or rediftilled from fresh quantities of the herb. This management feems to have been taken from Boerhaave, who has a very high opinion of the water thus prepared; he fays, he has experienced in himself, extraordinary effects from it taken on an empty flomach; that it has scarce its equal in hypochondriacal and hyflerical cases, the chlorosis, and palpitation of the heart as often as these diseases proceed from a disorder of the spirits rather than from any collection of morbific matter.

For my own part, I have already given my opinion with regard to the cohobation of thefe liquors; and shall here only obferve, that whatever virtues are lodged in balm, they may be much more perfectly and advantageously extracted by cold infusion in aqueous of spirituous menstrua; in this process, the liquor saffers no injury from being returned on fresh parcels of the herb; a few repetitions will load it with the virtues of the subject, and render it very rich. (See page 269.) The impregnation here is almost wnlimited; but in distilled waters, it is far otherwise.

AQUA

#### AQUA MENTHÆ.

Mint mater. Edinb.

Take of

ż

.

Spearmint leaves, freih, any quantity;

Water, three times as much. Diftil as long as the liquor which

Diffil as long as the liquor which comes over has any tafte or fmell of the mint.

#### AQUA MENTHÆ VULGARIS SIMPLEX.

Simple spearmint water.

Lond.

Take of

Spearmint leaves, dried, a pound and a half;

Water, as much as is sufficient to prevent burning.

Draw off by distillation one gallon.

THESE waters smell and taste very strong of the mint; and prove in many cases useful stomachics. Boerhaave commends them (cohobated) as a present and incomparable remedy, for strengthening a weak stomach, and curing vomiting proceeding from cold viscous phlegm; as also in lienteries.

# AQUA MENTHÆ PIPERITIDIS SIMPLEX. Simple pepper-mint water. L. E.

Take of

Pepper-mint leaves, dry, a pound and a half;

Water, as much as will prevent an empyreuma.

Draw off by diffillation one gallon.

This is a very elegant and

This is a very elegant and useful water: it has a warm, pungent taste, exactly resembling that of the pepper-mint itself. A spoonful or two, taken at a time, warm the stomach, and give great

relief in cold, flatulent colics. Some have substituted a plain insusion of the dried leaves of the plant, which is not greatly different in virtue from the distilled water.

### AQUA PETROSELINI. Parsley water.

This is distilled from the fresh leaves of parsley, after the same manner as the aqua mentha.

This water is fcarce ever called for, or kept in the shops. Parsley yields little virtue in distillation; and the leaves are not the part that yield most. The seeds give a considerable share of slavour, which is not disagreeable.

#### AQUA PIPERIS JAMAICENCIS.

Water of Jamaica pepper.

Lond.

Take of

Jamaica pepper, half a pound; Water, as much as will prevent burning.

Diftil off one gallon.

This diffilled water is a very elegant one, and has of late come pretty much into use: the hospitals employ it as a succedaneum to the more costly spice waters. It is, however, inserior in gratefulness to the spirituous water of the same spice hereaster directed.

## AQUA PULEGII SIMPLEX. Simple penny-royal water. -Cond.

Take of

Penny-royal leaves, dry, a pound and a half:

Water, as much as will prevent burning.

Draw off by distillation one gallon.

**AQUA** 

AQUA PULEGII.
Water of penny-royal,
Edinb.

Take of

Penny-royal leaves, fresh, any quantity;

Water, three times as much.

Diffil as long as the water comes
off well flavoured of the herb.

THESE waters possels, in a confiderable degree, the smell, taste, and virtues of the penny-royal. They are frequently taken in hysteric cases, and not without good effects.

AQUA ROSARUM DAMASCENARUM. Damask rose water. Lond.

Take of

Damaik roses, fresh gathered, six pounds; Water, as much as will keep

them from burning.

Distil off a gallon of the water.

Edinb.

Take three parts of water to one of the freih roses; and distil as long as the water which comes over has any smell of the flowers.

This water is principally valued on account of its fine flavour. which approaches to that generally admired in the rose itself. The purgative virtue of the roses remains entire in the liquor left in the still, which has therefore been generally employed for making the folutive honey, and fyrup, instead of a decoction or infusion of fresh roses prepared on purpose: and this piece of frugality the college have now admitted. A distilled water of red roses has been fometimes called for in the shops; and supplied by that of damask roses, diluted with common water: this is a very venial

fubilitation; for the water drawn from the red rose has no quality which that of the damask does not possess in a far superior degree; neither the purgative virtue of the one, nor the astringency of the other, arising in distillation.

AQUARUTE.
Rue water.

This is to be diffilled from the fresh leaves of rue, and cohobated on fresh parcels of them, after the same manner as the aqua melissæ.

Rue gives over in this process the whole of its sinell, and great part of its pungency. The distilled water stands recommended in epileptic cases, the hysteric passion, for promoting perspiration, and other natural secretions; see page 211.

AQUA SABINA. Savin water.

This is distilled from the fresh leaves of savin, after the same manner as aqua angelica.

This water is by some held in considerable esteem for the same purposes as the distilled oil of savin. Boerhaave relates, that he has sound it (when prepared by cohobation) to give an almost incredible motion to the whole nervous system, and that when properly used, it proves eminently serviceable for promoting the menses, and the hamorrhoidal slux.

AQUA SAMBUCI.

Elder flower water.

This is distilled from fresh elder flowers, after the same manner

as the aqua angelica.
This water imells confiderably
of the flowers; but is rarely made

of.

SECT.

Ľ.

: 5

1

I I

ï

į:

1

3

Í

#### S E C T. III.

#### Spirituous distilled waters and spirits.

HE flavour and virtues of distilled waters are owing, as observed in the preceding section, to their being impregnated with a portion of the essential oil of the subject from which they are drawn. Spirit of wine, considered as a vehicle for these oils, has this advantage above water, that it is their proper menstruum, and keeps all the oil, that rises with it, persectly dissolved into an

uniform limpid liquor.

Nevertheless, many substances, which, on being distilled with water, impart to it their virtues in great perfection; if treated in the same manner with spirit of wine, scarce give over to it any fmell or tafte. This difference proceeds from hence; that spirit is not susceptible of so great a degree of heat as water. Liquids in general, when made to boil, have received as great a heat as they are capable of fustaining: nrow, if the extent of heat between freezing and boiling water, as measured by thermometers, be taken for a standard, spirit of wine will be found to boil with less than four-fifths of that heat, or above one-fifth less than the heat of boiling water. It is obvious therefore, that substances may be volatile enough to rife with the heat of boiling water, but not with that of boiling spirit,

Thus, if cinnamon, for instance, be committed to distillation with a mixture of spirit of wine and water, or with a pure proof spirit, which is no other than a mixture of about equal parts of the two; the spirit will arise sirth, clear; colourless, and transparent, and

almost without any taste of the spice; but as soon as the more ponderous watery sluid begins to arise, the oil comes freely over with it, so as to render the liquor highly odorous, sapid, and of a milky hue.

The proof spirits usually met

with in the shops are accompanied with a degree of ill slavour; which, though concealed by means of certain additions, plainly discovers itself in distillation. This nauseous relish does not begin to arise, till after the purer spirituous part has come over; which is the very time that the virtues of the ingredients begin, also, most plentifully to distill; and hence the liquor receives an ungrateful taint. To this cause principally is ow-

diffiller, the latter being extremely curious in rectifying or purifying the spirits (when designed for what he calls sine goods) from all ill slavour.

SPIRITUS VINI

ing the general complaint, that

the cordials of the apothecary are less agreeable than those of the same kind prepared by the

RECTIFICATUS.

Rectified spirit of wine.

Edinb.

Take any quantity of French brandy, and with a very gentle heat distill it to one half.

This rectified spirit, being digested for two days with one-fourth its quantity of dry salt of tartar in powder, and then distilled in a glass cucurbit, with a very gentle heat, becomes AL-COHOL:

Spirits

Spirits distilled from malt liquors, or other fermented substances, after being rectified in the above method, require further purification; namely, repeated distillation from an equal quantity of spring water.

FRENCH brandy is rather too dear an article in this country, for distillation; nor is the spirit obtained from it any ways preferable to one procurable from cheaper liquors. The coarser inflammable spirits may be rendered perfectly pure, and sit for the nicest purposes, by the following method.

If the spirit is exceedingly foul, mix it with about an equal quantity of water, and distil with a flow fire; discontinuing the operation as foon as the liquor begins to run milky, and discovers, by its nauseous taste, that the impure and phlegmatic part is arising. By this treatment, the spirit leaves a considerable portion of its foul oily matter behind it in the water, which now appears milky and turbid, and proves highly difagreeable in taste. If the spirit was not very foul at first, this ablution is not necessary; if extremely fo, it will be needful to repeat it once, twice, or oftener.

As vinous spirits arise with a less degree of fire than watery liquors, we are hence directed to employ, in the distillation of them, a heat less than that in which water boils, and if due regard be had to this circumstance, very weak spirits may, by one or two wary distillations, be tolerably well freed from their aqueous phlegm; especially if the distilling vessels are of such a height, that the spirit, by the heat of a waterbath, may but just pass over them: in such case, the phlegmatic va-

pours, which arise for a little way along with the spirit, will condense and fall back again before they can come to the head. Very pompous instruments have been contrived for this purpofe, and carried in a fpiral or ferpentine form, to an extraordinary height. The spirit, ascending through The spirit, ascending through these, was to leave all the watery parts it contained, in its passage. and come over perfectly pure and free from phlegm. But these instruments are built, upon erroneous principles, their extravagant height defeating the end it was deligned to answer: if the liquor made to boil, a confiderable quantity of mere phlegm come over along with the spirit : and if the heat is not railed to this pitch, neither phlegm nor spirit The most convenient will distil. instrument is the common still, betwixt the body of which, and its head, an adopter or copper tube may be fixed.

The spirit being washed, as above directed, from its foul oil. and freed from the greatest part of the phlegm, by gentle diftillation in a water-bath; and to every gallon of it, a pound or two of pure, dry, fixt alkaline falt. Upon digesting these together for a little time, the alkali, from its known property of attracting water and oils, will imbibe the remaining phlegm, and fuch part of the disagreeable unctuous matter as may still be lest in the spirit, and sink with them to the bottom of the vessel. the spirit be now again gently drawn over, it will arise entirely free from its phlegm and nauseous flavour; but some particles of the alkaline salt are apt to be carried up with it, and give what the workmen call an urinous relish: this may be prevented by adding, previous

ŧ

•

ı

8

1

5

5

previous to the last distillation, a small proportion of calcined vitriol, allum, or sal catharticus amarus; the acid of these salts will unite with, and naturalize the alkali, and effectually prevent it from arising; while no more of the acid of the salts is extricated than what the alkali absorbs.

The spirit obtained by this means is extremely pure, limpid, perfectly slavourles, and sit for the finest purposes. It may be reduced to the strength commonly understood by proof, by mixing twenty ounces of it (by weight) with seventeen ounces of water. The distilled cordials made with these spirits, prove much more elegant and agreeable, than when the common recisited, or proof in the oils.

fpirits of the shops are made use of.

If the rectified spirit be distilled astresh from dry alkaline salt, with a quick sire, it brings over a considerable quantity of the salt, and in this state is supposed to be a more powerful menstruum, for certain substances, than the pure spirit. This alkalized spirit is called TARTARIZED SPIRIT OF

The general virtues of vinous spirits have been already mentioned in the preceding part: the spirits impregnated with the volatile oils of vegetables, to be treated of in this chapter, have joined to those, the aromatic, cordial, or other virtues which reside in the oils.

#### ARTICLE I. Distilled Spirits.

AQUA MELISSE COMPOSITA.

Compound balm water, commonly

called Eau de carmes.

Take of

Balm in flower, fresh gathered and cleared from the stalks, two pounds;

Lemon peel, fresh, as foon as pared from the fruit, four ounces:

Coriander feeds, eight ounces; Nutmegs,

Cloves,

Cinnamon, each, bruised, two ounces;

Angelica roots, dried and bruifed one ounce;

Spirit of wine, highly reclified, ten pints.

Steep the feveral ingredients in the spirit, four or five days; and then draw off; in the heat of a water bath, ten pints. Rectify the distilled liquor by a second distillation in a water bath, drawing off only about eight pints and three quarters.

This process is taken from the Elemens de pharmacis of M. Baume, who observes, that all the aromatic spirit ought to be prepared in the same manner. When the common spirits of this kind are rubbed on the hands, &c. they leave, after the more volatile parts have exhaled, a difagreeable empyreumatic fmell; and when diluted with water, and taken medicinally; they leave in like manner a naufeous flavour in the mouth. To remedy these impersections, he made many experiments, which shewed, that in order to obtain these liquors of the defireable qualities, the spirit must not only be perfectly pure at first, but that the liquor ought also to be rectified after it has been distilled from the subjects. In this rectification, only the more volatile, subtile, aromatic parts of the ingredients arise: there remains behind a white liquor, acrid, bitter, loaded only with the groffer

384

oil, and deprived of all the specific flavour of the subjects. Indeed the very imperfection complained of naturally points out this second distillation for the remedy; as it shews the spirit to contain a grateful and ungrateful matter, the first of which exhales while the other is left behind. The author fays, that when the aqua melissa is prepared as above directed, it has something in it more perfect, than any of the odoriferous spirits whose excellence is cried up, and which have the reputation of being the beft.

Aromatic spirituous waters have in general less smell, when newly distilled, than after they have been kept about fix months: M. Baume suspects that the preparations of this kind, which have been most in vogue, were such as had been thus improved by keeping; and found that the good effects of age might be produced in a short time by means of cold. He plunges quart bottles of the liquor into a mixture of pounded ice and fea falt : the spirit, after having suffered, for fix or eight hours, the cold hence refulting, proves as grateful as that which has been kept for feveral years. Simple waters also, after being frozen, prove far more agreeable than they were before, though they are always less so than those, which have been drawn with spirit, and exposed to a like degree of cold. This melioration of distilled waters by frost was taken notice of by Geoffroy, Hift. Acad. 1713.

SPIRITUS RORISMARINI.

Spirit of rosemary.

Anna

Take of Rolemary tops, fresh gathered, a pound and a half; Proof spirit, one gallon.

Distil in the heat of a water-bath, till five pints are come over.

SPIRITUS RORISMARINI, vulgo AQUA REGINÆ HUNGARIÆ.

Hungary water [E.]

Take of

Rosemary flowers, just gathered, two pounds;

Rectified spirit of wine, one gallon.

Put them together, and immediately diffil in a water-bath.

It is generally brought to us from abroad.

THIS spirit is very fragrant, infomuch as to be in common ufe as a perfume: that brought from abroad is superior in fragrance to fuch as is generally made among In order to prepare it in perfection, the vinous spirit should be extremely pure; the rolemary tops gathered when the flowers are full blown upon them, and committed immediately to diffillation, particular care being taken not to bruise or press them. The best method of managing the distillation, is that formerly recommended for the distillation of the more volatile essential oils and simple waters, viz. first to place the spirit in the still, and then fet in, above the liquor, either an iron hoop, with a haircloth stretched over it, upon which the flowers are to be lightly spread, or rather a basket, supported on three pins, reaching down to the bottom. A gentle heat being applied, just sufficient to raise the spirit, its vapour, lightly percolating through the flowers, will imbibe their finer parts, without making that disagreeable alteration, which liquors applied to fuch tender subjects, in their groffer

form, generally do. Probably the superiority of the French Hungary water, to that prepared among us, is owing to some skilful management of this kind, or that recommended for the foregoing preparation, and employing a persectly pure spirit.

In the Wirtemberg pharmacopoia, fome fage and ginger are added, in the proportion of half a pound of the former, and two ounces of the latter, to four pounds of

the rofemary.

#### SPIRITUS LAVENDULÆ SIMPLEX.

Simple spirit of lavender. Lond.

Take of

Lavender flowers, fresh gathered, a pound and a half;
Proof spirit, one gallon.
Draw off, by the heat of a waterbath, sive pints.

The fame cautions are to be observed here, as in the distillation of the foregoing spirit. Both of them, when made in perfection, are very grateful and fragrant: they are frequently rubbed on the temples, &c. under the notion of refreshing and comforting the nerves; and likewise taken internally, to the quantity of a tea-spoonful, as warm cordials.

### SPIRITUS LAVENDULÆ COMPOSITUS.

Compound spirit of lavender.

Lond.

Take of
Simple spirit of lavender, three
pints;
Spirit of rosemary, one pint;
Cinnamon,
Nutmegs, each half an ounce;
Red saunders, three drachms.

Digest them together, and then strain out the spirit for use.

The compound spirit of lavender of the former London pharmacopæia is as follows.

Take of Lavender flower, one gallon; Sage flowers, Rosemary flowers, Betony flowers, each one handful; Borage flowers, Bugloss flowers, Lillies of the valley, Cowslips, each two handfuls; Balm leaves, Feverfew leaves. Orange tree leaves, Orange flowers, Stechas flowers, Bay berries, each one ounce; French Brandy, four gallons.

Pour the brandy on the other ingredients fresh gathered, and after suitable digestion, draw off in a water-bath two gallons and a half. To this spirit add the following ingredients:

Citron peel,
Yellow faunders, each fix
Cinnamon, (drachms;
Nutmegs,
Mace,

Leffer cardamom feeds, Cubebs, each half an ounce; Aloes wood, one drachm.

Digest these together for twentyfour hours; then filter the spirit, and suspend in it the sollowing ingredients (where they are judged proper) tied up in a thin linen cloth; viz. of Musk,

Ambergris,
Saffron, each half a scruple;
Red roses dried,
Red saunders, each half an ounce.

In the edition of the Edinburgh pharmacoposia, preceding C c the the present, this spirit is thus directed.

Take of the diffilled oils of Lavender, an ounce and a half; Rosemary, an ounce; Marjoram, six drachms; Lemon peel, half an ounce; Nutmegs, three drachms; Cloves, two drachms; Cinnamon, one drachm.

Gradually drop these oils into three gallons of French Brandy, occafionally stirring them together. Distil the mixture in balneo mariæ to two-thirds; and in the spirit which comes over, sufpend the sollowing ingredients, tied up in a linen cloth; viz. of Red saunders, two ounces; English saffron,

Cochineal, each half an ounce; To which, if you would have the spirit perfumed, add of Ambergris, two scruples; Musk, one scruple.

In the present pharmacopæia of Edinburgh, this medicine stands as follows.

Take of

Fresh lavender slowers, two pounds;

Fresh rosemary flowers, one pound;

Fresh yellow rind of lemons, three ounces;

Rectified spirit of wine, one gallon and a half.

Diffil in the heat of a water-bath to dryness, and in the diffilled spirit macerate the following ingredients for three days, viz. of Cinnamon, three ounces;

Cloves, Cubebs, each one ounce; Shavings of red saunders, two

Strain out the spirit for use.

The red faunders is of no farther use in these compositions, than as a colouring ingredient. If a yellow spirit was liked, the yellow saunders would be an excellent article, as it not only communicates a fine colour, but likewife a confiderable share of medicinal virtue. A fpirit distilled from the flowers of lavender and fage, in due proportion, and digested in the cold for a little time with fome cinnamon, nutmegs, and yellow faunders, proves a very elegant and grateful one. Where essential oils are employed, as in the third of the above processes, particular care must be had in the choice of them : for on their goodness that of the medicine depends. The digestion of the spirit with the spices, &c. should be performed without heat, otherwise the flavour of the medicine will be injured.

All these spirits are grateful reviving cordials: the first, though considerably the most simple, is not inserior in elegancy to either of the others. This medicine has long been held in great esteem, under the name of PALSY DROPS, in all kinds of languors, weakness of the nerves, and decays of age: for which reason, we have given the different forms of preparing it that have been followed for some time past. It may be conveniently taken upon sugar, from ten to eighty, or an hundred drops.

AQUA ODORIFERA.

An odoriferous spirit, called fweet
boney-water.

Take of
Coriander feeds;
Honey, each one pound;
Cloves, an ounce and a half;
Nutmegs,
Benzoine,
Storaz, each an ounce:
Vanelloes,

Vanelloes, in number four; Yellow rind of three lemons; French brandy, one gallon.

Digest these ingredients together for forty-eight hours; and then distil off the spirit in balneo mariæ. To one gallon of this spirit, add

Orange flower water,

Rose water, of each one pound and a half;

Ambergris,

1

Musk, of each five grains.

First grind the musk and ambergris, with fome of the water,

gris, with fome of the water, and afterwards put all together, in a large matras; shake them well, and let them circulate for three days and nights in a gentle heat; then suffer them to cool, filter the liquor, and keep it close stopt up for use.

#### Another.

Take of
Coriander feeds, one pound;
Lemon peel, fresh,
Nutmegs, each four ounces;
Ambergris,
Musk, each five grains;
Clean melasses spirit, two gallons.

Bruise the nutmegs and coriander feeds, and put them, with the lemon peel and the spirit, into a small still placed in balneo mariz: tie a thin cloth over the mouth, and sprinkle thereon the ambergris and must, reduced into fine powder; lute on the head, let the whole stand in digestion for twelve hours, and then distil as much as a boiling heat of the bath can force over.

To this add, of Rose water, one pint; Orange flower water, half a pint.

THESE compositions are designated rather as presumes than as

medicines; though for fuch as can bear their fragrance, they might be used to advantage in this last intention. The musk and ambergris do not communicate so much of their smell as might be expected; and ferve chiefly to heighten the flavour of the other ingredients: which these perfumes excellently do, when employed in very small proportion, to all the odoriferous fimples, without imparting any thing perceptible of their own. Both the foregoing spirits are very agreeable; a few drops of either give a fine flavour to a large quantity of other liquor. Mr. Wilson, from whom the first is taken (Prad. Chem page 354.) tells us, that he often made it for King James II, and that it gives one of the most pleasant scents that can be smelt to. The other is formed on the same plan, by omitting fuch articles as appeared superfluous in the first.

## SPIRITUS COCHLEARIÆ. Spirit of Scureygrass. Edinb.

Take of

Fresh survygrass, bruised, ten pounds;

Rectified spirit of wine, five pints.

Steep the herb in the spirit for twelve hours; then with the heat of a water-bath, distil off five pints.

This spirit is very strong of the scurvygrass, and may be given in those cases where the use of this herb is proper, from twenty to an hundred drops. The virtues of scurvygrass reside in a very subtile, volatile oil, which arises in distillation both with water and pure spirit; and if the liquors are exposed to the air, soon exhales from both. The spirit, newly distilled,

Cc2

is extremely pungent, but if long kept, even in close vessels, becomes remarkably less so.

The makers of this spirit have frequently added to the scurvygrass a quantity of horseradish root, and sometimes substituted to it one drawn entirely from the horseradish; the flavour of these two simples being fo much alike, that their distilled spirits are scarce distinguishable from one another. Here it may be observed, that though arum and dracunculus are usually ranked in the same class with the two foregoing vegetables, and looked upon as fimilar to them; this process discovers a remarkable difference: whilst the former yield all their pungency in distillation both to water and fpirit; the latter give over nothing to either, and yet their virtues are destroyed in the operation.

SPIRITUS COCHLEARIM AU-

Golden, or purging spirit of scur-

Take of

Spirit of scurvygrass, one pound. Gamboge, one ounce.

Dissolve the gamboge in the spirit,

and if any fediment falls to the bottom, carefully decant the tinged liquor from it.

This spirit is otherwise made with scammony, or refin of jalap, instead of gamboge.

This has been in great efteem among the common people, and firongly recommended by the venders, in all kinds of scorbutic diforders. It is nevertheless a very indifferent medicine, and little deferves the pompous title given it. It may be taken from twenty to sixty drops, either upon sugar or mixed with syrup.

AQUA ANHALTINA. ... Anhalt water.

Take of Turpentine, fix ounces; Olibanum, one ounce: Aloes wood, three ounces; Cloves, Cinnamon. Cubebs, Rosemary flowers, Galangal, Masticb. Nutmegs, each fix drachms: Saffron, two drachms and a half. Bay berries, Fennel feeds, each half an ounce. Spirit of wine, five pints. Pulverize those ingredients which

Pulverize those ingredients which require such treatment, and digest the whole with the spirit for six days; then distill with an exceeding gentle heat, in balneo maria; the liquor which runs clear is to be separated from the turbid, and kept by itself.

Where the addition of musk is required, fifteen grains thereof are to be tied in a bag, and suspended in the head of the still.

We have inferted this composition from the Brandenburgh pharmacoponia, on account of its being held, in some places, in great esteem. It is rubbed on weak or paralytic limbs, against catarrhs, old pains and aches, &c. and likewife given internally, in doses of half an ounce, for strengthening the stomach, discussing flatulencies, relieving colicky pains, and promoting the uterine purgations. It is very unpleasant to the palate; the aromatics, though sufficiently numerous, and in confiderable quantity, not giving over near enough to cover the strong slavour of the turpentine; there are not many of them, indeed, that give over any thing confiderable at all. A more elegant Ì

elegant spirit of this kind might be prepared from turpentine, rosemary, lavender and sage flowers; or by distilling the spirit sirst from the turpentine alone, and then diffolving in it a proper quantity of any fuitable effential oils.

### ARTICLE II. Distilled spirituous waters.

By distilled spirits are underflood fuch as are drawn with a fpirit that has been previously rectified, or which is reduced nearly to that strength in the operation; by spirituous waters, those in which the spirit is only of the proof strength, or contains an admixture of about an equal measure of wa-These last have been usually called compound waters, even when diffilled from one ingredient only: as those, on the other hand, which are drawn by common water, tho' from a number of ingredients, are named simple; the title fimplex, here, relating not to simplicity in respect of composition, but to the vehicle being plain water. Edinburgh pharmacopæia denominates those waters simple which are drawn from a fingle ingredient, whether the vehicle becommon water, or spirituous water, and all those compounds which are distilled from more than one.

General rules for the distillation of spirituous waters, from the Edinburgh pharmacopæia.

I.

The plants and their parts ought to be moderately and newly dried, except such as are ordered fresh gathered.

II.

After the ingredients have been fleeped in the spirit for the time prescribed, add as much water as will be sufficient to prevent an empyreuma, or rather more.

The liquor which comes over first in the distillation, is by some kept by itself, under the title of spirit; and the other runnings, which prove milky, fined down by art. But it is better to mix all the runnings together, without fining them, that the waters may possess the virtues of the plant entire; which is a circumstance to be more regarded than their sineness or sightliness.

If the distillation is skilfully managed, the heat equable; and all along gentle, and no more drawn off than the quantity directed, most of the waters will appear sufficiently bright and fine: fome of them, which look turbid just after they are drawn, will, on standing for a few days, become clear and trans-parent. The practice here forbid, of faving some of the first runnings apart, is certainly very injurious to the composition; the water being not only robbed by it of some of the more volatile parts of the ingredients, but likewise rendered permanently milky, as wanting the fpirit which, by diffolving the oil of the ingredients that gives this appearance, would make the liquor transparent. Nor is the method of fining the turbid waters by alum, &c. less culpable; for these additions produce their effects only by separating from the liquor what it had before gained from the ingredients.

IV.

In the distillation of these waters, the genuine brandy obtained from wine is directed. Where this C c 3

is not to be had, take instead of that proof spirit, half its quantity of a well rectified spirit prepared from any other fermented liquors: in this steep the ingredients, and then add spring water enough, both to make up the quantity ordered to be drawn off, and to prevent burning.

By this method more elegant waters may be obtained, than when any of the common proof spirits, even that of wine itself, are made use of. All vinous spirits receive some flavour from the matter from which they are extracted; and this slavour, which adheres chiefly to the phlegm or watery part, they cannot be divested of without separating the phlegm, and reducing them to a rectified state.

### AQUA ABSINTHII COMPO-

Compound wormwood waters.

Take of

Calamus aromaticus,
Orange-peel, fresh,
Cinnamon, each four ounces;
Roman wormwood, half a pound;
Mint, three ounces;
Leffer cardamoms,
Mace, each one ounce;
French brandy, two gallons.
Having bruised the seeds and spices, and cut the other ingredients, pour on them the brandy, and after seeping them together for the space of four days, distil off two gallons.

This water was formerly prefcribed as a stomachic, along with bitter infusions; and for this purpose it is the least unsit (as being the most elegant and least unpleasant) of all the wormwood waters that the shops were sur nished with. It is nevertheless too ungrateful an addition to the sine bitters of our new pharmacopæias; and cannot be supposed to contribute any thing to their virtue, which more agreeable spirituous waters would not equally do. Some have expected wormwood water to be itself a bitter; but only the smell and slavour of the wormwood arises in this process, those parts in which its bitterness resides remaining behind in the still.

Is former editions of the London pharmacopæia there were two wormwood waters, which by some are still held in esteem, and were proposed by the committee of the college to be continued at the late revisal, with some amendments.

## AQUA ABSINTHII MINUS COMPOSITA.

Wormwood water less compounded. Take of

Common wormwood leaves, dried, two pounds; Leffer cardamom feeds, two ounces;

Coriander feeds, half a pound; French brandy, four gallons. Let them fleep together for fome

Let them theep together for some time, and then distil off four gallons.

#### AQUA ABSINTHII MAGIS COM-POSITA.

Wormwood water more compounded.
Take of

Sea wormwood,
Common wormwood, each dried, one pound;
Sage,
Mint,
Balm, each dried, two handfuls;
Galangal,
Ginger,
Calamus aromaticus,
Elecampane roots,
Sweet fennel feeds,

Coriander feeds, each three drachms;

Cinnamon,

Cloves,

n:

, 1

ď

۲ĭ

: 0

71

ďE

:

1

g 1

11

Ė

ģί

31

3

10

Cloves, Nutmegs, each two drachms; Leffer cardamom feeds, Cubebs, each one drachm; French brandy, twelve pints.

Having cut or bruised the ingredients, which require that treatment, steep them for some time in the brandy, and afterwards draw off by distillation twelve pints.

### AQUA ALEXETERIA SPIRI-TUOSA.

Spirituous alexeterial mater.

Lond.

Take of
Spearmint leaves, fresh, half a
pound;
Angelica leaves, fresh,
Sea wormwood tops, fresh, each
four ounces;
Proof spirit, one gallon;
Water, as much as will prevent
burning;
Diffil off one gallon.

This is a telerably pleasant water; it is looked upon as an alexipharmic and stomachic, and in these intentions is not unfrequently made use of in juleps, &c.

AQUA ALEXETERIA
SPIRITUOSA cum ACETO.
Spirituous alexeterial water with
winegor.
Lond.

Take of
Spearmint leaves,
Angelica leaves, each half a
pound,
Sea wormwood tops, four ounces;
Proof fpirit, one gallon;
Water, as much as is sufficient
to prevent burning;
Vinegar, one pint.

Distil the fresh herbs with the spirit and water, drawing off one

gallon; to which add the vinegar.

ANGELICA, after trial of fundry other materials, has been found the most effectually to remove the disagreeable flavour which the vinegar would otherwise communicate, and therefore this plant is ordered in a larger proportion here than in the other alexeterial waters. Perhaps it would be more eligible to add the vinegar occasionally; for when mixed with the liquor at first, it is apt to throw down, upon keeping, some of the more valuable parts which the water received from the herbs.

This water is given in the room of the AQUA THERIACALIS, or treacle water, a medicine of fome importance, which in the former Edinburgh pharmacopæia is thus directed.

Take of

Butterbur roots, one pound;
Angelica roots,
Mafterwort roots, each half a
pound;
Zedoary, four ounces;
Scordium leaves,
Rue leaves, each fix ouncess;

Theriaca, one pound;

French brandy, three gallons; Distilled vinegar, half a gallon. Let the roots, leaves and theriaca be steeped in the spirit for four days; then distill off two gallons and a half; to which add the distilled vinegar.

This water is ordered not to be drawn so low as the other distilled waters, and with great judgment; for the addition of the vinegar confiderably weakens it, and if drawn low, renders it very unsightly. It is lest to the choice of the operator, to employ either Andromachus's, or the Edinburgh treaches the latter is the best of the two, but neither of them are proper subjects for distillation; for besides, Cc 4

**4** l

that three parts in four are honey which yields nothing, they contain several other ingredients that afford as little.

The AQUA THERIACALIS of the former London pharmacopæia is as follows.

Take of

Juice of green walnuts, four pints; Rue, three pints;

Carduus,

Balm, each two pints; Butterbur roots, fresh, a pound and a half;

Burdock roots, fresh, one pound;

Angelica roots,

Masterwort roots, fresh, each half a pound;

Scordium, fresh, four handfuls; Venice treacle and

Mithridate, kept for some time, each eight ounces:

Lemon juice, two pints; French brandy, a gallon and a

half.

Draw off by distillation three gallons and a half, then add half a gallon of distilled vinegar.

The predominant flavour of this water is from the rue and angelica; the rest contribute only enough to render the whole more offensive. What qualities it can receive from the numerous ingredients of the imagined all powerful theriaca may be estimated by this, that the whole species of that electary, employed in half an ounce of the water, its usual dose, amounts not to a fingle grain; the mithridate, with which our pharmacopæia, by the advice of Sir Theodore Mayerne, had the honour of enriching the composition, being also just of the same importance in it.

The college of Edinburgh has now given the following amendment of this water, under the title of AQUA EPIDEMIA.

Plague water.

Edinb.

Take of

Masterwort roots, a pound and a half;

Angelica feed,

Elder flowers, each half a pound;

French brandy, three gallons. Digest for two days, then distil off two gallons and a half; to which add half a gallon of distilled vinegar.

THE foregoing compositions are the only distilled waters in which the heat of the spirit is tempered by the addition of vinegar, an ingredient which renders them ferviceable in many cases where spirituous liquors alone would be im-The treacle water has proper. long been held in great esteem as a fudorific and alexipharmac; and those which the London and Edinburgh colleges have now directed in the room of it, though far more fimple and elegant, are not inferior in efficacy.

### AQUA SEMINUM ANISI COM-POSITA.

Compound anifeed water.

Lond.

Take of

Anisceds,

Angelica feeds, each half a pound;

Proof spirit, one gallon;

Water, as much as is sufficient to prevent burning.

Draw off by distillation one gailon.

This is a very elegant anifeed water, the angelica feeds greatly improving the flavour of the anife: it is apt to turn out milky, if drawn fo low as here ordered.

AQUA

AQUA CORTICUM AURAN-TIORUM SPIRITUOSA. Spirituous orange peel water.

Lond.

Z

1

Take of Outer rind of Seville orange peel, dried, half a pound: Proof spirit, one gallon; Water, as much as is fufficient to prevent an empyreuma. Distil off one gallon.

THIS is confiderably stronger of the orange peel than the simple wa-It is used as a cordial, stomachic, and carminative.

AQUA BRYONIE COMPOSITA. Compound bryony water.

Take of

Bryony roots, one pound; · Wild valerian root, four ounces; Pennyroyal, Rue, each half a pound; Mugwort leaves, Feverfew flowers, Savin tops, each one ounce: Orange peel, fresh, Lovage feed, each two ounces: French brandy, two gallons and a half.

Having cut or bruifed those ingredients which require such treatment, steep them in the brandy four days; then draw off by distillation two gallons and a half of liquor.

This composition, designed as an antihysteric, is liable to considerable objections, not only in regard to its particular ingredients, but to the medicinal intention of the whole. Many, by the use of this and other. like waters, under the notion of medicines, have been betrayed into the pernicious habit of drinking drams: whereas, however spirituous liquors may give a temporary relief to the languors of hysterical and hypochondriacal perfons, none fuffer to foon the ill effects attending the constant use of

The unpleasant flavour of them. this water renders it exceptionable also as a vehicle of other antihysteric medicines, which in general are of themselves sufficiently ungrateful: a finall augmentation in the dose of the medicines themselves (as the London committee observe) would abundantly compensate any assistance to be expected from this water, and leave room for the use of a more agreeable vehicle.

The college of London has therefore wholly omitted this water, without giving any thing of fimilar intention in its place. That of intention in its place. Edinburgh still retains it, but has improved the composition, and rendered it more simple, by the rejection of the more exceptionable ingredients. The bryony root, from which the water receives its name, is the most exceptionable of them all: this being therefore now omitted, the water is distinguished by the name of another of its ingredients, and is directed as follows.

### AQUA VALERIANÆ COMPO-SITA.

Compound valerian water.

Take of

Wild valerian root, a pound and

Lovage seed, half a pound; Pennyroyal leaves, four ounces: Savin tops, two ounces; French brandy, two gallons. Digest for two days, and then draw off by diffillation two gallons.

> AQUA SEMINUM CARDAMOMI. Cardamom seed water. Lond.

Take of

Lesser cardamom seeds, freed. from the hulks, four ounces; Proof

Proof fpirit, one gallon;
Water, as much as is sufficient
to prevent burning.
Diffil off one gallon.

This water is a grateful cordial and carminative, the cardamom feeds giving over in this process the whole of their flavour. It is not perhaps very necessary to be at the trouble of separating the huses, for these communicate nothing disagreeable: the only difference is, that if employed unhusked, a proportionably larger quantity of them must be taken.

# AQUA SEMINUM CARUI. Caraway water. Lond.

Take of
Caraway feeds, half a pound;
Proof spirit, one gallon;
Water, as much as will prevent
burning.
Distil off one gallon.

This is a cordial in common use: it contains the flavour of the caraway seeds in perfection.

## AQUA CINNAMOMI SPIRITUOSA.

Spirituous cinnamon water.

Land

Take of
Cinnamon, a pound;
Proof spirit, a gallon;
Water, so much as will prevent
burning.
Draw off by distillation one gallon.

### AQUA CINNAMOMI CUM VINO.

Cinnamon water with wine.

Edinb.

Take of
Cinnamon, one pound;
French brandy, one gallon.

Let them steep together for two days, and then distil off one gallon.

This is a very agreeable and useful cordial water, but not fo strong of the cinnamon as might be expected; for very little of the virtues of the spice arise till after the pure spirituous part has distilled. Hence in the former editions of the London pharmacopæia, the distillation was ordered to be protracted till two pints more, than here directed, were come over. By this means, the whole virtue of the cinnamon was more frugally than judiciously obtained, for the difagreeable flavour of the feints of proof spirits, and the acidulous liquor arising from cinnamon as well as other vegetables, when their distillation is long continued, gave an ill relish to the whole; at the same time that the oil which was extracted from the spice, was by this acid thrown down.

In the pharmacoposia reformata, it is proposed to make this water, by mixing the aqua cinnamoni fimplex with somewhat less than an equal quantity of rectified spirit: on shaking them together, the liquor loses its milky hue, soon becomes clear, and more elegant than the water distilled as above: it is equally strong of the cinnamon, and free from the nauseous taint which the common proof spirits are impregnated with.

AQUA JUNIPERI COMPOSITA. Compound juniper water. Lond.

Take of
Juniper berries, one pound;
Sweet fennel feeds,
Caraway feeds, each an ounce
and a half;
Proof fpirit, one gallon;
Water, as much as is fufficient
to prevent burning.
Diffil off one gallon.

THIS

This water mixed with about an equal quantity of the rob of juniper berries, proves an useful medicine in catarrhs, debility of the stomach and intestines, and difficulty of urine. The water by itself is a good cordial and carminative: the fervice which this and other spirituous waters do in these intentions, is too commonly known; though the ill consequences that soliow their constant use, are too little regarded.

AQUA MENTHÆ PIPERITI-DIS SPIRITUOSA.

Spirituous pepper-mint water, Lond.

Take of
Pepper-mint leaves dry, a pound
and a half;
Proof spirit, a gallon;
Water, as much as is sufficient
to prevent an empyreuma.
Draw off by distillation one gallon.

THIS water is made use of in flatulent colics and other like diforders; in which it oftentimes gives immediate relief. it smells and takes strongly of the pepper-mint.

### **▲QUAMENTHÆ VULGARIS** SPIRITUOSA.

Spirituous Spearmint water.

Lond.

Take of
Spearmint leaves, dry, a pound
and a half;
Proof spirit, a gallon;
Water, as much as will prevent
burning.
Distil off one gallon.

THIS water, if the spirit be good, turns out a very elegant one, and preserable, in weakness of the stomach, reaching to vomit, and the like, to many made elaborate preparations. Where the disorder is not accompanied with heat

or inflammation, half an ounce of this water may be given diluted with fomeagreeable aqueous liquor-

AQUA MIRABILIS.

Take of
Cinnamon, two ounces;
Lemon peel, an ounce;
Angelica feeds,
Leffer cardamom feeds,
Mace, each half an ounce;
Cubebs, two drachms;
Balm leaves, fix ounces;
French brandy, one gallon.

Pour the brandy on the other ingredients bruifed; and after digelting them for four days, draw off by distillation one gallon.

THE above composition of this celebrated water is that which was formerly followed. At the late reformation it has received a considerable improvement: the cardamoms, cubebs, and balm, are omitted, and an addition of peppermint introduced. The formula at present is as follows.

### AQUA AROMATICA, vulgo MIRABILIS.

Aromatic water, commonly called aqua mirabilis. Edinb.

Take of
Cinnamon, two ounces;
Fresh yellow rind of lemons,
Angelica seeds, each one ounce;
Mace, half an ounce;
Pepper-mint, three ounces;
French brandy, one gallon.
Digest for two days, and then distil off one gallon.

THIS water is very rich of the spices; and proves a pleasant, warm cordial and carminative. In those who have not, by frequent use, deprived themselves of the benefit of these kinds of liquors, it often gives present relief in languors,

guors, flatulencies, colicky pains,

and other like complaints.

The spices in these two compositions being rather too dear for the purposes of a common cordial water, the wholesale dealers, as I have been informed, generally substitute to them a cheaper spice from our own plantations, Jamaica pepper. A very elegant water is prepared also from that spice by itself in the following proportions.

Aqua piperis Jamaicensis spirituosa.

Spirituous Jamaica pepper water. Take of

Jamaica pepper, half a pound; Proof spirit, three gallons; Water, a sufficient quantity to

prevent an empyreuma.

Draw off by distillation three gal-

This water is far more agreeable than a fimple water drawn from the fame spice; and has long had a place among the cordials both of the distiller and apothecary; though it has not yet been received into any public pharmacopæia.

AQUA NUCIS MOSCHATÆ.

Nutmeg water.

Lond.

Take of
Nutmegs, two ounces:
Proof (pirit, a gallon;
Water, as much as will prevent
burning.
Draw off by diffillation one gallon.

This water (with the addition only of fome hawthorn flowers, an article of very little fignificance) was formerly celebrated in nephritic diforders, under the name of AQUANEPHRITICA. At prefent, it is regarded only as an agreeable spirituous liquor, lightly impregnated with the nutmeg slavour.

AQUA POEONIE COMPOSITA.

Compound peony water.

Take of

Peony roots, two ounces; Wild valerian roots, an ounce and a half;

White dittany root, an ounce; Peony feeds, fix drachms; Lilies of the valley, fresh, four

ounces; Lavender flowers,

Rosemary flowers, each two ounces:

Betony, Marjoram, Rue,

Sage, tops of each, one ounce; French brandy, a gallon and a half.

Cut or bruife those materials that require such treatment, steep them four days in the brandy, and then distil over a gallon and a half of liquor.

This water was formerly distinguished by the title of Aqua ANTIEPILEPTICA; and recommended in all kinds of epilepses and nervous complaints. For fome time past, it has had little regard paid to it, having rarely been prefcribed any otherwife than as a vehicle, and as fuch not often. ingredients from which it receives its name, the peony roots and feeds, communicate little or nothing to the water; whatever virtues these are possessed of, remain behind in the decoction; nor are thefe the only exceptionable articles; the dittany, betony, and some others, though of the aromatic kind, afford so little, as not to deserve a place among more powerful mate-

The above formula is taken from the edition of the Edinburgh pharmacopæia preceding the prefent. It is here inferted for the fake of those, who may still have fome

some regard for forms so long received; and for the same reason, the peony water of the last London pharmacopæia is also subjoined. The committee endeavoured to amend it at the late reformation. by retaining only those ingredients to which a star is affixed.

Take of

Lilies of the valley \*, fresh gathered, one pound; Lime flowers\*, half a pound; Peony flowers \*, four ounces; Male peony root\*, two ounces and a half: White dittany root, Long birthwort, each half an ounce : Milletoe of the oak, Rue , each two handfuls: Peony feeds, husked, ten drachms; Rue seeds, three drachms and a half; Russia castor, Cubebs \*. Mace \*, each two drachms; Cinnamon , an ounce and a Rosemary flowers, six pugils: Stæchas flowers. Lavender flowers, each four pugils ; Betony flowers, Clove-july flowers, Cowslips, each eight pugils: Juice of black cherries, four pints : French brandy , two gallons and After proper maceration, distil off four gallons.

### AQUA PULEGII SPĪRITUOSA.

Spirituous penny-royal water. Lond.

Take of Penny-royal leaves, dry, a pound and a half; Proof spirit, a gallon;

Water, as much as will prevent burning. Distil off one gallon.

THIS water has a good share of the flavour of the penny-royal, and is pretty much in use as a carminative and antihysteric.

AQUA RAPHANI COMPOSITA. Compound borseradish water. Lond.

Take of Garden scurvygrass leaves, fresh, four pounds; Horseradish root, fresh, Orange peel, fresh, each two pounds; Nutmegs, nine ounces; Proof spirit, two gallons; Water, a sufficient quantity to prevent burning. Draw off by distillation two gallons.

#### Edinb.

Take of Horseradish root. Garden scurvygrass, fresh, each three pounds; Orange peel, fresh, Juniper berries, Canella alba, each four ounces: French brandy, three gallons. Steep the juniper berries and canella alba in the spirit, for two days: then add the other ingredients, and draw off three gallons.

Born these waters are very elegant ones, and as well adapted for the purposes of an antiscorbutic, as any thing that can well be contrived in this form. The committee of the London college observe, with regard to the first, that the horseradish and scurvygrass join very well together, giving a similar flavour, though not a little difagreeable; that the nutmeg sup-

preffes this flavour very successfully, without superadding any of its own; and to this, orange peel (no incongruous ingredient to the intention of the medicine) adds a flavour very agreeable. Arum root has generally had a place in this water, but is here deservedly thrown out; for it gives nothing of its pungency over the helm, notwithstanding what is afferted, by fome dispensatory-writers, to the contrary. Mustard seed, though not hitherto, that I know of, employed in these kinds of compositions, should feem to be an excellent ingredient; it gives over the whole of its pungency, and is likewife less perishable than most of the other substances of this class: this feed wants no addition, unless some aromatic material to furnish an agreeable flavour.

Aqua vulneraria, seu Aqua CATAPULTARUM. Arquebusade water. Pharm. Argent.

Take of Comfry, leaves and roots. Mugwort, Bugloss, each four handfuls: Betony, Sanicle, Ox-eye daify, Common daify, Greater figwort, Plantane, Agrimony. Vervain, Wormwood,

Fennel, each two handfuls: St. John's wort, Long birthwort, Orpine. Veronica, Lesser centaury: Milfoil, Tobacco. Mouse-ear. Mint. Hyssop, each one handful : Wine, twenty-four pounds.

Having cut and bruised the herbs, pour on them the wine, and let them fland together in digestion, in horsedung, or any other equivalent heat, for three days ; afterwards distil in an alembic with a

moderate fire.

This celebrated water has been for some time held in great esteem, in contusions, for resolving coagulated blood, discussing the tumours that arise on fractures and dislocations, for preventing the progress of gangrenes, and cleanfing and healing ulcers and wounds, particularly gun-shot wounds. Mr. Lemery has been at the pains of writing a whole treatife on it; in which he confiders each of the ingredients fingly, and supposes the water to possess their united virtues. In this, however, he is mistaken; for the virtues of most of the herbs, admitting them to be as great as he would have them. refide in fuch parts as are not capable of being elevated in this process.

### CHAPTER VI.

Concentration of the medicinal parts of juices and infusions by evaporation.

WHEN vegetable juices, or watery or spirituous decoctions or infusions, are exposed to a continued heat; the fluid, gradually evaporating, carries off with it such volatile matters as it was impregnated with, and leaves the more fixed united together into one mass. As the object of the preceding chapter was the collection of the volatile principle which exhales along with the fluid; that of the prefent is this re-union and concentration of the fixt matter. The mass which remains from the eva-

poration of the expressed juice of a plant, is called an inspissated juice; from watery decoctions or insusions, an extract; from spirituous timetures, a resin, or essential extract. The term extract is frequently used also as a general appellation of all the three kinds. Inspissated juices and watery decoctions, particularly the former, when evaporated no further than to the consistence of oil or honey, are called rob or sapa; and spirituous tinctures, reduced to a like consistence, are called bal-sam.

### SECT. I.

### Inspissated juices.

WHAT relates to the expression of juices, has already been delivered in chap, ii. with the most effectual means of preserving them in their liquid state, and a general account of what substances do, or do not, give out their virtues with their juices. In the inspissation of juices, there is further to be considered the volatility or fixity of their medicinal parts: if a plant loses its virtue, or part of its virtue, in being dried, it is obvious that the juice must lose as much in being inspissated to dryness; how gentle soever the heat be, with which the inspissation is performed, It is likewise to be observed, that the medicinal parts of some juices

are kept in a state of perfect folution by the watery sluid, so as to be completely retained by it after the liquor has been made sine by settling, straining, or other means; while the medicinal parts of others, not dissoluble by watery menstrua, are only dissured through the liquor in the same manner as the seculencies are, and separate along with these on standing.

ROB BACCARUM SAMBUCI.

Rob of elder berries.

Lond.

Let the depurated juice of elderberries be inspissated with a gentle heat.

Edinb.

Edinb.

Take two quarts of the juice of ripe elder-berries, and half a pound of white fugar. Evaporate over a gentle fire, or in a water-bath, to the confishence of honey.

This preparation, made with or without fugar, keeps well, and proves a medicine of confiderable importance, as an aperient, generally promoting the natural excretions by ftool, urine or sweat. The dose is from a drachm or two to an ounce or more. A spoonful, diluted with water, is usefully taken, in common colds, at bed time.

SUCCUS PRUNORUM SIL-VESTRIUM, five ACACIA GERMANICA.

Inspissated juice of sloes, or German acacia. Edinb.

Let any quantity of the juice of unripe floes be inspiffated over a gentle fire.

This juice is infpissated nearly to dryness, care being taken to prevent its burning, as directed in the following section for making extracts with water. It is a moderately strong astringent, similar to the Egyptian acacia, for which it has been commonly substituted in the shops. (See page 73.) It is given, in sluxes and other diforders where styptic medicines are indicated, from a scruple to a drachm.

EXTRACTUM PLANTAGINIS.

Extract of plantane. Edinb.

Let any quantity of the juice of plantane leaves be depurated; either by fuffering it to fettle,

and then decanting off the clear liquor; or by straining; or by clarification with whites of eggs. Afterwards evaporate the juice in a sand-heat, to the confishence of honey.

After the same manner, extracts may be made from all acid, cooling, styptic, juicy plants.

THIS is a method of treating plants very little practifed, but which promifes, if duly profecuted, to afford medicines of confiderable power. There are many common and neglected herbs, as plantane, chickweed, chervil, &c. whose juices in their dilute fate, as well as the herbs in substance, feem to be altogether infignificant, but which, when the juice is well depurated from the feculent matter, and concentrated by the evaporation of the fluid, yield extracts, which discover to the take no small activity. These extracts, like those prepared from the juices of most of the summer fruits, if inspissated to dryness, grow moult again in the air.

Extract um cicuta.

Extract of bemlock.

Take fresh hemlock leaves, gathered just before the plant begins to flower; which it commonly does in July, or about the latter end of June. Press out the juice; and immediately, without suffering it to fettle, put it into a shallow glazed earthen pan, over a very gentle fire; keeping it continually stirring, to prevent its burning, till it is reduced to a thick greenith brown This mass may be formed into pills with a little of the powder of the dried leaves of the plant.

THIS is the preparation of hemlock lately published at Vienna by

Dr. Storck; who recommends it as a high resolvent in many obstinate disorders, where the common remedies avail nothing. He observes, that fmall doses should always be begun with, as two grains, made into a pill, twice a day; and that by gradually increasing the dose, it may be given to two, three, or even four drachms a day, and continued in such quantities for feveral weeks: that it may be used with safety, in infancy, old age, and pregnancy: that it neither accelerates or disturbs the circulation; neither heats nor cools; nor affects the animal functions: that it increases the secretions, and renders the mouth moift; seldom purges; very rarely vomits; fometimes augments perspiration; often produces a copious discharge of viscid urine; but in many patients does not increase any of the fensible evacuations: that it removes obstructions and their consequences; relieves rheumatic pains, though of long continuance: discusses schirrous tumours, both internal and external, and cures dropfies and confumptions proceeding from schirrosities: that it often dissolves cataracts, or stops their progress, and has sometimes removed the gutta ferena; that inveterate Cutaneous eruptions. scald heads, malignant ulcers, cancers, the malignant flour albus, and gonorrhoea of long standing, obstinate remains of the venereal diseate, and caries of the bones generally yield to it; that for the most part it is necessary to continue this medicine for a very considerable time, before the cure is effected, or much benefit perceived from it: that in some cases it failed of giving any relief, and that he met with some persons who could not bear its effects; and that consequently there must be some

latent difference in the habit, the diagnostic signs of which are at present unknown: that though it is by no means infallible, any more than other medicines in their ref-.peclive intentions, yet the great number of deplorable cases that have been happily cured by it is fufficient to recommend it to further trials. The efficacy of this medicine is confirmed by many eminent practitioners abroad; tho' the trials hitherto made of it in this country have not been attended with much success. Somewhat perhaps may depend upon the time of the plant's being gathered, and the manner of the preparation of the extract. Dr. Storck himself takes notice of some mistakes committed in this respect: some have lest the herb in a heap for several days, whence part of it withered, part rotted, and the juice became thick and mucilaginous; others have taken a very large quantity of the juice, and boiled it down in copper vessels with a great heat, by which means a strong fetor was diffused to considerable distance, and the most efficacious parts dislipated; others, with officious care, have clarified the juice, and thus obtained a black tenacious extract, retaining but a finall degree of the specific smell of the plant: the extract duly prepared, according to the above prescription, is of a greenish brown colour, and a very disagreeable smell, like that of mice. But though there is reason to believe, that much of the extract used here had been ill prepared, we can by no means adunit that its general inefficacy was owing to this cause; for though there are few instances of its discocovering any valuable medicinal powers, there are several of its having activity enough, even in Dd

finall dofes, to produce alarming lymptoms.

#### ELATERIUM. Lond.

Slit ripe wild cucumbers, and having very lightly pressed out the juices, pass it through a fine hair sieve into a glazed earthen vessel. After standing for some hours, the thicker part will fall to the bottom; from which the thinner is to be poured off, and what liquid matter is ftill left is to be feparated by filtration. The remaining thick part is to be covered with a linen cloth, and exposed to the fun, or other gentle heat, till grown thoroughly dry.

WHAT happens in part in the foregoing preparation, happens in this completely, the spontaneous feparation of the medicinal matter of the juice on flunding for a little time: and the case is the fame with the juices of several other vegetables, as those of arum root, iris root, and bryony root. Preparations of this kind have been commonly called FOECULE. The filtration above directed, for draining off such part of the watery fluid as cannot be feparated by decantation, is not the common filtration through paper, for this does not fucceed here: the groffer parts of the juice, falling to the bottom, form a viscid cake upon the paper, which the liquid cannot pass through. The feparation is to be attempted in another manner, so as to drain the fluid from the top: this is effected by placing one end of fome moistened strips of woollen cloth, fkains of cotton, or the like, in the juice, and laying the other

end over the edge of the veffel, fo as to hang down, lower than the furface of the liquor: by this management the separation succeeds in perfection.

The Edinburgh pharmacopæia, directs the wild cucumbers to be gathered before they have grown fully ripe; and no more of the juice to be taken, than that which issness spontaneously upon flitting them. After fettling, the fluid part is ordered to be poured away; and the thick refiduum, without any further draining or filteration, to be exficcated in the fun.

THE juice of the unripe fruit is faid to operate with greater violence than of that which is ripe. The foregoing prescriptions do not perhaps differ so much in regard to the degree of maturity, as in the manner of expressing it; both feeming to intend the fruit to be taken just before it has grown fo thoroughly ripe, as to built and shed its juice on being touch If any pressure is used, it should be exceeding gentle, otherwife some of the inactive pulpy matter of the fruit will be forced out with the juice, and render the ftrength of the elaterium precarious; a point of primary confequence to be avoided, in a medicine of fuch powerful operation, and limitted to fo imall a dofe.

Elaterium is a strong irritating, cathartic, and oftentimes operates also as an emetic. It is never to be ventured on but in indolent phlegmatic habits, as in dropties. in which it is by some particularly Two or recommended. grains are in general a fufficient dofe.

### SECT. II.

### Extracts with water.

HESE extracts are prepared, by boiling the subject in water, and evaporating the strained decoction to a thick consistence.

This process affords us some of the more active parts of the plants, free from the useless, indissoluble, earthy matter, which makes the largest share of their bulk. There is a great difference in vegetable fubstances, with regard to their fitness for this operation; some yielding to it all their virtues, and others scarce any. Those parts in which the fweet, glutinous, emollient, cooling, bitter, auftere, aftringent virtues relide, are for the most part totally extracted by the boiling water, and remain almost entire upon evaporating it: whilst

those which contain the peculiar odour, flavour, and aromatic quality, are either not extracted at all, or exhale along with the menstruum. Thus, gentian root, which is almost simply bitter, yields an extract possessing, in a small volume, the whole taste and virtues of the root: wormwood, which has a degree of warmth and ftrong flavour joined to the bitter, loses the two first in the evaporation, and gives an extract not greatly different from the foregoing the aromatic quality of cinnamon is diffipated by this treatment, its aftringency remaining; whilft an extract made from the flowers of lavender and rolemary, discovers nothing either of the taste, smell, or virtues of the flowers.

### General rules for making extracts with water.

1. It is indifferent, in regard to the medicine, whether the subject is used fresh or dry: fince nothing that can be preserved in this process, will be lost by drying. In regard to the facility of extraction, there is a very considerable difference; vegetables in general giving out their virtues more readily, when moderately dried, than when fresh.

2. Very compact dry substances should be reduced into exceeding small parts, previous to the effusion of the menstruum.

3. The quantity of water ought to be no greater than is necessary for extracting the virtues of the subject. A difference herein will sometimes occasion a variation in the quality of the product; the

larger the quantity of liquor, the longer fire will be requisite for evaporating it, and consequently the more of the volatile parts of the subject will be dissipated. A long continued heat likewise makes a considerable alteration in the matter which is not volatile: sweet substances, by long boiling with water, become nauseous; and the drastic purgatives lose their virulence; though without any remarkable separation of their parts.

4. The decoctions are to be depurated by colature; and afterwards suffered to stand for a day or two, when a considerable quantity of sediment is usually sound at the bottom. If the liquor, poured off clear, be boiled down a little, and afterwards suffered to

Dd 2 . coo

cool again, it will deposite a fresh sediment, from which it may be decanted before you proceed to finish the evaporation. The decoctions of very resinous substances do not require this treatment, and are rather injured by it; the resin subsiding along with the inactive dregs.

5. The evaporation is most conveniently performed in broad shallow vessels; the larger the surface of the liquor, the sooner will the aqueous parts exhale: this effect may likewise be promoted by agi-

tation.

6. When the matter begins to grow thick, great care is necessary to prevent its burning. This accident, almost unavoidable if the quantity is large, and the fire applied as usual under the evaporating pan, may be effectually fecured against, by carrying on the inspiffation after the common manner, no farther than to the confiftence of a syrup, when the matter is to be poured into shallow tin, or earthen pans, and placed in an oven, with its door open, moderately heated; which acting uniformly on every part of the liquid, will foon reduce it to any degree of This may confistence required. likewise be done, and more securely, in balneo mariæ, by fetting the evaporating vessel in boiling water; but the evaporation is here exceeding flow and tedious.

7. Extracts are to be sprinkled with a little spirit of wine, to prevent their growing mouldy [L.] They should be kept in bladders moistened with sweet oil [E.]

# EXTRACTUM ABSINTHII. Extrad of wormwood. Edinb.

Boil dried wormwood leaves in water, supplying fresh water occasionally, till the herb has given out all its virtues to the liquor. Strain the decoction through a, woollen cloth, and evaporate it, in a fand-heat, to the confistence of honey.

THIS extract is almost simply bitter: the peculiar flavour of the wormwood being diffipated in the evaporation. The chemists usually prepare the extract of wormwood from the decoction which remains in the still after the diftillation of the effential oil: and. provided the fill has been perfectly clean, and the liquor not flood too long in it after the distillation, this piece of frugality is not to be disapproved of; since, whether we catch the exhaling vapour; or fuffer it to be diffipated in the air. the remaining extract will be the same.

## EXTRACTUM CENTAURII MINORIS.

Extract of leffer centaury, Edinb.

This is directed to be prepared in the same manner as the preced-It is the oldest extract we have any account of: its preparation is very accurately and circumstantially fet down, in a book usually ascribed to Galen, de vir-tute centaureæ. The author of that treatife recommends the extract as a medicine of excellent fervice in many cases; and looks upon cen-. taury as a specific against the bite of a mad dog and other venemous animals. It is doubtless an useful bitter, possessing the general virtues of the substances of that class. but cannot well be supposed to have any others.

# EXTRACTUM CHAMÆMELI. Extract of chamomile. Edinb.

This extract is prepared from the flowers of chamomile, in the fame manner

manner as those of the leaves of the two preceding plants. Nor is it greatly different from those extracts in quality; the specific flavour of the chamomile exhaling in the evaporation. The chemiss commonly prepare it, like that of wormwood, from the decoction remaining after the distillation of the essential oil.

## EXTRACTUM ENULÆ CAMPANÆ.

Extract of elecampane.

Lond.

Boil the roots of elecampane in water, press out and strain the decoction, and set it by to settle. Then pour off the clear liquor, and boil it down to a pilular consistence; taking care, towards the end, to prevent its burning to the vessel.

This extract retains a confiderable share of the virtues of the root: its talle is somewhat warm, and not ungratefully bitterish. It is given, from a scruple to a drachm, in a lax state of the sibres of the stomach, and in some disorders of the breast.

# EXTRACTUM GENTIANÆ. Extract of gentian. L. E.

This extract is prepared from the roots of gentian, in the same manner as the foregoing extracts. It is of a reddish brown colour, and an intensely bitter taste, being one of the strongest of the vegetable bitters.

#### EXTRACTUM GLYCYRRHIZÆ. Extract of liquorice. Lond.

Lightly boil fresh liquorice roots in water, press the decoction through a strainer, and after the feces have subsided, evaporate it until it no longer sticks to the singers, taking care, towards the end of the operation, to prevent an empyreuma.

IT is convenient, before boiling the root, to cut it transversely into finall pieces, that it may more readily give out its virtues by light coction: if the boiling is long continued, the rich sweet taste, for which this preparation is valued, will be greatly injured. For the same reason, the quantity of water ought to be no larger than is absolutely necessary to extract the virtues of the root: a quart, or at most three pints, will be fully sufficient for a pound of liquorice. It would be of considerable advantage to the preparation, and probably (when made in quantity) less expensive to the preparer, to use instead of the decoction, juice of liquorice, pressed out betwixt iron rollers, after the manner practifed abroad for obtaining the juice of the fugar-cane.

Large quantities of extract of liquorice have been usually brought to us from Spain, and other foreign countries; but it is very rarely met with in the shops in perfection; the makers of this commodity, both at home and abroad, being either very flovenly in its preparation, or defignedly mixing it with fand, and other impurities. When made with due care, it is exceedingly sweet, not at all bitterish, or nauseous, more agrecable in taste than the root itself; of a pleasant smell, a redbrown colour, and when drawn out into strings, of a bright golden colour; totally foluble in water, without depositing any fe-

This preparation would be very convenient for many purposes in D d 2 the

the shops, if kept in a somewhat softer consistence, than that of an extract. The only inconvenience attending this soft form is, its being apt in a short time to grow mouldy: this may be effectually prevented, by the addition of a small portion of spirit of wine.

#### EXTRACTUM HELLEBORI NIGRI.

Extract of black bellebore.

This extract is prepared from the roots of black hellebore, in the fame manner as that of elecampane roots above described. It purges with confiderably less violence than the hellebore in substance; and appears to be one of the best preparations of that root when intended to act only as a cathartic. The dose is from eight or ten grains to fifteen or more.

#### EXTRACTUM LIGNI CAMPÉCHENSIS. Extrad of logwood. Lond.

Take of logwood, reduced to powder, one pound. Boil it in a gallon of water till half the liquor is confumed; repeating the coction with fresh water four times, or oftener: the several decoctions are to be mixed together, passed through a strainer, and evaporated to a due consistence.

This wood very difficultly yields its virtues to watery menstrua, and hence the reducing it into fine powder is extremely necessary. The Edinburgh dispensatory directs spirit of wine to be called in aid; see the following section.

The extract of logwood has been used for a considerable time in some of our hospitals, but is now sirst received into the pharmaco-

pocia. It has an agreeable fweet taste, with some degree of astringency; and hence becomes ferviceable in diarrhoas, for blunting the acrimony of the juices, and moderately constringing the intestines and orifices of the smaller vessels; it may be given from a scruple to half a drachm, and repeated five or fix times a day to advantage. During the use of this medicine, the flools are frequently tinged red by it, which has occafioned some to be alarmed, as if the colour proceeded from blood: the prescriber therefore ought to caution the patient against any surprize of this kind.

### EXTRACTUM CORTICIS PE-RUVIANI.

molle et durum.

Extract of Peruvian bark, foft and
bard.

Lond.

Boil a pound of powdered bark in five or fix quarts of water, for an hour or two, and pour off the liquor, which whilst hot will be red and transparent, but on growing cold becomes yellow and turbid. The remaining bark is to be boiled again in the same quantity of water as before, and this process repeated till the liquor remains transparent when cold. All the decoctions, strained and mixed together, are to be evaporated over a very gentle fire to a due confiftence, care being taken to prevent the matter from burn-

This extract is directed to be kept in the shops, both in a soft and hard form; the sirst of a proper consistence for making into pills: the other fit for being reduced into pow-

der.

PERUVIAN

1

Z,

٠.,

Z

Ť

I

;

ġ:

**,** 

LI

5:

1

, 13

1:

: 2

...

12

1

5

Ľ

K

,1

٤

5

5

ç

ŕ

,

5

;

3

۶

PERUVIAN bark is a refinous drug; the refin melts out by the heat, but is not perfectly disfolved by the water; hence, in cooling, it separates, renders the liquor turbid, and in part falls to the bottom, as appears manifeltly upon examining the fediment by spirit of wine (see the account of this article in page 197.) This extract might be made to better advantage by the affistance of spirit of wine, after the same manner as that of jalap; and this method the Edinburgh college have directed. But, as the committee observe, all the spitits which can be expected to be employed for this process amongst us, are accompanied with some degree of a bad flawour: this adheres most strongly to, the phlegmatic part of the spirit, which evaporating last, must communicate this ill flavour to the extract; a circumstance of very great consequence; as this medicine is defigned for fuch, whose stomachs are too weak to bear a due quantity of bark in substance. Ten or twelve grains of the hard extract are reckoned equivalent to about half a drachm of the bark itself.

EXTRACTUM ligni GUAIACI, molle et durum.

Extract of guaiacum wood, foft and bard.

Lond.

Boil a pound of shavings of guaiacum in a gallon of water, till half the lequor is wasted, repeating the operation four times, or oftener; with the same quantities of fresh water. The seweral decoctions; passed through a strainer, are to be mixed and inspissated together; when the aqueous parts are almost entirely exhaled, a little rectified spirit of wine is to be added, that the whole may be reduced into an uniform and tenacious mass. This extract is to be prepared as the foregoing, in a soft and hard form.

HERE the refinous parts of the wood which were boiled out with the water, are apt to separate towards the end of the inspissation: hence an addition of spirit becomes necessary, to keep them united with the rest of the matter. The extract agrees in virtue with the wood; see page 148.

# EXTRACTUM RUTÆ. Extra@ of rue. Lond.

This is prepared from the leaves of rue, in the same manner as that of elecampane roots already described. It retains a considerable share of the warmth and pungency of the rue; for though the principal virtues of rue reside in an essential oil, yet the oil of this plant, as formerly observed under the head of those preparations (see page 364) is not of a very volatile kind.

## EXTRACTUM SABINÆ. Extraæ of Savin. Lond.

This extract is prepared from the leaves of favin, in the fame manner as the preceding. It does not retain so much, as that extract does, of the virtues of its subject, the oil of favin being more volatile than that of rue.

# GUMMI et RESINA ALOES. Gum and refin of aloes. Lond.

Boil four ounces of Socotorine aloes in two pints of water, till as much as possible of the aloes is dissolved. The solution suffered to rest for a night, will deposite the resin to the bottom of D d 4 the vessel; after which, the remaining liquor, strained, if needful, is to be evaporated; that the gum may be left.

The gum of aloes is somewhat less purgative, and considerably less disagreeable than the crude juice. This alteration is not owing, as might be supposed, to the feparation of the refin; for the pure refin of aloes is still less difagreeable, and lefs purgative, even than the gum; fome have denied that it has any purgative virtue at all, and others ascibe to it an astringent quality. I have exhibited this refin, divided by trituration with the testaceous powders, in the dose of a scruple, without observing any effect from it (see page 81.) The gum seems to be the only part here intended for medicinal use: if the refin is required, it ought to be further purified by folution in spirit of wine; for as it is obtained by precipitation from an aqueous folution of impure aloes, all the impurities of the drug, that are not foluble in water, will precipitate along with it.

PILULÆ, feu EXTRACTUM, RUDII.

The pills or extract of Rudius. Edinb.

Take of

Black hellebore roots,

Colocynth,

Socotorine aloes, each two

ounces:

Scammony, one ounce;
Vitriolated tartar, two drachms;
Distilled oil of cloves, one drachm.
Bruise the colocynth and hellebore, pour on them two quarts of water, and boil to the con-

fumption of half the liquor: pass the decoction through a strainer, and evaporate it to the consistence of honey, adding the aloes and scammony, reduced into fine powder: when the mass is taken from the sire, mix into it the vitriolated tartar, and distilled oil.

This preparation is a medicine of great importance as a eathartic, similar to one described hereaster in page 413, under the title of Extractum catharticum. Water appears to be a better mentruum, than spirituous liquors, both for the colocynth and the hellebore; the watery extracts being much tess irritating than the spirituous, though not perhaps less effectual purgatives.

ROB BACCARUM JUNIPERIOR Rob of juniper berries.

Let juniper berries, thoroughly bruifed, be boiled in a fufficient quantity of water, the liquor strained, and inspissated to the consistence of honey.

This preparation may be made also from the decoction that remains after the distillation of the essential oil of the berries. It has a sweet balsamic taste, accompanied with a greater or less bitterness, according as the feeds of the berry were more or less thoroughly bruised. This elegant preparation, though not received in our pharmacopæias, feems not unworthy of a place in the shops. Hossman has a great opinion of it in debilities of the stomach and intestines, and in the difficulties of urine familiar to persons of an advanced age.

### ["409]

#### ECT. III.

### Extracts with rectified spirit.

DECTIFIED Spirit of wine disfolves the effential oils and refins of vegetables, and does not readily carry off the oil in its exhalation; the heat sufficient to exhale pure spirit being much less than that in which water confiderably evaporates, or most effential oils diffil. Hence a refinous or spirituous extract of wormwood, contrary to that made with water, contains the warmth and flavour, as well as bitterness, of the herb; one made from cinnamon possesses its aromatic virtue, as well as its aftringency; and one from lavender and rofemary flowers retains great part of their flavour and virtues; the volatile parts, which are carried off by water in its evaporation, being left behind by spirit.

fpirit employed for this The purpole should be perfectly free from any ill flavour; which would be communicated, in part, to the preparation; and from any admixture of phlegm or water, which would not only vary its diffolving power, but likewise, evaporating towards the end of the inspissation, would promote the diffipation of the volatile parts of the subject. Hence also the subject itself ought always to be dry: thole substances, which lose their virtue by drying, lofe it equally on being submitted to this treatment with the purest spirit.

The inspissation should be performed, from the beginning, in the gentle heat of a water-bath. is not needful to fuffer the spirit to evaporate in the air; greatest part of it may be recovered by collecting the vapour in the common distilling vessels. (See chap. v.) If the distilled spirit is found to have brought over any flavour from the subject, it may be advantageoully referved for the same purposes

again.

It is observable, that though reclified spirit is the proper menstruum of the pure volatile oils, and of the groffer refinous matter of vegetables, and water of the mucilaginous and saline; yet these principles are, in almost all plants, so intimately combined together, that whichever of these liquors is applied at first, it will take up a portion of what is directly foluble only in the other. Hence fundry vegetables, extremely refinous, and whose virtues consist chiefly in their refin, afford nevertheless very useful extracts with water, though not equal to those which may be obtained by a prudent application of Hence, also, the extracts foirit. made from most vegetables by pure spirit are not mere refins; a part of the gummy matter, if the subject contained any fuch, being taken up along with the refin, an admixture of great advantage to it in a The spirituous medicinal view. extracts of several vegetable substances, as mint leaves, rhubarb. faffron, dissolve in water as well as in spirit.

Pure refins are prepared by mixing, with spirituous tinctures of very refinous vegetables, a quan-tity of water. The refin, incapable of remaining dissolved in the watery liquor, feparates and falls to the bottom; leaving in the

menstrum such other principles of the plant, as the spirit might have extracted at first along with it.

# RESINA JALAPPÆ. Refin of jalap. Edinb.

Take any quantity of jalap root very well bruised. Pour upon it fo much rectified spirit of wine as will cover it to the height of four fingers; and digest them together in a fand-heat, that the Spirit may extract the virtue of Filter the tincture the root. through paper, put it into a glass cucurbit, and distil off one half of the spirit. Add to the remainder a proper quantity of water, and the refin will precipitate to the bottom: divide it into little cakes, and dry it with a very gentle heat.

This preparation is a pure refin; fuch gummy parts as the spirit might have taken up, remaining suspended in the liquor. indiffolubility in any aqueous fluid, and its tenacious quality, by which it adheres to the coats of the inteftines, and occasions great irritation and gripes, forbid its being evergiven by itself. It is fitted for use, by thoroughly, triturating it with testaceous powders; by grinding it with almonds or powdered gum, and making the compound into an emuliion with water; or by diffolving it in spirit of wine, and mixing the folution with a proper quantity of fyrup, or of mucilage. Six or eight grains, managed in either of these ways, prove powerfully cathartic, and generally without griping or greatly difordering the body.

It has been faid, that refin of jalap is frequently adulterated with

common rosin; and that this abuse may be discovered by spirit of wine, which dissolves the sormer, without touching the latter. This criterion, however, is not to be relied on; for there are many cheap resins which are soluble in spirit of wine as well as that of jalap; and there is not any one which may not be artfully rendered so.

# RESINA SCAMMONII. Refin of scammony. • Edinb.

This resin is prepared in the same manner as the preceding; with which it agrees also in its general qualities; occasioning vehement gripes if taken by itself, and operating generally with sufficient safety when properly divided. Scammony is doubtless a valuable purgative; but what advantage there is in thus separating the purgative resin from its natural corrector, the gummy part, is not so clear.

# RESINA GUAIACI. Refin of guaiacum. Edinb.

This resin is prepared in the same manner as the two preceding. either from the wood of guaiacum, or from what is called gum guaiacum: is is obtained most commodiously from the latter.

The virtue of guaiacum confifts wholly in its refin; and the refin of the wood, and of the gum so called, is perfectly one and the same; the gum being the natural exudation from the tree. If this exudation could be had pure, there would be no occasion for any artificial preparation of this kind; but it always contains a large proportion of earthy matter, so as to stand greatly in need of this method.

Ġ

Ċ

1

Ì

t

¢

5

ļ

ŝ

¢

ĮĴ

Ę

0

ı

ţ

5

į

f

ŝ

thod of purification. Sixteen ounces of the best gum guaiacum do not yield above twelve ounces of pure resin. The same quantity of the wood yields about three ounces, more or less according to its goodness. The bark is somewhat less resinous than the wood.

# RESINA CORTICIS PERUVIANI. Refin of Peruvian bark. Edinb.

This refin is made in the fame manner as the foregoing, and proves an elegant preparation of the bark, much stronger in taste than the watery extract described in the preceding section; it is nearly equivalent to about ten times its quantity of the bark in There does not, howfubstance. ever, appear to be any advantage in separating the pure refin by the addition of water, either in this or in the other articles. In regard to the bark particularly, it is more adviseable to endeavour to unite into one compound all that can be extracted from it by watery and spirituous mentirua; and accordingly the Edinburgh college has received a preparation of this kind, which is described in the following fection.

Extract of Saffron.

Extract of Saffron.

Pharm. Brandenburg.

Digest saffron in such quantities of pure spirit of wine, so long as the spirit extracts any colour from it. Mix the several tinctures together, and distil off the spirit, in a tall glass wessel, by the heat of a water-bath, till the residuum appears of the consistence of oil or bassam.

This is a general process, for the preparation of extracts from aromatic and other odorous fubstances: which extracts have been commonly distinguished by the name of effential, for the same reason that the volatile oils are fo called, their retaining the specific odour and flavour of the subjects. In making the extracts of this class, the inspissation should never be carried much lower than the confisence above directed: for when the matter has become thick, the spirit exhales more difficultly than before, and is more apt to carry off with it some of the volatile parts. If the preparation is wanted in a folid or consistent form, it is more advisable to mix with it a fuitable quantity of any appropriated powdery matters, than to hazard the the loss of its virtue by a further evaporation. If any addition is wanted for giving confishence to the extract of fastron, fastron in substance appears to be the best.

The essential extract of saffron is an elegant and high cordial. Boerhaave says, it possesses such exhilarating virtues, that, if used a little too freely, it occasions an almost perpetual and indecent laughing: he observes, that it tinges the urine of a red colour: and that it mingles with water, spirit, and oils, but is most conveniently taken in a glass of Canary or other rich wine. A few drops are a sufficient dose. The distilled spirit contains also some share of the virtue of the faffron, though far less than the extract: it is said to have an advantage above most other cordial spirits, of disposing the patient to fweat; it may be taken, properly diluted, from a drachm to half an ounce.

SECT.

### SECT. IV.

### Extracts with spirit and water.

THERE are fundry vegetables, particularly those of a refinous nature, which are treated, to better advantage, with a mixture of water and spirit, than with either of them fingly. virtues of refinous woods, barks, and roots, may indeed be in great part extracted by long boiling in fresh portions of water; but at the same time they suffer a confiderable injury from the continued heat necessary for the extraction, and for the subsequent evaporation of so large a quantity of the fluid. Rectified spirit of wine is not liable to this inconvenience; but the extracts obtained by it, from the substances here intended, being almost purely refinous, are less adapted to general use than those in which the refin is divided by an admixture of the gummy matter, of which water is the direct menstruum.

There are two ways of obtaining these compound or gummy-refinous extracts: one, by using proof spirit, that is, a mixture of about equal parts of spirit and water, for the menstruum : the other, by digefting the fubject first in pure spirit and then in water, and afterwards uniting into one mass the parts which the two menstiua have separately extracted. In some cases, where a fufficiency of gummy matter is wanting in the fubject, it may be artificially supplied, by inspiffating the spirituous tincture to the confistence of a balfam, then thoroughly mixing with it a thick folution of any simple gum, as mucilage of gum Arabic, and

exficcating the compound with a gentle heat. By this method are obtained elegant gummy-refins, extemporaneously miscible with water into milky liquors.

EXTRACTUM JALAPII.

Extract of jalap.

Lond.

Upon powdered jalap pour some rectified spirit of wine, and with a gentle heat extract a tincture: boil the remaining jalap in fresh parcels of water. Strain the first tincture, and draw off the spirit, till what remains begins to grow thick: boil the strained decortion also to a like thickness: then mix both the inspissance in the spissance of the with a gentle fire reduce the whole to a pilular consistence.

#### Edinb.

Take any quantity of jalap root, very well bruifed, pour upon it as much rectified spirit of wine as will cover it to the height of four fingers, and digett them together in a fand-heat : pour off this tincture, and put to the remaining magma a fufficient quantity of water : boil them together for an hour : then pass the decoction through a strainer, and afterwards evaporate it to the confishence of honey, mixing in, toward the end of the evaporation, the spirituous tincture, and keeping them continually flirring, that the whole may be reduced into an uniform mass free from From the spirituous tincture great part of the spirit

may be recovered by diffillation, before its admixture with the watery decoction.

This extract is an uleful purgative, preferable to the crude root, as being of more uniform ftrength, and as the dose, by the rejection of the woody parts, is rendered smaller: the mean dose is twelve grains. If the spirituous tincture was inspissated by itself, it would afford a refinous mass, which, unless thoroughly divided by proper admixtures, occasions violent griping, and yet does not prove sufficiently cathartic; the watery decoctions yield an extract, which operates exceeding weakly: both joined together, as in this preparation, compose an effectual and fafe purge. This method of making extracts might be advantageously applied to fundry other rolinous substances, as the dry woods, roots, barks, &c. A fmall quantity of spirit takes up the refin, and much less water than would otherwise be necesfary extracts all the other foluble Darts.

In the preceding edition of the Edinburgh parmacopæia, a little fixt alkaline falt was ordered to be added to the water in which the jalap is boiled after the action of fpirit; on a supposition, that this would enable the water to extract more from the root than it could by itself. But, so far as the quantity of the alkaline falt could go, it had the opposite effect; impending the action of the water. The refinous parts of the jalap are dissolved by the spirit: and little other than the gummy matter remains for water to extract. Now, if pure gum Arabic be put into water along with any alkaline falt, the falt will render the water incapable of dissolving.

the gum: if the gum be diffolved first, the addition of any alkaline falt will precipitate it.

## EXTRACTUM CORTICIS PERUVIANI.

Extract of Peruvian bark.
Edinb.

The college of Edinburgh has directed the extract of bark to be made with water and spirit in the same manner as the preceding. In the bark we may distinguish two kinds of taftes, an aftringent and a bitter one; the former of which feems to refide in the refinous matter, and the latter chiefly in the gummy. The watery extract (described in page 404) is moderately strong in point of bitterness, but of the affringency it has only a small degree. pure resin, on the other hand, (page 411.) is strong in astringency, and weak in bitterness. Both qualities are united in the present extract; which appears to be the best preparation of this kind that can be obtained from this valuable drug.

### EXTRACTUM LIGNI CAM-PECHENSIS.

Extrad of logwood. Edinb.

This extract is directed in the Edinburgh pharmacoposia to be prepared as the foregoing; and the same treatment is judiciously ordered for all the resinous drugs in general.

### EXTRACTUM CATHARTI-CUM.

Cathartic extract.

Take of
Socotorine aloes, an ounce and
a half:

Colocynth, fix drachms;

Scam -

Scammony, Leffer cardamoms, husked, each half an ounce;

Proof spirit, one pint.

Having cut the colocynth fmall, and bruised the seeds, pour on them the vinous spirit, and digest with a gentle heat for four days. Press out the tincture, and dissolve therein the aloes and scammony, first seperately reduced to powder; then draw off the spirit, and inspissate the remaining mass to a pilular confiftence.

THIS composition answers very effectually the intention expressed in its title, so as to be relied on in cases where the patient's life depends on its taking place: the dose is from fifteen grains to half The proof spirit is a a drachm. very proper mentiruum for the purgative materials; dissolving nearly the whole substance of the aloes and scammony, except the impurities; and extracting from the colocynth, not only the irritating refin, but great part of the gummy matter. The purgative virtue of this last article appears indeed to be fufficiently got out by water; and the watery extract to operate with greater mildness than that with proof spirit, though in general effectually: the Edinburgh college have accordingly preferred water, in making a preparation, of the same intention with this described in page 408. In our former pharmacopæias, three spices were employed in this composition, cinnamon, mace, and cloves: the cardamom feeds, now introduced, are preferable, on account of their aromatic matter being of a less volatile nature; though a considerable part of the flavour, even of these, is disfi-

the phlegmatic part of the proof ipirit.

CONFECTIO CARDIACA. Cordial confection. Lond.

Take of

Rosemary tops, fresh, Juniper berries, each one pound: Leffer cardamom seeds, husked, Zedoary,

Saffron, each half a pound. Extract a tincture from these ingredients with about a gallon and a half of proof spirit: the tincture be strained off, and reduced by a gentle heat to the weight of about two pounds and a half; then add the following ingredients very finely pulverized, and make the whole into an electary.

Compound powder of claws, fixteen ounces; Cinnamon, Nutmegs, each two ounces;

Cloves, one ounce; Double-refined fugar, two pounds.

This confection is composed of the more unexceptionable ingredients of a composition formerly held in great efteem, and which was called, from its author, ConfectioRaleighana. The committee, appointed for reforming the London pharmacopæia, observe, that the original consection is composed of no less than five and twenty particulars; each of which they examined apart, except one, ros folis, the flower of which is too fmall to be gathered in sufficient quantity for the general use of the medicine, and the plant is possessed of hurtful qualities, as is experienced in cattle that feed where it grows. In this examination, many of the extracts pated during the evaporation of came out so very nauseous, that

1

Ī

1 : • it was impossible to retain them, tention from eight or ten grains confistent with any due regard to the taste of the composition. But some few of equal efficacy with any of the rest, being of a tolerable taste and flavour, were compounded in different proportions; and when after many trials, a compolition was approved, the quantity of each material, that would yield the proportion of extract which entered that composition, was calculated, and from thence the proportions collected as now fet down: after which the compound extract was made, and found to answer expectation.

The confection, as now reformed, is a sufficiently grateful, and moderately warm cordial; and frequently given in that into a scruple or upwards, in boluses and draughts. The extract retains a confiderable share of the flavour and virtue of the ingredients, though not near fo much as if a rectified spirit had been employed. The operator should be particularly careful to extract as much from the ingredients as the fpirit will take up; otherwise the inspissated matter turns out, so thin. and of so little tenacity, that the powders are apt to feparate and fublide from it in keeping. The crabs claw powder does not appear to be very necessary, and is inserted rather in compliance with the original, than from its contributing any thing to the intention of the medicine.

### EECT. V.

### Extracts by long digeftion.

N the foregoing part of this a particular account of an extract chapter it has been observed, of opium prepared on this principle; that the virtues of vegetable decoctions are altered by long boiling. Decoctions or infusions of drastic vegetables, by long continued boiling or digestion, lose more and more of their virulence; and at the same time deposite more and more of a gross sediment, resulting probably from the decomposition of their active parts. On this foundation it has been attempted to obtain safe and mild preparations from fundry virulent drugs, and some of the chemists have strongly recommended the process, though without specifying, or giving any intimation of, the continuance of boiling requifite for producing the due mildness in different subjects. M. Baume, in his Elemens de pharmacie lately published, has given

the substance of which is as follows.

Extract of opium prepared by long digestion.

Let five pounds of good opium, cut in pieces, be boiled about half an hour, in twelve or fif-teen quarts of water: strain the decoction, and boil the remainder once or twite in fresh water, that so much of the opium as is diffoluble in water may be got out. Evaporate the strained .. decoctions to about fix quarts, which being put into a tin cucurbit, placed in a sand-bath. keep up such a fire, as may make the liquor nearly boil, for three months together if the fire is continued day and night, and for fix months, if it is intermitted in the night; filling up the veffel with water in proportion to the evaporation; and scraping the bottom with a wooden spatula from time to time, to get off the sediment which begins to precipitate after some days digestion. The sediment needs not be taken out till the boiling is sinished, at which time the liquor is to be strained when cold, and evaporated to an extract, of a due consistence for being formed into pills.

THE author observes, that by keeping the liquor strongly boiling, the tedious process may be confiderably expedited, and the fix months digestion reduced to four months; that in the beginning of the digestion, a thick, viscous. oily matter rifes to the top, and forms a tenacious skin as the liquor cools; this is supposed to be analagous to effential oils, though wanting their volatility: that the oil begins to disappear about the end of the first month, but still continues fensible till the end of the third, forming oily clouds as often as the liquor cools: that the refin at the same time settles to the bottom in cooling, preserving for a long while its refinous form. but by degrees becoming powdery, and incapable of being any longer foftened, or made to cohere by the heat; that when the process is finished, part of it still continues a perfect resin, dissoluble in spirit of wine, and part an indisfoluble powder: that when the digested liquor is evaporated to about a quart, and fet in the cold till next day, it yields a brownish earthyfaline matter, called the effential falt of opium, in figure nearly like the fedative falt obtained from borax, intermingled with small

needled crystals. He gives an account of his having made this preparation fix or feven times, veffel he made use of was about two inches and a half diameter in the mouth: the quantity of water evaporated was about twenty-four ounces a day, and from a hundred and thirty to a hundred and forty quarts during the whole digestion. Out of fixty-four ounces of opium, seventeen ounces remained undiffolved in the water: the quantity of refinous matter, precipitated during the digestion, was twelve ounces: from the liquor, evaporated to a quart, he obtained a drachm of effential falt, and might, he fays, have separated more: the liquor being then further evaporated to a pilular confistence, the weight of the extract was thirtyone ounces.

It is supposed, that the narcotic virtue of opium refides in the oily and refinous parts; and that the gummy extract, prepared by the above process, is endowed with the calming, fedative, or anodyne powers of the opium, divefted of the narcotic quality as it is of the fmell, and no longer productive of the disorders, which opium itself, and the other preparations of it, frequently occasion. A case is mentioned, from which the innocence and mildness of the medicine are apparent; fifty grains having been taken in a day, and found to agree well, where the common opiate preparations could not be born. But what there it possesses of the proper virtues of opium, is not so clear; for the cure of convultive motions of the stomach and vomitings, length happened after extract had been continued daily in the above doses for several years (plusieurs annees) cannot perhaps

perhaps be ascribed fairly to the medicine.

If the theory of the process, and of the alteration produced by it in the opium, is just; a preparation equivalent to the above may be obtained in a much shorter time. If the intention is to separate the refinous and oily parts of opium, they may be separated, by means of pure spirit of wine, in as many hours as the digestion requires months. The separation will also be as complete, in regard to the remaining gum, though some part of the gum will in this method be loft. a little of it being taken up by the spirit along with the other principles.

In what particular part of opium

its peculiar virtues refide, has not perhaps been incontestably ascertained; but thus much seems clear from experiment, that the pure gum, freed from all that spirit can dissolve, has little, or rather nothing of its soporific power.

There are grounds also to prefume, that by whatever means we destroy or diminish what is called the narcotic, soporific, virulent quality of opium, we shall destroy or diminish likewise its salutary operation. For the ill effects, which it produces in certain cases, seem to be no other, than the necessary consequences of the same power, by which it proves so beneficial in others.

### [ 418 ]

### CHAPTER VII.

### Empryeumatic Oils.

TEGETABLE and animal fubstances, and mineral bitumens, on being urged with a red heat, have their original properties destroyed, and are resolved or changed into products of a different nature from what præexisted in the subject. By burning them in the open air, a part is changed into ashes, a part into foot, and a part is diffolved by the air. posed to the fire in close veffels (as in those called retorts, having receivers adapted to them for detaining the volatile parts) they are refolved into fetid oils, and different kinds of faline fubitances, which rise into the receiver : and a black coal, which remains behind, and which, though no farther alterable in close vessels, on admitting air burns into white ashes. The oils. called from their fetid burnt smell, empyreumatic, are the objects of the prefent chapter. Some of these however being obtained in the fame process with certain saline bodies of more importance than themselves, are referred to the head of faline preparations.

OLEUM BUXI.

Oil of box.

Lond.

Diffil pieces of box wood, in a retort, with a fand-heat gradually increased: the oil will come over along with an acid spirit, which is to be separated by a funnel.

OLEUM GUAIACI.
Oil of guaiacum.
Edinb.

Put any quantity of chips of guaiacum into an earthen longneck, or a glass retort, and distil either in a sand bath or an open fire, increasing the heat by degrees. At first an acid liquor will come over, afterwards a light red oil, and at length, in the utmost degree of fire, a thick black oil, which sinks through the other liquors to the bottom of the receiver.

Oils may be obtained after the fame manner from every kind of wood.

THE retort may be filled almost up to the neck with chips or small pieces of box or guaiacum, the refuse of the turner. Lute on a glass receiver with a paste made of linfeed meal and water: fet the retort on the bottom of a deep iron pot, with a little fand under it; and fill up the space, betwixt it and the fides of the pot, with more fand. Apply at first a gentle fire, and gradually increase it to the utmost that the furnace is capable of giving. Particular care must be had not to raise the heat too fast when the first reddish oil begins to come over; for at this time, a large quantity of elastic vapour is extricated from the wood, which, if the fire is urged, or if it is not allowed an exit, will burft the veffels: when the distillation is finished, and the veffels grown cool, unlute the receiver, and separate the oil from the acid liquor; the method of performing this by the funnel, as directed in the first of the above processes, is as follows. Pour the leveral liquors into a glass funnel, whose stem is stopt by the singer, the ponderous black oil finks lowermoft >

ţ

ermost: suffer this to run out: then close the stem again, and afterwards separate the acid siquor from the lighter oil in the same manner. They are more perfectly separated, by pouring them into a hollow cone of siltering paper moistened with water, and placed in a funnel: the acid siquor passes thro, and the oil remains on the paper.

The oils obtained by this treatment from different woods and plants are nearly of the same qualities: they have all a very difagreeable acrid taste, and a burnt stinking smell; without any thing of the peculiar slavour, taste, or virtues of the subject which afforded them. The present practice rarely employs those oils any otherwise than sor external surposes, as the cleansing of soul bones, for the tooth-ach, against some kinds of cutaneous eruptions, old pains and aches, and the like; and for these not very often.

# OLEUM LATERITIUM. Oil of bricks. Lond.

Heat bricks red hot, and quench them in oil olive, till they have foaked up all the oil: then break them into pieces small enough to he conveniently put into a retort; and diffil with a sand-heat gradually increased: an oil will arise, together with a spirit, which is to be separated from it as in the foregoing process.

THIS preparation has had a place in most dispensatories, under the pompous names of oleum philosophorum, fanctum, divinum, benedictum, and others, as improper as that under which it stands above. It is really oil olive, rendered strongly empyreumatic by heat: the spirit, so called, is no more than phlegm, or water, tainted with the burnt slavour of the oil. It has been celebrated for fundry external purposes, particularly against gouty and rheumatic pains, deafness and tingling of the ears, &c. and sometimes likewise given inwardly. But common practice seems to have now entirely rejected this loathsome remedy; and the college of Edinburgh have expunged it from their book.

### OLEUM PETROLEI BARBA-DENSIS.

Oil of Barbadoes tar.

Lond.

Diffil Barbadoes tar with a fandheat; as oil will arife, together with a fpirit, which is to be feparated from it.

Dr. Pemberton observes, that this oil will be more or less thin, according to the continuance of the distillation: that the tar will at last be reduced to a black coal, and then the oil will be pretty deep in colour, though perfectly fluid: that this oil has a property limitar to that of the tincture of nephritic wood in water, appearing blue when looked upon, but of an orange colour, when held betwixt the eye and the light. By long keeping I have observed it to lose this property. It is somewhat less disagreeable than the foregoing oils, tho' very acrid and flimulating.

OLEUM TEREBINTHINÆ ÆTHEREUM; & empyreumaticum five BALSAMUM. The ethereal oil of turpentine, and the empyreumatic sil or balfam.

Distil the effential oil of turpentine in a retort, with a very gentle fire, until what remains has ac-E e 2 quired quired the confistence of a bal-

Balsam of turpentine may likewise be obtained from the yellow resin left after the distillation of the essential oil: upon distilling this in a retort, at first a portion of thin oil arises, which is to be kept by itself, and asterwards a thick balsam: there remains in the retort a blackish resin, called colophony.

#### Edinb.

Melt any quantity of turpentine, over a gentle fire, and pour it into a glass retort, of which it may fill one half: then lute on a receiver, and distil in a fandbath. Apply at first a gentle heat, upon which an acid spirit will come over, and on gradually increasing the fire, a limpid oil, commonly called ethereal spirit of turpentine; at length a yellow oil will arise. In the bottom of the retort, there remains a refinous mass, called colophony : which, if still farther urged with successive degrees of heat to the highest, gives first a red oil, and afterwards a darker coloured one, which finks through the other liquors to the bottom of the receiver.

THESE processes are tedious, and accompanied with a good deal of danger; for unless the luting is very close, some of the vapour will be apt to get through, which if it catches fire, will infallibly burft The oil here called the vessels. ethereal does not confiderably differ in specific gravity, smell, taste, medical qualities, from the cheaper one obtained by the addition of water in the common still: nor are the empyreumatic thin oil and balfam of any great efteem in practice.

OLEUM COPAIVÆ COMPOSI-TUM,

Compound oil of balfam of Copaiva.

Lond.

Take two pounds of balfam of Copaiva, and four ounces of gum guaiacum. Diftil them in a retort, continuing the operation till a pint of oil is come over.

This mixture, undistilled, proves a medicine of confiderable efficacy in rheumatic cases, &c. In diftillation, the guaiacum gives over little, serving chiefly for the same purpose that bricks do in the oleum fateritium. The balfam diftilled in a retort, with or without the gum, yields first a light coloured oil, smelling confiderably of the subject; this is immediately followed by a darker coloured oil, and afterwards by a blue one, both which have little other smell than the empyraumatic one that diffinguishes the oils of this class: their tafte is very pungent and acrimonious. This balfam distilled with water yields as much essential oil, as above of empyreumatic.

# OLEUM CERÆ. Oil of wax. Edinb.

Melt yellow bees wax with twice its quantity of fand, and distil in a retort placed in a fand furnace. At first an acid liquor arises, and afterwards a thick oil, which sticks in the neck of the retort, unless it be heated by applying a live coal. This may be reclined into a thin oil, by distilling it several times, without addition, in a fand heat.

BOERHAAVE directs the wax, cut n pieces, to be put into the retort first, so as to fill one-half of it; when as much fand may be poured there-

thereon as will fill the remaining half. This is a neater, and much less troublesome way, than melting the wax, and mixing it with the fand, before they are put into the The author above-mentioned greatly commends this oil against roughness and chaps of the ikin, and other like purposes; the college of Strasburg speak also of its being given internally, and fay it is a powerful diuretic (ingens diureticum) in doses of from two to four and more drops; but its difagreeable smell has prevented its coming into use among us.

BALSAMUM ANODYNUM, vulgo GUIDONIS. The anodyne, commonly called Gui-

'I be anodyne, commonly called Gusdo's balfam.

*Edinb*. Take of

Take of
Tacamahaca, in powder,
Venice turpentine, each equal
parts.

Put them into a retort, whereof they may fill two-thirds, and diftil with a fire gradually increafed. Separate, according to art, the red oil, or balfam, from the liquor that swims above it.

THES oil is supposed to be anodyne and discutient. In foreign pharmacopæias, and in the preceding editions of the London and Edinburgh, oils are directed to be distilled in the same manner from different refinous and gummy refinous bodies separately, as tacamahaca, storax, ammoniacum, galbanum, sagapenum, &c. but it does not appear that they are materially different from one another in regard to their external use, which is the only intention in which they have been employed. The above compolition has loft one of its former ingredients, galbanum, without the least injury to its virtue.

OLEUM ANIMALE DIPPELII.

Dippel's animal oil.

Take any quantity of the empyreumatic oil distilled from animal substances, as that of hartshorn, (the preparation of which is dedescribed along with that of the volatile salt and spirit, in the following chapter.) Put it into a glass retort, and having fitted on a receiver, distil in a sand-heat : the oil will arise paler coloured and less fetid; and a black coaly matter will remain behind. Repeat the distillation in fresh retorts, till the oil ceases to leave any feces, and till it loses its ill fmell, and acquires an agreeable

THE quantity of oil employed in this process should be considerable: for it leaves fo much black matter behind in the several distillations, that it is reduced, at last, to a small portion of its original quantity. The distillation must be repeated, at least, twelve times, and frequently the requisite subtilization will scarcely be obtained with less than twenty distillations. It is faid, that the effect may be expedited, by mixing the oil with quicklime into a fost paste; the lime keeping down more of the gross matter, than would remain without fuch an addition.

Animal oils thus rectified, are thin and limpid, of a subtile, penetrating, not disagreeable smell and taste. They are strongly recommended as anodynes and antispassmodics, in doses of from sitteen to thirty drops. Hossman reports, that they procure a calm and sweet sleep, which continues often for twenty hours, without being followed by any languor or debility, but rather leaving the patient more alert and chearful

E e 3 than

than before: that they procure likewife a gentle fweat, without increasing the heat of the blood: that given to twenty drops or more, on an empty stomach, six hours before the accession of an intermittent sever, they frequently remove the disorder, and that they are likewife a very generous remedy in inveterate and chronical epilepsies, and in convulsive motions, especially if given before the usual time of the attack, and preceded by proper evacuations.

The empyreumatic oils of yegetables, rectified in the fame manner by repeated diffillations, fuffer a like change with the animal: their losing their dark colour and offensive smell, and becoming limpid, penetrating, and agreeable: in this fact they are supposed, like the animal oils, to be anodyne, antispassmodic, and disphoretic, or sudorific. It is observable, that more all the empyreumatic oils dissolve ness.

in spirit of wine, and that the oftener they are rectified or redistilled, they dissolve the more readily; a circumstance in which they differ remarkably from effential oils, which, by repeated distillations, become more and more difficult of solution.

How far these preparations really possess the virtues, that have been ascribed to them, has not yet been sufficiently determined by experience; the tediousness and trouble of the reclification having prevented their coming into general use, or being often made. They are liable also to a more material inconveniency in regard to their medicinal use, precariousness in their quality: for how perfeelly foever they be reclified, they gradually lose, in keeping, the qualities they had received from that process, and return more and more towards their original fetid-



Salts and Saline preparations.

### SECT. I.

### Fixt alkaline salts

HE ashes of most vegetables, steeped or boiled in water, give out to it a saline substance, separable in a solid form by evaporating the water. This kind of salt never præexists in the vegetable, but is always generated during the burning. It is called sixt alkaline salt.

# SA TARTARI. Salt of tartar. Lond.

Let any kind of tartar be wrapt up in strong brown paper, first made wet, or included in a proper vefsel, and exposed to the fire; that its oil may be burnt out: then boil it in water, filter the solution, and evaporate it, till there remains a dry salt, which is to be kept in a vessel closely stopt.

#### Edinb.

Wrap up any quantity of white tartar in wetted paper, and calcine it in a reverberatory furnace till it becomes exceedingly white. Then diffolve it in warm water, filter the folution, and evaporate it in a clean iron veffel, till a falt is left behind, perfectly dry, and white as fnow; observing towards the end of the operation to keep the matter continually fitring with an iron ladel, to prevent its sticking to the bottom of the vessel.

If a fironger sait of tartar is required, let the white sait be melted in a crucible, with the most intense degree of heat, and reverberated for some hours, till it has acquired a greenish or blue colour.

THE white and red forts of tartar are equally fit for the purpose of making fixt salt; the only difference is, that the white affords a somewhat larger quantity than the other; from fixteen ounces of this sort, upwards of sour ounces of fixt alkaline salt may be obtained. The use of the paper is to prevent the smaller pieces of the tartar from dropping down into the ash hole, through the intestices of the coals, upon first injecting it into the surrance.

The calcination of the falt (if the tartar was sufficiently burnt at first) does not increase its strength, fo much as is supposed; nor is the greenish, or blue colour any certain mark either of its strength, or of its having been long exposed to a vehement fire; for if the crucible is perfectly clean, close covered, and has stood the fire without cracking, the falt will turn out white, though kept melted and reverberated ever so long; whilst, on the other hand, a flight crack happening in the crucible, or a spark of coal falling in, shall in a few E e 4 minutes

minutes give the falt the colour ad-The colour, in effect, is mired. a mark rather of its containing some inflammable matter, than of its strength.

This falt has a pungent hery taste; and occasions in the mouth a kind of urinous flovour, probably from a resolution, which it produces in the faliva. It readily difsolves in water, and deliquiates in the air, but is not acted upon by pure vinous spirits. Instead of being dissolved by vinous spirits, if a faturated folution of it in water be dropt into the pure spirit, it will not mix therewith, but fall distinct to the bottom; if water be mixed with the spirit, the addition of fixt alkaline falt will imbibe the water, and form with it, as in the other cafe, a distinct fluid at the bottom: this property affords a commodious method of dephlegmating vinous fpirits, or separating their watery part, as we have already feen in page 382.

Salt of tartar, or folutions of it in water, raile an effervelcence on the admixture of acid liquors, and destroy their acidity, the alkali and acid uniting together into a compound of new qualities called neutral: earthy fubstances, and most metallic bodies, previously disfolved in the acid, are precipitated from it by the alkali. The alkaline falt changes the colours of the blue flowers of plants, or their infusions, to a green: it has the same effect on the bright red flowers and on the colourless infusions of white ones; but in many of the dark red, as those of the wild poppy, and of the yellow ones, it produces

no fuch change.

Solutions of this falt liquely all the animal juices, except milk;

corrode the fleshy parts into a kind of mucous matter; concrete with animal fats and vegetable oils, into foap : and dissolve sulphur into a red liquor; especially if affished by a boiling heat, and mingled with quicklime, which greatly promotes their activity. On pure earths and stones, these liquors have no fensible action; but if the earth or stones be mixed with four or five times the weight of the dry falt, and urged with a strong fire, they melt along with it, and become afterwards perfectly soluble both in water and by the moisture of the air: with a smaller proportion of the salt, as an equal weight, they run into an in-

dissoluble glassy matter.

The medical virtues of this falt are, to attenuate the juices, resolve obstructions, and promote the natural secretions. A dilute solution of it, drank warm in bed, generally excites fweat; if that evacuation is not favoured, its sensible operation is by urine. It is an excellent remedy in costive habits. especially if a few grains of aloes be occasionally interposed; with this advantage above other purgatives and laxatives, that when the complaint is once removed, it is not apt to return. Where acidities abound in the first passages, this falt absorbs the acid, and unites with it, into a mild aperient neutral falt. As one of its principal operations, is to render the animal fluids more thin, it is obvious, that where they are already colliquated, as in survies, and in all putrid diforders in general, this medicine is improper. The common dose of the falt is from two or three grains to a scruple; in some circumstances it has been extended to a drachm, in which case it must always be largely diluted with watery liquors.

SAL ABSINTHII. Salt of wormwood. Edinb.

Let any quantity of wormwood. either fresh gathered, or moderately dried, be put into an iron pan, and, with a gentle fire, reduced into white ashes. these with a fufficient quantity of fpring water, filter the liquor, and evaporate it till a dry salt is left behind; this proves of a brown colour; by repeated folution, filtration, and inspissation, it becomes at length pure and white.

IT is generally expected of a brown colour in the shops, and distinguished by this mark from the purer alkali of tartar. If required to be white, the means above recommended will scarcely render it fo; the remains of the oil of the plant, on which the brown colour depends, not being effectually separable without strong calcination. If the ashes have been fully calcined before the affusion of water, the falt will turn out white at once.

#### Lond.

Let the ashes of wormwood [which the shops are usually supplied with from the country] be put into an iron pot, or any other convenient vessel; and kept redhot over the fire for fome hours. after flirring them, that what oily matter remains may be burnt out; then boil the ashes in water, filter the ley through paper, and evaporate it till a dry falt remains; which is to be kept in a vessel close stopt.

After the same manner a fixt alkaline falt may be prepared from all those vegetables which yield this kind of falt [L.] as bean stalks, broom, &c.  $\{E_i\}$ 

THESE faits are obtained to greater advantage from dry plants than from green ones; they must not however be too dry, or too old: for in fuch case, they afford but a finall quantity of falt. The fire should be so managed, as that the subject may burn freely, yet not burst into violent flame; this last circumstance would greatly lessen the yield of the falt, and a very close smothering heat would have this effect in a greater degree: hence the ashes of charcoal scarce yield any falt, whilft the wood it was made from, if burnt at first in the open air, affords a large quantity.

If the ashes are not calcined after the burning, a confiderable portion of the oil of the subject remains in them unconfumed; and hence the falt turns out impure, of a brown colour, and somewhat saponaceous. Tachenius, Boerhaave, and others. have entertained a very high opinion of these oily salts, and endeavour as much as possible to retain the oil in them. They are nevertheless liable to a great inconvenience, uncertainty, in point of strength, without promoting any advantage to counterbalence it : if the coinmon alkalies are required to be made milder and less acrimonious (which is the only point aimed at in the making of these medicated salts as they are called) they may be occafionally rendered to by fuitable additions. Pure alkalies, united with a certain quantity of expressed oil, compose (as we shall see hereaster) a perfect foap, in which the pungent tafte of the alkaline falt is totally suppressed: it is obvious, therefore, that on the same principle the pungency may be covered in part. and this proportionably to the quantity of oily matter combined. But we may obtain more elegantly, by a process described in the following page (under the title of Sal alkalinus falis marini) a perfectly pure white alkaline falt, of all the mildness that can be wished for.

. The shops were formerly burthened with a great number of these falts, which are now very judicioully rejected: those here retained being abundantly sufficient to anfwer all the useful purposes that can be expected from these kinds of preparations. All fixt alkaline falts, from whatever vegetable they may be obtained (those of certain marine plants excepted, which partake of fea falt or its alkali) are nearly one and the same thing, and not distinguishable from each other, at least in their effect as medicines; and hence the college of London. in most of the compositions wherein these forts of salts are ingredients, allow any fixt alkaline fait to be made use of.

Some differences indeed are obferved in them as usually prepared; but these depend upon the manner in which the process for obtaining them is conducted, or on some faline matters of a different kind, which either przezisted in the vegetable, or were produced in the burning, remaining mixed with the alkali. A variation in the heat by which the plant is burnt or calcined, occasions a difference in the acrimony of the produce: the more vehement and lasting the fire (to a certain degree) the more acrid is the falt. The circumstances of using the ashes fresh burnt, or after they have been long exposed to the air, and of applying the water hot or cold to the ashes, likewise make a confiderable variation. By long exposure to the air, even the alkalies that have been made cauftic by quicklime, lofe all the adventitious acrimony which they had received from that treatment; the chemists affirm, that they imbibe

alfo from the air, in a length of time, a portion of vitriolic acid, by which a part of them is converted into a neutral falt, the same with the tartarus vitriolatus of the shops; and it is certain, that such a falt is often found among the ashes of vegetables; though it does not, perhaps, arife from that origin. Boiling water takes up this neutral falt from the ashes; whilst cold water extracts from them only the pure alkaline falt, unless it be used in too large quantity, or suffered to fland too long upon them. Boiling water dissolves also more, than cold, of the oily parts of the subject, if any remained unconfumed.

# NITRUM FIXUM.

Take of

Powdered nitre, four ounces; Charcoal in powder, five drachms. Mix them thoroughly together, by rubbing them in a mortar, and inject the mixture, by a little at a time, into a red-hot crucible. A deflagration, or a bright flame with a hilling noise, happens on each injection; the whole quantity being thus deflagrated, continue the fire strong for half an hour.

NITRE is composed of the common vegetable fixt alkaline falt, and a peculiar acid. In this procels, the acid is destroyed or changed to another nature; and the remaining falt proves merely alkaline. not different in quality from the fal tartari, except that a very minute portion of the nitre generally remains unchanged; the falt is purified by folution in water, filtration, and evaporation. It may be observed, that the falt receives no fenfible addition from the vegetable coal employed for the deflagration; for the athes of charcoal have exceeding

ceeding little faline matter; and the quantity of charcoal above directed, yields only a grain or two of ashes.

SAL ALCALINUS SALIS MA-RINI.

The alkaline salt of sea salt.
Take of

Cubical nitre (prepared as hereafter described in sect. vi. of this chapter) four ounces; Charcoal, five drachms.

Mix and deflagrate as in the preceding process.

CUBICAL nitre is composed of the nitrous acid united with the alkaline basis of sea salt: the acid being here separated in the deflagration, that alkali remains nearly pure. It possesses the general properties of the foregoing preparation; changing blue flowers, green; diffolving oils, salts, and sulphur; bringing earths and stones into fusion, and forming with them, according to its quantity, either a vitreous, or a foluble compound; effervescing with acids, precipitating earths, and metals dissolved in them, and uniting with the acid into a neutral falt. It differs from the foregoing alkalies, in being much milder in taste; not so readily disfolving in water; not at all deliquiating in the air; cafily affuming, like neutral salts, a crystalline form; and yielding with each of the common acids, compounds very sensibly different, both in their form and qualities, from those which refult from the coalition of the vegetable alkalies with the respective acids. The crystals of this falt itfelf are prismatic, greatly refembling those of the salt called fal mirabile. (See the fection of neutral falts.) Exposed to a warm air, they fall into a porous, friable mass, and lose above two-thirds of their weight.

How far this falt differs in medical virtue from the other alkalines, is not well known. It apparently possesses the same general virtues; and, as it is far milder, may be given in more considerable doses.

A falt of the same nature with this, but less pure, as containing an admixture of the common vegetable alkali, is prepared at Alicant, and some other places, from the ashes of certain marine plants, called kali; which plants are supposed to have given rife to the name al-The falt of the kali plants is called foda or bariglia: it has been long used medicinally in France. and begins now to be introduced into practice in this country; but the above pure alkali extracted from sea salt is doubtless preferrable to it.

LIXIVIUM TARTARI [L.]
Liquamenfalis tartari; vulgo Oleum
tartari per deliquium [E.]

Ley of tartar.

Or oil of tartar per deliquium.

Lond.

Let tartar, calcined to whiteness, be set by in a moist place, that it may liquefy.

Edinb.

Put any quantity of falt of tartar in a flat glass dish, and expose it to the air, for some days, in a moist place; it will run into a liquor, which is either to be filtered thropaper, or separated from the seces by decantation.—The higher the falt has been calcined, the more readily will it relent in the air.

THE folutions of fixt alkaline falts, effected by exposing them to a most air, are generally looked upon as being purer than those made by applying water directly: for though the falt be repeatedly disfolved

dissolved in water, filtered and ex- ficcated: yet on being liquefied by the humidity of the air, it will still deposite a portion of earthy matter: but it must be observed, that the exficcated falt leaves always an earthy matter on being dissolved in water, as well as on being deliquiated in the air. Whether it leaves more in one circumstance than in the other, I have not examined. The deliquiated lixivium is said to contain nearly one part of alkaline falt to three of an aqueous fluid. It is indifferent, in regard to the lixivium itself, whether the white athes of tarrar, or the falt extracted from them, be used; but as the ashes leave a much greater quantity of earth, the separation of the ley proves more troublesome.

# LIŽIVIUM SAPONARIUM.

Lond.

Take of

Russia potash,

Quicklime, each equal weights. Gradually sprinkle on them as much water as will slake the lime; then pour on more water, stiring the whole together, that the falt may be dissolved : let the ley settle, pour it off into another vessel, and, if there is occasion, filter it. A wine pint of this ley, measured with the greatest exactness, ought to weigh just fixteen ounces troy. If it proves heavier, for every drachm that it exceeds this weight, add to each pint of the liquor an ounce and a half of water by measure: if lighter, boil it till the like quantity is wasted, or pour it upon tresh lime and ashes.

QUICKLIME greatly increases the strength of alkaline salts; and hence this ley is much more acrimonious, and acts more powerfully as a men-

ftruum on oils, fats, &c. than a folution of the potash alone: the lime should be used fresh from the kiln; by long keeping, even in close vessels, it loses of its strength: such should be made choice of as is thoroughly burnt or calcined, which may be known by its comparative lightness.

All the inftruments employed in this process should be either of wood, earthen ware, or glass: the common metallic ones would be corroded by the ley, so as either to discolour, or communicate disagreeable qualities to it. If it should be needful to filter or strain the liquor, care must be taken that the filter or strainer be of vegetable matter: woollen, filk, and that fort of filtering paper which is made of animal substances, are quickly corroded and dissolved by it.

The liquor is most conveniently weighed in a narrow-necked glass bottle, of such a size, that the measure of a wine pint may arise some height into its neck: the place where it reaches to, being marked with a diamond. A pint of the common leys of our soft soapmakers weighs more than sixteen ounces; it has been sound that their soapley will be reduced to the standard here proposed, by mixing it with something less than an equal measure of water.

LAPIS SEPTICUS, feu CAUTE-RIUM POTENTIALE.

The septic stone, or potential cautery. Edinb.

Take of potash,

Quicklime, each equal parts;

Water, three times the weight of both.

Macerate for two days occasionally firring them, then filter the ley,

and evaporate it to dryness. Put the dry mass into a crucible, and urge it with a strong fire, till it flows like oil; then pour it out upon a flat plate made hot and while the matter continues foft, cut it into pieces of a proper fize and figure, which are to be kept in a glass vessel closely stopt.

THIS preparation is a strong and a sudden caustic. It has an inconvenience of being apt to liquely too much upon the part to which it is applied, fo that it is not easily confined within the limits in which it is intended to operate: and indeed the suddenness of its action depends on this disposition to liquefy.

### CAUSTICUM COMMUNE FORTIUS.

The stronger common caustic. Lond.

Boil any quantity of the soap-leys above described, to one-fourth part: then, whilft it continues boiling, some lime, that has been kept for several months in a glass vessel stopt with a cork, is to be sprinkled in by little and little, till it has absorded all the liquor, to as to form a kind of paste; which keep for use in a vessel very closely flopt.

HERE the addition of lime in substance renders the preparation

less apt to liquely than the foregoing, and confequently more eafily confineable within the intended limits, but proportionably slower in its operation. The design of keeping the lime is, that its acrimony may be somewhat abated.

It is observable, that both these caustics, and the soap leys, that is, alkaline falts increased in their power by quicklime, do not effervesce or emit air bubles, at least in any confiderable degree, on the admixture of acids; though this effervescence has been commonly reckoned one of the principal diffinguishing characters of alkaline falts. Exposed long to the air, they gradually resume their power of effervefcence, and lofe proportionably of the additional activity which the quick-lime had produced in them.

### CAUSTICUM COMMUNE MITIUS.

The milder common caustic. Lond.

Take of Fresh quicklime, Soft foap, of each equal parts, Mix them well together at the time of uling.

THIS caustic, notwithstanding the lime is used fresh, proves much milder than the former; the acrimony of the falt being here covered by the oil and tallow by which it is reduced into foap.

### SECT.

## Volatile alkaline salts.

S fixt alkalies are produced in the burning of vegetables, and remain behind in the ashes; volatile ones are produced by a like degree of heat from animal ter. These falts are obtainable also InbRances, and rife in distillation from some vegetable matters; and

along with the other volatile principles; the admission of air, necesfary for the production of the former, is not needful for the lat-

from vegetable and animal foot. Though a strong fire is requisite for their production, yet when once completely formed, they are dissipated by the gentlest warmth: in distillation, they rise fooner than the most highly rectified spirit of wine. They are produced in urine, by putresaction, without fire; and without fire also they exhale from it.

# SPIRITUS, SAL & OLEUM CORNU CERVI.

Spirit, salt and oil of bartsborn.

Lond.

Diffil pieces of hartshorn by a fire gradually raised almost to the highest: a spirit, falt and oil will ascend.

If the oil be separated, and the spirit and salt distilled again together, with a very gentle heat, they will both arise more pure. If this be carefully repeated several times, the salt will become exceedingly white, the spirit limpid as water, and of a grateful odour.

The falt, feparated from the spirit, and sublimed first from an equal weight of pure chalk, and afterwards from a little rectified spirit of wine, becomes the sooner pure.

Calcined hartshorn is generally made by burning the horns left after this distillation.

After the same manner, a spirit, salt, and oil, may be obtained from every kind of animal sub-stance.

Edinb.

Put pieces of hartshorn into a large iron pot furnished with an earthen head; and having fitted on a capacious receiver, and luted the junctures, diffil in an open fire gradually increased. At first a phlegin arises, then a spirit, and afterwards a volatile falt, accompanied with an oil: the oil that comes over first is of a yellowish colour, but on protracting the distillation, there succeeds a reddish one verging to black. In the bottom of the iron pot there remains a black coal, which being burnt to whiteness in the open air is called calcined hartshorn.

Having poured out of the recipient all the different matters which have come over into it. they may be separated from one another into the following manner: the oil separates from the phlegm and spirit in filtration: the two latter will pass through, and the oil remain on the filter. The phlegm may be separated from the spirit by distillation is a tall vessel, with a gentle heat: the spirit will come over into the recipient, and the phlegm remain at the bottom of the diftilling vessel.

The spirit may be divided into a volatile salt and phlegm, by distilling it in a very tall and narrow cucurbit; the salt will arise, and adhere to the head in a dry form; the phlegm remaining behind

The falt may be freed from the .

oil, by fubliming it from twice its quantity of potash, for the oil is kept down by the potash, whilst the salt arises.

The spirit also is rendered pure, by adding, to every pint, two ounces of potash, and distilling in a glass retort.

The remaining potash may be again purified for use, by calcining it in an open fire, so as to burn out the oil it had absorbed from the falt or spicit.

A fpirit, falt, and oil, may be obtained in the fame mounts from all the folid parts of assemble.

Tes

The wholefale dealers have very large pots for the distillation of hartshorn, with earthen heads almost like those of the common still: for receivers, they use a couple of oil-jars, the mouths of which are luted together; the pipe that comes from the head enters the lowermost jar, through a hole made on purpose in its bottom. large quantity of the subject is to be distilled, it is customary to continue the operation for feveral days fuccessively; only unluting the head occasionally, to put in fresh materials.

When only a small quantity of spirit or salt is wanted, a common iron pot, such as is usually fixed in sand surnaces, may be employed; an iron head being sitted to it. The receiver ought to be large, and a glass, or rather tin, adopter inserted betwixt it and the pipe of the head.

The distilling vessel being charged with pieces of the horn, a moderate fire is applied, which is flowly increased, and raised at length almost to the utmost degree. At first, a phlegmatic liquor arises; the quantity of which will be less or greater, according as the horns were more or less dry; this is succeeded by the falt and oil; the falt at first dissolves, as it comes over, in the phlegm, and thus forms what is called spirit: when the phlegm is faturated, the remainder of the falt concretes in a folid form to the fides of the recipient. If it is required to have the whole of the falt solid and undisfolved, the phlegm should be removed as foon as the falt begins to arife, which may be known by the appearance of white fumes; and that this may be done the more commodioully, the receiver should be left unluted, till this first part of the process is finished. The white

vapours which now arife, fometimes come with such vehemeace, as to throw off or burst the receiver: to prevent this accident, it is convenient to have a small hole in the luting; which may be occasionally stopt with a wooden peg, or opened, as the operator shall sind proper. After the salt has all surifen, a thick, dark coloured oil comes over: the process is now to be discontinued, and the vessels, when grown cold, unluted.

All the liquid matters being poured out of the receiver, the falt which remains adhering to its fides, is to be washed out with a little water, and added to the rest. It is convenient to let the whole stand for a few hours, that the oil may the better disengage itself from the liquor, so as to be first separated by a sunnel, and asterwards more perfectly by siltration through wetted paper. The salt and spirits are then to be farther purified as a-

bove directed.

The spirit of hartshorn met with in the shops is extremely precarious in point of strength; the quantity of falt contained in it (on which its efficacy depends) varying according as the distillation, in rectifying it, is continued for a longer or shorter time. If after the volatile salt has arisen, so much of the phlegm or watery part be driven over after it, as is just sufficient to diffolve it, the spirit will be fully faturated, and as strong as it can be made; if the process is not at this instant stopt, the phlegm, continuing to arife, must render the spirit continually weaker and weak-The distillation therefore ought to be discontinued at this period, or rather whilst some of the falt still remains undissolved; the spirit will thus prove always equal, and the buyer be furnished with a certain criterion of its strength. Very

few

few have taken any notice of the above-mentioned inconvenience of these kinds of spirits; and the remedy is first hinted at in the pharmacopteia reformata. The purity of the spirit is easily judged from its clearness and grateful odour.

VOLATILE alkaline salts, and their folutions called spirits, agree, in many respects, with fixt alkalies and their folutions or leys; as in changing the colour of blue flowers to a green; effervescing with and 'neutralizing acids; liquefying the animal juices, and corroding the fleshy parts, so as when applied to the skin, and prevented from exhaling by a proper covering, to act as caustics: diffolving oils and fulphur, though less readily than the fixt alkalies, on account, probably, of their not being able to bear any confiderable heat, by which their activity might be promoted. Their principal difference from the other alkalies seems to consist in their volatility: they exhale, or emit pungent vapours, in the coldest state of the atmosphere; and by their stimulating smell they prove ferviceable in languors and faintings. Taken internally, they difcover a greater colliquating as well as stimulating power; the blood drawn from a vein, after their use has been continued for some time, being found to be remarkably more fluid than before; they are likewife more disposed to operate by perspiration, and to act on the nervous system. They are particularly useful in lethargic cases; in hysterical and hypochondriacal diforders, and in the languors, headachs, inflations of the stomach, flatulent cholics, and other fymptoms which attend them: they are generally found more ferviceable to aged persons, and in phlegmatic habits, than in the oppolite circumstances. In some fevers, particularly those of the low kind, accompanied with a cough, hoarseness, redundance of phlegm, and siziness of the blood, they are of great utility; liquefying the viscid juices, raising the vis vitz, and exciting a falutary diaphorefis; but in putrid fevers, scurvies. and wherever the mass of blood is thin and acrimonious, they do harm. As they are more powerful than the fixt in liquefying fizy blood and tenacious humours, fo they prove more hurtful where the fluids are already in a colliquated state. In vernal intermittents, particularly those of the flow kind, and where the blood is dense or fizy, they are often the most efficacious remedy. Mr. Bisset observes, in his Estay on the medical constitution of Great Britain, that though many cases occur which will yield to no other medicine than the bark, yet he has met with a pretty many that were only suppressed from time to time by the bark, but were completely cured by alkaline fpirits: that these spirits will often carry off vernal intermittents, without any previous evacuation; but that they are generally more effectual, if a purge is premifed; and in pletboric or inflammatory cases, or where the fever personates a remittent, venæsection.

These salts are most commodiously taken in a liquid form, largely diluted; or in that of a bolus, which should be made up only as it is wanted. The dose is from a grain or two to ten or twelve. Ten drops of a well made spirit, or saturated solution, are reckoned to contain about a grain of the salt.

In intermittents, fifteen or twenty drops of the spirit are given in a tea-cup full of cold spring water, and repeated five or six times in each intermission.

THE volatile salts and spirits prepared from different animal substances, have been supposed capable of producing different effects upon the human body, and to receive specific virtues from the sub-The falt of vipers has been ject. esteemed particularly serviceable in the diforders occasioned by the bite of that animal: and a falt drawn from the human scull, in diseases of the head. But modern practice acknowledges no fuch different effects from these preparations, and chemical experiments have shewn their identity. There is indeed, when not sufficiently purified, a very perceptible difference in the smell, taste, degree of pungency, and volatility of these salts; and in this state, their medicinal virtues vary confiderably enough to deserve notice: but this difference they have in common, according as they are more or less loaded with oil, not as they are produced from this or that animal Substance. As first distilled, they may be looked upon as a kind of volatile foap, in which the oil is the prevailing principle: in this state, they have much less of the proper alkaline acrimony and pun--gency, than when they have undergone repeated distillations, and fuch other operations as disengage the oil from the falt; for by these means, they lofe their faponaceous quality, and acquiring greater degrees of acrimony, become medicines of a different class. These preparations, therefore, do not differ near so much from one another, as they do from themselves in different states of purity. To which may be added, that when we consider them as loaded with oil, the virtues of a distilled animal oil itself are likewise to be brought into the account.

These oils, as first distilled, are highly fetid and offensive, of an extremely heating quality, and of fuch activity, that, according to Hoffman's account, half a drop, dissolved in a drachm of spirit of wine, is sufficient to raise a copious sweat. By repeated recufications, they lofe their offensiveness, and at the same time become mild in their medicinal operation: the rectified oils may be given to the quantity of twenty or thirty drops, and are said to be anodyne and antispasmodic, to procure a calm fleep and gentle sweat, without heating or exagitating the body. (See page 421) It is obvious therefore, that the falts and spirits must differ, not only according to the quantity of oil they contain, but according to the quality of the oil itself in its different states.

The volatile salts and spirits, as first distilled, are of a brown colour, and a very offensive smell: by repeated rectification, as directed in the processes above set down, they lose great part of the oil on which these qualities depend, the salt becomes white, the spirit limpid as water, and of a grateful odour; and this is the mark of suf-

ficient reclification.

It has been objected to the repeated rectification of these preparations, that, by separating the oil, it renders them similar to the pure salt and spirit of sal ammoniac, which are procurable at an easier rate. But this is by no means the case. The intention is not to purify them wholly from the oil, but to separate the grosser part, and to subtilize the rest, so as to bring it towards the same state as when the oil is rectified by itself. I have repeated the rectification of spirit of hartshorn twenty times successively, and sound it still to participate of oil, but of an oil very different from what it was in the first distillation.

The rectified oils in long keeping, become again fetid. The falts and spirits also, however carefully rectified, suffer, in length of time, the same change; resuming their original brown colour and ill smell; a proof that the rectification is far from having divested them of oil.

SPIRITUS, SAL, et OLEUM FULIGINIS.

Spirit, salt, and oil of soot.

Lond.

Distil soot after the same manner as directed above for hartshorn: but here more labour is required to render the spirit and salt pure.

THE volatile falt and spirit of foot are, when sufficiently purified, not different in quality from those of animal substances; though some have preserved them in nervous complaints, particularly in epileptic cases.

SPIRITUS et SAL VOLATILIS SALIS AMMONIACI. The volatile falt and spirit of sal ammoniac.

Lond.

Take a pound and a half of any fixt alkaline falt, a pound of fal ammoniac, and four pints of water. Distil off with a gentle heat, two pints of spirit.

The volatile falt is made from a pound of fal ammoniac mixed with two pounds of pure chalk, and fet to sublime in a retort,

with a strong fire.

Edinb.

Take equal parts of fal ammoniac and falt of tartar; grind them separately to powder, then mix, and put them into a glass retort, pouring on gradually as much water as will dissolve the salts. Distil, with a gradual fire, in a fand-bath: the falt rifes first, and concretes in the receiver. If the falt is wated in a dry form. remove the receiver before any water comes over: if a spirit is wanted, continue the distillation, till so much water has arisen as is sufficient to dissolve the falt, taking care to protract it no longer.

SAL ammoniac is a neutral falt, composed of volatile alkali and marine acid. In these processes, the acid is absorbed by the fixt alkali or chalk; and the volatile alkali is of course set at siberty.

The fixt alkali begins to act upon the fal ammoniac, and extricates a pungent urinous odour, as foon as they are mixed. Hence it is most convenient not to mix them till put into the distilling vessel; the two salts may be dissolved separately in water, the solutions poured into a retort, and a receiver immediately fitted on. An equal weight of the fixt salt is fully, perhaps more than, sufficient, to extricate all the volatile.

Chalk does not begin to act upon the fal ammoniac, till a confiderable heat is applied. Hence
these may be without inconvenience, and indeed ought to be,
thoroughly mixed together, before
they are put into the retort: the.
surface of the mixture may be covered with a little more powdered
chalk, to prevent such particles of
the sal ammoniac, as may happen
to lie uppermost, from subliming
unchanged. Though the fire must

here

here be much greater than when fixt alkaline falt is used, it must not be too firong, nor too fuddenly raised; for if it is, a part of the chalk (though of itself not capable of being elevated by any degree of heat) will be carried up along with the volatile falt. M. du Hamel experienced the justness of this observation: he relates, in the Memoirs of the French academy of sciences for the year 1735, that he frequently found his volatile falt, when a very strong are was made use of in the sublimation, amount to more, fometimes one half more, than the weight of the crude fal ammoniac employed; and that, though it is certain that not threefourths of this concrete are pure volatile falt, the fixt earthy matter, thus once volatilized by the alkali, arose along with it again upon the gentlest resublimation, dissolved with it in water, and exhaled with it in the air.

When all the falt has sublimed, and the receiver grown cool, it may be taken off, and luted to another retort charged with fresh materials: this process may be repeated, till the recipient appears lined with volatile salt to a conconsiderable thickness; the vessel must then be broken in order to get out the salt.

The volatile salt and spirit of sal ammoniac are the purest of all the medicines of this kind. They are somewhat more acrimonious than those produced directly from animal substances, which always contain a portion of the oil of the subject, and receive from thence some degree of a sponaceous quality. These last may be reduced to the same degree of purity, by combining them with acids into ammoniacal salts; and afterwards recovering

the volatile alkali from these compounds by the processes above directed.

The matter which remains, in the retort, after the distillation of the spirit, and sublimation of the falt, of fal ammoniac, is found to confift of marine acid united with the fixt alkali or chalk em-When fixt alkaline falt ployed. has been used as the intermedium. the residuum, or caput mortuum as it is called, yields, on folution and crystallization, a salt exactly fimilar to the spiritus salis marini coagulatus hereafter described : and hence we may judge of the extraordinary virtues, formerly attributed to this falt, under the names of sal antiby stericum, antiby pochondriacum, febrifugum, dige stivum Sylvii, &c.

The caput mortuum of the volatile falt, where chalk is employed as an intermedium, exposed to a moist air, runs into a pungent liquor, which proves nearly the fame with a folution of chalk made directly in the marine acid: it is called by some, oleum cretæ, oil of chalk. If calcined shells or other animal limes be mingled with fal ammoniac, a mass will be obtained, which likewise runs in the air, and forms a liquor of the fame kind. This liquor feems to be the fecret of some pretenders to a dissolvent of the calculus.

SPIRITUS VOLATILIS CAUS-TICUS.

Volatile caustic spirit.

Take of fal ammoniac, one pound;

Quicklime, a pound and a half; Water, four pints.

Quench the lime in the water; and having put this mixture into a retort, add to it the powdered falt. Immediately adapt a re-F f 2 cipient cipient, and with a very gentle heat draw off two pints.

THIS spirit is commonly called, from the intermedium, spirit of sal ammoniac with quicklime. The effect of the quicklime on the sal ammoniac, is very different from that of the chalk and fixt alkali in the foregoing process. Immediately on mixture, a very penetrating vapour exhales; and in distillation, the whole of the volatile falt arises in a liquid form; no part of it appearing in a concrete state, how gently soever the liquor be redistilled. The spirit is far more pungent than the other both in smell and taste; and, like fixt alkalies rendered caustic by the same intermedium, it raises no effervescence on the admixture of acids.

This spirit is held too acrimonious for internal use, and has therefore been chiefly employed for smelling to in faintings, &c. though, when properly diluted, it may be given inwardly with safety. It is an excellent menstruum for some vegetable substances, as Peruvian bark, which the other spirit extracts little from.

Some have mixed a quantity of this with the officinal spirits both of fal ammoniac and of hartshorn: which thus become more pungent,

so as to bear an addition of a confiderable quantity of water, without any danger of discovery from This abuse the take or smell. would be prevented, if what has been formerly laid down as a mark of the strength of these spirits. (some of the volatile falt remaining undissolved in them) was complied with. It may be detected by adding to a little of the suspected foirit about one-fourth its quantity or more of rectified spirit of wine; which, if the solatile spirit is genuine, will precipitate a part of its volatile salt, but occasions no visible separation or change in the caustic spirit, or in those which are sophisticated with it.

Others have substituted, to the fpirit of fal ammoniac, a folution of crude fal ammoniac and fixt alkaline falt mixed together. mixture deposites a saline matter on the addition of spirit of wine, like the genuine spirit; from which however, it may be distinguished, by the falt which is thus separated, not being a volatile alkaline, but a a fixt neutral salt. The abuse may be more readily detected by a drop or two of folution of filver made in aquafortis; which will produce no change in the appearance of the true spirit, but will render the counterfeit turbid and milky.

## S E C T. III.

Combination of alkalies with oils and inflammable spirits.

SAPO AMYGDALINUS.

Almond foap

Lond.

AKE any quantity of freshdrawn oil of almonds, and thrice its quantity by measure of the foregoing soap leys. Digest them together in such a heat, that they may but just boil or simmer, and in a sew hours they will unite: after which, the liquor, in boiling, will soon become ropy, and in good measure transparent; a little of it suffered

fuffered to cool, will appear like When this happens, throw in by little and little fome common falt, till the boiling liquor loses it ropiness; and continue the coction, till, on receiving some drops on a tile, the foap is found to ccagulate, and the water freely separates The fire being then from it. removed, the foap will gradually arise to the surface of the liquor: take it off before it grows cold, and put it into a wooden mould a frame, which has a cloth for its bottom: afterwards take out the foap. and fet it by till sufficiently dried.

After the same manner, a soap may likewise be made with oil olive; but the purest oil must be used, that the soap may be as little ungrateful as possible either to the palate or stomach.

. This process is so fully described, as to render any farther directions unnecessary. The general virtues of foaps have been already delivered in page 219; that prepared after this manner is not different in quality from the hard fort there mentioned. The strength of foaps varies confiderably with their age, and the manner in which they have been kept; fresh foap, though apparently of a good confiftence, loses, upon being thoroughly dried, near one-third of its weight; the whole of which loss is mere water; a circumstance to particularly attended to, in the exhibition of this medicine.

Soap is decompounded (or the alkaline falt and oil, of which it is composed, separated from one another) by all acids; and hence it does not lather with waters that

contain any acid unneutralized. In pure water, it dissolves into a milky liquor, which on dropping in some oil of vitriol forms a kind of coagulum: on adding more of the acid, the liquor becomes clear, the oil of the foap arifes to the furface, its alkali uniting with the acid, and forming saline concretions at the The oil, carefully colbottom. lected, proves remarkably purer than when it first entered the composition of the soap; and, like the effential oils of vegetables, dissolves in spirit of wine: it may possibly be applicable to some useful purpoles, as it feems to be freed from its groffer matter, extremely pure, and is void of the pungency of effential oils.

It follows from the above experiments, that no kind of acid ought to be used along with soap; all acids absorbing the alkaline salt of the soap from the oil. Neutral salts have not this effect, their acid being already satisfied with an alkali; but salts composed of an acid and an earthy or metalic body, as the purging bitter salt, vitriol, &c. decompound the soap equally with pure acids; acids quitting an earth or metal to unite with an alkali brought in contact with them.

. Soap diffolves likewise, but in small quantity, in pure spirit of wine: it is observable of this solution, that if exposed to a degree of cold, a very little greater than that in which water begins to freeze, it congeals into a solid pellucid mass.

The menstruum which dissolves soap most persectly, and in greatest quantity, is a pure proof spirit. The common proof spirits have a slight acidity, not indeed dissinguishable by the taste or by the usual ways of trial, but sufficient

F f 3

to give somewhat of a milky hue to solutions of soap made in them. This may be corrected by the addition of a little alkaline falt: Mr. Geoffroy observes, in the Memoirs of the French academy, that twenty-eight parts of good proof spirit, with the addition of one part of falt of kali (see page 427) will diffolve ten parts of good hard foap into a perfectly limpid liquor. common alkaline salts, as that of tartar, answer equally in this respect with that of kali; but the latter, being much less acrimonious,
SAFO TARTAREUS.
feems preferable where the foSoap of tartar.
lution is intended for medicinal Take any quantity of falt of taruſe.

This facility of the decomposition of foap by acids renders it an useful criterion of low degrees of unneutralized acidity in waters, &c. The limpid folution of foap in proof spirit, dropt into any liquor, that contains either a pure acid, or a falt composed of an acid with an earth or metal, renders the liquor immediately milky, more or less, in proportion to the quantities it is impregnated with.

> SAPO PURIFICATUS. Purified soap.

Slice one pound of dry, hard, Genoa, Alicant, or any other oilfoap, into a clean pewter vessel, and pour upon it two gallons of reclified spirit of wine. Place the vessel in a water-bath, and apply such a degree of heat as may make the spirit boil, when it will foon dissolve the foap. Let the vessel stand close covered, in a warm place, till the liquor has grown perfectly clear: if any oily matter swim upon the furface, carefully scum it off. Then decant the limpid liquor from the feces, and distil off from it all the spirit that will arise in the heat of a water-bath. Expose the remainder to a dry air for a few days, and it will become a white, opake, and fomewhat friable mass. Prad. chem.

SOAP thus purified has little or no fmell, and proves, upon examination, not in any degree acrimonious, but quite mild and fost, and consequently well fitted for medicinal purposes.

tar, very well calcined and reduced into powder whilst hot: immediately pour upon it, in a broad glass vessel, twice its quantity of oil of turpentine; and let them stand together in a cellar for some weeks, till the oil has penetrated the falt: then add more oil by degrees, till the falt has absorbed thrice its own quantity, and both appear united into a foap; which, if the matter is every day flirred, will happen in a month or two. The effect succeeds sooner, if the containing veffel be fixed to the fail of a windmill, or any other machine that turns round with great velocity.

This tedious process, which is taken from the preceding edition of the Edinburgh pharmacopæia, might be finished in a very little time, by duly attending to a circumstance which our chemists, and the pharmaceutical writers, have in general overlooked; and which many have supposed to be a means even of preventing fuccels. the oil be poured upon the pulverized falt whilst very hot, they will immediately unite, with hilling histing noise; and by rubbing for a few minutes in a hot mortar, form a truly saponaceous mass, the medicine here intended. If the

falt is suffered to grow cold before the addition of the oil, it is scarce possible to unite them, as the committee of the London college obferves, without the addition of

a little water, which in this case promotes the effect. The regular, uniform motion above recommended, does not answer so well as

agitation or rubbing in a mortar; the different degrees of centrifugal force which the oil and falt acquire, when moved circularly, tending to keep them apart. The falt does not retain so much of the oil as might be expected; far the greatest part of this volatile fluid being diffipated in the pro-

cess. Mr. Baume relates, in his Manuel de chemie lately published, that experiments have convinced him, that the soap confects of only the refinous part of the oil united

with the alkali; that the more fluid and well rectified the oil is, the less soap is obtained; and that by adding a little turpentine in substance to the mixture, the

preparation is considerably accelerated. This medicine has been greatly

celebrated as a diuretic, in nephritic complaints, and as a corrector of certain vegetable substances, particularly opium; it was for fome time a great fecret in the hands of its first preparer, Starkey, under the names of philosophic foap, the vegetable correc-tor, &c. Its virtues, however, have not been fufficiently warranted by experience; nor does the prefent practice pay any regard to it. Accordingly both the London and Edinburgh colleges have rejected

it at the late reformation of their pharmacopecias.

LOTIO SAPONACEA. Saponaceous lotion.

Lond.

Take of

Damask rose water, three quarters of a pint;

Oil olive, one quarter of a pint: Ley of tartar, half an ounce by measure.

Grind the ley of tartar and the oil together, until they unite; then gradually add the rose wa-

This is defigned for external use, as a detergent wash; and, like other foapy liquors, answers this purpose very effectually. Where it is required to be more deterfive, it may be occasionally rendered so, by the addition of a fmall quantity of a folution of any fixt alkaline falt.

> LINIMENTUM SAPONACEUM. Saponaceous liniment. Lond.

Take of

Spirit of rolemary, one pint; Hard Spanish soap, three ounces; Camphor, one ounce.

Digest the soap in the spirit of rosemary, until it is dissolved; then add the camphor.

BALSAMUM SAPONACEUM, vulgo OPPODELDOCH. Saponaceous balfam, commonly called opodeldoc. Edinb.

Take of Spanish soap, ten ounces: Camphor, two ounces; Essential oil of rosemary, Effential oil of origanum, each half an ounce; Rectified spirit of wine, four pints. Digeft F f 4

Digest the soap in the spirit of wine, with a gentle heat, till it is diffolved; then add the camphor

and the oils, and shake the whole well together, that they may be perfectly mixed.

These compositions also are employed chiefly for external purposes, against rheumatic pains, sprains, bruises, and other like complaints. Soap acts to much better advantage, when thus applied in a liquid form, than in the folid one of a plaster.

ABLSAMUM ANODYNUM. vulgo BATEANUM. Anodyne balsam, commonly called Bate's balfam. Edinb.

Take of White foap, two ounces; Crude opium, half an ounce: Camphor, fix drachms; Essential oil of rosemary, one drachm: Reclified spirit of wine, eighteen

ounces. Digest the spirit with the soap and opium, in a gentle fand-heat, for three days; then strain the liquor, and add to it the camphor and effential oil.

THIS composition is greatly commended for allaying pains, and is said to have been sometimes used with benefit even in the gout; a cloth dipt in it being laid on the part. It is fometimes likewise directed to be taken inwardly, in the same disorder, as also in nervous colics, jaundices, &c. from twenty to fifty drops or more; though furely, in gouty cases, the use of opiate medicines requires great caution. One grain of opium is contained in about ninety drops of the ballam.

LINIMENTUM VOLATILE. Volatile liniment. Lond.

Take of Oil of almonds, one ounce by measure: Spirit of fal ammoniac, two drachms by weight. Stir them together in a wide-mouthed phial, until they perfectly unite.

### EPITHEMA VOLATILE. Volatile epithem. Lond.

Take of Common turpentine, Spirit of fal ammoniac, each equal weights. Stir the turpentine in a mortar, gradually dropping in the spirit, until they unite into a white mass.

### EMPLASTRUM VOLATILE. Volatile plaster. Edinb.

Take of Venice turpentine, Spirit of fal ammoniac, each one ounce. Pour the spirit gradually into the turpentine, stirring them diligently together in a mortar.

THE three foregoing are very acrid, stimulating compositions, and are principally applied against rheumatic and ischiadic pains. epithem or plaster was formerly made of a stiffer consistence, and more adhesive, by an addition of tacamahaca; which is here judicioully omitted, as it prevented the application from being fo expeditioully got off from the part, as its great irritating power made sometimes necessary.

SPIRITUS

Chap. 8. Combination of Alkalies wish Spirit, &c. 441

SPIRITUS SALIS AMMONIACI DULCIS.

Dulcified spirit of sal ammoniac. Lond.

Take half a pound of any fixtalkaline falt, four ounces of fal ammoniac, and three pints of proof fpirit of wine. Distil off, with a gentle heat, a pint and a half,

This spirit has lately come much into esteem, both as a medicine and a menstruum. It is a solution of volatile salt in rectified fpirit of wine; for though proof spirit is made use of, its phlegmatic part does not arife in the distillation, and ferves only to facilitate the action of the pure spirit upon the ammoniacle falt. Rectified spirit of wine does not dissolve volatile alkaline falts by fimple mixture: on the contrary, it precipitates them, as has been already observed, when they are previously diffolved in water: but by the present process, a considerable proportion of the volatile alkali is combined with the spirit. It might perhaps, for some purposes, be more advisable, to use in this intention the volatile spirit made with quicklime: for this may be mixed at once with rectified spirit of wine, in any proportions, without the least danger of any separation of the volatile alkali.

SPIRITUS VOLATILIS
FOETIDUS.
The volatile fetid spirit.

Lond.

Take of
Any fixt alkaline falt, a pound and a half;
Sal ammoniac, one pound;
Afafærida, four ounces;
Proof fpirit of wine, fix pints.

Draw off with a gentle heat, five pints.

THIS spirit is designed as an antihysteric, and is undoubtedly a very elegant one. Volatile spirits, impregnated for these purposes with different setids, have been usually kept in the shops: the ingredient here made choice of, is the best calculated of any for general use, and equivalent in virtue of them all. The spirit is pale when newly distilled, but acquires a considerable tinge in keeping.

SPIRITUS VOLATILIS
AROMATICUS.

Volatile aromatic spirit.

Lond.

Take of
Effential oil of nutmeg,
Effence of lemons, each two
drachms;
Effential oil of cloves, half a
drachm;

Dulcified spirit of sal ammoniac, one quart.

Distil them with a very gentle fire.

SPIRITUS VOLATILIS OLE-OSUS, vulgo SALINUS AROMATICUS.

Folatile oily Spirit, commonly called faline aromatic Spirit.

Edinb.

Take of
Volatile falt of fal ammonic,
eight ounces;
Effential oil of rolemary, one
ounce and a half;
Oil of amber, one ounce;
Effence of lemon peel, half an
ounce;
French brandy, one gallon and
a half.
Draw off hy difillation in the

Draw off by distillation, in the heat of a water-bath, near one gallon.

Volatile

Volatile falts, thus united with aromatics, are not only more agreeable in flavour, but likewise more acceptable to the stomach, and less acrimonious, than in their pure state. Both the foregoing compositions turn out excellent ones, provided the oils are good, and the distillation skilfully performed. The dose is from five or fix drops to fixty or more.

Medicines of this kind might be prepared extemporaneously, by dropping any proper effential oil into the dulcified spirit of sal ammoniac, which will readily dissolve the oil without the assistance of distillation, as in the following compositions; in which Jamaica pepper is chosen for the aromatic ma-

ently elegant one, and very well adapted to general use.

SPIRITUS VOLATILIS OLE-OSUS EXTEMPORANEUS. Extemporaneous volatile oily spirit. Take of

terial, as being a cheap and suffici-

Dulcified spirit of sal ammoniac, one pint;

Essential oil of Jamaica pepper, two drachms.

Mix them together, that the oil may be diffolved.

Or,

Take of
Spirit of wine, highly rectified,
Spirit of fal ammoniac; each
half a pint;
Effential oil of Jamaica pepper,
two drachms.

Diffolve the oil in the spirit of wine, and mix this solution with the spirit of sal ammoniac: a white coagulum will be immediately formed, which, in a warm place, soon resolves into a transparent liquor, depositing a quantity of a volatile oily salt.

By either of the above methods, a volatile oily spirit may be made occasionally, and adapted, at pleasure, to particular purpofes, by chusing an essential oil proper for the intention. Thus, in hysterical disorders, where the uterine purgations are deficient, a preparation of this kind made with the oils of rue, favin, pennyroyal, or other like plants, proves an useful remedy; for weakness of the stomach, oil of mint may be taken; where a cephalic is required, oil of marjorum, lavender, or rolemary; in coldness and faintings, oil of cinnamon; in cases of flatulencies, the oils of annifeeds and fweet fennel feeds. last greatly cover the pungency of the volatile spirit, and render it supportable to the palate. The spirits thus made by simple mixture, are no wife inferior, in medicinal efficacy, to those prepared by distillation, though the tinge, which they receive from the oil, may render them to some persons less fightly.

SPIRITUS VOLATILIS SUC-CINATUS.

Succinated volatile spirit.

Take of

Rectified oil of amber, from twelve to fixty drops:

Rectified fpirit of wine, one ounce:

Volatile spirit of fal ammoniac prepared with quicklime, twelve ounces.

Mix them together, and distil in a retort with a moderate fire.

This composition is extremely penetrating, and has lately come into esteem, particularly for smelling to in lownesses and faintings, under the name of Eau du luce. It has been hitherto brought from France. It is not quite lim

Ρια

pid, for the oil of amber diffolves ftrong, fcaronly imperfectly in the spirit; if be imbibed. the volatile spirit is not exceedingly

ftrong, fcarcely any of the oil will be imbibed.

### SECT. IV.

## Acid Spirits.

SPIRITUS VITRIOLI tenuis, et fortis (oleum dictus E.)
atque COLCOTHAR,

Weak foirit, and the firong foirit or

Weak spirit, and the strong spirit or oil, of witriol, and colcothar.

Lond.

ET calcined vitriol be diffilled in earthen vessels, with a reverberatory fire, for three days without intermission. What remains in the vessels is called colcothar of vitriol.

Put the diffilled liquor into a glass retort, and place in it a sand furnace: the weak spirit will come over, the strong (improperly called oil of vitriol) remaining behind.

#### Edinb.

Take any quantity of green vitriol, calcined to a flight yellow colour, and reduced into powder. Fill therewith one half of an earthen tetort, place it in a reverberatory furnace, fit on a very large receiver, and lute well the junctures: then proceed to distillation, gradually increasing the fire to the utmost degree, which is to be kept up as long as any vapours arise.

The phlegm, spirit, and oil improperly so called, may be separated from each other, by committing the whole to distillation in a retort placed in a fand surface. The phlegm (which will be in little quantity if the vitriol has been duly calcined)

will arife with a small degree of heat, and the spirit with a stronger, leaving the oil behind.

THE vitriol should be calcined till it acquires a yellowish colour inclining to red: if calcined only to whiteness, as has been commonly directed, it will change in the distilling vessels into a hard compact mass, from which the due quantity of acid can never be obtained, though urged with the most vehement fire for a great length of time. A retort is an inconvenient instrument for performing the distillation in; it requires an extraordinary expence of fuel and time to elevate the ponderous acid of vitriol, so high as the figure of this veffel demands: the vessels usually employed are fo contrived, that the vapour passes out laterally, without any afcent: these are called long-necks: the junctures of them with the receivers may be luted with Windfor loam, moistened with a folution of any fixt alkaline falt, and then beaten up with a small quantity of horse-dung. If the fire is fufficiently strong, the distillation will be finished in much less than three days, though vapours will not cease to appear long after this period; when the process has been continued for a certain time, which Boerhaave limits to eighteen hours, the spirit that arifes will not pay the expence:

expence: regard however must be had herein to the size of the furnace, the quantity of vitriol in each distilling vessel, and the degree of heat employed; those who make this commodity in quantity, continue the operation no longer than till the sumes which issue from the long-necks, at the greatest distance from the fire, begin to lessen, and the recipients grow somewhat clear.

This process is not practicable to advantage without a very large ap-Hence it is become a diflinct branch of the chemical bufiness; and considerable works have been erected for it, in fuch parts of the kingdom as fuel can be most easily procured in; some of the furnaces are so large as to contain an hundred earthen long-necks, or distilling vessels, at once. metallic part of the vitriol, or colcothar, which remains after the distillation, is ground down in mills. edulcorated with water, and employed as a pigment; in medical virtue, it is not different from some of the calces of iron, to be spoken of hereafter.

The acid spirit, as it arises in the first distillation, appears of a dark or blackish colour, and contains a confiderable portion of phlegm. In the second distillation, the phlegmatic parts arise first, together with the lighter acid, which are kept apart under the name of weak spirit: at the same time, the remaining strong spirit, or oil as it is called, loses its black colour, and becomes clear; and this is the ufual mark for discontinuing the distillation. Methods of farther purifying this acid for the nicer uses are described in Practical Chemistry, page 144:

The spirit of vitriol is the most ponderous of all the liquids we are acquainted with; and the most powerful of the acids. If any other acid be united with a fixt alkaline salt or earth; upon the addition of the vitriolic, fuch acid will be dislodged, and arise on applying a moderate heat, leaving the vitriolic in possession of the alkali; though without this addition. it would not yield to the most vehement fire. Mixt with water, it instantly conceives great heat, infomuch, that glass vessels are apt to crack from the mixture unless it is very flowly performed; exposed to the air, it imbibes moisture, and foon acquires a notable increase of weight. In medicine, it is employed chiefly as subservient to other preparations; it is likewise not unfrequently mixed with juleps and the like, in fuch quantity as will be sufficient to give the liquor an agreeable tartness in the intentions of a cooling antifeptic, restringent, and stomachic. page 325.

### SPIRITUS SULPHURIS

per campanam.

Spirit (commonly called Oil) of fulphur by the bell.

Let the fulphur be fet on fire, under a glass vessel fitted for this use, called a bell; and let the acid spirit, which trickles down from the sides of the bell, be received in a glass dish placed underneath.

#### Edinb.

Take any quantity of sulphur; melt it in an earthen dish, and dip into it twisted strips of linen, so as to form a sulphurated match. Fasten this in the mouth of a vial, which is to be set in the bottom of a glass or earthen dish, in a moist place screened from the wind: then kindle the sulphur with a red-hot iron; and hang

OACL

over it a glass bell, at such a distance, that the slame may not touch it. The vapour of the sulphur will condense in the bell by the cold, and drop down from its sides, like water, into the vessel placed underneath.

THE glass usually employed for this purpole by the chemists differs considerably from the bell shape; its belly is spherical, and has a rim at the bottom turned inwards a little: the upper part ends in a long open stein: a large receiver, with a hole cut in its bottom, and a long tube inferted into its mouth, would answer as well. If the fulphur happens to burn dull, the glass is taken off, and the matter firred with an iron wire, or a clean tobacco-pipe; as it confumes, fresh quantities are supplied, till all the fulphur designed for this use is The condensation of the burnt. fumes depends in great measure upon their imbibing aqueous moilture: hence in wet weather; or a damp place, the operation succeeds best. In dry weather, it is customary to moisten the bell, by sufpending it for a little time over the steam of boiling water.

This process is sufficiently troublesome, and the yield of acid spirit obtained by it extremely small; greatest part of the sumes escaping into the air, partly at the bottom, and partly through the upper aper-

ture of the bell.

Several contrivances have been made for preventing these inconveniencies. One of the best commonly known, is that described in vol. v. art. 14. of the Edinburgh essays; instead of the best, a large retort is employed, having a tubulated receiver (with the pipe turned uppermost) adapted to its neck; in-

stead of the large aperture in the bottom of the bell, a small one is made in the bottom of the retort; and thus by diminishing the aperture, enlarging the capacity of the vessels, and lengthening the passage of the sume, a considerably larger quantity of the sumes are detained than in the common instruments.

This apparatus may be greatly improved, by cutting the bole in the fide of the retort, and pouring into the bottom an ounce or two of warm water, in the middle of which is placed a shallow stone cup containing the fulphur. heat of the burning fulphur is foon communicated to the water, so as to keep it continually rifing in steam; with this aqueous vapour, the fumes of the brimftone are effectually blended as they ascend; and detained in confiderable quantity, in a much less proportion of phlegm than when the common methods are pursued; for here, the business of rectification or dephlegmation is carrying on, at the same time that the acid is collecting.

This affair is capable of being much farther improved. In the common method by the bell, in the most favourable circumstances. fcarce above two drachms of acid spirit are obtained from sixteen ounces of fulphur; by the fecond apparatus, an ounce may be obtained from the same quantity; and by the other, about two ounces. It appears however, from experiments related by Stahl, and others, that out of fixteen ounces of fulphur, at least fifteen ounces are pure acid, of such strength as to require being diluted with above an equal weight of water, to reduce it to the pitch of common spirit of fulphur. It follows therefore, that if we could contrive a method of burning fulphur, fo as to preferve all the fumes, we might obtain from it much more than its own weight, of an acid of the or-

dinary strength

The acid obtained from sulphur is in all respects similar to that of The acid of fulphur, united with iron or copper, forms a true vitriol; and the acid of vitriol, combined with inflammable matters, produces fulphur, not diftinguishable from pure common brim-ftone. The identity of these acids is well known to some particular persons: who supply us with almost all that is now fold under the name of oil of vitriol, prepared from the fumes of burning fulphur. method by which they obtain the acid so plentilully, and at so cheap a rate, from this concrete which bas hitherto yielded it so sparingly, differs from the processes above defcribed. Instead of an open bell, or a retort with the mouth open. they use for burning the sulphur in, very large spherical glass vessels blown on purpose, of the size of a hogshead or more, with only one aperture, through which the fulphur is introduced, and which is afterwards immediately closed, till the fumes have fubfided and incorporated with the vapour of the warm water placed in the lower part of the vellel.

AQUA SULPHURATA.
Sulpburated water, usually called
gas sulpburis.
Lond.

Take a quart of water, and half a pound of fulphur. Let part of the fulphur be fet on fire in an iron ladel, and fuspended over the water in a close vessel; as soon as the sumes subside, some more of the sulphur is to be fired in the fame manner; and this repeated till the whole quantity is burnt.

A convenient way of managing this process is, to put the water into a glass receiver, placed on its side; and to have the ladel, containing the burned sulphur, fixed to a plug, made to go freely into the neck of the vessel; the use of the plug is to keep the ladel from dipping into the water: the sumes which issue betwixt it and the glass, may be confined by a cloth thrown round the neck.

The water is impregnated, in this process, with a subtile volatile acid, different in many respects from the foregoing spirits of sulphur and of vitriol. The acid-may likewife be obtained in the fame volatile state, both from vitriol and fulphur, without water. If the retort or long-neck, during the distillation of oil of vitriol, happens to crack in the fire, all the acid that rifes afterwards is found to be thus volatilized. If cloths, moistened with a folution of fixt alkaline salt, be suspended over burning brimftone, the acid fumes will be imbibed by the alkali, and form with it a neutral falt : if this neutral falt be rubbed off from the cloths, and some common oil of vitriol poured upon it, the volatile acid it had imbibed from the fulphur will be immediately extricated again, and may be collected by distillation. The acid proves in all these casas so volatile, as to diftil in a heat scarcely greater than that which the hand can support: it has a pungent fuffocating fmell, like that of the fumes of burning brimstone, bot discovers to the taste very little acidity or corrofiveness. Exposed for some time to the air, it loses these properties, and becomes comes a fixt acid, and corrofive like common oil of vitriol.

The aqua fulphurata is liable to great uncertainty in point of strength, partly on account of the water being impregnated with a greater or less quantity of the fumes, according as the process is more or less skilfully managed; and partly on account of the above change of the acid from a volatile to a fixt When newly prepared, it is highly volatile and pungent, fmelling like burning brimstone, but in tafte rather bitterith and austere than acid; in keeping, the volatility and smell are lost, and the liquor (sooner or later, according as the air is more or less admitted to it) becomes in all respects the fame as water acidulated with a little common oil of vitriol.

This preparation is faid to give relief in fits of the convultive althma. It is taken to the quantity of a fpoonful or half an ounce two or three times a day, in any fuitable

vehicle.

# SPIRITUS NITRI Glauberi. Glauber's spirit of nitre. Lond.

Take three pounds of nitre, and one pound of the strong spirit, or oil of vitriol. Mix them cautiously and gradually together, under a chimney; and then distil, at first with a gentle, and afterwards with a stronger heat.

#### Edinb.

Put two pounds of nitre into a glass retort; and add by degrees one pound of oil of vitriol diluted with an equal quantity of warm water. Distil in a sand heat, gradually increased, till the matter remains dry.

This spirit is reclified by a second

distillation, with the heat of a water-bath, in a glass cucurbit with its head and receiver: the phlegm arises, leaving the spirit behind.

HERE the vitriolic acid expels that of the nitre, in red corrolive vapours, which begin to issue immediately upon mixture, and which the operator ought cautiously to avoid. A pound of oil of vitriol is fufficient to expel all the acid from about two pounds of nitre, not from more : some direct equal parts of the two. The spirit, in either case, is in quality the same: the difference in this respect affecting only the residuum. When two parts of nitre are taken to one of oil of vitriol, the remaining al-kaline basis of the nitre is completely faturated with the vitriolic acid, and the refult is a neutral falt, the same with vitriolated tartar, as we shall see hereaster; if more nitre is used, a part of the nitre in substance will remain blended with this vitriolated falt; if less nitre, it cannot afford alkali enough to faturate the vitriolic acid, and the residuum will be not a neutral falt, but a very acid one. In this last case there is one conveniency; the acid falt being readily diffoluble in water, so as to be got out without breaking the retort, which the others are not.

The acid of nitre is next in strength to the vitriolic, and dislodges all but that from alkaline salts or earths. It differs from all the other acids in deslagrating with inflammable matters: if a solution of any inflammable substance, as hartshorn, &c. in this acid be set to evaporate; as soon as the matter approaches to dryness, a violent detonation ensues. The chief use of this acid is as a menstruum for

certain

certain minerals, and as the basis of some particular preparations, of which hereafter. It has been given likewise diluted with any convenient vehicle, as a diuretic, from ten to sifty drops.

### SPIRITUS SALIS MARINI Glauberi

Glauber's spirit of sea salt. Lond.

Take two pounds of sea salt, and the same quantity of strong spirit or oil of vitriol. Dilute the acid spirit with a pint of water, and pour this mixture, by little and little, on the salt under a chimney; then distil, at first with a gentle, and afterwards with a stronger sire.

Edinb.

Put into a glass retort two pounds of sea falt decrepitated (that is, dried over the fire till it ceases to crackle, and pour thereon, by little and little, one pound of oil of vitriol diluted with an equal quantity of warm water. Place the retort in fand; and, with a fire gradually increased, distil to dryness. This spirit is rectified by a fecond distillation, in a glass cucurbit, with a head and receiver adapted to it; by the heat of a water-bath the phlegm will arise, leaving the spirit behind.

THE marine acid arifes, not in red sumes like the nitrous, but in white ones. The addition of water is more necessary here than in the foregoing process; the marine vapours being so volatile, as scarce to condense without some adventitious humidity; and hence the rectification, directed in the second process, does not succeed so well as that of the nitrous acid, a part of the marine spirit arising along with the phlegm. The oil of vitriol

is most conveniently mixed with the water in an earthen or stoneware vessel; for unless the mixture is made exceedingly slowly, it grows so hot as to endanger breaking a glass one.

THE spirit of sea salt is the weakest of the mineral acids, but stronger than any of the vegetable: it requires a greater sire to distil it than that of nitre, yet is more readily dissipated by the action of the air. It is used chiestly as a menstruum, for the making of other preparations: sometimes likewise it is given, properly diluted, as an antiphlogistic, aperient, and diuretic, from ten to sixty or seventy drops.

Spirit of falt.

Take a pound of sea salt thoroughly dried, and three pounds of powdered bricks. Mix and put them into an earthen retort, of fuch a fize that thefe may fell only one half of it. Place the retort in a reverberatory furnace, adapt to it a large receiver, and lute well the junctures. the fire be applied, at first very sparingly, and afterwards increased by degrees, until all the spirits are driven over in the form of clouds. When the vessels are grown cold, pour out the distilled liquor into a glass cucurbit, and gently abstract from it the phlegin, which will leave the fpirit pure.

Instead of brickdust, some have used bolar earths and clays. It has been supposed, that these substances ast by discontinuing and dividing the particles of the salt, so as to enable the fire to expel the spirit: if this was true, glass or sand would prove equally serviceable, and the same intermedium would

would answer as well for a number of times as at first, the reverse of which, experiments shew to be. true. Brick earth, and other substances of this kind, contain a small quantity of vitriolic acid, whose known property it is to disengage the acid of fea falt, and which is the only part of them of use in this process. The quantity of spirit, therefore, obtained by these intermedia, is only in proportion to that of the acid contained in them, which is extremely fmall. This has occasioned some to make use of vitriol, as containing a latger quantity of the vitriolic acid; but though vitriol is in this respect greatly preferable to brickduft, or the argillaceous earths; yet in another, it is found lefs eligible; its metallic part fo strongly adheres to the marine acid, as to keep it down after it is separated from its basis, or elfe arifes along with it, and defiles the product. Thefe methods therefore of extracting the spirit of salt have been for some time laid aside, the foregoing, in which the pure vitriolic acid itself is used, being in all respects more convenient and advantageous.

# AQUA FORTIS.

Take of
Nitre,
Green vitriol uncalcined, each
three pounds,
The fame vitriol calcined, one
pound and a half.

Mix them well together, and diffil with a very firong fire, as long as any red vapour arises.

# AQUA FORTIS SIMPLEX. Single aquafortis. Edinb.

Take two parts of vitriol calcined to whiteaels, and one part of powdered nitre. Mix them very well together, and fill therewith an earthen retort to two-thirds; then fit on a large receiver, and proceed to distillation; which is to be performed in the same manner as directed for spirit of fait.

THE vitriol here is not liable to the inconvenience mentioned in the foregoing remark; it only occasions a greater heat to be necessary than when the pure vitriolic acid is afed. for the acid of the vitriol must be extricated before it can act on the nitre; the fire, however, must not be extremely firong, otherwise some of the metallic parts of the vitriol will be forced over along with the nitrous acid: the direction of theroughly mixing the ingredients ought to be well attended to, for if this is neglected, or but flightly performed, the due quantity of aeid will not be obtained. The produce of these processes is a spirit of nitre containing fo much more phlegm, or watery moisture, than Glauber's spirit, as the vitriol employed in its preparation does more than an equivalent quantity of oil of vitriol.

# AQUA FORTIS DUPLEX. Double aquaforsis. Edinb.

Take of
Green vitriol calcined to whitenefs.
Clay dryed and powdered,

Powdered nitre, of each equal parts.

Mix them well together, and diffil in an earthen retort as above.

This process is an unartful one. The clay appears to be of very little use, though the contrivers of the process seems, from the reduction of the vitriol, to have G g laid

laid confiderable stress on it: all it can do is to kinder the melting of the salts. It would doubtless be better to omit the clay, and increase the quantity of the vitriol; which, in order to make the aquafortis of the strength here intended, should undergo a farther degree of calcination.

The great demand which there is in fundry bulinesses for aquafortis has occasioned the preparation of it to become a trade by itself. Hence larger and less expensive instruments than those mentioned above, have been contrived. common distilling vessel is a large iron pot, with an earthen, or stoneware still-head, to which is adapted a large glass globe, or else a jar made of the same kind of clay as The workmen are notthe head. at the trouble either of drying the vitriol, or pounding the nitre, but throw them both promiscuously into the pot, where the fire foon. liquefies, and mixes them together. The aquafortis, prepared after this manner, is extremely impure, and utterly unfit for many purposes, fuch in particular are the folution of mercury and of filver: the violence of the fire, employed in the operation, never fails to elevate some of the metallic parts of the vitriol; the nitre is used rough or unrefined, which containing a portion of sea salt, sends over some of the marine along with the nitrous acid: nor are the ingredients free from bits of wood, or other vegetable matters, which burning in the process foul the spirit with an empyreumatic oil, giving it, at the fame time, an high colour. Ιf therefore common aquafortis be employed in any medicinal preparation, it ought to be previously purified; the most effectual method of doing which is the following.

AQUA FORTIS PURIFICATA.

Purified aquafertis.

Drop into the aquafortis a drop or two of folution of filver. If it becomes milky or cloudy, drop in a little more of the folution, till a fresh addition occasions no further change; allowing proper intervals for the white matter to fettle, that the effect of a new addition may be the better perceived. Then pour the liquor into a glass retort, and diffil in a fand-heat to drynes.

THE milkiness produced by the folution of filver is a certain mark of marine or vitriolic acid in the aquafortis; the filver absorbing those acids, and forming with them a concrete which the liquor is incapable of holding diffolved. If the aquafortis is not made at all cloudy by this folution, we may be certain of its having been previously free from the least admixture of those heterogeneous acids; and when it ceases to become milky from a fresh addition, we may be equally cartain, that how much foever it might have contained of them at first, they are now perfectly separated.

The folution of filver is to be made in aquafortis already purifi-Where this cannot be had, the little quantity generally sufficient for the present purpose, may be made in the common impute fort of aquafortis, which will be purified during the diffolution itself. Put a thin bit of filver into a little of the aquafortis, and fet the vial in a fand-heat; if the aquafortis is pure, numerous minute bub-bles will issue from the silver on all fides, and the metal will gradually diffolve without altering the transparency of the hquor: but if the aquafortis contains marine or vitriolic acid, it will quickly become milky, those

acids uniting with the filver, as in the above process, as fast as the nitrous acid dissolves it. As the white matter precipitates upon, and adheres to, the furface of the filver, so as to impede the further action of the mentruum; the liquor must be filtered and treated in the same manner with a bit of fresh filver: if any milkiness still ensues, the operation must be repeated with another piece of the metal, till all the foreign acids are separated, and the filver is found to dissolve clear. Good aquafortis takes up about half its own weight of filver.

The filver may be recovered from the white fettlings, without any confiderable lofs, by the following

method.

Let the matter be thoroughly dried, then mixed with a little potash, and the mixture made into
a passe with oil. Put this passe
into a crucible, surrounding it
every where with a little more
potash. Set the crucible in a
proper surrace, and gradually
raise the fire so as to bring the
whole into suspense. When the
crucible is grown cold, a lump
of sine silver will be sound in the
bottom.

# AQUAFORTIS COMPOSITA. Compound aquafortis, Lond.

Take fixteen ounces of aquafortis, and one drachm of fea falt. Distil them to dryness.

THIS is defigned as a mentiruum for quickfilver, for the preparation of the red mercurial corrofive, or red precipitate as it is called; which the marine acid in this compound liquor renders of a more fparkling appearance, and more beautiful to the eye, than when made with the nitrous acid alone.

# AQUA REGIA.

Put an ounce of powdered (al ammoniac into a large cucurbit, and add to it, by little and little at a time, four ounces of spirit of nitre, or double aquafortis. Let them stand together in a sandheat, till the salt is entirely disolved.

THE glass in which the mixture is made should be placed under a chimney (to carry up the offensive vapour) and its orifice by no means floot till such time as the falt is perfeetly diffolved, and the fumes cease to arise with impetuosity. These cautions are extremely necessary, if the process be conducted according to the directions above. But if the fal ammoniac, finely powdered, be gradually added to the acid spirit (which ought to be of a middle degree of strength between lingle aquafortis and strong fpirit of nitre) the folution will proceed without any inconvenience; and may be finished in a reasonable compass of time, provided the mixture be now and then stirred. - The only use of aqua regia and the aquæ fortes, is as menstrua for certain mineral substances.

### ACETUM DISTILLATUM, vel SPIRITUS ACETI. Distilled winegar, or spirit of

vinegar. Lond.

Let vinegar be distilled with a gentle heat as long as the drops fall free from an empyreuma.

If fome part of the spirit which comes over first be thrown away, the rest will be the stronger.

#### Edinb.

Put any quantity of the best vinegar into a large, shallow, glass vessel, and with a gentle heat, G g 2 in in a fand-bath, evaporate about one-fourth part of it: then distil the remainder in an alembic, with a glass head, gradually increafing the fire, as long as the spirit comes off clear.

This process may be performed either in a common still with its head, or in a retort. The better kinds of wine vinegar should be made use of; those prepared from malt liquors, however fine and clear they may feem to be, contain a large quantity of a viscous substance, as appears from the sliminess and ropiness to which they are very much subject; this not only hinders the acid parts from arising freely, but likewise is apt to make the vinegar boil over into the recipient, and at the same time dispoles it to receive a disagreeable impression from the fire. And indeed, with the best kind of vinegar, if the distillation be carried on to any great length, it is extremely difficult to avoid an empyreuma. The best method of preventing this inconvenience is, if a retort be made use of, to place the sand but a little way up its fides, and when somewhat more than half the liquor is come over, to pour on the remainder a quantity of fresh vinegar equal to that of the liquor drawn off; this may be repeated three or four times; the vinegar supplied at each time being previously made hot: the addition of cold liquor would not only prolong the operation, but also endanger breaking the retort. If the common still is employed, it should likewise be occasionally supplied with fresh vinegar, in proportion as the spirit runs off; and this continued, until the process can be conveniently carried no farther: the distilled spirit must be rectified by a second distillation in a retort, or glass alembic, for though the head and receiver be of glass or stone ware, the acid will contract a metallic taint from the pewter worm.

The refiduum of this process is commonly thrown away as useless. though if skilfully managed, it might be made to turn to good account; the most acid parts of the vinegar still remaining in it. Mixed with about three times its weight of fine dry fand, and committed to distillation in a retort, with a well regulated fire, it yields an exceeding strong acid spirit; together with an empyreumatic oil, which taints the spirit with a disagreeable odour. This acid is nevertheless. without any rectification, better for some purposes (as a little of it will go a great way) than the pure fpirit; particularly for making the fal diureticus of the London dispensatory; for there the oily matter, on which its ill flavour depends, is

The spirit of vinegar is a purer and ftronger acid than vinegar itfelf, with which it agrees in other respects. The medical virtues of these liquors may be seen in the fection of acids, page 61, and under the article ACETUM, page. 74, their principal difference from the mineral acids confifts in their being milder, less stimulating, less dispossed to affect the kidneys, and promote the urinary fecretions, or to coagulate the animal juices. The matter, left after the distillation in glass vessels, though not used in medicine, would doubtless prove a serviceable detergent, saponaceous acid; and in this light it stands recommended by Boerhaave.

burnt out by the calcination.

### SECT. V.

## Combination of acid with vinous spirits.

L L the mineral acids, on being mixed with spirit of wine, raise a great ebullition and heat. If the acid is in small quantity, it unites intimately with the vinous spirit, so as to arise with it in distillation. The taste and all the characters of acidity, are destroyed; and the mixture acquires a grateful slavour, which neither of the ingredients had before.

# SPIRITUS VITRIOLI DULCIS. Dulcified spirit of vitriol. Lond.

Take of the strong spirit or oil of vitriol, one pound; of rectified spirit of wine, one pint. Cautiously mix them together by little and little at a time, and distil the mixture, with a very gentle heat, till a black froth begins to arise: then immediately remove the whole from the fire, lest this froth should pass over into the recipient, and frustrate the operation.

Edinb.

Into four pints of rectified spirit of wine, drop cautiously one pound of oil of vitriol. Digest in a close stopt matrass, with a very gentle warmth in sand, for eight days; and then distil with a soft fire, till oily drops begin to appear in the neck of the retort.

Add to the diffilled spirit, an equal measure of water, in which some salt of tartar has been dissolved, in the proportion of a drachm of the salt to a pint of the water: mix them together, by shaking the vessel; and draw

off the spirit again by a gentle heat.

THE different proportions of the acid spirit to the vinous in these processes, make no variation in the quality of the produce, provided the distillation is duly conducted; all the redundant acid being left in the residuum.

A good deal of caution is requifite in mixing the two liquors. Some direct the spirit of wine to be put first into the retort, and the oil of vitriol to be poured upon it all at once; a method of procedure by no means adviseable, as a violent heat and ebullition always ensue, which not only dissipate a part of the mixture, but hazard also the breaking of the vessel, to the great danger of the operator. Others put the oil of vitriol into the retort first, then by means of a funnel, with a long pipe that may reach down just to the surface of the acid, pour in the spirit of wine: if this is done with sufficient caution, the vinous spirit spreads itself on the furface of the oil of vitriol, and the two liquors appear distinct: on flanding for a week or two, the vinous spirit is gradually imbibed. wishout any commotion, and the vessel may then be safely shaken, to complete the mixture; but if the spirit is poured in too hastily at first, or if the vessel is moved before the two liquors have in some degree incorporated, the same effect enfues as in the foregoing case. The only secure way is, to add the oil of vitriol to the spirit of wine by a little quantity at a time, waiting till the first addition is incor-Gg 3 porated

porated before another quantity is put in: by this management, the heat that enfues is inconfiderable, and the mixture is effected without

any inconvenience.

The distillation should be performed with an equable and very gentle heat, and not continued fo long as till a black froth begins to appear; for before this time a liquor will arise of a very different nature from the spirit here intend-The feveral products are most commodioully kept apart by using a tubulated receiver, so placed, that its pipe may convey the matter which will come over, into a vial set underneath : the juncture of the retort and recipient is to be luted with a paste made of linseed meal, and further fecured by a piece of wet bladder; the lower juncture may be closed only with some soft wax, that the vial may be occasionally removed with ease.

The true dulcified spirit arises in thin subtile vapours, which condense upon the sides of the recipient in streight strize. It is colour-less as water, very volatile, inslammable, of an extremely fragrant smell, in taste somewhat aromatic.

After the fire has been kept up for some time, white sumes arise, which either form irregular strize, or are collected into large round drops like oil: on the first appearance of these, the vial (or the receiver, if a common one is made use of) must be taken away. If amother be substituted, and the distillation continued, an acid liquor comes over, of an exceeding pungent smell, like the sumes of burning brimstone. At length a black froth begins hastily to arise, and prevents our carrying the process further.

On the furface of the sulphureous spirit is found swimming a small quantity of oil, of a light yellow colour, a strong, penetrating, and very agreeable smell. This oil seems to be nearly of the same nature with the essential oils of vegetables. It readily and totally disfolves in rectified spirit of wine, and communicates to a large quantity of that menstruum the taste and smell of the aromatic or dulcified solution.

spirit.

The matter remaining after the distillation is of a dark blackith colour, and still highly acid. Treated with fresh spirit of wine, in the fame manner as before, it yields the same productions; till at length, all the acid that remains unvolatilized being fatiated with the inflammable oily matter of the spirit, the compound proves a bituminous, suphureous mass: which exposed to the fire in open vessels, readily burns, leaving a confiderable quantity of fixt ashes; in close ones, explodes with violence; and with fixt alkaline falts, forms a compound, nearly similar to one composed of alkalies and sulphur.

Dulcified spirit of vitriol has been for some time greatly esteemed both as a menstruum and a medicine. It dissolves some resinous and bituminous substances more readily than spirit of wine alone, and extracts elegant tinctures from fundry vegetables: especially if rectified, as in the second of the above processes, from a little fixt alkaline falt, to feparate any redundant acidity. a medicine, it promotes perspiration and the urinary secretion, expels flatulencies, and in many cafes abates spasmodic strictures, cases pains and procures fleep: the dofe is from ten to eighty or ninety drops in any convenient vehicle. It is not effentially different from the celebrated anodyne liquor of Hoffman; to which it is, by the author himself, not unfrequently directed as a fuccedaneum.

Liquos

LIQUOR ANODYNUS MINERA-LIS HOPPMANNI.

Hoffman's mineral anodyne liquor. Into half a pound of concentrated oil of vitriol, placed in a large glass retort, pour by little and little, through a long-stemmed funnel, one pint and a half of highly rectified spirit of wine. Stop the mouth of the retort, digest for some days, and then distil with a very gentle heat. At first a fragrant spirit of wine will arise; and after it, a more fragrant volatile spirit, to be caught The receivin a fresh receiver. er being again changed, a fulphureous, volatile, acid phlegm comes over; and at length a fweet oil of vitriol, which should be immediately separated, lest it be absorbed by the phlegm. Mix the first and second spirits together, and in two ounces of this mixture dissolve twelve drops of the fweet oil. If the liquor has any fulphureous smell, redistil it from a little falt of tartar. Paril.

WHETHER this is the exact preparation, fo much recommended and so often prescribed by Hossman, as an anodyne and antispasmodic, we cannot determine. We learn from his own writings, that his anodyne liquor was composed of the dulcified spirit of vitriol, and the aromatic oil which arises after it : but not in what proportions he mixed them together. The college of Wirtemberg feem to think that all the oil was mixed with all the spirit obtained in one operation, without regard to the precise quantities.

### AQUA RABELLIANA. Eau de Rabel.

Take four ounces of oil of vitriol, and twelve ounces of rectified spirit of wine. Pour the vinous spirit gradually into the acid, and digest in a close matrass. Parif. This liquor has been greatly celebrated in France as a reftringent, and for the same purposes as the dulcified spirit; from which it differs in having a considerable acidity.

# SPIRITUS ÆTHEREUS. Etbereal Spirit. Edinb.

Take dulcified spirit of vitriol, rectified as in the second of the foregoing processes under that title, page 453.

Water in which falt of tartar has been diffolved, in the proportion there prescribed; of each equal parts.

Mix them together, then fuffer them to reft, and separate the ethereal spirit which rises to the top.

Great part of the dulcified spirit of vitriol may be recovered from the remaining liquor by distillation.

The preparation of this fingular fluid, now first received into a public pharmacopæia, has hitherto been confined to a few hands: for though several processes have been published for obtaining it, the fuccess of most of them is precarious, and some of them are accompanied also with danger to the operator. The principal difficulty confifts in the distillation of the dulcified spirit of vitriol; where that spirit itself only is wanted, the method above directed for it succeeds in perfection: but when it is made with a view to the ether, a little variation is necessary, for only a fmall quantity of ether can be feparated from the spirit so prepared. There, the distillation is performed with an equable and gentle heat: here the fire should be hastily raised, so as to make the liquor boil, for on this circumstance the produce of ether principally depends. (See a paper on this sub-Gg4 ject

ject by Dr. Morris, in the second volume of the Medical Observations and inquiries, published by a fociety of physicians in London.) The heat may be kept up to this degree till the black froth begins to appear: but it is more adviscable, when the ebullition has continued for a little time, to abate the heat. The most fecure way will be, as foon as the mixture boils with large bubbles, to entirely remove the fire : the retort should be furrounded with fand in an iron pot, and the heat, which these have by this time acquired, will be fufficient for keeping up the ebullition a little longer, and for completing the diffillation. The diffilled spirit is to be drawn over again from some fixt alkaline fult, but the addition of water, in this rectification, does not appear to be necessary. ether is to be separated from the rectified spirit in the manner above preferibed.

THE ether or ethereal spirit is the lightest, most volatile, and inflammable, of all known liquids. It is lighter then the most highly rectified spirit of wine, in the proportion of about 7 to 8: a drop, let fall on the hand, evaporates almost in an instant, scarcely rendering the part moift. It does not mix, or only in a fmall quantity, with water, fpirit of wine, alkaline lixivia, volatile alkaline spirits, or acids, but is a powerful diffolvent for oils, balfams, refins, and other analogous fubitances. It has a fragreat odour, which, in confequence of the volatility of the fluid, is diffused through a large space. Its medical effects are not as yet much known, though it is not to be doubted that a fluid of so much fubtility must have considerable ones. It has often been found to give eafe in violent head achs, by being applied externally to the part, and to relieve the tooth-ach, by being laid on the afflicted jaw. It has been given also internally, with benefit, in whooping coughs, and hyfterical cases, from two or three drops to five and twenty, in a glass of wine or water; which should be swallowed as quick as possible, as the ether so speedily exhalos.

# SPIRITUS NITRI DULCIS. Dulcified spirit of nitre. Lond.

Take a quart of rechified spirit of wine, and half a pound of Glauber's spirit of nitre. Mix them, by pouring the nitrous spirit into the other; and distil with a gentle heat, as long as the liquor which comes over does not raise any effervescence with lixivial salts.

#### Edinb.

Put three parts of rectified spirit of wine into a large bolt-heat, and gradually add thereto one part of spirit of nitre. Digest them together for two days; and then distil in a sand-heat, according to art; taking care, towards the end of the operation, that the retort break not from too great a heat.

Here the operator must take care not to invert the order of mixing the two liquors, by pouring the vinous spirit into the acid; for if he should, a violent effervescence and heat would ensue, and the matter be dispersed in highly noxious red sumes. The most convenient and safe method of performing the mixture seems to be, to put the inflammable spirit into a large glass body with a narrow mouth, placed under a chimney, and to pour upon it the acid, by means of a glass

a glass funnel, in very small quantities at a time : shaking the vessel as foon as the effervescence ensuing upon each addition ceases, before a fresh quantity is put in; by this means, the glass will heat equally, and be prevented from breaking. During the action of the two spirits upon one another, the veffel should be lightly covered; if close stopt, it will burft, and if left entirely open, some of the more valuable parts will exhale. Lemery directs the mixture to be made in an open veffel: by which unscientifical procedure, he usually loft, as he himfelf observes, half his liquor; and we may prefume that the remainder was not the medicine here intended.

The liquors, mixed together, fhould be suffered to rost for at least twelve hours, that the fumes may entirely subside, and the union be in some measure compleated. distillation should be performed with a very flow and well regulated fire a otherwise the vapour will expand with fo much force as to burft the veffels. Wilson seems to have experienced the justness of this obfervation; and hence directs the juncture of the retort and receiver not to be luted, or but flightly: if a tubulated recipient, with its upright long pipe, be made use of, and the distillation performed with the heat of a water-bath, the veffels may be luted without any danger; this method has likewife another advantage, as it afcertains the time when the operation is finished: examining the distilled spirit every now and then with alkaline falts, as directed above, is

fufficiently troublesome: whilst in a water-bath, we may safely draw over- all that will arise, for this heat will elevate no more of the acid than what is dulcified by the vinous spirit.

Dulcified spirit of nitre has been long held, and not undefervedly, in great esteem. It quenches thirst, promotes the natural fecretions, expels flatulencies, and moderately ftrengthens the stomach: it may be given from twenty drops to a drachm, in any convenient vehicle. Mixed with a small quantity of spirit of hartshorn, the spiritus volatilis aromaticus, or any other alkaline spirit, it proves a mild yet efficacious diaphoretic, and often notably diuretic; especially in some febrile cases where such a salutary evacuation is wanted. A fmall proportion of this spirit added to malt spirits, gives them a flavour approaching to that of French brandy.

# SPIRITUS SALIS DULCIS. Dulcified spirit of salt. Edinb.

This is made with spirit of salt, after the same manner as dulcified spirit of nitre.

THE dulcification of the spirit of falt does not succeed so perfectly, as that of the two foregoing acids, only a minute portion of it uniting with the spirit of wine, and unless the process is skilfully managed, scarce any. Some have held this spirit in great esteem against weakness of the stomach, indigestion, and the like, following from hard drinking; at present it is not often made use of or kept in the shops.

### E C T. VI.

## Neutral Salts.

WHEN any acid and any alkakaline falts are mixed together, in such proportion that neither of them may prevail, they form by their coalition a new compound, called NEUTRAL. In all the combinations of this kind (except some of those with vegetable acids) the alkali and acid are so strongly retained by one another, that they are not to be disunited by any degree of fire. How volatile soever the acid was by itself, if combined with a fixt alkali, it proves almost as fixt as the pure alkali : if the alkali is of the volatile kind, the compound proves also volatile, subliming in its whole substance, without any separation of its parts. There are, however, means of procuring this disunion by the intervention of other bodies, as we have already feen in the feparation of the volatile alkali of sal ammoniac, and of the acids of nitre and fea falt: but in all cases of this the neutral falt can possibly be obthis happening in virtue of the super- neutral salts. added body uniting with the Other:

There is another kind of compound falts, formed by the coalition of acids with earthy and metallic bodies. These salts differ from the true neutral ones in feveral obvious properties: some of them change blue vegetable juices to a green like alkalies, and others to a red like acids, while neutral falts make no change in the colour: mixed with boiling milk, they coagulate it, while neutral falts rether prevent its coagulation: from most of them, the acid is disunited by fire, without the intervention of any additional matter, of which we bave seen an instance in the distillation of the acid of vitriol: but the most distinguishing, and universal, character of these salts is, that folutions of them, on the addition of any fixt alkali, grow turbid, and deposit their earth or metal. It were to be wished that custom had appropriated some particular name to the falts of this kind, only one of the ingredients of class, to prevent their being confounded, which several of them tained by itself, the separation of have often been, with the persect

	VITRIOLIC ACID.	ACED	MARINE ACID.	ACETOUS
COMMON fixt ALKALI	Vitriolated tartar	Common	Kegenerat- ed fea falt.	Sal diure- ticus.
ALKALI OF SEASALT.	Glauber's falt.	Cubical nitre	Sea falt.	A fait himlar
VOLATILE ALKALI	Philotophic fal ammon.	Volatile nitre	niac.	Spiritus Mindereri
CALCARE- OUS EARTH.	ociciities.	Calcareous nitre	Calcareous muriaric falt	A jubaitrin- gent falt.
MAGNESIA	Sal catharti- cus amarus	Purging taits, not diftinguished by any particular name.		
Soluble earth of CLAY.	Alum.	Attringent falts, not diffinguished by any particular name.		

The preceding table exhibits, atone view, the several compound falts resulting from the union of each of the pure acids with each of the common alkalies and foluble earths; the acids being placed on the top, the alkalies and earths on the left hand, and the compound falts in the respective intersections; and is thus to be understood. In the upright columns, under each of the acids, are feen the feveral compound falts resulting from the union of that acid with the respective alkalies and earths on the left side. In the transverse columns, opposite to each particular alkali and earth, are feen the compound falts refulting from the union of that alkali or earth with the respective acids on the top; and, conversely, of each of the compound falts expressed in the table, the component parts are found on the top of the upright column, and on the left fide of the transverse column, in whose intersection that particular falt is placed. Some of these salts have been already treated of in the Materia Medica: but it was thought proper to unite them here into one view, for the greater perspicuity in regard to their composition, and the different properties which their component parts assume in different combinations.

### Crystallization of falts.

This is a general operation on neutral and most of the other compound falts. It depends upon these principles: that water, of a certain degree of heat, dissolves, of any particular salt, only a certain determinate quantity: that on increasing the heat, it dissolves more and more (except only in one instance, common salt) till it comes to boil, at which time both its heat

and dissolving power are at their height: that in returning to its first temperature, it throws off again all that the additional heat had enabled it to dissolve: that independently of any increase or diminution of heat, a gradual evaporation of the fluid itself will occasion a proportional separation of the falt: and that the particles of the falt, in this separation from the water, unless too hastily forced together by sudden cooling, or strong evaporation, or disturbed by external causes, generally concrete into transparent and regularly figured masses, called crystals. The several salts assume, in crystallization. figures peculiar to each: thus the crystals of nitre are hexagonal prisms; those of sea falt, cubes: those of alum, octohedral masses; while fal ammoniac shoots into thin fibrous plates like feathers.

459

The use of preparing salts in a crystalline form is not merely in regard to their elegance, but as a mark of, and the means of fecuring. their purity and perfection. From fubstances not dissoluble in water. they are purified by the previous folution and filtration: by crystallization, one falt is purified from an admixture of fuch other faline bodies as dissolve either more easily or more difficultly than itself. For if two or more falts be diffolved together in a certain quantity of hot water, the falt which requires the greatest heat for its solution in that quantity of water, will first begin to separate in cooling: and if the water is kept evaporating, in an uniform heat, the falt which requires most water in that heat will be the first in crystallizing. In all cases of this kind, if the procels is duly managed, the first shootings are generally well figured and pure: the fucceeding ones. fooner or later according to the quantity quantity of the other falts in the liquor, retain an admixture of those falts, which they betray by their smallness and by their

figure.

In order to the crystallization of faline folutions, it is customary to boil down the liquor, till fo much of the fluid has exhaled, as that the falt begins to concrete from it even while hot, forming a pellicle upon the furface exposed to the air: when this mark appears, the whole is removed into a cold place. This method seldom affords perfect crystals: for when water is thus faturated with the falt in a boiling heat, and then fuddenly cooled; the particles of the falt run hastily and irregularly together, and form only a confused semitransparent mafs. It is by flow concretion that most falts assume their crystalline form in perfection. The evaporation should be gentle, and continued no longer, than till some drops of the liquor, in a heat below boiling, being let fall upon a cold glas plate, discover crystalline filaments: the liquor is then immediately to be removed from the fire into a less warm, but not a cold place: and the vessel covered with a cloth to prevent the access of cold air, and the formation of a pellicle, which falling down through the fluid, would difturb the regularity of the crystallization. This is the most effectual method for most falts; though there are fome, whose crystallization is to be effected, not by an abatement of the heat, but by a continued equable evaporation of the fluid; fuch in particular is sea salt.

Salts retain in crystallization a portion of the aqueous sluid, without betraying any marks of it to the eye; on this their crystalline form appears in great measure to depend. The quantity of phlegm or water varies in different falts: dry crystals of nitre were found, on several careful trials, to contain about one-twentiethof their weight; those of alum, one fixth; sea salt, one-fourth; borax, green vitriol, and the purging salts, no less than one half. The same salt appears always to retain nearly the same quantity.

Some falts diffolve in spirit of wine; and here also, as in water, the solution is limited, though the salt is not easily recovered in a crystalline form. Such in particular are, combinations of the nitrous acid with volatile alkalies, and with calcareous earths; of the marine acid with all the soluble earths; of the acetous with faxt and volatile alkalies. Scarce any of the compound salts, whose acid is the vitriolic, are affected by

vinous spirits.

Salts differ greatly in their dispofition to assume and retain a crystalline form. Many, even of the compound kind, imbibe humidity like fixt alkalies, so as to crystallize with difficulty; and when crystallized, or exsiccated by heat, to deliquiate again in the air. Such are the combinations of the nitrous and marine acid with all the foluble earths, and of the acetous both with earths and alkalies. vitriolic acid, on the other hand, forms, with all the fubftances it dissolves, permanent crystals; as do likewise the other mineral acids with all alkalies.

The crystallization of those salts, which are not dissoluble in spirit of wine, is generally promoted by a small addition of that spirit; which, absorbing the water, or weakening its dissolving power on the salt, disposes the salt to part from it more freely. The operator must be careful however not to add

t00

too much of the spirit, especially where the fait is composed of an earthy or metallic body united with the acid; lest it absorb the acid as well as the water, and inflead of a gradual and regular crystallization, hastily precipitate the earth or metal in a powdery form.

Mr. Rouelle, of the French academy of sciences, has examined with great attention the phænomena of the crystallization of salts, and published the result of his observations in different volumes of the Memoirs of that academy. Among other curious particulars, he has given a general distribution of falts, in regard to their crystallization, which will be of prac-

tical utility to the artist.

He divides evaporation into three degrees; insensible evaporation, or that effected by the natural warmth of the atmosphere, from freezing up to the heat of the fummer's fun; mean evaporation, commencing with the fun's heat, and extending to that in which the exhaling stream is visible to the eye, and the liquor too hot to be endured by the hand; and frong evaporation, reaching from this period to boiling. He divides falts into fix classes; the distinctions of which are taken from the degree of evaporation in which they crystallize most perfectly, from the figure of their crystals, their disposition to remain single unite into clusters, receiving an increase from a continuance of the crystallization.

I. The first class confists of falts which crystallize into small plates or very thin scales. The crystals are single. They are, of The crystals all falts, those which crystallize most frequently on the furface of their folutions, which retain least water in their crystals, and require most They to dissolve in. crystallize most perfectly by insensible evaporation.

Salts whose crystals are cubes, cubes with the angles truncated, or pyramids of four or fix sides. They form single, and change their figure by new accretions. By infentible evaporation they crystallize at the bottom, by mean evaporation at the furface, and by both kinds they prove perfect and regular: by ftrong evaporation, the liquor contracts a pellicle, and in cooling yields few crystals, and those ill figured:

III. Salts whose crystals are ] tetrahedral, pyramidal, parallelopipeds, rhomboidal, and rhomboidal parallelopipeds; with the angles variously truncated according to different circumitances. They form fingle (except that fome few unite by the bases) and change their figure by new accretions. They crystallize at the bottom, most perfectly by infensible evaporation: by mean and strong evaporation, the liquor contracts a pellicle, and in cooling the crystals adhere to the pellicle, and prove confuled and ill formed. They retain a large quantity of water.

ġ,

Green Cubical

Salts whose crystals? are flattened parallelopipeds, with the extremities terminating in two furfaces inclined to one another, so as to form a point and acute angles with the large fides. They cluster together, uniting by the bases, The crystals into tufts. are largest and most regular by insensible evaporation: by mean and artar umited hasty evaporation, a pellicle is formed, and in cooling the crystals prove tartar. They revery small. tain a large quantity of water in crystallization, and require little to diffolve in.

Salts whose crystals? are very long, in form of needles, prisms, or columns of different sur-They shoot at the bottom, and cluster together into tufts of regular figures. By inevaporation **fensible** they scarce ever crystallize well. By mean and strong evaporation, they give a pellicle, and in flow cooling, if the evaporation was not carried too far, they yield perfectly well formed crystals, which at first fwim, but foon fall to the bottom. If the evaporation was too long continued, the crystals prove confused and ill formed.

al ammon. Philof. Jal ammon. Nitre. Vol. nitre. Glauber's Jalt. Salt of amber. Vineg. united with chalk. Volat. vitri. acid united with fixt alk. VI. Salts whose crystals are in very small needles. or of other indeterminate None of them figures. crystallize by insensible evaporation, and few of them by the mean degree. They require to be reduced, by strong evaporation, to a thick confistence; they then contract a pellicle, and crystallize with confufion. If the crystals are wanted regular, spirit of wine must be used, or some other medium if the falt is soluble in spirit. They readily disfolve in water, and liquefy in the air.

Sal diureticus. Marine acid united with absorbent earths. Nitrous acid united with absorbent earths.

### NITRUM PURIFICATUM. Purified nitre. Lond.

Boil nitre in water till it is diffolved; filter the folution through paper; and then, after due evaporation, fet it by in a cold place, that the nitre may shoot into crystals. Edinb.

The liquor, which remains after the crystallization, may be further evaporated, and set to shoot as before; but this process must not be too long protracted.

COMMON nitre contains usually a considerable proportion of sea falt, which in this process is separated, the sea falt remaining dissolved after greatest part of the nitre has crystallized. The crystals which shoot after the sirst evaporation, are large, regular, and pure: but when the remaining liquor is surther evaporated, and this repeated a second or third time, the crystals prove at length small.

fmall, imperfect, and tipt with appears to be the very fame spelittle cubical glebes of sea salt.

When rough nitre, in the state wherein it is first extracted from the earths impregnated with it (fee page 183) is treated in this manner, there remains at last a liquor, called mother-ley, which will no longer afford any crystals. This appears to participate of the nitrous and marine acids, and to contain an earthy matter dissolved by those acids. On adding alkaline lixivia, the earth is precipitated, and when thoroughly washed with water, proves infipid. the liquor be evaporated to dryness, a bitterish saline matter is left, which being strongly calcined in a crucible, parts with the acids, and becomes, as in the other case, insipid.

This earth has been celebrated as an excellent purgative, in the dose of a drachm or two; and, in fmaller doses, as an alterant in hypochondriacal and other disorders. This medicine was for fome time kept a great secret, under names of magnefia alba, nitrous panacea, count Palmer's powder, il polvere albo Romano, poudre de Seneinelli, &c. till Lancifi made it public in his notes on the Metallotheca Vaticana. It has been supposed that this earth is no other than a portion of the lime commonly added in the elization of nitre at the European nitre-works: but though the specimens of magnefia examined by Neumann, and some of that which has lately been brought hither from abroad, gave plain marks of a calcareous nature; yet, the true magnesia must be an earth of a different kind, calcareous earths being rather aftringent than purgative. The earthy basis of the sal catharticus amarus is found to have the properties ascribed to the true magnetia of nitre, and

cies of earth: from that falt therefore this medicine is now prepared, as will be seen hereaster.

#### SAL AMMONIACUS PURIFI-CATUS.

Purified sal ammoniac Lond.

This falt is purified by folution in water, filtration, and crystallization, after the manner above directed for nitre.

Edinb.

The liquor remaining after the crystallization is to be further evaporated, and the cryftallization repeated, fo long as any falt will shoot from it.

THE impurities of fal ammoniac are commonly such as will not disfolve in water: and hence the purification is effected by the folution and filtration. The very last crystals seldom betray an admixture of any other falt.

#### FLOS SALIS AMMONIACI. Flowers of fal ammoniac. Edinb.

Take any convenient quantity of dry sal ammoniac in powder: put it into an earthen cucurbit; and having fitted on a blindhead, sublime the falt with a fire gradually increased.

This process seems to be intended with a view to the further purification of the falt. As falammoniac, however, carries up with it substances which of themselves are not volatile, as it is originally prepared by a fimilar process, and may possibly suffer some alteration in its quality from repetitions of it; the fublimation does not appear to be either needful or Neumann expedient. observes. that by repeated fublimations, it acquires at length a yellowish

tinge

tinge, and a particular fmell, of which it discovered nothing at first, and which he attributes to the extrication of the oily or inflammable matter of its volatile animal falt; for that sal ammoniac participates of an inflammable principle, appears from its deslagration with nitre.

#### VITRIOLUM PURIFICA-TUM, vulgo GILLA VITRIOLI.

Purified white vitriol, commonly called gilla of vitriol.

Edinh.

Diffolve white vitriol in a sufficient quantity of warm water, filter the solution, and evaporate it to the consumption of two-thirds: fet the remainder in a cold place, that the salt may shoot into crystals upon the sides of the wessel, and afterwards dry the crystals in the sun. The remaining liquor is to be surther evaporated, and set to crystallize as before; and this process repeated till no more salt will shoot.

SOLUTIONS of white vitriol deposite, on standing, a yellow ochery fubstance; which, if not suffered to separate before the liquor is exhaled and fet to shoot, will foul the crystals. Wilson directs the vitriol to be dissolved in just as much water as will keep it from crystallizing, viz. two pounds or two pounds and a half of water to one of the vitriol; and the filtered folution kept warm, to settle, for twenty-four hours: being then evaporated to a proper pitch for crystallization, a yellow matter is still frequently found at the bottom, from which the liquor must be decanted before it is fet by to shoot. It may be observed, that the separation is by far the most plentiful and speedy while the liquor boils: solutions, which had stood in the cold for some days, and appeared perfectly clear, on being made to boil, became immediately turbid, and threw off a yellow other.

### SAL VITRIOLI. Sal of vitriol. Lond.

Take of

White vitriol, one pound; Strong spirit (called oil) of vitriol, one ounce by weight;

Water as much as is sufficient. Boil them together till the vitriol is diffolved, then filter the liquor, and after due evaporation set it by in a cold place to cryftallize.

HERE the intention is not to feparate the othery matter of the vitriol, but to prevent its separating and colouring the crystals. This is effectually answered by the addition of the acid, by which it is kept diffolved.

### ALUMEN USTUM. Burnt alum. Land.

Let alum be calcined in an iron or earthen veffel, so long as it bubbles and swells up.

THE bubbling or bliftering proceeds from the phlegm retained in the crystals; after that is expelled, the salt cannot be made liquid by any degree of fire. Alum is composed of vitriolic acid and an earth; and it is remarkable, that combinations of that acid with all earths, with most metals, and even with vegetable fixt alkalies, are unfassible.

The alum thus deprived of its phlegm, proves confiderably ftronger, and more acrid, infomuch .as to be fometimes employed for confuming fungous flesh: it is faid to have an inconvenience of leaving. a hardness upon the part.

## VITRIOLUM CALCINATUM. Calcined vitriol. Lond.

Let green vitriol be calcined in an earthen vessel, with an open fire, till it becomes thoroughly dry: then breaking the vessel, take out the vitriol, and set it by for use, well closed from the air. The vitriol is sufficiently calcined, if it has acquired a red colour at the sides and bottom of the vessel.

This process succeeds tolerably well for small quantities, but does not answer so perfectly for larger. As the action of the fire is exerted first on the external parts of the mass, these will be calcined first, and, where the quantity is large, exhibit the mark of sufficient calcination, whilst the internal part remains almost unchanged: and even if the process is still farther continued, the effect required will not be produced; for the outlide growing first hard, prevents the evaporation of the aqueous parts from within.

Edinb.

Expose any quantity of powdered green vitriol, in an unglassed earthen vessel, to the action of a moderate fire, till it becomes white: keeping the matter continually stirring, to prevent its sticking to the vessel, and acquiring a stony hardness. If this be urged with a more vehement fire, it passes into a deep red substance, called colcothar of vitriol.

This method is sufficiently troublesome: for unless the heat be

very gentle, and the matter spread very thin over the bottom of a broad shallow vessel, it is almost impossible to avoid melting it, which makes it adhere to the sides of the pan, and renders the previous pulverisation an useless labour.

The method usually practised by the chemists is, to place a deep earthen pan, with some vitriol in it, upon a gentle fire; the vitriol foon liquefies, boils up, and by degrees incrustates to the sides of the vessel: some more vitriol is then thrown in and suffered to incrustate in the same manner, and this procedure repeated till the pan is nearly full of the concretted matter, which proves of a whitish colour, except on the outfide next the pan (which must be broken to take it out) where it appears yellowish or reddish, according to the continuance and degree of fire. If the vitriol be delired still farther dephlegmated, this may be commodiously effected, by reducing the mass into a gross powder (which will now no longer melt) and then calcining it over a strong fire, in a shallow iron pan, till it has gained the degree of dryness required, which may be known from its colour. - The principal use of calcined vitriol is for the distillation of the spirit of vitriol: if employed for this purpose uncalcined, it would melt in the distilling vessel, and running into a lump, scarce give out any spirit, and the little obtained would . be very weak.

#### TARTARUM VITRIO-LATUM, Vitriolated tartar. Lond.

Diffolve eight ounces of green vitriol in four pints of boiling water: and whilft the liquor conti-H h nues nues boiling, throw into it salt of tartar, or any other alkaline salt, till no farther effervescence arises upon a fresh addition; which generally happens when some some some salt the more, of the salt have been used. Filter the liquor through paper, and aster due evaporation set it by to crystallize.

HERE the acid of the vitriol forfakes the iron which it was before in possession of, to unite with the alkaline falt: particular care ought to be had that the quantity of alkali be sufficient to fully saturate the acid, otherwise it will It is not deposit all the metal. convenient, even after the faturation seems, from the effervescence ceafing, to be compleated, to throw in a little more of the alkali; for by this means the preparation is fecured from containing any metallic matter; whilst the superfluous quantity of alkali can do no prejudice, as it remains uncrystallized.

It is remarkable, that although the vitriolic acid and fixt alkaline falt do each readily unite with water, and strongly attract moisture even from the air; yet the neutral falt refulting from the combination of these two, vitriolated tartar is one of the falts most difficult of folution, very little of it being taken up by cold water. Hence some have directed the liquor in this process to be filtered whilst very hot, suspecting, that if it was suffered to cool, great part of the falt would be thrown off and left upon the paper. The college, however, have avoided this inconvenience, by ordering a quantity of water which is found to be sufficient for keeping the falt dissolved in the cold, or at least in a moderate warmth.

Edinb.

Take oil of vitriol diluted with equal its quantity of warm water; put it into a large glass vessel, and gradually drop into it oil of tartar per deliquium till the effervescence ceases. Then filter the liquor, evaporate it till a pellicle appears upon the surface, and set it by in a cold place to crystallize.

This is an elegant, and one of the least troublesome ways of preparing this falt. The Edinburgh college, in their former editions, ordered the acid liquor to be dropt into the alkaline: by the converse procedure, now received, it is obviously more easy to secure against a redundance of acidity: for the greater certainty in this point, it may be expedient, as in the foregoing process, to drop in a little more of the alkaline ley than the cessation of the effervescence seems to require.

But though the manner of preparation, here directed, appears to be the most commodious, there is one imperfection in the process, a deficiency in the quantity of wa-There is not near water enough to keep vitriolated tartar difsolved, and of consequence, as fast as the alkaline salt is neutralized by the acid, great part falls to the bottom in a powdery form. the Leyden pharmacopæia, this inconvenience is judiciously provided against: the oil of vitriol is diluted with four times, its quantity of water, and the alkaline ley being gradually dropt into it till the point of faturation is obtained, four times the quantity more of water is added, and the mixture boiled, that fuch part of the falt as had precipitated, may be dissolved: the liquor is then filtered while hot, and fet by to crystallize. In order to

obtain

obtain perfect and well formed cryftals, the liquor should not be set in the cold, but continued in the moderate heat, such as the hand can scarcely bear, that the water may flowly evaporate.

Vitriolated tartar, in small dofes, as a scruple or half a drachm, is an useful aperient; in larger ones, as four or five drachms, a mild cathartic, which does not pass off so hastily as the fal catharticus amarus, or sal Glauberi, and seems to extend its action further. The wholesale dealers in medicines have commonly substituted to it an article otherwise almost useless in their shops, the residuum of Glauber's spirit of nitre: this may be looked upon as a venial fraud, if the spirit has been prepared as formerly directed, and the reliduum dissolved and crystallized; but it is a very dangerous one if the vitriolic acid has been used in an over proportion, and the caput mortuum employed without crystallization; the falt in this case, instead of a mild neutral one of a moderately bitter taste, proving highly acid. The purchaser ought therefore to insist upon the salt being in a crystalline form. crystals, when perfect, are oblong, with fix flat fides, and terminated at each end by a fix-fided pyramid: fome appear composed of two pyramids joined together by the bases, and many, in the most perfect crystallizations I have They seen, are very irregular. decrepitate in the fire, somewhat like those of sea salt, for which they have sometimes been mistaken.

### NITRUM VITRIOLATUM. Vitriolated nitre. Lond.

Dissolve in warm water the mass

,

which remains after the diffillation of Glauber's spirit of nitre: filter the solution through paper, and crystallize the salt.

This salt is not different from the tartarum vitriolatum, being composed of the vitriolic acid, and the alkaline basis of nitre, which alkali is no other than the common vegetable fixt alkaline salt, as salt of tartar or potash: it is, in effect, from the ashes of vegetables, that the nitre prepared in Europe receives its alkaline basis. If any unchanged nitre remains in the mass, it is left dissolved in the water while the vitriolated alkali crystallizes.

### SAL POLYCHRESTUM. Salt of many virtues. Edinb.

Take

Nitre in powder,

Flowers of sulphur, of each equal parts.

Mingle them well together, and inject the mixture, by little and little at a time, into a red-hot crucible: after the deflagration ceases, keep the crucible in the fire for an hour. The falt may be purished by dissolving it in warm water, filtering the solution, and exhaling it to dryness; or by crystallization.

This is another method of uniting the vitriolic acid with the common vegetable fixt alkali. Both the nitre and the fulphur are decompounded in the operation: the acid of the nitre, and the inflammable principle of the fulphur, detonate together, and are dissipated; while the acid of the fulphur, (which, as we have already seen, is no other than the vitriolic acid) remains combined with the alkaline bass of the nitre. The shops, accordingly

cordingly, have substituted to the sal polychrest the foregoing preparation.

### SAL PRUNELLÆ. Edinb.

Take of

Pure nitre reduced to powder, two pounds,

Flowers of sulphur, one ounce.

Melt the nitre in a crucible, and sprinkle into it the sulphur by little at a time. When the deflagration is over, pour out the melted salt upon a clean, dry, and warm brass plate, so as to form it into cakes.

THOSE who prepare sal prufiellæ in large quantities, make use of a clean iron pot instead of a crucible; and when the nitre is melted, and the sulphur designated, take out, the salt with an iron ladel, and pour it into brass moulds kept for this purpose. The previous pounding of the nitre, directed above, may be as well omitted, as occasioning a needless trouble.

This preparation was formerly in great esteem, and is sometimes still ordered in prescription. It is nevertheless built upon an erroneous foundation, which supposed, that the nitre was purified by the deflagration it undergoes upon injecting a little fulphur on it; from proper experiments it appears, that the fulphur is so far from depurating the nitre, or tending to its improvement as a medicine, that it really alters some part of it into a salt, which has quite different properties. The real effect of this process will be easily under-Rood from the preceding one: there, nearly all the nitre is decompounded, and a falt, not different from vitriolated tartar, is found in its place: here, only about one twenty-fourth part of it suffers this change. Boerhave instead of deflagrating the nitre with sulphur, orders it to be only well purified after the common method, and then melted by itself and poured into moulds: the sussing the falt into a less compass, by evaporating the aqueous moisture which had concreted with it in its crystallization; though even in this intention it is not of much use, the quantity of water, which nitre retains, not being very considerable.

#### SAL CATHARTICUS GLAU-BERI.

The cathartic salt of Glauber, commonly called sal mirabile.

Lond.

Diffolve in warm water, the mass which remains after the diffillation of spirit of sea falt: filter the solution, and crystallize the salt.

Edinb.

If the crystals (obtained as above) prove too sharp, dissolve them again in water, filter the liquor, and cautiously evaporate it to such a pitch only as may dispose the falt to crystallize.

THERE is no great danger of the crystals proving too sharp, even when the spirit of salt is made with the largest proportion of oil of vitriol directed under that process. The liquor which remains after the crystallization is indeed very acid; and with regard to this preparation, it is convenient it should be so; for otherwise, the crystals will be very small, and likewise in little quantity. Where a sufficient proportion of oil of vitriol has not been employed in the distillation of the spirit, it is necessary to add some to the liquor,

in order to promote the crystallization of the falt.

The title of this salt expresses Taken from its medical virtues. half an ounce to an ounce, or more, it proves a mild and ufeful purgative; and in finaller doses. largely diluted, a ferviceable aperient and diuretic. The shops frequently substitute to it the fal catharticus amarus (see page 214.) which is nearly of the same quality, but somewhat more unpleasant, and as is faid, less mild to opera-They are very easily diffinguifhable from one another, by the effect of alkaline salts upon solutions of them. The folution of Glauber's saits suffers no visible change from this addition, its own basis being a true fixt alkali: but the folution of the fal catharticus amarus grows instantly white and turbid, its balis, which is an earth, being extricated copiously by the alkaline fait; as in the following process.

#### MAGNESIA ALBA. White magnefia. Edinb.

Diffolye sal catharticus amarus in a sufficient quantity of water. Filter the folution, and add to it a filtered ley of potash, so long as a fresh addition continues to occasion any milkiness. A white powder will precipitate; which, being separated from the liquor, is to be carefully washed in fresh portions, both of hot and cold water, and afterwards dried.

This powder appears to be the same species of earth with that obtained from the mother-ley of nitre (see page 463.) which was for several years a celebrated secret in the hands of fome particular persons abroad. Hoffman, who describes the preparation of the

nitrous magnelia, gives it the character of an useful antacid, a safe and inoffensive laxative in doses of a drachm or two, and a diaphoretic and diuretic, when given in smaller doses of fifteen, or twenty grains. Since this time, it has had a considerable place in the practice of foreign physicians, and now begins to come into effeem among us, particularly in heartburns, and for preventing or removing the many disorders which children are so frequently thrown into from a redundance of acid humours in the first passages: it is preferred, on account of its laxative quality, to the common absorbents, which (unless gentle purgatives are given occasionally to carry them off) are apt to lodge in the body, and occasion a costiveness very detrimental to infants.

Though the preparation of this medicine is now commonly known. its nature and properties are very little understood: whilst some suppose it to possels uncommon virtues, others affirm, that, when duly edulcorated, it is in no refpect different from calcined hartshorn, or any other simple animal, or vegetable earth. The following observation of its real properties will be sufficient to determine

this point.

Magnesia alba, when prepared in perfection, is a white and very subtile earth, perfectly void of fmell or tafte, of the class of those which dissolve in acids. It diffolves freely, even in the vitriolic acid: which, in the common way of making folutions, takes up only an inconsiderable portion of other earths. Combined with this acid, it forms a bitter falt, very easily soluble in water, while the common absorbents form with the same acid almost insipid concretes,

НЬз

very difficult of folution. Solutions of magnefia in all acids are bitter and purgative; while those of the other earths are more or less austere and astringent. A large dose of the magnefia, if the stomach contains no acid to dissolve purge or it, does not duce any sensible effect : a moderate one, if an acid is lodged there, or if acid liquors are taken after it, procures feveral stools, whereas the common absorbents, in the same circumstances, instead of loofening, bind the belly. It is obvious, therefore, that magnefia is specifically different from the other earths, and that it is applicable to useful purposes in medi-

### NITRUM CUBICUM. Cubical nitre.

Dissolve chalk or lime in purified aquafortis, and add the solution by degrees to a solution of Glauber's salt in water, so long as a fresh addition produces any milkiness: a white powder will precipitate; after which the liquor is to be siltered, and, after due evaporation, set to crystallize.

In this process, both the solutions are decompounded, and two new compounds produced. The vitriolic acid of the Glauber's falt unites with the chalk, and forms with it an indiffoluble selenitic concrete, which of course precipitates; while the alkali of the Glauber's falt, and the nitrous acid, unite into a neutral falt, which is separated from the liquor by crystallization: the crystals are rhomboidal, of a cooling tafte, greatly resembling that of common nitre. How far this salt differs from common nitre in its medical virtue, is not known. The process is here inserted, partly, as be-

ing a very instructive one in regard to the transpositions which happen on the mixture of different faline bodies, and partly as affording the most convenient means of obtaining the pure alkaline basis of sea falt. In the distillation of spirit of falt, that basis was disunited from its own acid, and combined with the vitriolic: it is here transferred from the vitriolic to the nitrous: and in page 427, we have given a method of diffipating or destroying the nitrous acid, and leaving the alkali, that was combined with it. pure.

#### SPIRITUS SALIS MARINI COAGULATUS.

Spiris of Sea Salt coagulated.

Lond.

Drop, into Glauber's spirit of sea falt, a ley of any fixt alkaline falt, till all effervescence ceases: then evaporate the mixture to dryness.

This preparation is inferted; under the fame title, in the Wirtemberg pharmacopæia. It has been commonly called regenerated fea falt, though with little propriety, as it differs from that falt in its basis; the common vegetable alkali being here substituted to the mineral alkali of fea falt. How far it differs from fea falt in its medical qualities, I cannot take upon me to determine: it is manifestly sharper in taste, and somewhat more difficult both of solution in water and of susion in the fire.

#### TARTARUS REGENERA-TUS.

Regenerated tartar. Edinb.

Put any quantity of dry fait of tartar, powdered, into a large glass vessel; and pour thereon, by by little and little, as much diffilled vinegar as is necessary to saturate it. Filter the siquor, and exhale it, over a very gentle fire, to dryness, taking great care that the matter contract not an empyreuma. On the falt which remains, pour as much more spirit of vinegar as will saturate it; then depurate the liquor again, and carefully exsicate it into a dry salt.

Ir the common alkalies are made use of for this process, they should be previously purified, by solution and crystallization, from the neutral salt which they generally contain. The distilled vinegar must be perfectly free from any empyreumatic taint: it is not necessary to dephlegmate it, or throw away the first runnings in the distillation, since these contain a portion of the acid (the part here wanted) as well as the phlegm.

It is difficult to hit the point of faturation betwixt the acetous acid. After aand the alkaline falt. bout fourteen parts of strong distilled vinegar have been gradually poured upon one of the fixed falt, the addition of a little more of the acid will not occasion any further effervescence in the cold: but if the mixture be now firongly ftirred and well heated, the effervescence will appear afresh; upon which fome more vinegar is to be added, till it again ceases. faturation is not as yet complete; for upon exhaling the aqueous parts, the remaining falt still effervesces with fresh vinegar. When fo much of the acid has now been added, that no marks of fermentation any longer appear, a little more of the vinegar may be poured in before you proceed to the last evaporation; by this means,

the faturation of the alkali will be fecured, whilft, if the acid prevails, the superfluous quantity of it will exhale.

The falt thus prepared, is of a dark brown colour, a peculiar, not ungrateful odour, a penetrating, saponaceous, saline taste, in no wise alkaline or acid. Its brown colour, and saponaceous quality, proceed from the oily parts of the vinegar; the depuration of the salt from this oil, is not in the foregoing process insisted on. In the London pharmacopæia, the salt is ordered to be purified to perfect whiteness, under the title of

### SAL DIURETICUS. Diuretic falt. Land.

Take a pound of any fixt alkaline falt, and boil it, with a very gentle heat, in four or five times its weight of distilled vinegar. When the fermentation ceases, add more distilled vinegar; and proceed with fresh additions thereof, until the vinegar being almost evaporated, fresh vinegar will no longer raife any fermentation; which generally happens by the time that twenty pounds of distilled vinegar have been Then flowly exhale to uſed. dryness.

Melt the remaining impure falt for a little time, but not too long, over a gentle fire; then diffolve it in water, and filter the folution through paper. If the melting has been duly performed, the filtered liquor will be limpid and colourless as water; but if otherwise, of a brown colour.

Evaporate the limpid folution, with an exceeding gentle heat, in a shallow glass vessel, occafionally stirring the salt as it H b 4 dries,

dries, that its moisture may be the fooner exhaled. Afterwards keep it for use in a vessel very closely stopt; for it will liquely

by the air.

This falt ought to be of perfect whiteness; and should totally dissolve both in water and in spirit of wine, without leaving any feces. If the salt, though ever so white, deposits any feces in spirit of wine; the whole of it must be dissolved in that spirit, the solution siltered, and exsiccated again.

THE purification of this salt is not a little troublesome. The operator must be particularly careful in melting it, not to use too great a heat, or to keep it liquefied too long; a little should be occasionally taken out, and put into water: and as foon as it begins to part freely with its black colour. the whole is to be removed from the fire. In the last drying, the heat must not be so great as to melt it; otherwise it will not prove totally soluble. If the solution in spirit of wine be exsiccated, and the remaining falt liquefied with a very fost fire, it gains the leafy appearance, which has procured it the name terra foliata.

In the fourth volume of the Memoirs of the correspondents of the French academy lately published, Mr. Cadet has given a method of making the falt white at the first evaporation, without the trouble of any further purification. He obferves, that the brown colour depends upon the oily matter of the vinegar being burnt by the heat commonly employed in the evaporation; and his improvement confifts in diminishing the heat at the time that this burning is liable to happen. The process he recommends as is follows.

Dissolve a pound of falt of tartar in a sufficient quantity of cold water, filter the folution, and add by degrees as much distilled vinegar as will faturate it, or a little more. Set the liquor to evaporate in a stone-ware vessel. in a gentle heat not fo strong as to make it boil; when a pellicle appears on the surface, the rest of the process must be finished in a water-bath. The liquor acquires by degrees an oily confiftence, and a pretty deep brown colour, but the pellicle or fcum on the top looks whitish, and when taken off and cooled, appears a congeries of little brilliant filverlike plates. The matter is to be kept continually stirring, till it is wholly changed into this white flaky matter, the complete drying of which is most conveniently effected in a warm oven.

We shall not take upon us to determine whether the pure or impure salt are preferable as medicines; observing only, that the latter is more of a saponaceous nature, the former more acrid, though fomewhat more agreeable to the stomach. Mr. Cadet reckons the falt prepared in his method superior both to the brown and white forts made in the common way, as possessing both the oily quality of the one, and the agreeableness of the other, and as being always uniform, or of the same power, whereas the others are liable to warv confiderably, according to the degree of heat employed in the evaporation. They are all medicines of great efficacy, and may be so dosed and managed as to prove either mildly cathartic. or powerfully diuretic: few of the faline deobstruents come up to them in virtue. The dose is from half

half a fcruple to a drachm or two. A bare mixture of alkaline falt and vinegar without exficcation, is not perhaps much inferior as a medicine to the more elaborate falt: I have known two drachms of the alkali, faturated with vinegar, occasion ten or twelve stools, in hydropic cases, and a plentiful discharge of urine, without any inconvenience.

# SPIRITUS MINDERERI. Spirit of Mindererus. Edinb.

Take any quantity of the volatile alkaline falt of fal ammoniac, and gradually pour upon it diftilled vinegar, till the effervefcence ceafes; occasionally flirring the mixture, to promote the action of the vinegar on the falt.

This is an excellent aperient faline liquor. Taken warm in bed, it proves commonly a powerful diaphoretic or sudorific; and as it operates without heat, it has place in febrile and inflammatory disorders, where medicines of the warm kind, if they fail of procuring sweat, aggravate the dis-Its action may likewife temper. be determined to the kidneys, by walking about in a cool air. common dose is half an ounce. either by itself, or along with other medicines adapted to the intention. Its strength is not a little precarious, depending in great measure on that of the vinegar: an inconvenience which cannot easily be obviated, for the faline matter is not reducible to the form of a concrete salt.

#### SECT. VII.

#### Anomalous salts.

# CRYSTALLI TARTARI. Cryftals of tartar. Edinb.

ET powdered white tartar be boiled in twenty times its quantity of water, till perfectly diffolved; and the folution, whilft it continues hot, passed through filtering paper, or a woollen cloth, and received in a wooden vessel; then expose it for a night or longer to the cold air, that crystals may form themselves, and shoot to the sides of the vessel; the water being now poured off, the crystals are to be collected and dried for use.

THE filtration of the folution of tartar through paper succeeds very slowly, and unless managed

with a good deal of address, not at all; for as foon as the boiling liquor begins to grow fensibly less hot, it deposites much of the tartar all over the furface of the paper, which hinders the remainder from passing through. Zwelffer, in his animadversions on this process in the Augustan pharmacopæia, directs the folution to be clarified with whites of eggs. and strained only through a linen cloth; he likewise judiciously orders the vessel to be close covered, and the crystallization performed a warm place; for if the folution be suffered to cool very fast, it is in vain to expect any appearance of crystals; the tartar will inevitably be precipitated to the bottom of the vessel in the form of sand. And indeed, the bulinels

Part III.

business of refining and crystallizing tartar is fo very troublefome, and requires fo large an apparatus, that scarce any of the apothecaries, or even of the trading chemists, are at the trouble of it; but either import it ready refined from Holland, or purchase it from some people here who make it their sole business. (See the article TARTAR, page 237.)

#### CREMOR TARTARI. Cream of Tartar. Edinb.

Take any quantity of folution of tartar, made as in the foregoing process, and passed through a filter. Boil it over the fire, till a thick cuticle appears on the furface, which is to be taken off with a wooden skimmer, bored full of holes; continue the boiling till a fresh cuticle arises, which is be taken off as the foregoing, and the operation repeated till the whole quantity of liquor is thus confumed. Afterwards dry all the cuticles together in the fun.

This process seems inserted on ly to retain a name long familiar to the shops; for the preparation itself in no respect differs from crystals of tartar reduced to powder. Indeed the purchaser ought always to prefer the crystals; for the powder is often sophisticated with faline substances of another kind.

The college of Edinburgh obferves, that both the crystals and cream are brought to us from abroad; that they are not different in quality from one another; and that good

white tartar, unrefined, is not inferior to either of them.

#### TARTARUM SOLUBILE. Soluble Tartar. Lond.

Dissolve a pound of any fixt alkaline falt in a gallon of boiling water; and gradually throw in crystals of tartar, as long as a fresh addition thereof raises any effervescence; which generally ceases before three pounds of the crystals have been used. Then filter the liquor, and after due evaporation, fet it by to cryftallize; or evaporate it to dryness, and keep the remaining saline mass for use.

#### Edinb.

Boil crystals of tartar, till they are perfectly dissolved, in ten times their quantity of water; and gradually drop into the folution, whilst it continues boiling, oil of tartar per deliquium, till the effervescence ceases. Filter the liquor whilft hot, and evaporate it till a pellicle appears on the furface, that when removed into a cold place, it may crystallize.

Common white tartar is perhaps preferable for this operation to the crystals usually met with (see the article TARTAR, page 237.) Its impurities can here be no objection; fince it will be fufficiently depurated by the subsequent filtration.

The preparation of this medicine by either of the above methods is very easy; though some chemists have rendered it sufficiently troublesome by a nicety that is not at all wanted. They infift upon hitting the very exact point of faturation betwixt the alkaline fair and the acid of the tartar: and caution the operator to be extremely careful, when he comes near this mark, left by imprudently adding too large a portion of either, he rendered the falt too acid, or too alkaline. If the liquor be suffered to cool a little before it is committed to the filter, and then properly exhaled and crystallized, no error of this kind can happen, though the faturation should not be very exactly hit : for fince crystals of tartar are very difficultly foluble even in boiling water, and when dissolved therein, concrete again upon the liquor's growing cold; if any more of them has been employed, than is taken up by the alkali, this superfluous quantity will be left upon the filter: and on the other hand, if too much of the alkali has been made use of, it will remain uncrystallized. crystallization of this salt indeed cannot be effected without a good deal of trouble: it is therefore most convenient to let the acid falt prevail at first, to separate the superfluous quantity, by suffering the liquor to cool a little before filtration, and then proceed to the total evaporation of the aqueous fluid, which will leave behind it the neutral falt required. The most proper vessel for this purpurpose is a stone-ware one : iron discolours the falt.

Soluble tartar, in doses of a scruple, half a drachm, or a drachm, is a mild cooling aperient: two or three drachms commonly loosen the belly; and an ounce proves pretty strongly purgative. Malouin says it is equal in purgative wirtue to the cathartic salt of Glauber. It is an useful addition to the purgatives of the resinous kind, as it promotes their operation, and at the same time tends

to correct their griping quality. But it must never be given in conjunction with any acid, for all acids decompound it; absorbing its alkaline salt, and precipitating the tartar.

SAL RUPELLENSIS.
Sel de Seignette, or Rochel falt.
Pharm. Parif.

Let the falt extracted from the ashes of the kelp or kali of Alicant be calcined till it melts. then dissolve in water, the folution filtered, and after due eyaporation fet by, that the salt may shoot into pure white crystals. Dissolve crystals of tartar in boiling water, and faturate the folution with the crystals of kali: the proportions necessary for this purpole will be, about fixteen ounces of the latter to twenty of the former. Duly exhale the liquor in the heat of a water-bath, and after filtration, fet in the cold to crystallize.

This is a species of soluble tartar, made with the salt of kali or soda, which is the same with the mineral alkali or basis of sea salt: (see page 427.) It crystallizes far more easily than the preceding preparation, and does not, like it, grow moist in the air. It is also considerably less purgative, but is equally decompounded by acids. It appears to be a very elegant salt, and begins now to come into esteem in this country, as it has long been in France.

SAL ESSENTIALE. ACETOSÆ. Esfential falt of forrel. Edinb.

Let the juice of forrel, after fettling and decantation from the feces, be evaporated, till only

one

one-third remains, then strained through a flannel bag, and exhaled again till a pellicle appears upon the surface. Put the liquor into a glass vessel, and, a little oil olive being poured upon the top, fet it by in a cellar till plenty of crystals are formed: thefe are to be gently washed with water, and afterwards dried. After the same manner, essential

falts are obtained from all acid, austere, aftringent, and bitterish plants that contain but a small

quantity of oil.

Herbs of a dry nature are to be juices. moistened, in the bruising, with a little water, that the juice may be the more easily pressed

The waters of these plants, which are in vain endeavoured to be drawn over by distillation, may be obtained by dissolving a suitable quantity of their effential falts in common water.

Some pharmaceutical writers direct the plants to be gathered early in the morning; but this is of very little moment. In order to make the subject yield its juice readily, it should be chopt to pieces, and well bruised in a marble mortar, before it is committed to the press: the magma which remains in the bag, still containing no inconfiderable quantity of faline matter, may be advantageously boiled in water, and the decoction added to the expressed juice. The whole may be afterwards depurated together, either by the method above directed, or by running the liquor leveral times through a linen cloth. In fome cases, the addition of a confiderable portion of water is necessary; that the juice, thus diluted, may part the more freely from its feculencies; on the separation of which, the success of the process in great measure de-

The evaporation should be performed either in shallow glass bafons, or in fuch earthen ones as are of a compact close texture; fuch are those usually called stone-ware. The common earthen vessels are subject to have their glasing corroded, and are so extremely porous, as readily to imbibe and retain a good quantity of the liquor: metallic vessels are particularly apt to be corroded by these acid kinds of

The directions for the time of discontinuing the second evapora-tion are not so easily observed as one could wish. These juices are fo viscid, and abound so much with heterogene matter, of a quite different nature from any thing faline, that a pellicle, or pure saline incrustation upon the surface, is in vain expected. Boerhaave therefore, and the more expert writers in pharmaceutical chemistry, with great judgment, direct the evaporation of the superfluous moisture to be continued until the matter has acquired the confiftence of cream. If it be now suffered to stand for an hour or two in a warm place, it will, notwithstanding the former depurations, deposite a fresh sediment, from which it should be warily decanted before it is put into the vessel in which it is designed to be crystallized.

Some recommend an unglazed earthen vessel, as preferable for this purpose to a glass one; the smoothness of the latter being supposed to hinder the salt from sticking thereto; whilst the juice easily infinuating itfelf into the pores of the former, has a great advantage of shooting its saline spicula to the fides. Others flightly incrustrate

the fides and bottom of whatever vessel they employ, with a certain mineral falt, which greatly disposes the juice to crystallize, which of itself it is very averse to: but as this addition is, with regard to its medical virtue, quite different from the falt here intended, we forbear to mention it.

The use of the oil is to preserve the juice uncorrupted, and to prevent it from running into fermentation or putrefaction, during the great length of time which this process requires: as much oil as will fully cover the furface of the liquor, is sufficient for this purpose. washing of the crystals is intended to cleanse them from the mucilaginous feculencies which adhere to them: it ought to be performed with the utmost caution, to prevent any of the falt itself from being dissolved. The liquor which remains after the crystillization, may be depurated by a gentle colature, and after due inspissation let to shoot again; when a farther yield of crystals will be obtained.

The process for obtaining these falts is very tedious, insomuch as scarce to be completed in less than feven or eight months; and the quantity of salt which the juices afford, is extremely small: hence they are hardly ever made or expected in the fhops. The chemists have contrived several methods for expediting the process, among which the two following feem the most

remarkable.

Take any quantity of wormwood, carduus benedictus, or the like plants, gently dried in the shade. Pour thereon a suitable portion of spirit of wine, and digest them together with a foft heat, till the menstruum has acquired a green colour. This tincture is

to be put into a glass cucurbit. and distilled with the heat of a water-bath, till so much of the spirit is come over, as that the remainder may be left of the confiftence of honey. The whole being now fuffered to remain unmoved till grown perfectly cold, beautiful pyramidal crystals will be found to have shot from the sides of the distilling vessel towards its center. Spiessius, in Miscell. Berolin. continuat. ii. p.

91, 92. This gentleman relates likewise, that having made an essence (that is, a faturated tincture) of elecampane roots, with spirit of wine, and kept it unmoved for a year, he found a great number of crystals shot from the bottom of the glass upwards, of the thickness of a quill, and about an inch long.—The crystals obtained by this method are faid to be of the nitrous kind, but of a more fubtile tafte than the common nitre, impressing only an agreeable cool-

ness upon the tongue.

The fecond process is from the celebrated Dr. Stahl:

Take wormwood, brooklime, pellitory, mercury, foapwort, or any other plants of the same kind, dried quick in a shady place. Cut the herb small, and pour thereon a sufficient quantity of highly-rectified spirit of wine: digest them together till the menstruum becomes saturated with the oil, or refinous parts of the plant; then pour off the tinged liquor, add a fresh parcel of spirit, and digest as before, continuing to add more of the menstruum, till such time as it no longer extracts any colour from the vegetable. The plant thus freed from its oily matter, is to be gently exficcated,

and boiled in water, till the liquor has taken up its faline parts: the decoction being then passed through a silter, afterwards evaporated to a due consistence, and set by in a cool place, will shoot into saline crystals, which, on examination, prove manifestly nitrous. Stablii fund. chem. pag. 68, et alibi.

THE two foregoing processes agree but ill with one another: how far they are adequate to the purpofes intended by them, has not yet been sufficiently examined. It is certain, that spirit of wine diffolves the fubtile oils and the refins of vegetables, which prove a great impediment to the crystallization of falts; from whence it should feem that the falt might afterwards be prepared by water from the residuum to much better advantage. But it is certain also. that this menstruum dissolves some of the native vegetable falts themfelves; and that if the tincture is fufficiently loaded with the foluble parts of the subject, the salt separates, while the oily and refinous matter remain dissolved. Thus manna, an effential falt of the sweet kind, dissolves totally in rectified fpirit, and, however foul before, is recovered white as snow, its oily impurities being left in the menftruum; and thus spirituous tinctures of celery, beet-roots, and other plants of the sweet kind, deposite, on standing, true saccharine concretions. It is probable that one process is best adapted to some plants, and the other to others: the first doubtless is for those of the fweet kind, and the fecond for acid herbs, as forrel and woodforrel.

The virtues of the effential falts have not been sufficiently deter-

mined from experience. Thus much, however, is certain, that they do not, as has been supposed, possess the virtues of the subjects entire, excepting only the acids and fweets. The others feem to be, almost all of them, nearly similar, whatever plant they were obtained from. In watery extracts of wormwood, carduus, chamomile, and many other vegetables, kept for some time in a soft state, I have often observed fine saline efflorescences on the furface, which had all nearly the same taste, somewhat of the nitrous kind. They are supposed by some to be at bottom no more than an impure species of volatile nitre (that is, a salt composed of the nitrous acid and volatile alkalies:) those which were examined by the chemists of the French academy, deflagrated in the fire, and, being triturated with fixt alkalies, exhaled an urinous odour; plain marks of their containing those two ingredients.

SACCHARUM LACTIS.

Sugar of milk.

Pharm. Parif.

Take common whey of cows milk, made with calves rennet. Clarify it with whites of eggs; and, if it is not perfectly limpid, pass it through a filter. Then evaporate it, in a glass vessel, in the heat of a water-bath, and set it by in a cellar to crystallize. The crystals are to be washed with cold water.

This preparation has been greatly celebrated in diforders of the breaft, but is far from answering what has been expected from it. It has little sweetness, and is difficult of folution in water. A faline substance, much better deserving the name of sugar, may

be obtained by evaporating new milk, particularly that of the afs, to drynefs, digesting the dry matter in water till the water has extracted its soluble parts, and then inspissating the filtered liquor. This preparation is of great sweetness, though neither white nor crystalline: nor is it perhaps in the pure crystallizable parts of milk, that its medicinal virtues lie.

### FLORES BENZOINI. Flowers of Benzoine. Lond.

Put fome powdered benzoine into an earthen pot placed in fand; and with a gentle heat fublime the flowers into a conical paper cap fitted to the pot.

Or, the sublimation may be performed in a retort; the slowers will arise with a soft heat, into the neck.

If the flowers have any yellow tinge, mix them with tobaccopipe clay, and fublime again.

#### Edinb.

The fublimation is to be performed in a glazed earthen pot, and repeated in the fame inftruments with fresh parcels of benzoine, till the paper cap becomes foul with oil.

BENZOINE, exposed in a retort to a gentle fire, melts and sends up into the neck white, shining crystalline slowers, which are followed by an oily substance. On raising the heat a little (a recipient being applied to the neck of the retort) a thin yellowish oil comes over, intermingled with an acid liquor, and afterwards a thick buty-raceous substance; this last, liquefied in boiling water, gives out to it a considerable quantity of saline matter (separable by sitration and proper exhalation) which

]

۶

\$

appears in all respects similar to the flowers.

It appears therefore, that the whole quantity of flowers which benzoine is capable of yielding, cannot be obtained by the above processes, since a considerable portion arises after the time of their being discontinued: that greatest part of the flowers arises with a less degree of heat than what is necessary to elevate the oil: but that if the operation is hastily conducted. or if the fire is not exceeding gentle, the oil will arise along with the flowers, and render them foul. Hence in the way of trade, it is extremely difficult to them of the requisite whiteness and purity; the heat which becomes necessary, when large quantities of the benzoine are employed, being so great as to force. over some of the oil along with

In order therefore to obtain these flowers in perfection, only a fmall quantity of benzoine should be put into the vessel at a time: and that this may not be any impediment to the requifite dispatch. a number of shallow, flat-bottomed, earthen dishes may be employed, each fitted with another vesfel inverted over it. With these you may fill a fand-furnace: having fresh dishes charged in readiness to replace those in the furnace, as foon as the process shall appear finished in them: the refiduum of the benzoine should be scraped out of each of the vessels. before a fresh parcel is put in.

These flowers, when made in persection, have an agreeable taste and fragrant smell. They totally dissolve in spirit of wine; and likewise, by the assistance of heat, in water; but separate again from the latter upon the liquor's growing

cold,

cold, shooting into saline spicula, which unite together into irregular masses. By the mediation of fugar, they remain fuspended in cold water, and thus form an elegant balsamic syrup. Some have held them in great efteem, as pectoral and sudorific, in the dose of half a scruple or more: but the present practice rarely makes use of them, on account of the offenfive oil which, as usually pre-pared, they are tainted with, and from which a fresh sublimation from tobacco-pipe clay does not free them fo effectually as might be wished. The observations above related, point out a method of depurating them more perfectly, viz. by folution, filtration, and crystallization.

SAL SEDATIVUS. Salt of borax, called sedative falt. Put eight ounces of powdered borax into a wide-necked retort: pour thereon three ounces of water: and then add three ounces of oil of vitriol. Place the retort in a proper furnace, adapt to it a receiver, and increase the fire till the vessel becomes red hot. The fedative falt will arise into the neck, in form of thin shining plates, which are to be fwept out with a feather: and a little liquor will pass into the receiver. When the matter in the retort is grown cool, pour back upon the distilled liquor, and fublime again. Repeat this procels to long as the borax continues to yield any considerable quantity of faline flowers.

Or,
Diffolve the borax in a fufficient
quantity of warm water, and
add thereto the oil of vitriol.

Evaporate this mixture, till thin plates begin to appear upon the furface; then fuffer the fire to decay, and let the veffel fland unmoved, till plenty of crystals are formed, which are to be well rinsed with cold water, and then dried for use.

In the preparation of this falt by fublimation, the fire must be expeditionly raised when the matter begins to grow dry, for it is only at this period that the falt The sublimed falt itself, fublimes. in a perfectly dry state, proves fixt in the fire: if moiltened with water. and then exposed to a smart heat. part of it continues to rife, till the moisture is wholly exhaled a after which, nothing more can be forced up by heat, till the falt is again moistened. Hence the use of returning the distilled liquor, and repeating the sublimations. Lemery fays, he found flowers continue to rife till the thirty-fixty sublimation : and that the quantity obtained by all these sublimations amounted to half an ounce and thirtyfive grains from two ounces of borax.

The part of the borax which does not sublime, appears to be the fame (when the common refined borax of the shops is made use of) with the alkaline salt of fea fait: the fedative fait, united with that alkali, recomposes borax again. The extrication of the sedative salt from the borax happens on the same principle as that of the marine acid from fea salt, viz. the vitriolic acid uniting with the alkali; and the residuum is in both cases the fame, viz. the falt called fal mirabile, or Glauber's falt: the fedative falt may be extricated also from borax by other acids, but but most commodiously and effec-

tuelly by the vitriolic.

The progress by crystallization is less troublesome than that by sublimation; but the falt proves generally less white, and is apt likewise to retain a part of the Glauber's salt, especially if the evaporation is too long protracted.

The sedative salt appears to the tuste a neutral salt; but examined with alkalies has the properties of an acid, effervescing, uniting, and crystallizing with them, and destroying their alkaline quality. It dissolves both in water and in spirit of wine; though not very readily in either. As to its virtues, it is supposed to be a mild anodyne, (whence its name) to calm the heat of the blood in burning fevers, to prevent or remove delirious fymptoms, and allay spasmodic affections, whether hypochondriacal or hysterical, at least for a time. The dose is from two to eighteen grains, in any proper liquor.

### SPIRITUS, SAL, ET OLEUM SUCCINI.

Spirit, Salt, and oil of amber. Lond.

Diftil amber in a fand heat gradually increased; there will come over a spirit, an oil, and a salt fouled with the oil.

The oil distilled again by itself, is divided into a thinner oil which arises; and a thicker part that remains behind, called bal-sam of amber.

The falt is to be boiled in the diffilled spirit, or in common water, and set to crystallize; by this means it is freed from its adhering oil. The oftner this is repeated, the purer it will be.

Edinb.

Mix powdered white amber with

thrice its weight of clean fand. and put them into a glass retort. of which the mixture may fill one half; then adapt a large receiver, and distil in a sandfurnace, with a fire gradually increased. At first a spirit will come over, with fome yellow oil: then more yellow oil, along with a little falt; and upon railing the heat, more of the falt, with a reddish coloured oil. When the distillation is finished, empty the liquor out of the receiver; and having collected together the falt which adheres to the fides, dry it by gentle preffure between the folds of some spongy paper.

The oil may be separated from the spirit by siltration: and afterwards received by distilling it

from brine of sea salt.

The falt is to be rectified in the following manner. Grind it well with twice its quantity of sea salt, and put the mixture into a tall and narrow glass cucurbit; sit on a blind-head, and proceed to sublimation in a sand heat, taking care that the oil does not rise. When the vessels are grown cold, sweep out the salt with a seather.

In the distillation of amber, the fire must for some time be continued gentle, fcarce exceeding the degree at which water boils, till the aqueous phlegm and thin oil have arisen: after which it is to be flowly increased. If the fire was urged hastily, the amber would swell up, and rife in its whole substance into the receiver, without undergoing the required decomposition or separation of its parts. When fand or other like intermedia are mixed with it, it is less subject to this rarefaction, and the fire may be . Tailed

raifed fomewhat more expeditioufly; though this little advantage is perhaps more than counterbalanced by the room which the fand

takes up in the retort.

Our chemists generally leave the receiver unluted, that it may be occasionally removed as the salt riles and concretes in the neck of the retort, from whence it is every now and then scraped out to prevent the oil from carrying it down into the receiver. When a gross thick oil begins to arise, and no more salt appears, the distillation is stopt, though it might, perhaps, be continued longer to advantage.

Pott informs us, (in a Mr. ... curious differtation on the falt of amber puplished in the volume of the Memoirs of the academy of sciences of Berlin) that the Prussian workmen, who prepare large quantities of the falt for exportation, from cuttings and fmall pieces of amber, perform the distillation without any intermedium, and in an open fire: that sweeping out the salt from the neck of the retort being found too troublesome, they suffer the oil to carry it down into the receiver, and afterwards separate it by means of bibulous paper, which imbibes the oil, and leaves the falt dry; which paper is afterwards fqueezed and distilled: that they continue the distillation till all that can be forced over has arisen, with care only to catch the last thick oil in a separate receiver; and that from this they extract a confiderable quantity of falt, by shaking it in a strong vessel with three or four fresh portions of hot water, and evaporating and crystallizing the filtered wa-

The spirit of amber so called, is no more than a solution of a

fmall portion of the falt in phlogen or water; and therefore is very properly employed for diffolying the falt in order to its cryftallisation.

The last, freed from as much of the oil as fpongy paper will imbibe, retains fo much as to uppear of a dark brown colour, Mr. Pott fays, the method he has found to succeed best, and with least loss, is, to dissolve the falt in hot water, and put into the paper, through which the foliation is to be filtered, a little cotton flightly moistened with oil of amber; this, he fays, detains a good deal of the oil of the falt, and the folution paffes through the more pure, The liquor being evaporated with a very gentle fire, as that of a water-bath, and fet to shoot, the first crystals prove transparent, with a flight yellowish tinge, but those which follow are brown, oily, and bitter, and are therefore to be further depurated in the fame manner. The whole quantity of cryftals amounts to about onethirtieth of the weight of the crude amber employed. By fublimation from sea salt, as directed above. the falt is more perfectly and more expeditiously purified: Mr. Pott objects to sublimation, that a part of the falt is decomposed by it, a coaly matter being left behind, even though the falt was previously purified by crystallization: it may be prefumed, however, that this coal proceeds rather from the burning of fome remains of the oily matter, than from the decomposition of any part of the true falt.

Pure falt of amber has a penetrating, subastringent acid taste. It disloves, both in water and in rectified spirit; though not readily in either, and scarcely at all in the latter without the assistance of

heat ;

heat: of cold water in fummer, it requires for its folution about twenty times its own weight, of boiling water only about twice its weight. Exposed in a glass vessel, to a heat a little greater than that of boiling water, it first melts, then rifes in a white sume, and concretes again in the upper part of the glass, into fine white flakes, leaving, unless it was perfectly pure, a little coaly matter behind. It effervesces with alkalies both fixt and volatile, and forms with them neutral compounds, greatly refembling those composed of the same vegetable acids, and Mixed with acid liquors, it makes no fenfible commotion. Ground with fixt alkaline falts, it does not exhale any urinous odour. By these characters, it is conceived, this falt may be readily distinguished from all the other matters that have been mixed with, or vended for it. With regard to its virtue, it is accounted aperient, diegretic, and, on account of its retaining some portion of the water.

oil, antihysteric: Boerhauve gives it the character of diureticorum et antihystericorum princeps. Its great price, however, has prevented its coming much into use; and perhaps its real virtues are not equal to the opinion generally entertained of them.

The rectified oil has a strong bituminous fmell, and a pungent, acrid tafte. Given in a dose of ten or twelve drops, it heats, stimulates, and promotes the fluid fecretions; it is chiefly celebrated in hysterical disorders, and in deficiencies of the uterine purgations. Sometimes it is used externally, in liniments for weak or paralytic limbs, and rheumatic pains. oil differs from all those of the vegetable kingdom, and agrees with the mineral petrolea, in not being soluble, either in its reclified or unrectified state, by spirit of wine. fixt alkaline lixivia, or volatile alkaline spirits; the oil, after long digeftion or agitation, separating as freely as common oil does from

#### CHAPTER IX.

#### Preparations of Sulpbur.

FLORES SULPHURIS. Flowers of Sulphur. Lond.

UBLIME fulphur in proper vef-I fels; and reduce the flowers, that concrete, into powder, either in a wooden mill, or in a marble mortar with a wooden pestle.

Edinb.

Put any quantity of yellow fulphur, grossly powdered, into an earthen cucurbit placed in a fand-furnace; and having fitted on a glass blind-head, or inverted upon it another earthen cucurbit, begin the sublimation with a gentle heat, which may be afterwards increased. The flowers will arise into the uppermost part of the veffels, from whence they are to be swept out and carefully washed with very hot water.

This process is rarely attempted by the apothecaries, a large apparatus being necessary for performing it to advantage. Those who prepare the flowers of brimstone in quantity, use for the subliming veffel a large iron pot, capable of holding two or three hundred weight; this communicates with an arched chamber, lined with glazed tiles, which ferves for the recipient.

This preparation of fulphur makes no change in its qualities; only separating its impurities, and at the fame time reducing it into a finer powder than it can easily be brought to by other means. At the bottom of the subliming vessel there remains a ponderous grey-coloured

ters, with a small portion of sur!phur that has escaped the subliming heat. This is usually broken in pieces, and vended in the shops under the name of fulphur vivum.

#### FLORES SULPHURIS LOTI. Washed flowers of Sulpbur. Lond.

Pour upon the flowers as much water as will arise to the height of four fingers above them, and boil them for fome time: then pouring off this water, let some cold water be added, and thoroughly wash the flowers; after which they are to be dried for use.

As the flowers of fulphur are generally sublimed into very capacious rooms, which contain a large quantity of air, or in vessels not perfectly close; some of those that arise at first, are apt to take fire, and thus are changed into volatile acid vapour, which mingling with the flowers that fublime afterwards. communicates to them a notable degree of acidity. In fuch case the ablution here directed is for the general use of the medicine absolutely necessary; for the flowers, thus tainted with acid, fometimes occasion gripes, and may, in other respects, be productive of effects different from those of pure sulphur. The Edinburgh college, as appears in the foregoing process, allow only the washed flowers to be kept in the shops: there are, however, some particular combinations, to which they are supposed to be better adapted when unwashmais, composed of fand, earth, ed, as their union with mercury stony, and sometimes metallic matinto ethiops mineral; and accordingly

ingly for that preparation the un- This is made after the same manwashed flowers are directed by the London college.

#### **BALSAMUM SULPHURIS** SIMPLEX.

Simple balfam of Sulpbur. Lond.

Boil flowers of fulphur, with four times their weight of oil olive, in a pot flightly covered, until they unite into the confistence of a balfam.

#### **BALSAMUM SULPHURIS** CRASSUM.

Thick halsam of sulphur. Edinb

Take a pint of linfeed oil, and four ounces of flowers of fulphur. Boil them together over a gentle fire, keeping them continually firring, till they come to the confishence of a ballam.

LINSEED oil more readily difsolves sulphur than oil olive, and the preparation made with it is reckoned fomewhat less disagreeable. The vessel they are boiled in ought to be capable of holding at least three times the quantity of As foon as the the ingredients, oil begins to act upon the fulphur, which happens nearly at the point of ebullition, the mixture rarifies very much, so as, if not prudently removed from the fire, to run over into the furnace; and as the matter is very susceptible of flame, dangerous consequences may enfue, especially if the quantity is The operator ought therelarge. fore to be upon his guard in the management of this process.

BALSAMUM SULPHURIS BARBADENSE. Balsam of sulphur with Barbadoes tar. Lond.

ner as the foregoing, by using Barbadoes tar inflead of the oil.

#### BALSAMUM SULPHURIS TEREBINTHINATUM.

Balfam of fulphur with oil of turpentine. Edinb.

Take two ounces of washed flowers of fulphur, and fix ounces of oil of turpentine.

Digest them together, in a sandheat, till the oil is saturated with the fulphur.

#### BALSAMUM SULPHURIS ANISATUM.

Balsum of sulphur with oil of aniseed. Edinb.

Take two ounces of washed flowers of fulphur; fix cunces of oil of turpentine; and four ounces of essential oil of anifeeds.

Digest them together as in the preceding process.

THESE preparations are more conveniently and fafely made in a tall glass body, with the mouth at least an inch in diameter, than in the circulatory or close vessels in which they have commonly been directed to be prepared; for when the fulphur and oil begins to as vehemently upon each other, they not only rarify into a large volume, but likewise throw out impetuously great quantities of an elastic vapour, which, if the vessels are closed, or the orifices not sufficient to allow it a free exit, infallibly burft them: Hoffman relates a very remarkable history of the effects of an accident of this kind. In the veilel above recommended, the process may be completed, without danger, in four or five hours, by duly managing the fire; which should be very gentle for li3

some time, and afterwards increas- the virtues ascribed to them when ed so as to make the oil just bubble or boil, in which state it should be kept till all the fulphur appears to

be taken up.

Effential oils employed as menftrua for sulphur, undergo a great alteration from the degree of heat, necessary for enabling them to disfolve the fulphur; and hence the balfams have not near so much of their flavour as might be expected. It should therefore seem more eligible to add a proper quantity of the effential oil to the simple balfam; there readily incorporate by a gentle warmth, if the veffel be now and their shaken. parts of essential oil, and fix of the balfamum fulphuris craffum compole a balfam more elegant than those made in the foregoing manner, and which retains so much of the flavour of the oil, as in some measure sufficient to cover the taste of the fulphur, and render it Tupportable.

The balfams of fulphur have been firongly recommended coughs, confumptions, and other diforders of the breaft and lungs. But the reputation which they have had, in these cases, does not appear to have been built upon any fair trial, or experience of their vir-They are manifestly hot, acrimonious and irritating; and therefore should be used with the They have freutmost caution. quently been found to injure the appetite, offend the stomach and viscera, parch the body, and occasion thirst and sebrile heats. The dose of the simple balsam is from ten to foty drops; those with esfential oils are not given in above half these quantities. Externally,. they are employed for cleanling and healing foul running ulcers; Boerhaave conjectures, that their which in these cases give occasion to

taken internally.

HEPAR SULPHURIS. Liver of sulphur. Edinb.

Take three ounces of flowers of fulphur; and one ounce and a half of powdered falt of tartar. Melt the fulphur in an earthen dish, under a chimney, and add to it by degrees the falt of tartar: keeping the matter constantly flirring with a spatula till it has acquired a red colour: care must be had to prevent its taking

IT is much more convenient to melt the fulphur first by itself, and add the falt of tartar by degrees, as here directed; than to grind them together, and afterwards endeavour to melt them as ordered in former editions; for in this last case, the mixture will not flow sufficiently thin to be properly united by Rirring; and the fulphur either takes fire, or sublimes in flowers, which probably has been the reason why so large a propor-tion of it has been commonly directed. Even in the prefent method a confiderable part of the fulphur will be dissipated; and if it was not, the hepar would not be of its due quality: for one part of fulphur requires two of the alkaline falt, to render it perfectly foluble in water, which this preparation ought to be.

The hepar sulphuris has a setid fmell, and a nauseous taste. Solutions of it in water, made with fugar into fyrup, have been recommended in the same intentions as the ballams above-mentioned: our pharmacopæias nevertheless have deservedly rejected this syrup, as common practice has almost done the balsams. The hepar, di-

gested in rectified spirit of wine, imparts a rich gold colour, a warm, somewhat aromatic taste, and a peculiar, not ungrateful smell: a tingure of this kind is kept in the shops, under the name of another suneral,

## SULPHUR PRÆCIPITATUM. Precipitated fulphur. Lond

Boil flowers of fulphur in water, with thrice their weight of quick-lime, till the fulphur is dissolved. Filter the folution, and drop into it some of the weak spirit of vitriol: this will throw down a precipitate, which is to be washed in fresh portions of water, till it becomes inspid.

### LAC SULPHURIS.

Boil the hepar fulphuris, reduced to powder, in four times its quantity of water for three hours; adding more water if there is occasion. Then filter the folution whilft hot, and drop into it spirit of vitriol, till the effervescence ceases; a powder will be precipitated to the bottom, which is to be washed with hot water, and afterwards dried for use.

THE method of preparing this lac, as it is called, with hepar fulphuris, is the most expeditious, and least troublesome, provided the hepar be well made: and, on the other hand, quicklime gives the preparation a more faleable whiteness. Some have been accustomed to add to the quicklime a portion of alkaline salt, with a view to promote its dissolving power.

The medicine is searly the fame in both cases. It would be exactly the same, if the precipitation

was performed with any other acid than the vitriolic: for this acid forms with the dissolved lime a selenitic concrete, which precipitates along with the fulphur, and is not afterwards separable by any ablation: whilft the neutral fait, which the acid forms with the fixt alkali of the hepar, may be totally diffolved, and washed off by repeated ablution with hot water; and the. combinations of all the other acids, both with the lime and alkali, are feparated by cold water. It is probably to the admixture of the white felenitic matter, refulting from the vitriolic acid and lime, that the finer colour of the preparation made with lime is owing.

Pure lac fulphuris is not different in quality from pure sulphur itself: to which it is preferred, in unguents, &c. only on account of its colour. The whitehes does not proceed from the sulphur having lost any of its parts in the operation, or from any new matter superadded: for if common sulphur be ground with alkaline salts, and set to sublime, it arises of a like white colour, the whole quantity of the alkali remaining unchanged; and if the lac be melted with a gentle fire, it returns into yellow

fulphur again.

It may be observed, that the name lac fulphuris, or milk of sulphur, applied among us to the precipitate, is by the French writers confined to the white liquor before the precipitate has fallen from it.

TINCTURA SULPHURIS VOLA-

Volatile tindure of fulpbur.

Take of

Flowers of fulphur, fix ounces; Sal ammoniac, one pound;

Quicklime, a pound and a half.

Sprinkle fome water on the lime,
and when flaked and fallen into
f i 4

powder,

ture in a retort, with a fire gradually increased. The distilled close stopt, for use.

THIS liquor has a strong offenfive smell, somewhat similar to that which arises in the precipitation of the virtues of this preparation. He lac sulphuris. The vapour in both cases spreads to a considerable diftincture with three of spirit of wine, tance, changes silver or copper in a dose of thirty or forty drops, attensils to a brown or blackish proves a most powerful diaphoretic; colour, and produces difagreeable and that a liquor composed of this alterations in many medicinal preparations: to this circumstance the gout, by bathing the feet with therefore due regard ought to be it. This tincture may be a powerhad in the performance of that pro- ful medicine, but it is certainly a cefs, and in the keeping of this very unpleafant one. tincure. If a piece of paper, writ-

powder, grind it first with the ten upon with a saturated solution fulphur, and afterwards with the of lead in vegetable acids, and fal ammoniac, in small quanti- gently dried, be placed in the midties at a time: then distil the mix- dle of a quire of paper, or of a pretty thick book, and brought near the unftopt orifice of the bottle conliquor is to be Rept, in a bottle taining this tincture, the vapour will quickly reach it, and change the colourless writing to a legible black.

Hoffman has a great opinion of fays a mixture of one part of the

Metallic preparations.

#### SECT. I.

### Preparations of gold.

OLD is the most ponderous and perfect of the metals: it abides fixt and unaltered in the strongest fire; and is not acted upon by alkaline, or any fimple acid menstruum. It dissolves in aqua regis glone, into a yellow transparent fluid: this folution. stains the fkin, &c. purple: the ethereal spirit of wine, and some effential oils, take up the gold from it: alkalies precipitate the metal in form of a yellowish mud, which exsiccated, and exposed to a small beat, violently explodes.

As to the medicinal virtues of this metal, experience has sufficiently shown; that it is not possessed of any valuable ones. In its metallic form, however finely comminuted, it proves inactive; when fatiated with acid, corrofive; and in the intermediate flates, either fignificant or unfafe. 

AURUM POTABILE. Potable gold. 19 11/11

111111

Diffolve with a moderate heat, half a drachm of fine gold, in two ounces of aqua regia; and add to the folution one nunce of the essential oil of rosemary. - Shake them regether, and then faffer them to reft: the acid lofes its gold yellow colour; and the oil, which arifes to the furface; becomes richly impregnated therewith. Separate the oil by decantation, and add to iv four or

five ounces of reclified spirit of wine: digest this mixture for a month, and it will acquire a purplish colour.

THERE have been many preparations of this kind contrived by the deligning pretenders to alchemy, and imposed upon the credulous and unwary, as cordials and diaphoretics of inestimable value. above feems to be one of the best and fafest of them; though it would be equally serviceable as a medicine, if made without the ingredient, which it receives its name from: The gold is indeed taken up from the acid, and kept for a time dissolved by the oil; but on standing it totally separates, in form of fine yellow films, like leaf-gold. The effect is the same, whether the oil or the vinous spirit be mixed with the folution of the gold in aqua regis : the only difference is, that the gold is thrown off from the oil so the lides of the glass; while the spirit revives it into such subtile films, as to float upon the furface of the liquor. No means have yet been found of permanently combining gold with either oils or vinous Spirks.

> · AURUM PULMINANS. Fulminating gold. Parif.

Put a drachm of filings of gold, , with half an ounce of aqua regia

newly made, into a matrass, placed in fand. When the menstruum ceases to act, pour off the folution; and, if any of the gold is left, add as much more aqua regia as shall be sufficient to diffolve it. Dilute the solution with ten times its quantity of warm water : and then drop in oil of tartar per deliquium till the effervescence and precipitation cease. The whole being now fuffered to fettle, the clear liquor is to be poured off, and the precipitated matter washed with warm water till it becomes inlipid, and afterwards exficcated.

This powder requires to be exficcated with the utmost precaution; for in a fmall heat it explodes with great wolence: the

same effect ensues likewise upon strongly rubbing it. This property has given name to the preparation; and is the only one on account of which it is at present taken any notice of. It has been recommended indeed, in fevers, as a diaphoretic, in the dose of a few grains: its more certain effect, however, is to operate downwards, and that not always with fafety; Konig and Ludovici relate, that in some sebrile cases, it has occasioned almost mortal diarrhoeas; and Stahl (de proexeucrisi medica, sect. vii.) reports, that the intellines have been found croded by it. The more thoroughly it is washed and edulcorated, the less corrolivé it is in the human body, and the less vielently it fulninates when heated.

### SECT. II.

# Preparations of silver.

SILVER is the most permanent: in the fire of all the metals after gold. It disloves in the pure nitrous acid, into a colousless staffinarent liquor, intensity bitter and corrosive. This folution exacested, furnishes the shops with an useful caustic; which has likewise; been taken internally in small doses, and mixed with other substances, as; an hydragogue: it stains the skim black.

## CAUSTICUM LUNARE. The lunar cauftic. Lond.

Let pure filver be diffolved in about twice its weight of aquafortis, upon warm fand; then gently increase the heat, until a dry mass it left. Melt this in a crucible, that it may be poured into proper moulds, cantilly

aroiding overmuch heat; left the matter should grow too thick,

CAUSTICUM LUNARE, fou,
LAPIS INFERNALIS.
The luner couplic, or informal flore.
Edinb.

Take any quantity of well-aupalled filver, flatted into plates and cut in pieces. Dissolve, it by the heat of a fanti-bath, in three ' \* pines its sneight of spirit of nitre. .. Evaporate the foliation, to:dryneika and put the remaining galx in a large crucible. Let the fire at first be gentle, and angment at by segrees, until the mais flows like oil, and confer to fume: then bonk it into it on hipes made for this purpate, previously mested and greated : lattle, let it be dried, and kept for use in a glass vessel close steps. STRONG

STRONG spirit of nitre will disfolve somewhat more than half its weight of pure filver; and the weaker of the aquæ forses, formerly deferibed, proportionably less, according to their quantity of pure nitrous acid. Sometimes this ipirit contains a portion of the vitriolic, or marine acids; which, however minute, renders it unfit for diffolving this metal, and should therefore be carefully separated before the solu-The method tion is attempted. which the refiners employ, for examining the parity of their aquafortis, and purifying it if necessary. is, to let fall into it a few drops of a perfect folution of filver already made: if the liquor remains clear, and grows not in the least turbid or whitish, it is fit for their use: otherwise, they add a smallquantity more of the folution, which immediately turns the whole of a milk white colour; the mixture. being then suffered to rest for some time, deposites a white sediment; from which it is warily decanted, examined afresh, and, if need be, farther purified, by a fresh addition of the folution. See page

The filver, flatted into thin plates as directed in the second of the above processes, needs not be cut in pieces; the folution will go on the more speedily, if they are only turned round into spiral circumvolutions, so as to be conveniently got into the glass, with care that the feveral furfaces do not touch one another: by this management,. a greater extent of the surface is exposed to the action of the menstruum, than when the plates are cut in pieces and laid above one another. Good aquafortis will diffolve about half its weight of ver. and it is not adviseable to use a greater quantity of the menitruum than is sufficient for effecting the is required, as in the laying open

folution; for all the furplus must be evaporated in the fubfequent fusion.

The crucible ought to be large enough to hold five or fix times the quantity of the dry matter; for it bubbles and fwells up greatly, fo as otherwise to be apt to run over: during this time, also, little drops are now and then spirted up, whose caudicity is increased by their heat. and which the operator ought therefore to be on his guard against. The fire must be kept moderate till this ebuilition ceases, and till the matter becomes confiftent in the heat that made it boil before; then quickly increase the fire till the matter flows thin at the bottom, like oil, on which it is to be imnvediately poured into the mould. without waiting till the fumes cense to appear, for when this happens, the preparation aroves not only too thick to can freely into the mould. but likewife less corrosive than it is expected to be.

In want of a proper iron mould, one may be formed of tempered tobacco-pipe clay, not too moift, by making in a lump of it, with a smooth slick first greafed, as many holes as there is occasion for; pour the liquid matter into these cavities, and when congealed, take a out by breaking the mould. Each piece is to be wiped clean from the greafe; and wrapt up in dry foft paper, not only to keep the air from acting upon them, but likewife to prevent their corroding or discolouring the fingers in hand-

This preparation is a strong cauflic, and frequently employed as fuch, for cooluming warts and other fleshy excrescences, keeping down fungous flesh in wounds or ulcers, and other like uses. It is rarely applied where a deep eschar

of

of imposshumations and tumours; for the quantity necessary for these purposes, liquefying by the moisture of the skin, spreads beyond the limits in which it is intended to operate.

PILULE LUNARES.

The lunar pills.

Dissolve pure silver in aquasortis, as in the foregoing process, and after due evaporation, set the liquor to crystallize. Let the crystals be again dissolved in common water, and mingled with a solution of equal their weight of nitre. Evaporate this mixture to dryness, and continue the exsiccation with a gentle heat, keeping the matter constantly stirring, till no more summer arise.

HERE it is necessary to continue the fire till the fumes entirely cease, as more of the acid is required to

be dissipated, than in the preceding process. The preparation is, nevertheless, in taste very sharp, intenfely bitter and nauseous; applied to ulcers, it acts as a caustic. but much milder than the forego-Boerhaave, Boyle, and others. greatly commend it in hydropic The former affures us, that two grains of it made into a pill. with a crumb of bread and a little fugar, and taken on an empty stomach, (fome warm water, fweetened with honey, being drank immediately after) purge gently without griping, and bring away a large quantity of water, almost without the patient's perceiving it: that it kills worms, and cures many inve-terate ulcerous disorders. He nevertheless cautions against using it too freely, or in too large a dole; and observes, that it always proves corrofive and weakening, especially to the flomach.

#### SECT. III.

### Preparations of iron.

RON calcines by fire the most easily, and melts the most difficultly of all the metals. Sulphur promotes its sussion, and changes it into a substance not greatly dissimilar to a combination of the metal with vitriolic acid. All acids dissolve this metal; even the air corrodes it into a rust or calx.

Iron, in its metallic form, or lightly calcined, or combined with vegetable or with mineral acids, acts in the human body in the fame manner, (but with different degrees of power) by constringing the fibres. In all these states, it promotes, or restrains secretions, where the deficiency or excess proceed from a laxity and debility of the vessels;

and, in general, raifes the pulfe, and quickens the circulation. The calces feem to be the least active preparations; the crude metal, duly comminuted, is more easily soluble in the animal sluids, and if acescent juices are lodged in the primæ viæ, soon manifests its operation by nidorous eructations, and the black colour of the alvine feces: if previously combined with saline bodies, it scarce ever fails of taking effect.

As the calces of iron are fearcely diffoluble in acids, it has been concluded that they are not foluble in the human body, and that therefore they are to be looked upon no otherwise than as a mere inactive

canto

But admitting the absolute indiffolubility of iron while it continues a calx, it must be observed, that the calces of this metal are remarkably easy of revival into their metallic state. Mr. Baume relates, that calx of iron, digested for an hour or two in oil olive, resumes its perfect metalic nature, so as to be attracted by the magnet, and totally foluble in acids: from whence he infers, that a like revival of the metal happens in the human It is matter of common obfervation, that calces of iron tinge the excrements black, a fure mark of their taking effect; though their effects appears to be neither so speedy not so great as that of iron in fome other forms.

#### CHALYBIS RUBIGO PRÆPARATA. Ruft of fleel prepared.

Expose filings of theel to the air, frequently moistening them with vinegar or water, until they change into rust; then grind them in a mortar, and pouring on water, wash over the more subtile powder. The remainder is to be exposed afresh to the air, and moistened as at first, then triturated and washed again, and the powders that have been washed over, dried, and kept for use.

#### MARTIS LIMATURA PRÆPARATA. Filings of Iron prepared. Edinb.

Set filings of iron, first cleansed by the magnet, in a moist place, that they may turn to rust, which is to be ground into an impalpable powder.

They may likewife be prepared by moistening them with vinegar.

THE cleanfing of iron filings by means of a magnet is very tedious, and does not answer so well as might be expected; for if they are rufty, they will not be attracted by it, or not sufficiently; nor will they, by this means, be entirely freed from brass, copper or other metallic fubitances which may adhere to them. It appears from the experiments of Henckel (Pyritolog. cap. vom eisem im kiess) that if iron be mixed by tufion with even its own weight of any of the other metals, regulus of antimony alone excepted, the compound will be vigourously attracted by the loadstone.—The rust of iron is to be procured at a moderate rate from the dealers in iron, free from any impurities, except such as may be washed off by water.

The rust of iron is preferable as a medicine to the calces, or croci, made by a strong fire. Hoffman relates, that he has frequently given it with remarkable success, in obflinate chlorotic cases, accompanied with excessive headaches, and other violent symptoms; and that he usually joined with it pimpinella. arum root, and falt of tartar, with a little cinnamon and fugar. dose is from four or five grains to twenty or thirty; some have gone as far as a drachm; but all the preparations of this metal answer best in small doses, which should rather be often repeated than enlarged.

### ÆTHIOPS MARTIALIS. Martial etbiops.

Put filings of steel into an unglazed earthen vessel, with so much water as will stand above them about four inches; the whole is to be well stirred every day, and more water supplied, as that in the vessel exhales, so that the filings may remain always covered: continue this procedure fore-

feveral months, till they lose their motalic aspect, and are reduced so a fine powder of an inky blackness.

THIS preparation is described by Lemery in the Memoirs of the French academy. It is faid, that if the filings are left unstirred for fome days, they unite together fo firmly that the mass is scarcely to be beaten into a powder by blows of a bankmer; if they are lest for a little while uncovered wish water, the deep black colour does not succeed, a part of them changing into ruft. Mr. Malouin observes, that this ethiops is better fitted for general use, than any other preparation of iron; that the metal is here in as subtile a state as in the croci of iron; and that it is no more decompounded, or changed in its nature, than the crude filings are. He therefore recommends fubilitating it to the filings and croci, in doles of from four grains to eighteen. The tedioninels of the process, however, has prevented its coming into use : especially as it does not promise any advantage above the common chalybeate preparations, to counterbalance that inconvenience.

### CHALYBIS CUM SULPHURE PRÆPARATUS.

Steel prepared with fulphur. • Lond.

Heat the steel with a very sierce fire to a strong white heat; and in this state apply it to a roll of salphur held over a vessel of water: the steel will melt and fall down in drops, which are to be picked out from the sulphur that runs down with them, and ground into an impalpable powder.

Ir has been supposed by many,

that this preparation is no other than common brimftone, and that it contains nothing of the steel. If the steel indeed is not made extremely hot, it will not melt on applying it to the sulphur, and the latter will run into the water by itself; but if the metal is heated to the degree above directed, it will readily melt and fall down in drops of a brown colour; whilst the sulphur runs into long yellow strings.

The heat requisite for this purpole, is not easily procurable in the common furnaces of the apothecary; and even if the steel is fusficiently heated at first, it will foon become too cool to be corroded by the fulphur. For this reason, and on account of the offensive fumes which arise very copiously, and which are not avoidable by the operator, this process has been long neglected. The shops have been generally supplied with a preparation of fleel with fulphur made at an easier rate in the following manner.

## MARS SULPHURATUS. Sulphurated iron. Edinb.

Mix filings of iron with twice their weight of powdered sulphur, and as much water as is sufficient to make them into a paste, which on standing at rest for six hours, will swell up. The matter is then to be pulverized, put by degrees into a hot crucible to designate, and kept continually stirring with an iron spatula till it falls into a deep black powder.

If the quantity of this mixture is confiderable, and strongly pressed down, it will not only swell on standing for some hours, but will heave up very weighty obstacles, and burst out into stame.

CRO

### CROCUS MARTIS APERIENS.

Opening crocus of iron This is made by keeping the foregoing preparation longer over the fire, till it assumes a red colour.

> :CROCUS MARTIS. ASTRINGENS. Aftringent crocus of iron. Edinb.

This is made from the opening crocas of iron, by reverbarating it for a long time in the most extreme degree of heat.

THESE preparations differ from one another in virtue; though the difference is not of fuch a kind as the titles, they have been usually distinguished by, import. All the preparations of steel act by an aftringent quality; that above. denominated aftringent, seems to have the least effect. They may be given in form of bolus, electary, or pills, from ax grains to a scraple.

In fome foreign pharmacopæias, the croci of iron are prepared from This strongly pure green vitriol. calcined (or the colcothar remaining after the distillation of oil of vitriol) is the aftringent crocus: when less calcined, it is called aperint. These preparations differ little, if at all, from those above diffinguished by the same appellatious: and accordingly the Edingburgh college has now allowed the substitution of colcothar of vitriol to both the croci.

#### MARS SOLUBILIS, seu CHALYBS TARTARIZATUS. Soluble, or tartarized steel. Edinb.

Mix equal parts of iron filings, and crystal of tartar, with as much them into a mais: this mais is to be dried in a fand heat a then powdered, moistenened and dried again; and this process repeated. till such time as the matter will easily grind into an impalpable powder.

This is a very elegant and useful preparation of steel, and will in many cases take effect after all the foregoing ones have failed; the falt here joined rendering the metal fufficient soluble in the animal fluids. It may be given either in a liquid form, or in that of a bolus, &c. in doles of four or five grains, or half a scruple. Dr. Willis is faid to have been the inventor of this preparation, and by his same it has been usually distinguished in the shops. The chemists have received another method of preparing iron with tartar; which is as fol-

MARS SOLUBILIS ALCALIZA-TUS.

Alkalized soluble steel.

Take equal quantities of filings of iron, and of white tartar. Grind them together, and put them into a crucible, which is to be fet in a fire strong enough to make the materials red hot; in this state, let them continue for some time, When grown cold, powder the matter in a mortar; and the part which will not pals through a fine fieve, heat and pulverize again; repeating this, until the whole was passed through. Mix the several liftings together, and keep them in a reffel close ftopt from the air.

This preparation is foluble like the foregoing. Exposed to the air, it deliquates like alkaline salts (the tartar being converted into an alkali by the fire) and therefore it. water as is fufficient to reduce is not to be prescribed in any

dry form. It is very rarely made use of.

#### FLORES MARTIALES. Martial flowers. Lond.

Take of

Colcothar of green vitriol washed, or filings of iron, one pound;

Sal ammoniac two pounds. Mix and fublime in a retort. Grind the flowers with the matter which remains in the bottom of the retort, and repeat the fublimation until the flowers arise of a beautiful yellowish colour. To the refiduum you may add half a pound of fresh sal ammoniac, and sublime as before; repeating this as long as the flowers arise well coloured.

THE success of this process depends principally upon the fire being hastily raised, that the sal ammoniac may not fublime before the heat is become strong enough to enable it to carry up a sufficient Hence glass quantity of the iron. vessels are not so proper as earthen or iron ones; for when the former are made use of, the fire cannot be raised quick enough, without endangering the breaking of them. The most convenient vessel is an iron pot : to which may be luted an inverted earthen jar, having a fmall hole in its bottom, to suffer the elastic vapours, which arise during the operation, to escape. It is of advantage to thoroughly mix the ingredients together, moisten them with a little water, and then gently dry them; and to repeat the pulverization, humectation, and exficcation, two or thee times, or oftener. If this method is follow-·· ed the fal ammoniac may be increased to three times the quantity of the iron, or farther; and a fingle sublimation will often times be fufficient to raile flowers of a ver v

deep orange colour.

This preparation is supposed to be highly aperient and attenuating: though no otherwise so than the rest of the chalybeates, or at most, only by virtue of the faline matter joined to the iron. It has been found of good service in historical and hypochondriacal cases, and in diffempers proceeding from a laxity and weakness of the solids as the rickets. It may be conveniently taken in the form of a bolus, from two or three grains to ten; it is nauseous in a liquid form (unless in spirituous tincure) and occasions pills to fwell and crumble, except fuch as are made of the gums.

#### LIXIVIUM MARTIS. Ley of iron. Lond.

Let the matter which remains after the sublimation of the martial flowers be fet by in a moist place: it will run into a liquor which is to be kept for use.

THIS liquor feems greatly to resemble a saturated solution of iron made in spirit of salt : its taste is highly astringent, and fomewhat sweetish. It may be given in doses of a drop or two in any convenient vehicle, for the fame intentions as the other chalybeates. It is called by some of the chemical writers, oleum martis per deliquium, and efsentia martis.

#### SAL MARTIS. Salt of fleel. Lond.

Take of Strong spirit or oil of vitriol, eight ounces. Iron filings, four ounces, Water, two pints.

Mix

Mix them together; and after the ebullition ceases, let the mixture stand for fome time upon warm fand: then pour off and filter the liquor; and after proper exhalation, fet it by to crystallize.

VITRIOLUM MARTIS, feu SAL CHALYBIS. Vitriol of iron, or falt of steel. Edinb.

Take of

Oil of vitriol diluted with equal its quantity of warm water, four ounces :

Filings of iron, three ounces.

Cautiously mix them together, and digest in a cucurbit for twelve hours, that the metal may be dissolved; filter the folution whilst hot, then evaporate it till a pellicle appears on the surface, and fet it in a cold place, until the vitriol has crystallized at the bottom of the vessel. The liquor poured off from the crystals is to be again evaporated till a pellicle forms on the top, and fet to shoot as before. Collect all the crystals together, and dry them on a paper in the shade.

During the dissolution of the iron, a strong sulphureous vapour arises, which on the approach of flame, catches fire; and explodes, to as fometimes to burit the veffel: to this particular therefore, the operator ought to have due rezard.

The chemists are seldom at the trouble of preparing this falt according to the directions above given; but in its stead substitute common green vitriol, purified by folution in water, filtration, and crystallization. The only difference betwixt the two is, that the comvitriol contains somewhat more metal in proportion to the acid; and hence in keeping, its

green colour is much sooner debased by a rusty brownish cast. The superfluous quantity of metal may be easily separated, by suffering the folution of the vitriol to stand for some time in a cold place. when a brownish yellow ochery sediment will fall to the bottom 2 or " it may be perfectly diffolved, and kept suspended, by a suitable addition of oil of vitriol. If the vitriol is suspected to contain any cupreous matter, (which it does not appear that the common English vitriol ever does, though almost all the foreign vitriols do) the addition of fome bright iron wire to the folution will both discover, and effectually separate that metal: for the acid quits the copper to dissolve a proportionable quantity of the iron : and the copper, in its separation from the acid, adheres to the undiffolved iron, and forms a skin of a true copper colour upon its surface. Even a vitriol of pure copper may, on this principle, be converted into a pure vitriol of iron.

But though the vitriolic acid appears, in this operation, to have for much stronger a disposition to unite with iron than with copper, that it totally rejects the latter upon prefenting the former for it to act upon; the operator may, nevertheless, give a dangerous impregnation of copper to the purest and most saturated solution of iron in the vitriolic acid, by the use of copper vessels. If the martial solution be boiled in a copper veffel, it never fails to dissolve a part of the copper, diftinguishable by its giving a cupreous strain to a piece of bright iron immersed in it. addition of the iron, the copper is separated; by boiling it again without iron, more of the copper is dissolved; and this may in like manner be feparated by adding

more iron. K k

The falt of Reel is one of the most efficacious preparations of this metal and not unfrequently made use of, in cachectic and chlorotic cases, for exciting the uterine purgations, strengthening the tone of the viscera, and destroying worms. It may be conveniently taken in a liquid form, largely diluted with aqueous fluids; Boerhaave directs it to be dissolved in an hundred times its quantity of water, and the folution to be taken in the dose of twelve ounces, on an empty fromach, walking gently after it: thus managed, he fays, it opens the body, purges, proves diuretic, kills and expels worms, tinges the excrements black, or forms them into a matter like clay, strengthers the fibres, and thus cures many differ-The quantity of ent distempers. vitriol in the above dose of the folution is fifty-feven grains and a half: but in common practice, such large doses of this strong chalybeate are never ventured on. Four or five grains, and in many cases half a grain, are sufficient, for the intentions in which chalybeate medicines are given. Very dilute folutions, as that of a grain of the falt in a pint of water, may be used as succedanea to the neutral chalybeate waters, and will in many cales. produce fimilar effects.

### SECT. IV.

### Preparations of copper.

topper is lefs eafy of folution than iron; and, in its metallic ftate, does not appear to be acted on by the animal fluids, or to have any confiderable effect in the body. Diffolved, it proves extersally an efcharotic; internally, a violent purgative and emetic. Acids of every kind diffolve it, and likewife volatile alkalies. With the vegetable and marine acids, it forms a green folution; with the vitriolic acid, and volatile alkalies, a blue.

Æs ustum.

Let thin copper plates be firatified in a crucible with fulphur; and calcined with a firong fire until they become pulverable.

PREPARATIONS of this kind, made with fulphur, nitre, and common falt, or mixtures of these, or by calcining the copper without addition, have been sometimes used in external applications, for drying and cleanling ulcers, and preventing the growth of fungous flesh; and sometimes likewise internally. They are still retained in some soreign pharmacopæias, but have not for a long time been taken notice of among us, for any medicinal intention.

CRYSTALLI VENERIS.

Cryftals of copper.

Diffolve pure copper in thrice its weight of aquafortis, adding the metal to the acid by little and little at a time. Evaporate the liquor by a gentle heat, till one half of it is wasted; then set the remainder in a cool place to crystallize: afterwards dry the crystals, and keep them in a vial close stopt from the air.

THESE crystals are strongly cauflic, similar to the causticum lunare; but are so much disposed to liquesy, that they are scarce ever made use of, and cannot easily be preserved.

Tinc-

Tinctura veneris volatilis.

Volatite tincture of copper.

Take of

Copper filings, one drachm; Spirit of fal ammoniac, twelve drachms.

Let them fland together in a close vessel, frequently shaking it, until the liquor is tinged of a beautiful violet colour.

This tincture, or folution, of copper has been given internally, in the dose of a few drops, as a diuretic. Beerhaave directs at first three drops to be taken in a morning fasting, with a glass of mead, and this dose to be daily doubled till it comes to twenty-four drops; which last quantity is to be continued for some days; he says, that by this means, he cured an hydropic person sabouring under a confirmed ascites; and that the medicine procured furprifing discharges of urine; that nevertheless, on trying it in another case of the same kind, it did not answer. See the article Cuprum, page 132.

### ENS VENERIS.

Take of

Colcothar of blue vitriol well edulcorated with water, and afterwards dried.

Sal ammoniac, of each equal parts.

Reduce them separately into powder; then mix, and put them into an earthen cucurbit, so as to fill two-thirds thereof. Place the cucurbit in an open fire, and having adapted to it a glass blind-head, apply at first a gentle heat, which is to be increased by degrees, and continued as long as the flowers arise of a yellow colour inclining to red: when the vessels are grown cold, let

the flowers be carefully fwept out with a feather.

If the blue vitriol be perfectly good, this process will not succeed in the manner here fet down: where it does succeed, that is, where the flowers prove of a reddish yellow colour (ex luteo rubentes) it is to be prefumed, that the fuccess is owing to the vitriol partaking largely of iron, and that the preparation is not greatly different from the flores martiales of the preceding fection. The colour of blue vitriol is undoubtedly owing to copper; but most of the common vitriols of this kind contain, also, no inconsiderable quantity of iron; and a reddish yellow colour of the flowers may be looked upon as a mark, that it is chiefly or folely the iron that the sal ammoniac has carried up. For this is the colour which iron always gives in its fublimates with fal ammoniac : whereas copper, in all its folutions, or foluble combinations with fal. ammoniac, or other faline bodies, gives a blue or green, or a colour compounded of these two.

The process is originally taken from Mr. Boyle, who tells us, that he and a chemist, endeavouring to imitate Butler's stone by a preparation of calcined virtiol, and finding the medicine upon trial, though far short of what Helmont ascribes to his, yet no ordinary one, they called it, for the mineral's sake it was made of, ens primum veneris.

The composition of vitriols was at that time but impersectly known; and this is not the only instance of an effect being ascribed to the cupreous part of a vitriol, which was owing to the ferugineous. Though Boyle looked on the preparation as proceeding from copper, and accordingly directs a good venereal vitriol to be used; yet, in the Goslarian and Dantzick vitriol, which he recommends as being very fit for

Kk 2 the

the purpose, iron is in the prevailing metal, the quantity of copper being very inconsiderable; and it appears from his own words, that sometimes at least, he used the English vitriol, which he scarcely ever found to contain any metallic matter besides iron. The yellow or reddish colour which he ascribes to this sub-limate, and its property of turning to an inky blackness with insusion of galls, are matks of its having been truly a chalybeate preparation.

In the preceding edition of the London pharmacopæia, agreeably to Boyle's opinion of the production of the fublimate, the process was inserted with blue vitriol; and those of Edinburgh and Paris followed the example. The London college, at the last revisal of their book, have corrected this error, and ordered green vitriol, or filings of iron itself, to be used; but the

mistake is still continued in the other pharmacopæias.

From good blue vitriol, or pure vitriol of copper, the sublimate here required cannot be obtained: and although it may be prepared from the common blue vitriol of the shops, as I have on trial found that it may, yet it is furely imprudent to endanger impregnating the preparation with that noxious metal; more especially as pure vitriols of iron are procurable at a much cheaper rate than the others. Those mixed vitriols in which the copper greatly prevails, give first a green or blue cupreous sublimate. and afterwards a yellow or reddish ferrugineous one; and those in which iron abounds most, give first the ferrugineous, and afterwards the cupreous flowers: though poffibly neither fublimate is entirely free from an admixture of the other.

### SECT. V.

### Preparations of lead.

EAD readily melts in the fire, and calcines into a dufky powder: which, if the flame is reverberated on it, becomes at first yellow, then read, and at length melts into a vitreous mass. This metal dissolves easily in the nitrous acid, difficultly in the vitriolic, and in small quantity in the vegetable acids; it is also soluble in expersed oil, especially when calcined.

Lead and its calces, whilst undissolved, have no considerable effects as medicines. Dissolved in oils, they are supposed to be (when externally applied) anti-inflammatory and desiccative. Combined with vegetable acids, they are notably so, and taken internally,

prove a powerful but dangerous styptic.

## PLUMBUM USTUM. Burnt lead. Edinb.

Melt lead with a gentle fire, and keep it continually stirring, with an iron spatula, till it changes into powder.

### MINIUM. Redlead. Edinb.

Let any quantity of lead be melted in an unglazed earthen veffel, and kept stirring with an iron spatula, till it falls into a powder, at first blackish, afterwards yellow, and at length of a deep

red

red colour, in which last state it is called minium; taking care not to raise the fire so high as to run the calx into a vitreous mass.

THE preparation of red lead is so troublesome and tedious, as scarce ever to be attempted by the apothecary or chemist; nor indeed is this commodity expected to be made by them, the preparation of it being a distinct branch of business. The maker melt large quantities of lead at once, upon the bottom of a reverberatory furnace built for this purpole, and fo contrived, that the flame acts upon a large surface of the metal, which is continually changed by the means of iron rakes drawn backwards and forwards, till the fluidity of the lead is destroyed; after which, the calx is only now and then turned. By barely ftirring the calx, as above directed, in a vessel over the fire, it acquires no redness; the reverberation of flame upon the furface being absolutely necessary for this effect. It is said, that twenty pounds of lead gain, in this process, five pounds; and that the calx, being reduced into lead again, is found one pound less than the original weight of the metal.

These calces are employed in external applications, for abating inflammations, cleansing and healing ulcers, and the like. Their effects, however, are not very considerable; nor are they perhaps of much farther real use, than as they give consistence to the plaster, unguent, &c.

CERUSSA. Cerusse, or white lead. Edinb.

Put some vinegar into the bottom of an earthen vessel, and sufpend over the vinegar very thin plates of lead, in such a manner that the vapour which arises from the acid, may circulate about the plates. Set the containing vessel in the heat of horsedung, for three weeks: if at the end of this time, the plates are not totally calcined, scrape off the white powder, and expose them again to the steam of vinegar, till all the lead is thus corroded into powder.

THE making of white lead also is become a trade by itself, and confined to a few persons, who have large conveniencies for this purpose. The general method which they follow, is nearly the same with that above described. See the Philosophical Transactions, No. 137.

In this preparation, the lead is fo far opened by the acid, as to discover, when taken internally, the malignant quality of the metal; and to prove externally, when fprinkled on running fores, or ulcers, moderately cooling, drying, and astrictive.

## SACCHARUM SATURNI. Sugar of lead. Lond.

Boil cerusse with distilled vinegar, in a leaden vessel, until the vinegar becomes sufficiently sweet: then filter the vinegar through paper, and after due evaporation set it to crystallize.

#### Edinb.

Put any quantity of ceruste into a cucurbit, and pour thereon distilled vinegar to the height of four inches. Digest them together for some days in a sand-heat, till the vinegar has acquired a sweetish taste, when it is to be suffered to settle, and then poured off. Add fresh vinegar to the remainder, and repeat this pro-K k 2

cess till the menstruum no longer extracts any fweet tafte. Let all the impregnated liquors rest for fome time; and after they have been poured from the feces, evaporate them, in a glass vessel, to the confidence of thin honey; so that, upon being fet in a cool place, the fugar may shoot into crystals, which are afterwards to be dried in the shade. Exhale the remaining liquor to a pellicle, fet it again in the cold, and more crystals will shoot; repeat this operation till no crystals can be any longer obtained.

CERUSSE (especially that sort called flake lead, which is not, like the others, subject to adulteration) is much preferable either to minium or litharge, for making the fugar of lead; for the corrosion, which it has already undergone from the steam of vinegar, disposes it to disfolve more readily. It should be finely powdered before the vinegar is put to it, and during the digeftion, or boiling, every now and then stirred up with a wooden spatula, to promote its diffolution, and prevent its concreting into a hard mass at the bottom. The strong acid obtained from the caput mortuum of vinegar (see page 452.) may be employed for this process to better advantage than the weaker, though purer acid, above directed. If a small quantity of rectified spirit of wine be prudently added to the folution as foon as it is duly exhaled, and the mixture fuffered to grow cold by flow degrees, the fugar will concrete into very large and transparent crystals, which are scarcely to be obtained by any other method.

The fugar of lead is much more efficacious than the foregoing preparations, in the several intentions which they are applied to. Some have ventured upon it internally, in doles of a few grains, as a styptic, in hæmorrhages, profuse colliquative sweats, seminal fluxes, the fluor albus, &c. nor has it failed their expectations. It very powerfully restrains the discharge; but almost as certainly as it does this, it occasions symptoms of another kind, often more dangerous than those removed by it, and sometimes fatal. Violent pains in the bowels, or through the whole body, and obstinate constipations, sometimes immediately follow, especially if the dose has been confiderable: cramps, tremors, and weakness of the nerves, generally, fooner or later, enfue.

Boerhaave is of opinion, that this preparation proves malignant only, in to far as its acid happens to be absorbed in the body; for in such case, he says, " it returns again "into ceruste which is violently poisonous." On this principle it would follow, that in habits where acidities abound, the fugar of lead would be innocent. But this is far from being the case. Lead and its preparations act in the body only in so far as they are combined with acid: cerusse possesses the qualities of the faccharum only in a low degree; and either of them freed from the acid, have little, if any effect at all.

### SECT. VI.

### Preparations of tin.

IN easily melts in the fire. and calcines into a dufky powder, which by a farther continuance of heat, becomes white. A mass of tin heated till it is just ready to melt, proves extremely brittle, so as to fall in pieces from a blow, and by dextrous agitation into powder. Its proper menstruum is aqua regia; though the other mineral acids also may be made to diffolve it, and the vegetable ones in small quantity. It crystallizes with the vegetable and vitriolic acids; but with the others, deliauiates.

The virtues of this metal are little known. It has been recommended as an antihysteric, antihectic, &c. At present, it is chiefly

used as an anthelmintic.

### STANNUM PULVERATUM. Powdered tin.

Lond.

Melt the tin, and pour it into a wooden box rubbed in the infide with chalk: then immediately let the box be nimbly shaken, and a part of the tin will fall into powder. The remainder is to be melted a second time, and treated in the same manner, till the whole of the metal is thus reduced into powder.

THIS preparation has been used for some time as a remedy against worms, particularly the flat kinds, which too often elude the force of other medicines. The general dose is from a scruple to a drachm; some confine it to a few grains. But Dr. Alston assures us, in the Edinburgh essays, that its success chiefly

depends upon its being given in much larger quantities: he gives an ounce of the powder on an empty stomach, mixed with four ounces of melasses; next day, half an ounce; and the day following, half an ounce more after which, a cathartic is administered: he fays the worms are usually voided during the operation of the purge, but that pains of the stomach occasioned by them are removed almost immediately upon taking the first dose of the tin. The experiments on tin related in page 231 of this work, account fufficiently for its being destructive to these animals; though not for its being safe to the patient.

CALX JOVIS.

Calx of tin.

Edinb.

Melt any quantity of tin in an unglafed earthen veffel, and keep it continually flirting with an iron spatula, until it falls into a

This process is not here intended to be carried so far as the pharmaceutical writers in general direct: it must be discontinued as soon as the metal is reduced into a dusky powder; if calcined to whiteness, the following operation would not well succeed. As to the virtues of the calx, they do not seem to be greatly different from those of the foregoing preparation.

SAL JOVIS.

Salt of tin.

Edinb.

Take one pound of the foregoing calx of tin; and four ounces of K k 4 aqua

aqua regia, diluted with fix times its quantity of spring water. Digest them together in a sandheat for two days: then shake the vessel, and after the more ponderous parts of the calx have subsided, pour off the turbid liquor, and evaporate it almost to drynes; the further exsiccation of the matter is to be performed on bibulous paper. On the calx which is lest, pour half as much of the dissolvent as was employed at first; and proceed in the same manner as before.

In former editions, the menstruum, after digestion upon the calx of tin, was ordered to be filtered, then evaporated till a pellicle appeared upon the furface, and fet by to crystallize. But the crystallization fucceeded very ill; and fuch crystalline matter, as was with difficulty obtained, proved to be little other than a nitrous ammoniacal falt afforded by the aqua regia; for this menstruum does not diffolve, or diffolves only an inconsiderable quantity of, the calx of tin. The process is now rendered more practicable, by allowing the finer parts of the calx to be mixed with the liquor in an undiffolyed state, and the whole to be inspissated and exficcated together. It is probable, however, that the preparation here intended, might be obtained in a manner still more commodious.

I cannot apprehend what advantage there is in calcining the tin. Tin, in its metallic state, dissolves freely in aqua regia, but calcination renders it almost indissoluble in that menstruum; the surther it is calcined, the more does it lose of its solubility. If tin and its calk were of equal solubility, it could scarcely be suspected that the solutions of the two would be different

in quality; for the phlogiston, or inflammable principle, which fire expels from metals in their calcination, is equally extricated by acids in their dissolution. A falt of tin with aqua regia may therefore be more advantageously prepared in the following manner.

Let melted tin be poured in small fireams into a vessel of cold water, that it may be reduced into grains. Drop these by little and little, as a grain at a time, into aqua regia, that the dissolution may go on slowly, without effervescence or the discharge of summer. When the metal is no longer acted on, pour off the solution, and evaporate it in a sand-heat, till a dry salt is lest.

This preparation feems intended chiefly for external use, as a mild escharotic and detergent. It is not so corrosive as might be expected, nor much disposed to liquefy in the air, though it is not easily made to assume a crystalline falt may be obtained from tin by the vitriolic acid, in the following manner.

Take two ounces of tin, reduced into grains of filings; and five ounces of oil of vitriol. them into a wide-necked glass, in a fand-heat, and increase the fire till the liquid boils and evaporates, and the matter remains almost dry. Then remove the vessel from the fire, and when the faline residuum has concreted, add a proper quantity of water, which, by the affiftance of a moderate heat, will diffolve nearly the whole. Filter the folution, and after due evaporation, fet it to crystallize.

SALT

SALT of tin for internal use, has been commonly directed to be prepared with distilled vinegar; by digesting the vinegar on calcined tin, and then evaporating and crystallizing. Several of the chemysts have denied that any crystals would by this means be obtained, or that the distilled vinegar would dissolve any part of the calx: and indeed when the tin is but moderately calcined, as above directed, it does not appear that any solution happens.

There are two states in which tin is confiderably acted upon by vegetable acids: its perfect metalic state, and that of a perfect calx. Plates of pure tin, put into common vinegar, are in a few hours corroded: by degrees the liquor becomes quite opake and turbid, and deposites great part of the corroded fin to the bottom in form of a whitish powder; but still retains a part exquisitely divided; for after flanding for many days, and after passing through a filter, so much remained suspended, as to give a whitishness and opacity to the fluid. Acid juices of fruits, substituted to vinegar, exhibited the same phoenomena. These experiments, though they do not shew that the tin is thus fufficiently dissolved to afford a perfect crystalline falt, prove, nevertheless, what is of more importance to be known, that tin or tinned vessels, however pure the tin be, will give a metalline impregnation to light vegetable acids fuffered to stand in them for a few hours.

In order to the obtaining a perfect folution, of tin for crystallization, the metal must be highly calcined; for though its folution in mineral acids is prevented by calcination, it is otherwise in regard to the vegetable. Some take the common calz of tin, and having fpread it thinly over the bottom of a proper vessel, continue the calcination in a gentle heat, frequently flirring the powder, for three or four days, in a furnace where the air may pass freely over the furface. Others mix the common calx or filings of tin with twice their weight of nitre, and inject the mixture by degrees into a vessel strongly heated, over which are fitted a number of aludels, or earthen pots with holes in their bottoms: the lowermost of these vessels has a hole also in the side. through which the matter is thrown in : during the deflagration which happens on each injection, a part of the tin is volatalized, and adheres to the pots in form of a fine white powder, which is fwept out and washed with water. Others obtain a calx of tin, perhaps not less perfect, more expeditionly and with less trouble; by dissolving the metal in aqua regia, (which as already observed, has, in this respect, nearly the same effect as fire,) and afterwards recovering the calx, by diluting the folution with about four times its quantity of water, and gradually adding to it spirit of fal ammoniac till the effervescence ceases: a white curdly matter precipitates, which is to be washed with water and dried.

Take of calx of tin, prepared in either of the above methods, one pound; of distilled vinegar, one gallon. Digest them together, occasionally stirring up the matter from the bottom, till the vinegar has required a rough sweetish taste: then evaporate the liquor to the consistence of a syrrup, add to it about one twentieth its weight of rectified spirit of wine, and suffer the heat slowly to decrease, that the salt may crystallize.

THE

THE crystals obtained by this method are hard, folid, colourless, transparent, void of acrimony. They have been recommended in the dose of a few grains, in uterine diforders: but it does not appear that experience has warranted the virtues attributed to them; nor are any of these salts at present made use of in the common practice, or kept in the shops.

The powder precipitated from aqua regis, either by volatile alkalies, or by water alone, is fometimes employed as a cosmetic, under the name of MAGISTERY OF A whiter and more elepreparation of this kind gant, might be obtained, by dissolving the metal in the vitriolic acid, and precipitating with volatile spi-

rits.

### AURUM MUSIVUM. Mosaic gold. Lond.

Take of

Tin, one pound: Flowers of fulphur, seven ounces; Sal ammoniac,

Purified quickfilver, of each half

a pound.

Melt the tin by itself, add to it the quickfilver and when the mixture is grown cold, reduce it into powder: mix this with the fulphur and fal ammoniac, and fublime in a matrafs: the mofaic gold will be found under the fublimed matter, with some dross at the bottom.

THE management of this procefs, fo as to give to the preparation the beautiful colour and appearance for which it is admired, has been held as a secret. chemists seem greatly divided as to the proportions which the ingredients ought to bear to each other, and in this some make the chief difficulty to confift; while others make the due regulation of the fire to be the principal There does not however point. appear to be any very great nicety in either respects. I have found the process to succeed equally with very different proportions of the materials; by mixing them thoroughly together; putting them into a wide necked matras upon a little fand in an iron pot; applying a gentle fire for fome time, till the white fumes, which arose copiously at first, and passed out at the neck of the glass, begun to abate; then gradually increasing the fire till the fand became red hot, and keeping it up in this flate for a confiderable while, according to the quantity of the mixture.

The mosaic gold is chiefly valued, and receives its name from its sparkling gold-like hue. As a medicine, it is at present little regarded; though formerly held in considerable esteem, in historical and hypochondriacal complaints. malignant fevers, and venereal disorders. In these last it has been recommended from a supposition of its being a mercurial; but on confidering the circumstances of the process, and the phenomena that occur in it, there will appear little probability of any of the mercury being retained in the pre-

paration. The matrass being broken when the process is finished, the mosaic gold is found in the bottom; and the sublimed substance, above it, confifts partly of fal ammoniac, partly of fulphur, and partly of a cinnabar refulting from the combination of part of the fulphur The aurum moand mercury. faicum is found to weigh more than the tin employed; but pure tin, in being calcined by itself,

gains

gains very nearly as much as it does in this process: the golden colour is probably owing to a minute portion of sulphur adhering to the tin. On roasting the aurum over a gentle fire, it smokes a little, and soon changes its golden hue to a dirty coloured one not unlike that of tin lightly calcined; being then mixed with a proper flux, and melted in a crucible,

it yields a lump of tin, not far fhort of the original weight of the metal.

The college of Edinburgh though they formerly divided this preparation into two processes, one for amalgamating the tin with the mercury, the other for the sublimation with the sulphur and sal ammoniac, have now entirely rejected it.

### SECT. VII.

### Preparations of Mercury.

ERCURY, or quicksilver, is a ponderous metalic fluid, totally volatile in a strong fire, and calcinable by a weaker one (though very difficultly) into a red powdery substance. It diffolves in the nitrous acid, is corroded by the vitriolic, but not acted on by the marine in its liquid state: it nevertheless may be combined with this last, if skilfully applied in the form of fume. Quickfilver unites, by trituration with earthy, uncluous, refinous, and other like substances. so as to lose its fluidity: triturated with fulphur, it forms a black mass, which by fublimation changes into a beautiful red one.

The general virtues of the mercurial preparations are, to fuse the juices, however viscid, in the minutest and remotest vessels; by this means they prove eminently serviceable in inveterate chronical disorders, proceeding from a thickness and sluggishness of the humours, and obstinate obstructions of the glands. Crude mercury has no effect this way. Resolved into sume, or divided into minute particles, and prevented from re-uniting by the interposition of other substances, it operates very power-

fully; unless the dividing body be sulphur, which restrains its action. Combined with a small quantity of the mineral acids, itacts effectually, though in general mildly; with a larger, it proves violently corrosive.

# ARGENTI VIVI PURIFICATIO, Purification of quickfilver. L. E.

Diftil quickfilver in a retort; and afterwards wash it with water and common salt, or with vine-

IF a glass retort is made use of for this operation, it ought to have a low body, and a long neck; and the neck should be considerably inclined downwards, so as to allow the elevated mercury a quick defcent: the receiver should be filled almost to the neck of the retort with water; the use of this is not to condense, but to cool, the distilling quickfilver, left falling hot upon the bottom, it should crack the glass. The distillation may be more conveniently performed in an iron retort, or an iron pot fitted with a head.

The fire should be raised no higher than is sufficient to elevate the

the mercury; for certain mineral fubstances, which are said to be sometimes mixed with it, prove in part volatile in a degree of heat, not much greater than that in which mercury distils. Mr. Boyle relates, that he has known quickfilver carry up with it a portion even of lead, so as to have its weight very sensibly increased thereby; and this happened though only a moderate fire was used.

### MERCURIUS ALCALIZATUS. Alkalized mercury.

Take of

Pure quickfilver, three drachms; Prepared crabs-eyes, five drachms. Grind them together in a glass mortar, till the mercurial globules disappear.

THIS preparation, which has never been received into the London pharmacopæia, and is now rejected from the Edinburgh, is inferted here on account of its being still now and then called for. and held by some in considerable efteem. It has never come much into common practice, the labour of making it having been a temptation to a grievous abuse in its preparation, viz. the addition of an intermedium, which facilitates the union of the mercury with the crabs eyes, but greatly abates its medicinal powers. The medicine, when duly prepared, is an useful alterative; and may be given in cutaneous or veneral cases, from two or three grains to a scruple.

### MERCURIUS SACHARA-TUS.

Sugared mercury.
Edinb.

Take of
Pure quickfilver,
Brown fugar-candy, of each half
an ounce;

Effential oil of juniper berries, fixteen drops.

Grind them together in a glass mortar until the mercury ceases to appear.

THE effential oil, here added, is faid to be a very useful ingredient: not only promoting the extinction of the quickfilver (which however is still not a little difficult and tedious) but likewise improving the medicine. The intention, in this and the foregoing process, is only to divide the mercury by the interpolitition of other bodies; for when thus managed (as already observed) it has very powerful effects; though whilst undivided it seems to be altogether inactive. alone apparently answers this in-tention; but on the commixture of aqueous fluids, the fugar diffolves by itself, leaving the mercury to run together again in its original form: the addition of the oil is faid in great measure to prevent this inconvenience. The dose of this medicine, as an alterative, is from two or three grains to a scruple.

## ÆTHIOPS MINERALIS. Ethiops mineral. Lond.

Take

Purified quickfilver, Flowers of fulphur, unwashed, of each equal weights.

Grind them together in a glass or stone mortar, until they are united.

Edinb.

Take of

Purified quickfilver, Flowers of sulphur washed, each

equal weights.

Grind them together in a glass mortar, with a glass peftle, till till the mercurial globules totally disappear.

Αn

An ethiops is made also with a double quantity of mercury.

The union of the mercury and fulphur might be greatly facilitated by the affiftance of a little warmth. Some are accustomed to make this preparation in a very expeditious manner, by melting the fulphur in an iron ladel, then adding the quickfilver, and flirring them together till the mixture is complet-The finall degree of heat here sufficient, cannot reasonably be supposed to do any injury to fubstances, which have already undergone much greater fires, not only in the extraction from their ores, but likewise in the purifications of them directed in the pharmacopæia. In the following process, they are exposed in conjunction to astrong fire, without suspicion of the compound receiving any ill quality from it. Thus much is certain, that the ingredients are more perfectly united by heat, than by the degree of triture usually bestowed upon them. From the ethiops prepared by triture, part of the mercury is apt to be spued out on making it into an electary or pills: from that made by fire, no separation is observed to happen.

Ethiops mineral is one of the most inactive of the mercurial preparations. Some practitioners have boldly afferted its poffetling extraordinary virtues; and most people imagine it a medicine of some efficacy. But what benefit is to be expected from it in the common doses of eight or ten grains, or a scruple, may be judged from hence, that it has been taken in doses of feveral drachms, and continued for a considerable time, without producing any remarkable effect. Sulphur eminently abates the power of all the more active minerals,

and feems to be at the same time restrained by them from operating in the body itself. Boerhaave. who is in general sufficiently liberal in the commendation of medicines, disapproves the ethiops in very strong terms. " It cannot " enter the absorbent vessels, the " lacteals or lymphatics; but paffes directly through the intestinal tube where it may happen to destroy worms if it operates " luckily. They are deceived " who expect any other effects " from it; at least I myself could never find them. I am afraid, " it is unwarily given, in fuch " large quantities, to children " and persons of tender constituti-" ons ; as being a foreign mals, " unconquerable by the body, the " more to be suspected, as it there continues long, fluggish, " inactive. It does not raise a fa-" livation, because it cannot come " into the blood. Who knows the effects of a substance, which fo long as it remains compound-" ed, seems no more active than " any ponderous infipid earth?" The ethiops with a double proportion of mercury, now received into the Edingburgh pharmacopæia, has a greater chance for operating as a mercurial; and probably the quantity of mercury might be still further increased to advantage.

## CINNABARIS FACTITIA. Artificial cinnabar. Lond.

Take of

Purified quickfilver, twenty five ounces;

Sulphur, seven ounces.

Melt the sulphur, and mix into it the quicksilver; if the mixture happens to catch slame, extinguish it by covering the vessel. The matter is afterwards to be reducreduced into powder, and fublimed.

#### Edinb.

Take of

Purified quickfilver, three pounds and a half:

Flower of fulphur, washed, one

pound.

Melt the fulphur in a large iron veffel, over a gentle fire, and add to it by degrees the quick-filver previously heated, stirring them constantly together with an iron spatula, that they may be perfectly mixed. Immediately fit upon the veffel a wooden cover, to prevent the mixture from taking fire: before the matter is grown cold, grind it into powder, and sublime according to art.

It has been customary to order a larger quantity of sulphur than here directed; but these smaller proportions answer better; for the less sulphur, the siner coloured is the cinnabar.

As foon as the mercury and fulphur begin to unite, a considerable explosion frequently happens, and the mixture is very apt to take sire, especially if the process is somewhat hastily conducted. This accident, the operator will have previous notice of, from the matter swelling up, and growing suddenly consistent: as soon as this happens, the vessel must be immediately close covered.

During the sublimation, care must be had that the matter rise not in the neck of the vessel, so as to block up and burst the glass: to prevent this, a widenecked bolf-head, or rather an oval earthen jar, coated, should be chosen for the subliming vessel. If the former is employed, it will be convenient to introduce, at

times an iron wire, fomewhat beated, in order to be the better affured, that the paffage is not blocking up, the danger of which may be prevented, by cautiously raising the vessel higher from the fire.

If the ingredients were pure, no feces will remain: in such case, the sublimation may be known to be over, by introducing a wire as before, and feeling therewith the bottom of the vessel, which will then be perfectly smooth: if any roughness or inequalities are perceived, either the mixture was impure, or the sublimation is not compleated; if the latter be the case, the wire will soon be covered over with the

rifing cinnabar.

The preparers of cinnabar in large quantity, employ earthen jars, which in thape pretty much These are of resemble an egg. different fizes, according to the quantity intended to be made at one fublimation, which fometimes amounts to two hundred weight. The jar is usually coated from the fmall end, almost to the middle, to prevent its breaking from the vehemence, or irregularity of the fire. The greater part, which is placed uppermoft, not being received within the furnace, has no oc-The whole cation for this defence, fecret with regard to this process, is (1) the management of the fire, which should be so strong as to keep the matter continually subliming to the upper part of the jar, without coming out at its mouth, which is covered with an iron plate; (2) to put into the subliming veffel, only small quantities of the mixture at a time.

A method is mentioned in the practical chemistry of making cinnabar without sublimation, by agitating or digesting mercury in the volatile tincture of sulphur,

already

I have found already described. a fulphureous liquor more eafily parable to have a like effect; the folution for lac sulpburis will, with fome address, succeed.

The principal use of cinnabar is It was formerly as a pigment. held in great esteem as a medicine, in cutaneous foulnesses, gouty and rheumatic pains, epileptic cases, &c. but of late, it has loft much of its reputation. It appears to be nearly similar to the etbioss, already spoken of. Cartheuser relates, that having given cinnabar in large quantities to a dog, it produced no sensible effect, but was partly voided along with the feces unaltered, and partly found entire in the stomach and intestines upon opening the animal. The celebrated Frederick Hoffman, after bestowing high encomiums on this preparation, as having, in many instances within his own knowledge. perfectly cured epilepsies and vertigoes from contufions of the head (where it is probable, however, that the cure did not so much depend upon the cinnabar, as on the spontaneous recovery of the parts from the external injury) observes, that the large repeated doses, necesfary for having any effect, can be borne only where the first passages are strong, and that if the fibres of the stomach and intestines are lax and flaccid, the cinnabar, accumulated and concreting with the mucous matter of the parts, occations great oppretion; which feems to be an acknowledgment that the cinnabar is not subdued by the powers of digestion, and has no proper medicinal activity. There are indeed fome instances of the daily use of cinnabar having brought on a falivation; perhaps from the cinnabar, made use of in those cases, having contained a less pro-

portion of fulphur than the forts commonly met with. The regulus of antimony, and even white arlenic, when combined with a certain quantity of common sulphur, seem to have their deletereous power destroyed: on separating more and more of the fulphur, they exert more and more of their proper virulence. It does not feem unreasonable to prefume, that mercury may have its activity varied in like manner; that when perfectly satiated with fulphur, it may be inert: and that when the quantity of fulphur is more and more lessened, the compound may have greater and greater degrees of the proper efficacy of mercurials.

Cinnabar is fometimes used in fumigations against venereal ulcers in the nose, mouth, and throat. Half a drachm of it burnt, the fume being imbibed with the breath. has occasioned a violent salivation. This effect is by no means owing to the medicine as cinnabar; when fet on fire, it is no longer a mixture of mercury and sulphur: but mercury resolved into sume, and blended in part with the volatile vitriolic acid; in either of which circumstances, this mineral, as already observed, has very powerful

effects.

### MERCURIUS CALCINATUS. Calcined mercury.

Lond.

Put purified quickfilver into a broadbottomed glass vessel, having a fmall hole open to the air; and keep it in a constant heat, in a fand-furnace, for feveral months. until it is calcined into a red powder.

This tedious process might, in all probability, be greatly expedited, by employing, instead of a

veffel with a small aperture, a very wide-mouthed, flat bottomed glass body, of such a height that the mercury may not escape: by this means, the air, which is effentially necessary to the calcination of all metallic substances, will be more freely admitted. A veffel might be so contrived, as to occasion a continual flux of air over the surface of the mercury.

This preparation is by some highly esteemed in venereal cases, and supposed to be the most efficacious and certain of all the mercurials. It may be advantageously given in conjunction with opiates: a bolus or pill, containing from half a grain to two grains of this calx, and a quarter or half a grain or more of opium, with the addition of some warm aromatic ingredient, may be taken every night. Thus managed, it acts mildly, though powerfully, as an alterative and disphoretic: given by itself in larger doses, as four or five grains, it proves a rough emetic and cathartic.

## MERCURII SOLUTIO. Solution of mercury. Edinb.

Take equal quantities of pure quickfilver, and double aquafortis. Digest them together, in a phial placed in a sand-surnace, that a limpid solution may be made.

AQUAFORTIS dissolves mercury more easily, and in larger quantity, than any other acid: fixteen ounces, if the menstruum is very strong and pure, will take up eleven or twelve. As the liquor grows cold, a considerable part concretes, at the bottom of the vessel, into a crystalline form. If the whole is wanted to remain suspended, a proper quan-

tity of water should be added after the solution is completed.

This process is given only as preparatory to some of the following ones. The solution is highly caustic, so as scarce to be safely touched. It stains the skin purple or black.

## CALX MERCURII. Calx of mercury. Edinb.

Take any quantity of the folution of mercury, and evaporate it over a gentle fire, till a white dry mass remains.

This calx, or rather falt, of mercury, is violently corrolive. It is rarely made use of any otherwise than for making the following preparation and the corrolive sublimate.

### MERCURIUS CALCINATUS, vulgo

PRÆCIPITATUS RUBER.

Red calx of mercury, commonly called, red precipitate.

Edinb.

Take any quantity of the calx of mercury, and reverberate it in a crucible, with fucceffive degrees of heat. Its white colour will change first into a brown, and afterwards a yellow; at length, upon increasing the fire, it passes into a deep red.

## MERCURIUS CORROSIVUS RUBER.

The red mercurial corrofive-Lond.

Take of
Purified quickfilver,
Compound aquafortis, described in page 451, of each equal weights.

Mix, and set them in a broad-

bottomed vessel, in a sand-heat.

till

till all the humidity is exhaled, and the mass has acquired a red colour.

The marine acid in the compound menstruum, ordered in this last process, disposes the mercurial calx to assume the bright sparkling look admired in it; which, though perhaps no advantage to it as a medicine, ought nevertheless to be infifted on by the buyer as a mark of its goodness and strength. foon as the matter has gained this appearance, it should be immediately removed from the fire, otherwife it will foon lose it again. The preparation of this red precipitate, as it is called, in persection, is supposed by some to be a secret not known to our chemists; infomuch that we are under a necessity of importing it from abroad. This reflexion feems to be founded on misinformation: we sometimes indeed receive considerable quantities from Holland; but this depends upon the ingredients being commonly cheaper there than with us, and not upon any secret in the manner of the preparation.

This precipitate is, as its title imports, an escharotic, and in this intention is frequently employed by the furgeons, with basilicum, and other dreffings, for confuming fungous flesh in ulcers, and the like purposes. It is subject to great uncertainty in point of frength; more or less of the acid exhaling, according to the degree and continuance of the fire. The best criterion of its strength, as already observed, is its brilliant appearance; which is also the mark of its genuineness: if mixed with minium, which it is sometimes said to be, the duller hue will discover the This admixture may be more certainly detected by means of fire: the mercurial part will totally evaporate, leaving the minium behind.

Some have ventured to give this medicine internally, in venereal, scrophulous, and other obstinate chronic disorders, in doses of two or three grains, and more. But certainly the milder mercurials, properly managed, are capable of answering all that can be expected from this; without occasioning violent anxieties, tormina of the bowels, and other ill consequences. which the best management can scarcely prevent this corrolive preparation from fometimes doing. The chemists have contrived sundry methods of correcting and rendering it milder, by divelting it of a portion of the acid; but to no very good purpose, as they either leave the medicine still too corrofive, or render it fimilar to others which are parable at an easier rate.

## MERCURIUS CORALLINUS. Coralline mercury. Lond.

Pour on the red mercurial corrofive, about thrice its weight of rectified spirit of wine, and digest them together, with a gentle heat, for two or three days, frequently shaking the vessel; then set fire to the spirit, keeping the powder continually stirring till all the spirit is burnt away.

Ir is supposed, that all the more violent preparations of this kind, composed of metallic bodies united with acids, are rendered milder by digestion in spirit of wine; the acid being dulcified, or in part absorbed by the spirit. This evidently happens in some cases, where the proportion of acid is large, or sufficient to render the compound soluble in water: but that it happens equally

in others, I cannot affirm. Thus much is certain, that the mercurius cotallinus, whether from this cause, or barely from some of the acid being dissipated by the heat of the burning spirit, proves considerably milder than the corrosive was at first. It is still, however, a medicine of great activity, and seems to be scarce sufficiently safe for internal use: a few grains of it generally prove cathartic or emetic, and sometimes occasion violent symptoms.

ARCANUM CORALLINUM.

The coralline fecret.

Take five ounces of the red mercurial corrolive, and eight ounces of spirit of nitre: distil off the fpirit in a retort; return it with four ounces of fresh spirit of nitre upon the residium, and draw it off again as before: repeat this process with four ounces of new spirit; and at last keep the fire up very strong, for at least two hours. powder, which remains in the retort, is to be put into a crucible, and kept of a worm-red heat for feven or eight minutes: then boil it for half an hour, in three pints of pure water; distil from it twelve ounces of tartarized spirit of wine, cohobating the spirit twice; digest it for forty-eight hours in a fund heat, with the same quantity of fresh tartarized spirit; raising the fire towards the end, so as to make the spirit simmer a little : afterwards suffer the whole to cool, decant off the fpirit, and dry the powder for ule.

Tries preparation, notwithstanding its pompous name, is a very unthrifty and injudicious one.

The cohobation of spirit of nitre upon the corrosive, answers no useful purpose; for whatever the acid communicates, is afterwards dissolved and separated by the water: if the direction of keeping up a strong fire for some time, after the last distillation, is not strictly complied with, all the mercury will dissolve in the water, and the solution will prove similar to the solution mercurii above described.

### Pulvis Principis. Prince's powder.

Grind eight ounces of the red mercurial corrolive into a fine powder; and digeft it with two quarts of water, in an almost boiling heat, for twelve hours, occasionally stirring up the powder from the bottom: then pour off the liquor, and digeft the powder in a fresh parcel of water as before; repeating this process a third time. The last water being poured off, grind the powder with double its weight of fixt alkaline salt, and digest it as at first in fresh waters, till it becomes infipid. Afterwards boil it in spirit of wine; and lattly, pouring off the spirit, dry the powder for use.

PANACEA MERCURII RUBRA.

Red panacea of mercury.

Digest the red mercurial corrosive with eight times its weight of water, for twenty-sour hours, shaking the vessel three or four times: pour off the water, dry the powder, and digest it with eight times its weight of spirit of wine, for fisteen days. The spirit being then decanted off, burn upon the calx twice its weight of tincture of sulphur: afterwards digest it two or three days longer in fresh spirit of wine;

wine; and in the last place, exsiccate it for use.

THE three foregoing preparations have been kept in particular hands as fecrets. At bottom they are all nearly the same, and much too trivial to deserve the pains taken about them. They are perhaps farther divested of acid, than the mercurius corallinus of the shops: but have this disadvantage, that the quantity of acid separated in the troublesome digestions, &c. must vary according to different circumstances in the operation. All the four fland recommended in small doses, two grains for instance, as excellent alterants and diaphoretics: in the larger ones, they prove emetic and cathartic.

MERCURIUS CORROSIVUS SUBLIMATUS, vel ALBUS. The white mercurial corresive, or corresive mercury sublimate. Lond.

Take of

Purified quickfilver, forty ounces; Sea-falt, thirty three ounces; Nitre, twenty eight ounces; Calcined green vitriol, fixty-fix ounces.

Grind the quickfilver in a wooden or stone mortar, with an ounce or more of corrolive mercury fublimate already made, until the quickfilver is divided into fmall grains: this mixture is to be ground with the nitre, and afterwards with the fea falt; then add the calcined vitriol. continuing the triture only for a little time longer, lest the quickfilver should run together again. Lastly proceed to sublimation, in a glass matras; to which you may adapt a head, in order to fave a little spirit that will come over.

IT has been supposed that corrofive sublimate participates of all the ingredients employed in this process; though it is certain, that it consists only of mercury and the acid of the sea salt united together The materials being mixed and exposed to the fire, first the vitriol parts with its acid; which, dislodging those of the nitre and marine falt, takes their place. The marine acid, refolved into fume and affifted by the nitrous, diffolyes the mercury now also strongly heated. This acid, though it very difficultly acts on mercury, yet when thus once united with it, is more strongly retained thereby than any other The nitrous spirit, therefore, having nothing to retain it (for its own basis, and that of the fea fal are both occupied by the vitriolic; and that which the vitriolic forfook to unite with thefe. is now scarcely combinable with it) arises leaving the mercury and marine acid to sublime together when the heat shall be strong enough to elevate them. Some small portion of the marine spirit arises along with the nitrous; and hence this compound acid has been usually employed insted of the aqua fortis composita, to which it is similar, for making the red corrolive.

It appears, therefore, that the vitriol and the bases of the nitre and sea salt, are of no farther use in this process, than as convenient intermediums for facilitating the union of the mercury with the marine acid. They likewise serve to afford a support for the sublimate to rest upon, which thus assumes the form it is expected in, that of a placenta or cake. The design of adding a little sublimate already made is to facilitate the extinction of the mercury, or its mixture with

the other materials.

Lla THERE

THERE are fundry other ways of making this preparation, or of combining mercury with the ma-rine acid. If mercury corroded by the vitriolic acid into a white mass (as for making the yellow mercurial emeric or turpeth mineral described hereafter) be mixed with an equal quantity of fea falt and fet to fublime: the vitriolic acid will quit the mercury to unite with the basis of the sea salt; and the acid of the sea falt, now set at liberty, will unite with the mercury, and fublime with it into the compound required. The discovery of this method is generally attributed to Boulduc; though it is found also in Kunckel's laboratorium chymicum.

If the mercury be corroded by the nitrous acid instead of the vitriolic, the event will be the same; that acid equally quitting the mercury, and setting loose the marine. This method the college of Edinburgh have received.

#### Edinb.

Take

Calx of mercury (that is, a folution of mercury in aquafortis, evaporated to a dry white mass,)

Decrepitated sea salt, of each equal quantities.

Powder, and mix these well together; and put them into a matras, of which they may nearly fill one half: place the vessel in a sand sturnace, and proceed to sublimation; applying at first a gentle heat, and afterwards increasing it, till all the sublimate has arisen in a white cristalline mass, to the upper part of the matrass: separate this from the red scoria, and purify it, if needful by a second sublimation.

THE sublimate made by this method is the same with the foregoing; but as the quantity of fixt matter is small, it difficultly affumes the form of a cake. It requires indeed fome skill in the operator, to give it this appearance when either process is followed. When large quantities are made. this form may be easily obtained, by placing the matrafs no deeper in the fand than the furface of the matter contained in it: and removing a little of the fand from the fides of the glass, as soon as the flowers begin to appear in the neck; when the heat should likewise be fomewhat lowered, and not at all raifed during the whole process. The fublimation is known to be compleated by the edges of the crystalline cake, which will form upon the furface of the caput mortuum, appearing fmooth and even, and a little removed from it.

Onr apothecaries rarely, and few even of the chemists, attempt the making of this preparation themselves; greatest part of what used among us comes from Venice and Holland. This foreign sublimate has been reported to be adulterated with arsenic. affirm that this dangerous fraud may be discovered by the sublimate turning black on being moiftened with alkaline ley: which by others is denied. As this point feemed of fome importance to be determined. I made fundry experiments with this view, which . convinced me of the infufficiency of alkalies for discovering are-Alkaline ley, poured into folution of pure fublimate, into a folution of pure arfenic, and into a mixture of the two solutions in different proportions. produced no blackness in any ; and though the pure sublimate.

and

and the mixtures of it with arfenic, exhibited some differences in these trials, yet these differences were neither so constant, nor so strongly marked, as to be laid down, universally, for criteria of the presence or absence of arsenic: different specimens of sublimate, known to be pure, differed considerably in this respect; probably from their holding a little more or less mercury in proportion to the acid, or from their retaining some small portion of those acids which were employed in the preparation as intermedia.

Some chemists deny the prasti-cability of this adulteration. There is a process common in books of chemistry, wherein sublimate and arfenic being mixed together and fet to fublime, they do not arise in one mass, or yield any thing similar to the preparation here intended; the arienic absorbs the acid of the sublimate, and is reduced thereby into a liquid or butyraceous confidence; while the mercury thus freed from the acid. diffils in its running form: if the quantity of arfenic is infufficient to decompound the whole of the fublimate, the remainder of the sublimate concretes distinct from the arfenical butter. From whence they conclude, that arfenic and fublimate cannot be united together, into a crystalline cake, the form in which this preparation is brought

The above experiment is not altogether decifive; for though arfenic and sulphur do not assume the required form by the common process, it is possible they may by some other management. It will therefore (though I have never found any reason to suspect that the abuse is practised) be proper to point out means for the satisfaction of those, who may be

defirous of convincing themselves of the genuineness of this important preparation. Let some of the sublimate, powdered in a glass mortar, be well mixed with twice its weight of black flux, (page 532.) and a little filings or shavings of iron : put the mixture into a crucible capable of holding four or five times as much; give a gradual fire till the ebullition ceases, and then hastily increase it to a white heat. If no fumes of a garlic smell can be perceived during the process; and if the particles of iron retain their form, without any of them being melted; I think we may be secure that the mixture contained no arsenic.

SUBLIMATE is a most violent corrosive, presently corrupting and destroying all the parts of the body it touches. A solution of it in water, in the proportion of about a drachm to a quart, is made use of for keeping down proud flesh, and cleansing soul ulcers, and a more dilute solution as a cosmetic, and for destroying cutaneous insects. But a great deal of caution is requisite even in these external uses of it.

Some have nevertheless ventured to give it internally, in the dose of one-tenth or one-eighth of a grain. Boerhaave relates, that if a grain of it be dissolved in an ounce or more of water, and a drachm of this solution, softened with syrup of violets, taken twice or thrice a day, it will perform wonders in many reputed incurable distempers; but particularly cautions us not to venture upon it, unless the method of managing it is well known.

Sublimate diffolved in vinous fpirits has of late been given internally in larger doses from a quarter of a grain to half a grain.

Ll 3

This method of using it was brought into vogue by baron Van Swieten at Vienna, particularly for venereal maladies; and several trials of it have been made in this kingdom also with success. Eight grains of the sublimate are dissolved in sixteen ounces of reclished spirit of wine or proof spirit; the reclified spirit dissolves it more perfectly, and feems to make the medicine milder in its operation, than the proof spirit of the original prescription of Van Swieten. Of this folution, from one to two spoonfuls, that is, from half an ounce to an ounce, are given twice a day, and continued till all the symptoms are removed: observing to use a low diet with plentiful dilution, otherwise the sublimate is apt to purge, and gripe fe-It generally purges more or less at the beginning, but afterwards feems to operate chiefly by urine and perspiration.

Sublimate consists of mercury united with a large quantity of marine acid. There are two general methods of destroying its corrosive quality, and rendering it mild; combining with it so much fresh mercury as the acid is capable of taking up, and separating a part of the acid by means of alkaline salts, and the like. On the sufficient on the latter, white pre-

cipitate.

# MERCURIUS DULCIS SUBLIMATUS. Dulcified mercury fublimate. Land.

Take of

Corrofive mercury fublimate, one pound;

Purified quickfilver, nine ounces. Having powdered the sublimate, add to it the quickfilver, and digest them together in a matras, with a gentle heat of fand, until they unite; then, increasing the heat, let the mixture be sublimed. The sublimed matter, freed from the accrimonious part at top and such mercurial globules as happen to appear dithinch in it, is to be reduced into powder, and sublimed again; and this sublimation repeated six times.

#### Edinb.

Take of

Corrofive mercury sublimate, reduced to powder in a glass mor-

tar, four ounces;

Pure quickfilver, three ounces. Mix them well together, by long trituration in a glass or marble mortar, until the quicksilver ceases to appear; taking care to avoid the finer powder that flies off. Put the powder into an oblong phial, of fuch a fize, that only one-third of it may be filled; and fet the glass in a sand furnance, fo as that the fand may reach up to one-half its height. By degrees of fire successively applied, almost all the mercury will sublime, and adhere to the upper part of the vessel. glass being then broken, and the red powder which is found in its bottom, with the whitish one that slicks about the neck, being thrown away, let the white mercury be sublimed again three or four times.

The trituration of corrolive sublimate with quicksilver is a very noxious operation: for is is almost impossible, by any care, to prevent the lighter particles of the former from arising, so as to affect the operator's eyes and mouth. It is nevertheless of the utmost consequence, that the ingredients be perfectly united before the sublimation is begun;

begun: this may be most commodiously effected, by the digestion ordered in the first of the above processes. It is indeed still necessary to pulverize the sublimate, before the mercury is added to it; but this may be fafely performed, with a little caution; especially, if during the pulverization, the matter be now and then sprinkled with a little spirit of wine; this addition does not at all impede the union of the ingredients, or prejudice the fublimation: it will be convenient not to close the top of the subliming vessel with a cap of paper at first (as is usually practifed) but to defer this till the mixture begins to -fublime, that the spirit may es-

The rationale of this process deferves particular attention; and the more so, as a mistaken theory herein has been productive of feveral errors, with regard to the operation of mercurials in general. It is supposed, that the dulcification, as it is called, of the mercurius corroficus, is owing to the spicula or sharp points, on which its corrolivenels depends, being broken and worn off by the frequent sublimations. If this opinion was just, the corrolive would become mild, without any addition, barely by repeating the fublimation; but this is contrary to all experience. The abatement of the corrofive quality of the fublimate is entirely owing to the combination of so much fresh mercury with it, as is capable of being united; and by whatever means this combination is effected, the preparation will be sufficiently dulcified. ture und digestion promote the union of the two, whilst sublimation tends rather to difunite them. The prudent operator, therefore, will not be folicitous about separating fuch mercurial globules as appear

distinct after the first sublimation; he will endeavour rather to combine them with the rest, by repeating the triture and digestion.

The college of Wirtemberg require their mercurius dulcis to be only twice sublimed; and the Augustan but once; and Neumann proposes making it directly, by a fingle fublimation, from the ingredients which the corrosive sublimate is prepared from, by only taking the quickfilver in a larger proportion. If the medicine, made after eitheir of these methods, should prove in any degree acrid, water, boiled on it for some time, will disfolve and separate that part in which its acrimony confifts. marks of the preparation being fufficiently dulcified, are, its being perfectly infipid to the taste, and indiffoluble by long boiling in wa-Whether the water, in which it has been boiled, has taken up any part of it, may be known by dropping into the liquor a ley of any fixt alkaline falt, or any volatile alkaline spirit: if the decoction has any mercurial impregnation, it will grow turbid on this addition: if otherwise, it will cantinue limpid. But here care must be taken not to be deceived by any extraneous faline matter in the water itself: most of the common spring waters turn milky on the addition of alkalies: and therefore, for experiments of this kind, distilled water, or rain water ought to be used.

Mercurius dulcis, seven times sublimed, has been commonly called Calomelas, and Aquila alba; names which are now dropt both by the London and Edinburgh colleges. Calomelas is indeed a very improper name for a white preparation, the word implying a black colour: by grinding mer-L14

curius dulcis with volatile spirits, it becomes blackish, and this per-

haps is the true calomel.

Mercurius dulcis appears to be one of the best and safast preparations of this mineral for general use, whether intended to act as a fialogogue, diaphoretic, or alterant. Many of the more elaborate processes are no other than attempts to produce from mercury such a medicine as this really is. The dose, for raising a fallyation, is ten or fifteen grains, taken in the form of a bolus or pills, every night or oftner, till the ptyalism begins. As an alterant and diaphoretic, it is given in doles of five or fix grains; a purgative being occasionally interposed, to prevent its affecting the mouth. It answers, however, much better, when given in fmaller quantities, as one, two, or three grains every morning and evening, in conjunction with fuch fubiliances as determine its action to the skin, as the extract or refin of guaiacum; the patient at the same time keeping warm, and drinking liberally of warm diluent liquors. By this method of managing it, obstinate cutaneous and venereal distempers have been successfully cured, without any remarkable increase of the fensible evacuations.

## PANACEA MERCURII. Mercurial panacea.

Take any quantity of levigated calomel, and four times as much spirit of wine. Digest them together in a sand-heat for twenty days, frequently shaking the vessel; then pour off the spirit, and dry the powder for use.

This preparation differs very little, if at all, from the foregoing; for, as Lemery observes, the spirit of wine does not dissolve any part

of the calomel. Some chemists have therefore recommended a proof spirit, or common water, as more fuitable for this purpose than rec-tified spirit: if any part indeed of the calomel remains not fufficiently dulcified, this will be diffolyed by boiling in water, and consequently the preparation becomes milder; but if the calomel is well made, even water will have no effect upon it; the mercury and spirit of salt being so closely united to each other, as not to admit of any feparation by the means here proposed. Nor indeed does good mercurius dulcis want any of its acid to be taken away, as being already sufficiently safe, and mild in its operation. The Edinburgh college therefore, who received this preparation in the former editions of their pharmacopæia, have now rejected it.

# MERCURIUS PRÆCIPITATUS ALBUS. White precipitate of mercury. Edinh.

Dissolve sublimate corrosive mercury in a sufficient quantity of hot water, and gradually drop into the solution some spirit of sal ammoniac, as long as any precipitation ensues. Wash the precipitated powder upon a silter, with several fresh quantities of warm water.

This preparation is used chiefly in ointments, in which intention its fine white colour is no small recommendation to it. For internal purposes, it is rarely employed, nor is it at all wanted: it is nearly similar to mercurius dulcis, but less certain in its effects. Corrosive sublimate, as we have already seen, consists of mercury united with a large proportion

portion of acid: it is there dulcified by adding as much fresh mercury as is sufficient to satiate all the acid; here, by feparating all the acid that is not fatiated. This last way seems an unfrugal one, on account not only of the loss of the acid, but of the volatile spirit necessary for absorbing The operator may, however, if it should be thought worth while, recover the volatile falt from the liquor, by adding to it after the precipitate has been foparated, a proper quantity of potash, and distilling with a gentle heat, in the same manner as for the spirit of the volatile salt of sal ammoniac: for a true fal. ammoniac is regenerated in the precipitation, from the union of the volatile spirit with the marine acid of the sublimate. It is by no means advisable to use the liquor itself as a solution of fal ammoniac, or to separate the fal ammoniac from it by evaporation and chrystallization as a part of the mercury might be retained, and communicate dangerous qualities but the volatile falt separated by distillation may be used without fear of its containing any mercury, none of which will arise with the heat by which volatile falts are distilled.

Fixt alkalies answer as effectually, for precipitating solutions of sublimate as the volatile; but the precipitate, obtained by means of the former, instead of being white as with the latter, is generally of a reddish yellow or orange colour. If sal ammoniac be dissolved along with the sublimate, the addition of fixt alkalies will now extricate the volatile alkali of the sal ammoniac, occasion as white a precipitation, as if the volatile alkali had been previously separated and employed

in its pure state: and this compendium is now allowed by the London college. The process is as follows. Lond.

Take

Sublimate corrofive mercury,
Sal ammoniac, of each equal
weights.

Diffolve them both together in water, filter the folution, and precipitate it with a folution of any fixt alkaline falt. Wash the precipitated powder, till it is perfectly sweet (that is insipid or void of acrimony.)

HERE the sal ammoniac, befides its use in the capital intention. promotes the folution of the alkali to make a white precipitation, promotes the folution of the fublimate: which, of itself, is difficultly, and scarce at all totally soluble by repeated boiling in water: for however skilfully it is prepared, some part of it will have an under proportion of acid, and confequently approach to the state of mercurius dulcis. A good deal of care is requifite in the precipitation, for if too large a quantity of the fixt alkaline folution be imprudently added, the precipitate will lose the elegant white colour for which it is valued.

A PRCIPITATE of a different nature from the preceding has been commonly diftinguished by the same name, MERCURIUS PEECIPITATUS ALBUS; the preparation of which, in the preceding edition of the Edinburgh pharmacopozia, is as follows.

Take any quantity of the folution of mercury (made in aquafortis) and pour into it, by little and little, fome very strong brine of fea falt, until all the quicksilver is precipitated in form of a very white powder; which is to be washed upon a filtre with warm water, till the water comes off without any acrimony. The powder is then to be put betwixt the folds of paper, and dried with a very gentle heat.

This is a very unfrugal preparation: for fea falt, in whatever proportion it be added, will not precipitate all the mercury: this evidently appears upon adding a fmall quantity of a folution of fixt alkaline falt, or volatile alkaline spirit to the liquor which remains after the precipitate is fallen, when it will again grow turbid, and let fall a confiderable of fresh precipitate. quantity Homberg observes, that if the acid spirit bears an over-proportion to the mercury in the folution, no precipitation at all will follow upon the affusion of the brine of sea salt. If the precipitate be washed too often with hot water, it will all dissolve and pass the filtre: the same accident will likewise happen, if the brine employed at first to throw down the mercury be suffered to stand too long upon the precipitate.

Some have been accustomed to fubstitute the above officinal white precipitate in the place of this: but very injudiciously: the first is fo mild, as not improperly to deserve the appellation by which it is distinguished in the former Edinburgh pharmacopæia, dulcis: whilst this last is so far corrosive. as to be employed by the farriers for the purposes of an escharotic. Internally, it is among us very rarely made use of; notwithstanding the character given of it by Boerhaave, of being "perhaps the best remedy hitherto af-" forded by mercury." Mercurius dulcis produces the good efrects which this is supposed to do, with a greater degree of certainty, and without disordering the constitution, occasioning vomitting,&c. which this precipitate, in a dose of two or three grains, frequently does.

MERCURIUS PRACIPITATUS
PUSCUS, vulgo WURTZII.
Brown, commonly called Wurtz's,
precipitate.

Take any quantity of a folution of mercury (made in aquafortis) and gradually drop into it oil of tarrar per deliquium, till the effervesscence ceases. A powder will precipitate, which is to be edulcorated as the foregoing.

THIS preparation was in confiderable efteem fome years ago, but at present is rarely or never made use of, and hence it is now rejected both by the London and Edinburgh colleges. It does not seem to differ in strength or effects from the sweet precipitate.

# MERCURIUS PRÆCIPITATUS VIRIDIS. Green precipitate of mercury. Edinb.

Diffolve four ounces of corrolive fublimate mercury (previously reduced to powder) in a quart of hot water.

Digest an ounce and an half of copper filings, with eight ounces of spirit of sal ammoniac, in a marrass, until a deep blue tincture is extracted.

Filter the tincture, and drop it by degrees into the mercurial folution: when the precipitate has fallen, evaporate in a fand heat to dryness.

THIS differs from the sweet precipitate, in containing an admixture of copper, which renders it an emetic too rough to be used internally with fafety: and hence the present practice has almost entirely

rejected it.

This preparation is confiderably different from the green precipitate of foreign pharmacopæias. There the proportion of copper, contained in the preparation when finished is much greater; for though the quantity directed to be taken is less, yet aquafortis being employed for the menstruum, the whole is dissolved; whereas the wolatile spirit, here employed, exgracts but a very small portion of it.

### MERCURIUS EMETICUS **FLAVUS**

The yellow mercurial emetic.

Lond.

Upon purified quickfilver, conpour tained in a glase vessel, double its weight of the strong spirit, or oil of vitriol. the liquor by degrees, so as at length to make it boil, till a white mass remains, which is to be thoroughly dried with a This mass on the strong fire. affusion of warm water, grows yellowish, and falls into powder, which is to be diligently ground with the water, in a glass mortar : then suffer it to settle, pour off the water and wash the powder in several parcels of fresh water, until it is sufficiently dulcified.

**MERCURIUS** PRÆCIPITATUS FLAVUS, feu

TURPETHUM MINERALE. Yallow precipitate of mercury, or turpeth mineral.

Edinb.

Take four ounces of pure quickfilver, and eight ounces of oil of vitriol. Cautioully mix them together and distil in a retort, placed in a fand furnace, to dryness; the white calx, which is left at the bottom, being ground to powder, and thrown into warm water, immediately grows of a yellow colour: wash this in fresh waters renewed several times, until it has lost all its acrimony; then dry it for uſc.

THE quantity of oil of vitriol, formerly directed was double to that in the above prescriptions: the reduction, now made in this article, greatly facilitates the process: and even less than the present quantity would fuffice:

Boerhaave directs this preparation to be made in an open glass, flowly heated, and then placed immediately upon burning coals; care being taken to avoid the fumes, which are extremely noxious. This method will succed very well, with a little address, when the ingredients are in small quantity: but where the mixture is large, it is better to use a retort, placed in a fand furnace, with a recipient, containing a finall quanty of water, luted to it. Great care should be taken, when the oil of vitriol begins to bubble, to steadily keep up the heat, without at all increasing it, till the ebullition ceases, when the fire should be augmented to the utmost degree, that as much as possible of the redundant acid may be expelled.

If the matter be but barely exficcated, it proves a caustic salt, which in the ablution with water will almost all dissolve, leaving only a little quantity of turpeth:

the more of the acid has been diffipated the less of the remaining mercury will dissolve, and consequently the yield of turpeth will be the greater; fire expelling only the acid (viz. fuch part of the acid as is not completely fatiated with mercury) while water takes up always along with the acid, a proportionable quantity of the mercury itself. Even when the matter has been strongly calcined, a part will still be soluble; this evidently appears upon pouring into the washings a little solution of fixt alkaline falt, which will throw down a considerable quantity of yellow precipitate, greatly refembling the turbith, except that it is less violent in operation.

From this experiment, it appears that the best method of edulcorating this powder is, by impregnating the water, intended to be used in its ablution, with determined proportion of fixt alkaline falt: for by this means, the washed turbith will not only turn out greater in quantity, but, what is of more consequence, always have an equal degree of strength: a circumstance which deserves particularly to be confidered, especially in making such preparations as, from an error in the process, may prove too violently corrofive to be used with any to-

lerable degree of fafety.

It is observable, that though the superfluous acid is here absorbed from the mrcrury by the alkaline salt; yet in some circumstances this acid forfakes that falt to unite with mercury. If tartarum vitriolatum, or nitrum vitriolazum (i. e. a combination of vitriolic acid with fixt alkali) be dissolved in water, and the solution added to a folution of mercury in aquafortis; the vitriolic acid

will unite with the mercury, and form with it a turbith, which falls to the bottom; leaving only the alkali dissolved in the aquafortis, and united with the acid thereof into a regenerated nitre. On this principle depends the preparation described by Wilson, under the title of An excellent precipitate of mercury; which is no other than a true turbith, though not generally known to be fuch. It is made by dissolving four ounces of nitrum vitriolatum in sixteen ounces of spirit of nitre; dissolving in this compound liquor four ounces of mercury; abstracting the menstruum in a fand heat; and edulcorating with water the gold-coloured mass which remains.

Turbith mineral is a strong emetic, and in this intention operates the most powerfully of all the mercurials that can be fafely given internally. Its action however is not confined to the primz viæ; it will fometimes excite a ptyalism, if a purgative is not taken soon after it. This medicine is used chiefly in virulent gonorrhæas, and other venereal cases, where there is a great flux of humours to the parts: it is faid likewise to have been employed with great success, in robust constitutions, against leprous disorders, and obstinate glandular obstructions: the dose is from two grains to fix or eight. It may be given in doses of a grain or two as an alterative and dia-

fooken of. This medicine has been of late recommended as the most effectual preservative against the hydropho-There are several examples of its preventing madness in dogs that had been bitten; and some,

phoretic, after the same manner

as the mercurius calcinatus alteady

of its performing a cure after the madness was begun: from six or seven grains to a scruple may be given every day, or every other day, for a little time, and repeated at the two or three succeeding sulls and changes of the moon. Some few trials have likewise been made on human subjects, bitten by mad dogs; and in these also the turbith, used either as an emetic or alterative, seemed to have good effects. See James's treatise on canine madness.

The washings of turbith mineral are used by some, externally, for the itch and other cutaneous soulnesses. In some cases mercurial lotions may be proper, but they are always to be used with great caution: this

is by no means an eligible one. as being extremely unequal in point of strength; more or less of the mercury being dissolved, as observed above, according to the degree of calcination. pharmacopæia of paris directs a mercurial wath free from this inconvenience, under the title of Aqua mercurialis, or Mercurius liquidus. It is composed of an ounce of mercury, dissolved in a sufficient quantity of spirit of nitre, and diluted with thirty ounces of distilled water. In want of distilled water, rain water may be used: but of spring waters there are very few which will mix with the mercurial folution, without growing turpid and precipitating a part of the mercury.

### SECT. VIII.

### Preparations of antimony.

ANTIMONY is composed of a metal, united with sulphur common brimstone.

If powdered antimony be exposed to a gentle fire, the sulphur exhales; the metallic part remaining in form of a white calx, reducible, by proper fluxes, into a whitish brittle metal, called regulus. This is readily distinguished from the other bodies of that class, by its not being soluble in aquasortis; its proper menstruum is aqua regis.

If aqua regis be poured upon crude antimony, the metallic part will be diffolved; and the fulphur thrown out, partly to the fides of the veffel, and partly to the furface of the liquor, in form of a greyish yellow substance. This, separated and purified by fublimation, appears on all trials the fame with pure common brimstone.

The metal, freed from the sulphur naturally blended with it, and afterwards sufed with common brimstone, resumes the appearance and qualities of crude antimony.

The antimonial metal is a medicine of the greatest power of any known substance: a quantity too minute to be sensible on the tenderest balance, is capable of producing virulent effects, if taken dissolved or in a soluble state. If given in such a form as to be immediately miscible with the animal sluids, it proves violently emetic; if so managed as to be more slowly acted on, cathartic; and in either case, if the dose

is extremely small, diaphoretic. Thus though vegetable acids extract so little from this metal, that the remainder seems to have lost nothing of its weight, the tinctures prove in no large doses strongly enteric, and in smaller ones powerfully diaphoretic. The regulus has been cast into the form of pillars, which acted as virulent cathartics, though without suffering any sensible diminution of weight in their passage through the body; and this repeatedly, for a great number of times.

This metal, divested of the inflammable principle which it has in common with other metalic bodies, that is reduced to a calx, becomes indisfoluble and inactive. The calx nevertheles, urged with a strong sire, melts into a glass, as easy of solution (partially) and as virulent in operation, as the regulus itself: the glass, thoroughly mingled with such substances as prevent its solubility, as wax, resins, and the like, is again render-

ed mild.

Vegetable acids, as aiready observed, dissolve but an extremely minute portion of this metal: the folution nevertheless proves powerfully emetic and cathartic. The nitrous and vitriolic acids only corrode it into a powder, to which they adhere fo flightly as to be separable in good measure by water, and totally by fire, leaving the regulus in form of a calx fimilar to that prepared by fire alone. The marine acid has a very different effect : this reduces the regulus into a violent corrolive, and though it difficultly unites, yet very closely adheres to it, infomuch as not to be separable by any ablution, nor by fire, the regulus ariting along with it.

The nitrous or viriolic acids expel the marine, and thus reduce the corrofive into a calx fimilar to the foregoing.

Sulphur remarkably abates the powder of this metal: and hence crude antimony. (in which the regulus appears to be combined with from one fourth to one half its weight of fulphur) proves altogether mild. If a part of the fulphur be taken away, by fuch operations as do not deftroy or calcine the metal, the remaining mass becomes proportionably more active.

The sulphur of antimony may be expelled by designation with nitre: the larger the quantity of nitre to a certain point, the more of the sulphur will be dissipated, and the preparation will be the more active. If the quantity of nitre is more than sufficient to confume the sulphur, the rest of it, designating with the insammable principle of the regulus itself ren-

ders it again mild.

The sulphur of antimony is likewise absorbed, in susion, by certain metals, and by alkaline salts. These last when united with sulphur, prove a menstruum for all the metals (zinc excepted) and hence if the susion is long continued, the regulus is taken up, and rendered soluble in water.

CROCUS ANTIMONII MEDCI-

Medicinal crocus of antimony.

Take of

Antimony, eight parts; Nitre, one part-

Mix, and throw them by little at a time, into a red hot crucible: when the deflagration ceases, take the crucible out of the fire, and reduce the matter into powder

THIS

THIS preparation is sufficiently mild, though confiderably more active than the crude mineral: eighteen or twenty grains will in some constitutions operate, though very gently, both upwards and downwards. It appears to be nearly fimilar to the medicinal regulus hereafter described.

In this and the following processes with nitre, the operator must observe to throw into the crucible only a little of the matter at a time, and to wait till the deflagration of one parcel is over before another is added; for if much was put in at once; the deflagration would be so violent, that great part of the matter would be thrown over the crucible. The powder is most conveniently introduced by means of a small iron ladle : care must be taken not to bring back with the ladle any spark of coal, which would fet fire to the rest of the mixture.

CROCUS ANTIMONII MITIOR. The milder crocus of antimony, Take of

Antimony, two parts; Nitre, one part.

Mix them together, and throw the powder by degrees into a red-As foon as the hot crucible. deflagration ceases, remove the matter from the fire (without fuffering it to melt) and reduce it into powder.

This preparation is called mitior, not in regard to the crocus above described, but to that which follows. It acts much more powerfully than the foregoing; the increase of the nitre occasioning a greater quantity of the fulphur of the antimony to The London combe dissipated. mittee received it in their first

draught, with the character of an antimonial of mild operation. which had proved a fuccessful medicine in numerous instances. without any one example of its being unsafe. Some trials however, afterwards reported to them. where the operation of this and the following crocus were compared, induced them to lay this preparation aside. It appears to differ from the other only in being less violent.

#### CROCUS' ANTIMONII.

Crocus of antimony, commonly called Crocus metallorum, and by foreign writers, Hepar antimonii, or Liver of antimony. Lond.

Take

Antimony,

Nitre of each equal weights. Reduce them separately into powder; then mix, and inject them into a crucible heated to a white heat, that the mixture (after deflagration) may melt. pour it out, separate the scoriz. and referve the matter underneath them for use: it proves different in colour, according to the continuance of the heat; the

longer it has been kept in fusion,

#### Edinb.

the yellower it will be.

The mixture of antimony and nitre, made as above, is to be injected into a red-hot crucible; when the detonation is over, separate the reddish metallic matter from the whitish crust, and edulcorate it by repeated washings with hot water.

Here the antimonial sulphur is almost totally confumed, and the metallic part left divested of its corrector. These preparations.

given

given from two to fix grains, act as violent emetics, greatly difordering the constitution. Their principal use is in maniacal cases; as the basis of some other preparations; and among the farriers, who frequently give to horses an ounce or two a day, divided into different doses, as an alterative: in these and other quadrupeds, this medicine acts chiefly as a dia-

phoretic.

The chemists have been accustomed to make the crocus with a less proportion of nitre than directed above; and without any farther melting, than what enfues from the heat that the matter acquires by deflagration, which when the quantity is large, is very considerable: a little common salt is added to promote the fusion. The mixture is put by degrees into an iron pot, or mortar, somewhat heated, and placed under a chimney: when the first ladleful is in, a piece of lighted charcoal is thrown to it, which fets the matter on fire: the rest of the mixture is then added by little and little: the deflagration is foon over, and the whole-seen ears in perfect fulion: when cold, a confiderable quantity of scoriz are found upon the furface; which scorize are easily knocked off with a hammer. The crocus prepared after this manner is of a redder colour, than that of the first of the above processes.

### CROCUS ANTIMONII.

Washed crocus of antimony.

Lond.

Reduce the crocus into a very fubtile powder, and boil it in water; then, throwing away this water, wash the powder several times in fresh warm water, until it becomes perfectly infipid.

THIS process is designed chiefly to fit the crocus for the preparation of emetic tartar, of which hereafter, and of the antinionial emetic wine, page 293. 'If the crocus was employed for those purpoles, without washing, alkaline salt, which it is in some degree impregnated with from the deflagration of the nitre, would in part satiate the acids of the tartar, and of the wine, and thus, impeding their actions on the metallic part of the antimony, render the medicines very precarious in strength: that uncertainties of this kind may be the more effectually guarded against, the glass, or rather the pure regulus of antimony, is by some preferred to the crocus, both for the emetic tartar and The Edinburgh college. as appears in the foregoing process, does not allow the crocus to be kept in its unwashed state : making the ablution a part of the preparation of it.

EMETICUM MITE ANTIMONII.

A mild antimonial emetic.

Take of

Antimony, one part; Nitre, two parts.

Grind them together, and throw them by little and little into a red-hot crucible: when the deflagration is over, the remaining matter which proves white, is to be washed for use.

THE quantity of nitre is here follarge, as to confume not only the fulphur of the antimony, but likewife great part of the inflammable principle of the regulus. Boerhaave, from whom this preparation is taken, informs us that it is fo mild, as often to occasion

only

only fome light nausea and gentle vomiting, with a large discharge of saliva, and thick urine. Its effects seem to be nearly the same with those of the regulus medicinalis and crocus medicinalis.

### CALX ANTIMONII.

Calx of antimony, commonly called
Diaphoretic antimony.

Lond.

Take of

Antimony, one part; Nitre, three parts.

Let the powdered antimony be well mixed with the nitre, and gradually injected into a crucible, heated to a light white heat; the matter being afterwards taken from the fire, is to be washed with water, both from the salt which adheres to it, and from the grosser part that is less perfectly calcined.

Edinb.

Take of

Antimony, half a pound; Nitre, a pound and a half.

Reduce them feparately into powder, then mix them together, and throw the mixture, by a finall ladleful at a time, into a red-hot crucible: when the detonation is over, let the white mass be calcined in the fire for half an hour longer; then powder and keep it in a glass vessel closely stopt.

This powder, unwashed, is called ANTIMONIUM D'APHORETI-CUM NITRATUM.

Nitrated diaphoretic antimony.

When the powder is washed with fresh quantities of water, till the water comes off insipid, it is called.

ARTIMONIUM DIAPHORETI-CUM LOTUM.

Washed diaphoretic antimony.

The feveral washings, mixed together, filtered, and evaporated over a gentle fire till a cuticle forms on the furface, yield, in the cold, crystals, called NITRUM STIBIATUM.

Antimoniated nitre.

The calx of antimony, when freed by washing from the saline matter, is extremely mild, if not altogether inactive. Hoffman, Lemery, and others, affure us, that they have never experienced from it any fuch effects as its usual title (that under which it stands in the last of the above processes imports: Boerhaave declares, that it is a mere metallic earth, entirely destitute of all medicinal virtue: and the committee of the London college admit that it has no fenfible operation. The common dose is from five grains to a scruple, or half a drachm: though Wilson relates, that he has known it given by half ounces, and repeated two or three times a day, for several days together.

Some report, that this calx, by being kept for a length of time, contracts an emetic quality: from whence it has been concluded, that the powers of the reguline part are not entirely destroyed; that the preparation has the virtues of other antimonials which are given as alteratives, that is, in fuch small doses as not to stimulate the prime viz; and that, therefore, diaphoretic antimony, as it is certainly among the mildest preparations of that mineral. may be useful for children, and fuch delicate constitutions where the stomach and intestines are easily affected. The observation. however, from which thele conclusions are drawn, does not appear to be well founded · Ludovici relates, that after keeping the powder powder for four years, it proved as mild as at first : and the Strasburgh pharmacopæia, with good reason, suspects, that where the calx has proved emetic, it had either been given in such cases as would of themselves have been attended with this symptom (for the great alexipharmac virtues, attributed to it, have occasioned it to be exhibited even in the more dangerous malignant fevers, and other disorders, which are frequently accompanied with vomiting) or that it had not been sufficiently calcined, or perfectly freed from fuch part of the regulus as might remain uncalcined. The uncalcined part being groffer than the true calx, the separation is effected by washing over with water, in the same manner as directed in page 257, for feparating earthy powders from their groffer parts.

It has been observed, that when diaphoretic antimony is prepared with nitre abounding with sea salt, of which all the common nitre contains some portion, the medicine has proved violently emetic. This effect is not owing to any particular rect is not owing to any particular quality of the sea salt, but to its quantity, by which the proportion of the nitre to the antimony is ren-

dered less.

The nitrum flibiatum is produced by the deflagration of the fulphur of the antimony with the nitre, in the same manner as the fall polycrest (page 467.) from which it differs no otherwise than in retaining some portion of the antimonial calx.

CERUSSA ANTIMONII.

Cerusse of antimony.

Take of

Regulus of antimony, one part; Nitre, three parts. Deflagrate them together, as in the foregoing process.

THE refults of both processes appear to be altogether the same. It is not necessary to use so much nitre here, as when antimony itself is employed; for the sulphur which the crude mineral contains, and which requires for its dissipation nearly an equal weight of nitre to the antimony, is here already separated. Two parts of nitre to one of the regulus are sufficient. It is better however to have an over-proportion of nitre than an under one, lest some parts of the regulus should escape being sufficients.

Ciently calcined.

It may be proper to observe, that though crude antimony and the regules yield the fame calces, yet the falts separated in washing the calces are very different. As erude antimony contains common fulphur, the acid of the fulpher unites with the alkaline bases of the nitre, and the refult is a neutral falt (page 467.) As the imflammable princible, but no fulphur, the nitre is alkalized, as it would be by charcoal or other inflammable bodies (page 426.) and is at the same time rendered more acrimonious than the common alkaline falts. If only equal parts of the regulus and nitre be employed, and the fire kept up strong for an hour or more, the falt will prove more caustic than even the potential cautery of the shops. But the caustiticity of the falt will still be far greater, if, instead of the simple regulus of antimony, the martial regulus be used.

REGULUS

REGULUS ANTIMONII MEDICINALIS.

Medicinal regulus of antimony. Edinb.

Take of
Antimony, five ounces;
Sea falt, four ounces;

Salt of tartar, one ounce. them into powder, and throw the mixture, by little at a time, into a red-hot crucible; occasionally breaking, with an iron rod, the crust that forms on the furface. When the fusion is compleated, pour out the matter into an heated cone, gently shaking it now and then, or striking it on the sides, that the regulus may fettle to the bottom; when grown cold, beat off the fcorize, and grind the regulus into a powder, which is to be kept in a close-stopt vial.

TRIS medicine is nearly similar in quality to one made with oneeighth of nitre, already described: in both processes, the antimony is freed from a fmall portion of its fulphur, which is diffipated in flame by the nitre, and absorbed This preby the alkaline falt. paration is greatly celebrated by Hoffman, and other German phyficians, in fundry obstinate chronieal diforders, and esteemed one of the best antimonials that can be given with fafety as alterants: it operates chiefly as a diaphoretic, and fometimes, though rarely, proves emetic. The dose is from three or four grains to twenty.

This regulus, reduced in a fubtile powder, is the genuine FEBRIFUGE FOWDER of Craanius (Pharm. Borusso-Brandenburg, edit 1734. pag. 107.) and has been greatly commended in all kinds of fevers, both of the intermittent and continual kind.

(Pharm. Argent. 1725. pag. 252.) It is faid that a dose or two have frequently removed these disorders, by occasioning either a salutary diaphoresis, or acting mildly by stool or vomit. The colour of the levigated powder is a purplish brown. The antimonial emetic of Boerhaave, already mentioned, which is white, is nearly similar to it in its medicinal effects.

The common falt feems to be of no further use in the process, than as it serves to promote the fusion; and even for this it is not necessary. The medicine is said to be rather more mild and certain in operation, if prepared without it.

REGULUS ANTIMONII.

Regulus of antimony.

Take of Antimony,

Nitre, Crude tartar, of each equal

parts. Grind them feparately into a powder, then mix, and rub thenr all Throw the powder, together. at several times, into a red-hot crucible, taking care to break the crust, which forms on the furface, with an iron rod: when the detonation is over, let a strong fire be made, that the matter may flow like water, theh pour it out into a warm greafed cone, which is to be gently ftruck on the fides, that the regulus may feparate and fall to the bottom; when grown cold. let the regulus be cleared from the scorize that lie a top of it.

In this proces, (which is taken from the edition of the Edinburgh pharmacopæia published in the year 1744) an alkaline salt is pro-M m 2 duced duced from the nitre and tartar, in fuch quantity, as entirely to absorb the fulphur of the antimony: the alkali, thus sulphurated, will take up more or less of the reguline part, according to its quantity, and the continuance of the fulion.

As the ingredients are above proportioned, the yield of regulus proves extremely finall, and if the fusion is long continued, scarce perceptible, almost the whole of it being taken up into the scoriæ: in order to obtain the largest quantity, the nitre ought to be diminished one half. It is convenient to rub the nitre and tartar together, and deflagrate them in an iron ladle or pan, before their mixture with the antimony; for by this means, the loss of some part of the antimony, which otherwise happens from the vehemence of the deflagration, will be prevented, a smaller crucible will serve, and less time and labour complete the process.

The mixture of nitre and tartar deflagrated together, will reduce any of the antimonial calces (as the diaphoretic antimony, cerusse, or antimony calcined by itself) into regulus; the oily matter of the tartar fupplying the inflammable principle, which all calces require for their revival into a metallic form; and the alkaline falt promoting their fulion. is the common reducing flux of the chemists; by whom it is called, from its colour, the black flux. The largest yield of regulus, hitherto obtained from antimony, has been got by calcining it without addition, as directed hereafter for making glass of antimony, and reviving the calk by fusion, with this, or other like compositions. Mr. Geoffroy, who first communicated this method to the French academy, feems to look upon

foap (the substance he happened to make use of himself) as the only one that will fucceed; but the effects of this are not different from those of the foregoing flux-Both confift of an alkaline falt, and an inflammable (not fulphureous) fubflance, which are the only materials here necessary. Upon the whole, the most advantageous process for obtaining the regulus, appears to be the following.

Let powdered antimony be calcined or roafted over a gentle fire, as directed hereafter for making the glass. Mix the calx with about equal its weight of fome reducing flux, such as the black flux above-mentioned. Melt the mixture in a crucible. with a quick fire, and when in thin fusion, pour it into a cone heated over a smoaky flame: the pure regulus will fall to the bottom, the scoriz floating on the top.

REGULUS ANTIMONII MAR-TIALIS.

Martial regulus of antimony. Take of

Antimony,

Nitre.

Crude tartar, of each one pound; Small pieces of iron, half a pound.

Heat the iron in a crucible to a white heat: then gradually add the other ingredients, first powdered and mixed together, and proceed in the fame manner as in the foregoing process.

THE nitre might here be diminished to one-fourth its weight, and the tartar to half that quantity. The pieces of iron may be small nails; the filings of the metal, lying closer together, are not so readily acted upon by the antimony.

REGULUS

REGULUS ANTIMONII STEL-LATUS. .

Stellated regulus of antimony.

This is made by melting the martial regulus feveral times with fresh mitre and tartar.

THE fimple regulus of antimony is more readily made to exhibit a starry appearance on its furface, than the martial; which it will also do by one, as well as by any number of fusions; the phænomenon entirely depends upon the regulus being pure, brought into extreme thin fusion, and cooled flowly in the cone, wirhout shaking or moving it. If the martial regulus is employed, it is convenient to add some fresh antimony (about one-fourth the weight of the regulus) to absorb such part of the iron as may be retained in it: when the whole is in persect fusion, inject, at times, about one-eighth of nitre, or fixt alkaline falt, previously dried, and made very hot.

The three foregoing reguli are at present rarely, if ever, made use of in medicine; the emetic cups, and perpetual pills, formerly made from them, have long been laid afide as precarious and unsafe. Hence the Edinburgh college, which retained them all in the edition of their pharmacopœia published in 1744, have at the late revifal rejected them. It, should seem, however, that the pure regulus, though greatly too virulent to be taken by itself, might be employed to advantage for the making of some other preparations, particularly the antimonial wine and emetic tartar: for the uncertainty in strength, which has often been complained of in those medicines, appears to proceed chiefly from faline or fulphureous matter in the antimonial preparation

made use of for communicating the impregnation to the wine or tartar; and (except the calces, which are divested of the proper antimonial virtues) the regulus is the only form in which we can expect to have the metallic part of the antimony free from such admixures, the only antimonial preparation which we can depend on heing always aqual in its own degree of power.

The scoriz produced in the foregoing processes, afford medicines less violent than the regulus itself, some of which are in considerable esteem. These scoriz consist of the sulphur of the antimony united with an alkaline salt, and a part of the regulus taken up by this compound, and rendered soluble in water.

#### Sulphur auratum antimonii.

Golden fulphur of antimony.

Let the scorize of regulus of antimony be reduced into powder, whilst warm, and then boiled for a considerable time in thrice their quantity of water. Filter the yellowish red solution, and drop into it a proper quantity of spirit of vitriol: a powder will precipitate, which is to be washed with water, till persectly edulcorated and freed from its ill smell.

# SULPHUR ANTIMONII PRÆCIPITATUM. Precipitated fulpbur of antimon? Lond.

Take of
Antimony, fixteen ounces;
Tartar, a pound;
Nitre, half a pound.

Let there be reduced feparately into powder, then mixed, thrown by degrees into a red-hot crucible, and melted with a strong M m 3 fire. fire. Pour out the matter into a conical mould; the metallic part, commonly called regulus of antimony, will fink to the bottom, the scorize swimming above it. Dissolve these scorize in water, filter the solution through paper, and precipitate the fulphur by dropping in some spirit of sea salt : lastly, wash the sulphur from the salts, and dry it for use.

#### SULPHUR AURATUM ANTI-MONII.

Golden sulphur of antimony. Edinb.

Boil, in an iron pot, four pints of foap leys, diluted with three pints of water, and throw in by degrees two pounds of powdered antimony; keeping them continually stirring, with an iron spatula, for three hours, over a gentle fire; and occasionally sup-The liquor, plying more water. loaded with the fulphur of antimony, being then ftrained thro' a woollen cloth, drop into it gradually, whilft it continues hot, fo much spirit of nitre, diluted with an equal quantity of water, as shall be sufficient to precipitate the sulphur, which is afterwards to be carefully washed with hot water. If the liquor remaining after the precipitation be purified by filtration, evaporated till a pellicle forms on the furface, and then fet to shoot, it will yield crystals of antimoniated nitre. (the same with that obtained from the washings of diaphoretic antimony.)

THE foregoing preparations are not strictly sulphurs; they contain a confiderable quantity of the metallic part of the antimony,

which is reducible from them by proper fluxes. That made by the first of the above processes contains greatest part of the metal; for as we have already seen, very little, sometimes scarce any at all, separates in the fusion. quantity of regulus taken up in the second also will be different, according to the degree of fire employed, and the length of time that the fusion is continued. These medicines, therefore, must needs be liable to great variation in point of strength, and in this respect they are, perhaps, the most precarious, though some have affirmed, that they are most certain, of the an-

timonial medicines.

The foregoing preparations prove emetic when taken on an empty stomach, in a dose of four, five, or fix grains; but in the present practice, they are scarce ever prescribed in this intention; being chiefly used as alterative doobstruents, particularly in cuta-Their emetic neous disorders. quality is easily blunted, by making them up into pills with refins or extracts, and giving them on a full stomach: with these cautions, they have been increased to the rate of fixteen grains day, and continued for a considerable time, without oc casioning any disturbance upwards or downwards. As their strength is precarious, they should be taken at first in very small doses, and increased by degrees according to their effect.

A composition of the sulphur auratum, with mercurius dulcis, has been found a powerful, yet safe, alterative in cutaneous disorders; and has completed a cure aster salivation had failed: venereal cases likewise, this medicine has produced excellent effects.

A mixture of equal parts of the fulphur and calomel (well triturated together, and made into pills with extracts, &c.) may be taken from four to eight or ten grains, morning and night; the patient keeping moderately warm, and drinking after each dole, a draught of the decoction of the woods, or other like liquors. This medicine generally promotes perspi-ration, scarce occasioning any tendency to vomit or purge, or affecting the mouth. See the Edinburgh essays. vol. i. and the Ada natur. curie∫. vol. v.

#### KERMIS MINERALIS. Kermis mineral.

Take of

Antimony, fixteen ounces: Any fixt alkaline falt, four ounces ;

Water, one pint.

Boil them together for two hours, then filter the warm liquor; as it cools, the kermes will precipitate. Pour off the water, and add to it three ounces of fresh alkaline falt, and a pint more of water: in this liquor boil the remaining antimony as before: and repeat the process a third time, with the addition of only two ounces of alkaline falt, and another pint of water; filtering the liquor as at first, and collecting the powders which subside from them in cooling.

This medicine has of late been greatly esteemed in some places, under the names of Kermes mineral. pulvis Carthussanus, poudre des Chartreaux, &c. It was originally a preparation of Glauber, and for some time kept a great secret. till at length the French king purchased the preparation from M. de la Ligerie, for a considerable sum, and communicated it to the public in the year 1720. In virtue. it is not different from the fulphurs above-mentioned; all of them owe their efficacy to a part of the regulus of the antimony, which the alkaline falt, by the mediation of the fulphur, renders foluble in water.

PANACEA ANTIMONII. Panacen of antimony.

Take of

Antimony, fix ounces: Nitre, two ounces: Common salt, an ounce and an half;

Charcoal, an ounce.

Reduce them into a fine powder, and put the mixture into a redhot crucible, by half a spoonful at a time, continuing the fire a quarter of an hour after the last injection: then either pour the matter into a cone, or let it cool in the crucible, which when cold must be broken to get it out. In the bottom will be found a quantity of regulus; above this a compact liver-coloured subflance; and on the top, a more spongy mass: this last is to be reduced into powder, edulcorated with water, and dried, when it appears of a fine golden colour.

That preparation is supposed to have been the basis of LOCKYER's PILLS, which were formerly a celebrated purge. Ten grains of the powder mixed with an ounce of white fugar-candy, and made up into a mais with mucilage of gum tragacanth, may be divided into an hundred small pills; of which one, two, or three, taken at a time, are said to work gently by stool and vomit. The compact livercoloured substance, which lies immediately above the regulus, ope-This last rates more churlishly.

appears Mm 4

appears to be nearly of the same nature with the crocus antimonit, and the former with the fulphur auratum.

#### VITRUM ANTIMONII. Glass of antimony. : Edinb.

Take of

Antimony, reduced to powder,

one pound.

Calcine it over a gentle fire, in an unglazed earthen vessel, keeping it continually stirring with an iron spatula, until the fumes cease, and the antimony is reduced into a grey powder. Melt this powder in a crucible, with an intense fire, and pour out the liquid matter, into a heated brais dish.

THE calcination of antimony, to fit it for making a transparent glass, succeeds very flowly, unless the operator be very wary and circumspect in the management of The most convenient vessel is a broad shallow dish, or a smooth flat tile, placed under a chimney. The antimony should be the purer fort, fuch as is usually found at the apex of the cones: this, grossly powdered, is to be evenly spread over the bottom of the pan, so as not to lie above a quarter of an inch thick on any part. The fire should be at first no greater than is just sufficient to raise a fume from ' the antimony, which is to be now and then stirred: when the fumes begin to decay, increase the heat, taking care not to raise it so high as to melt the antimony, or run the powder into lumps: after some time the vessel may be made red hot, and kept in this state, until the matter will not, upon being stirred, any longer, fume. If this part of the process be duly conducted, the antimony will appear

in an uniform powder, without any lumps, and of a grey colour.

With this powder, fill two-thirds of a crucible, which is to be covered with a tile, and placed in a wind-furnace. Gradually increase the fire, till the calx is in perfect fusion, when it is to be now and then examined by dipping a clean iron wire into it: if the matter, which adheres to the end of the wire, appears fmooth and equally transparent, the vitrification is completed, and the glass may be poured out upon a hot smooth stone, or copper plate, and fuffered to cool by flow degrees, to prevent its cracking and flying in pieces. It is of a transparent yellowish redcolour.

The glass of antimony usually met with in the shops, is said to be prepared with certain additions: which may perhaps render it not so fit for the purposes here defigned. By the method above directed, it may be easily made, in the requifite perfection, without any addition.

As we have feen, in a former process, antimony rendered nearly or altogether inactive by calcination; it might be expected that the calk and glass of the present process, would be likewise inert. But here the calcination is far less perfect than in the other case, where the inflammable principle of the regulus is totally burnt out by deflagration with nitre: there the calx is of perfect whiteness; and a glass made from that calx (with the addition of any faline flux, for of itself it will not vitrify) has little colour: but here fo much of the inflammable principle is left, that the calx is grey, and the glass of a high colour. The calcined antimony is said by Boerhaave to be violently emetic. Experience has shewn that the glass is so, insomuch \*s to be unfafe for internal use. It is employed chiefly, in the prefent practice, as being subservient to some other preparations, particularly the emetic tartar and antimonial wine; and in combination with wax, and some other substances, by which its power is obtunded.

# VITRUM ANTIMONII. CERATUM. Cerated glass of antimony. Edinb.

Take of

Yellow wax, a drachm; Glass of antimony, reduced into

powder, an ounce.

Melt the wax in an iron veffel, and throw into it the powdered glass: keep the mixture over a gentle fire for half an hour, continually flirring it; then pour it out upon a paper, and when cold grind it into powder.

THE glass melts in the wax, with a very foft heat; after it has been about twenty minutes on the fire, it begins to change its colour, and in ten more, comes near to that of Scots snuff, which is a mark of its being sufficiently prepared: the quantity set down above, loses about one draches of its weight in the process.

This medicine has for some time been greatly esteemed in dysenteties: several instances of its good effects in these cases, may be seen in the fifth realization of the fifth.

in the fifth volume of the Edinburgh effays, from which the above remarks on the preparation are taken. The dose is from two or three grains to twenty, according to the age and strength of the patient. In its operation, it makes fome persons sick and vomit; it purges almost every one; though it has sometimes effected a cure,

without occasioning any evacuation or sickness,

Mr. Geoffroy gives two pretty fingular preparations of glass of antimony, which feem to have fome affinity with this. One is made by digesting the glass, most subtilely levigated, with a solution of mastich made in spirit of wine. for three or four days, now and then shaking the mixture; and at last evaporating the spirit, so as to leave the mastich and glass exactly mingled. Glass of antimony thus prepared, is faid not to prove emetic, but to act merely as a cathartic, and that not of the violent kind. A preparation like this was first published by Hartmann under the name of chylistia.

The other preparation is made by burning spirit of wine upon the glass three or four times, the powder being every time exquisitely rubbed upon a marble. The dose of this medicine is from ten grains to twenty or thirty: it is said to operate mildly both upwards and downwards, and sometimes to prove

sudorific.

### Antimonium catharti-

The purging antimony of Wilson. Take four ounces of glass of antimony, finely powdered, and gradually pour thereon twelve ounces of oil of vitriol; diffil in a fand heat: and wash the powder, which remains in the bottom of the retort, till all its acrimony is loft; then dry it, and grind it with an equal weight of Glauber's cathartic falt, and a double quantity of vitriolated nitre. Let this mixture be kept a quarter of an hour in gentle fution in a crucible; and afterwards pulverized, washed, and dried for use.

Mr. Wilson, the inventor of this preparation, informs us, that it is the most certain antimonial purge

he ever met with; that it operates without nauseating the stomach; and that by the use of this powder only, he knew three confirmed poxes cured. His dose is from two

grains to ten.

We have already observed, that the glass of antimony contains a part of the regulus not fully divested of its inflammable principle. The vitriolic acid, and neutral salts, containing this acid, absorb the inflammable principle from fundry metallic and other bodies; and on this probably depends the mitigation of the glass in the prefent process.

# CAUSTICUM ANTIMONIALE. The antimonial caustic. Lond.

Take of

Crude antimony, one pound; Corrolive mercury sublimate,

two pounds.

Reduce them feparately into powder; then mix, and distil them in a wide-necked retort, with a gentle fand heat. The matter, which arises into the neck of the retort, is to be exposed to the air, that it may run into a liquor.

## BUTYRUM ANTIMONII. Butter of antimony. Edinb.

Take of

Crude antimony, one part; Corrolive mercury sublimate,

two parts.

Grind them first separately, then thoroughly mix them together, taking the utmost care to avoid the vapours. Put the mixture into a coated glass retort (having a short wide neck) so as to fill one, half of it; the retort being placed in a sand-surnace, and a receiver adapted to it, give first

a gentle heat, that only a dewy vapour may arife: the fire being then increased, an oily liquor will ascend and congeal in the neck of the retort, appearing like ice, which is to be melted down by a live coal cautiously applied. This oily matter is to be rectified, in a glass retort, into a pellucid liquor.

THESE processes are extremely dangerous, infomuch that even the life of the operator, though tolerably versed in common pharmacy. may be affected for want of taking due care herein. Boerhaave relates, that one, who from the title he gives him is not to be supposed inexpert in chemical operations. or unacquainted with the danger attending this, was fuffocated for want of proper care to prevent the bursting of the retort. The fumes which arise, even upon mixing the antimony with the fublimate, are highly noxious, and fometimes iffue fo copiously and suddenly, as very difficultly to be avoided. utmost circumspection therefore is necessary.

The caustic or butter, as it is called, appears to be a folution of the metallic part of the antimony in the marine acid of the fublimate; the fulphur of the antimony, and the mercury of the sublimate, remain at the bottom of the retort, united into an ethiops. This folution does not succeed with spirit of falt in its liquid flate, and cannot be effected, unless (as in the case of making sublimate) the acid is highly concentrated, and both the ingredients strongly heated. If regulus of antimony was added in the distillation of spirit of sea salt without water, a like folution would be made.

When the congealed matter that arises into the neck of the retort, is liquefied

liquefied by the moisture of the air, it proves less corrosive than when melted down and rectified by heat : though it feems, in either case, to be sufficiently strong for the purposes it is intended for, as the confuming of fungous flesh, and the callous lips of ulcers. It is remarkable, that though this faline concrete, readily and almost entirely dissolves by the humidity of the air, only a small quantity of white powder separating, it nevertheless will not dissolve on putting water to it directly: even when previously liquefied by the air, the addition of water will precipitate the folution.

## CINNBARIS ANTIMONII. Cinnabar of antimony. Lond.

Let the matter, which remains in the retort after the distillation of the caustic, be sublimed in a coated matrass, in an open fire.

Edinb.

As foon as the red vapours begin to appear in the distillation of the butter, change the receiver, without luting the junctures; and increase the fire until the retort becomes intensely red hot: in an hour or two, the whole of the black powder will be sub-hlimed and its colour changed into red. Then break the retort, and diligently separate the cinnabar, which will be found in the neck, from the black droffy matter.

THE cinnabar of antimony is composed of the sulphur of the antimony, and the mercury of the sublimate, which are perfectly the same with the common brimstone and quicksilver, of which the cinnabaris fastitia is made. The antimonial cinnabar therefore, whose ingredients are laboriously extract-

ed from other substances, is not different from the common cinnabar made with the same materials procured at a much chéaper rate. The former indeed is generally of a darker colour than the other, and has fomewhat of a needled appearance; like that of antimony itself; from whence it has been supposed to participate of the metallic part of that mineral. But it appears from experiment, that both the colour and needled form are entirely accidental, and owing to the mixture containing a larger portion of fulphur, and being fublimed in a more languid manner.

MERCRIUS VITE, feu PULVIS
ALGEROTHI.

Mercury of life, or Algeroth's powder. Take of

Rectified butter of antimony, as much as you pleafe.

Pour to it a sufficient quantity of spring water, and an exceeding white powder will be precipitated: edulcorate this by repeated assusping as flow fire.

THIS powder has not, as its name should seem to imply, any thing of mercury in it, but is folely composed of the reguline part of the antimony, corroded by the acid spirit of sea salt; which acid is so closely united, as not to be separated by any ablution with wa-Le Mort directs some alkaline salt to be dissolved in the water, in order to obund the acid: feveral other methods also have been contrived for correcting and abating the force of this violent emetic; but they either leave it still virulent, or render it inert. has therefore for a long time been laid aside by practitioners, and the Edinburgh college, who retained it in their preceding edition, have, at

the late revifal of their pharmacopæia, expunged it.

BEZOARDICUM MINERALE.

Bezoar mineral.

Take any quantity of butter of antimony newly rectified, and gradually drop into it spirit of nitre, till the efferve scence ceases. Draw off the spirit in a glass vessel, placed in a sand heat, till a dry powder remains behind: add to this a little fresh spirit of nitre, and again exsiccate it. Repeat this a third time; then commit the powder in a crucible to a naked sire, till it has received an almost white heat, and detain it in this state for half an hour.

This preparation may be easier made, and with greater fafety to the operator, by dropping the butter of antimony into three or four times its weight in spirit of nitre, and distilling the mixture in a retort, until a dry white mass is left behind, which is afterwards to be calcined, as above directed. may likewise be made by distilling spirit of nitre from the mercuries vitæ, and calcining the remainder: or by deflagrating the mercurius witze with thrice its weight of pure nitre. This last method, proposed by Wedelius, is followed by the Augustan college.

Bezoar mineral was formerly held in great esteem as a diaphore-tic, but its reputation is at present almost lost. It is not different in medical virtue, or in any sensible quality, from the calces of anti-mony made directly by destagration with nitre, some of which have generally supplied its place in the shops. It appears at first pretty extraordinary, that the violent caustic, butter of antimony, should be rendered indolent by the corrosive spirit of nitre: how this happens

will be easily understood, upon considering that the nitrous acid expels the marine (to which the caustic quality of the butter is owing) and is itself expelled from most metalic substances by sire.

### TARTARUM EMETICUM.

Emetic tartar, Lond.

Take of

Washed crocus of antimony, Crystals of tartar, each half a pound;

Water, three pints.

Boil them together for half an hour; then filter the liquor, and after due evaporation, fet it by to crystalize.

Edinb.

Take of

Cream of tartar, four ounces; Glass of antimony powdered, two ounces.

Boil them together in fix pints of water, for ten hours, ftirring them frequently with a spatula, and adding more water as there is occasion. Fiker the liquor while warm; and evaporateit, either to dryness, or only till a pellicle forms that it may shoot into crystals.

It may likewife be prepared, in the fame manner, from the crucus metallorum washed.

This preparation has been usually made with the unwashed crocus of antimony; by employing, as here directed the washed crocus, or the glass, it proves of a whiter colour, and likewise more certain is frength (see page 533;) though it will still be somewhat precanous in this last respect, if the crystallization is complied with: for some of the tartar, even though the operation is performed with a good deal of case, will be apt to shoot

by itself, retaining little or nothing of the antimony. It should seem therefore more eligible, as soon as the solution has passed the filter, to proceed to the total evaporation of the liquor, or at least to evaporate lower than is usual for crystallization, that the whole may shoot at once.

The title of this medicine expresses its principal operation. It is one of the best of the antimonial emetics, acting more powerfully than the quantity of crocus contained in it would do by itself, though it does not so much russe the constitution. And indeed antimonials in general, when thus rendered soluble by vegetable acids, are more safe and certain in their effects, than the violent preparations of that mineral exhibited by them-

selves; the former never varying in their action from a difference in the food taken during their use, or other like circumstances, which occasioning more or less of the others to be dissolved, make them operate with different degrees of force. Thus crude antimony, where acid food has been liberally taken, has sometimes proved violently emetic; whilst in other circumstances, it has no such effect.

The dose of emetic tartar, when designed to produce the full effect of an emetic, is from four to fix or eight grains. It may likewise be advantageously given in smaller doses, half a grain for instance, as a diaphoretic and alterative in cutaneous disorders; and added, in the quantity of a grain, as a stimulus to ipecacuanha, &c.

### SECT. IX.

### Preparations of bismuth.

THIS metal resembles in appearance the regulus of antimony; but differs greatly from it, in its pharmaceutical properties and medical qualities. It melts in a very small heat, long before ignition; and totally dissolves, with great effervescence, in aquafortis. which only corrodes the antimonial metal. As a medicine, it feems, when pure, to have little or no effect; though some preparations of it were formerly counted diaphoretic. At present, only one preparation comes under the notice of the apothecary or the chemist, and that designed for external use.

MAGISTERIUM BISMUTHI.

Magistery of bismuth.

Dissolve bismuth in a proper quantity of aquafortis, without heat,

adding the bismuth by little and

little at a time. Pour the folution into fixteen times its quantity of fair water: it will grow milky, and on ftanding for fome time deposite a bright white precipitate: the addition of spirit of wine will expedite the precipitation. Wash the powder in fresh parcels of water; and dry it in a shady place betwixt two papers.

This preparation is of some esteem as a cosmetic, which is the only use it is now applied to. The diaphoretic virtues, attributed to it when taken internally, have very little soundation, and by the present practice are not at all regarded. It was proposed to be received in our pharmacopæia at the late revisal, but was sound much too insignificant to be admitted there.

SECT.

### SECT. X.

### Preparations of zinc.

HIS metal melts in a red heat; and if the air is admitted, flames, and fublimes into light, white, downy flowers; if the air is excluded, it arifes by a strong fire, in its metallic form. Sulphur which unites with, or fcorifies all the other metals except gold does not act on zinc. Acids of every kind diffolve it.

Zinc, its flowers or calces, and folutions, taken internally, prove frong and quick emetics; in small doses, they are said to be diaphoretic. Externally they are cooling,

astringent, and desiccative.

Purification of zinc.

Melt zinc with a heat no greater than is just sufficient to keep it fluid. Stir it strongly with an fron rod, and throw in alternately pieces of sulphur and of tallow, the first in largest quantity. If any consistent matter, or scoria, forms on the top, take it off and continue the process, until the sulphur is sound to burn freely and totally away on the surface of the sluid zinc.

Zinc usually contains a portion of lead, which this process effectually separates. Sulphur united with lead forms a mass, which does not melt in any degree of fire that zinc is capable of sustaining.

FLORES ZINCI.

Flowers of zinc.

Let a large and very deep crucible,
or other deep earthen vessel, be

placed in a furnace, in an inclinedituation, only half upright. Put a small quantity of zinc into the bottom of the vessel, and apply a moderate sire, no greater than is necessary to make the zinc slame; white slowers will arise, and adhere about the sides of the vessel like wool. When the zinc ceases to slame, stir it with an iron rod, and continue this operation till the whole is sublimed.

THESE flowers should seem preferable for medicinal purposes, to tutty, and the more impure sublimates of zinc, which are obtained in the brass works; and likewise to calamine, the natural ore of this metal, which contains a large quantity of earth, and frequently a portion of heterogeneous metallic matter.

SAL, feu VITRIOLUM ZINCI.
Salt, or wittiol of zinc.

Diffolve purified zinc, by a gentle heat of fand, in a mixture of one part of oil of vitriol, and four of water. Filter the folution, and after due evaporation, fet it to crystallize.

This falt is an elegant white vitriol. It differs from the common white vitriol, and the fal vitrioli of the shops, only in being purer, and perfectly free from any admixture of copper, or such other soreign metallic bodies as the others generally contain.

SECT.

#### S E C T. XI.

### Compound metallic preparations.

LAPIS MEDICAMENTOSUS.

The medicinal flone.

Land.

AKE of
Litharge,
Bole armenic, or French bole,
Alum, each half a pound;
Colcothar of green vitriol, three
ounces;

Vinegar, a quarter of a pint.

Mix and dry them till they grow hard.

THIS preparation is employed externally as an aftringent, for faftening loofe teeth, preferving the gums, healing and drying up ulcers and wounds, and repreffing defluctions of thin acrid humours upon the eyes. It is fometimes used in injections for checking a gonorrhuea, after the virulence is expelled. A preparation much refembling this is said, in the Memoirs of the French academy, to be generally esteemed among the surgeons in the army as a vulnerary.

Speificum adstringens.

Maetzii.

An astringent preparation taken from Maetz, which has been sold under the name of

Colbatch's flyptic powder.

Take any quantity of iron filings, and as much spirit of salt as will rise above them three or sour inches. Digest them together with a gentle heat, till the spirit ceases to act on the metal; then pour off the liquor, evaporate it to one half, and add thereto an equal weight of sugar of lead. Continue the evaporation, with

a finall heat, until the matter remains dry, and assumes a red colour.

If the process is stopt as soon as it becomes dry, it has exactly the appearance of Colbatch's powder. It must be kept close from the air, otherwise it deliquiates.

THIS is said to be the styptic. with which fo much noise was made fome time ago, by the author of the novum lumen chirurgie; and for the fale of which. a patent was procured; only in that was used oil of vitriol, instead of the spirit of salt in this. a difference not very material. The preparation stands recommended in all kinds of hæmorrhages and immoderate fluxes, both internally and externally: the dole is from four grains to twelve. It is undoubtedly an efficacious styptic, but for internal use a dangerous one. See the article LEAD, and its preparations.

ANTIHECTICUM POTERII.

Poterius's antichectic.

Take of

Martial regulus of antimony, fix ounces;

Fine tin, three ounces.

Melt these together in a crucible; then pour them out into a warm greased mortar, and when the mass is grown cold, grind it into a powder. Add to this thrice its weight of pure nitre, and deflagrate the mixture in a crucible, throwing in only a spoonful at a time; then calcine it [that

is, keep in fusion for an hour: and having afterwards ground it into an impalpable powder, pour thereon a fufficient quantity of warm water: stir them well together with a peftle, till the water grows milky, which, thus loaded with the finer pasts of the powder, is to be poured off, and fresh water put to the remainder: repeat this operation till nothing but indisfoluble feces remain behind. Suffer all the milky liquor to reft : a powder will fall to the bottom, which is to be washed with repeated affusions of warm water, and laftly dried for use.

THE regulus of antimony should be melted before the tin is added to it: for if they both are put into the crucible together, a part of the tin will be diffipated by the heat requilite for the fulion of the

regulus.

The chemists have been greatly divided with regard to the proportion which these two ingredients ought to bear to one another. Some vary so much from the above prescription, as to order two parts of the antimonial regulus to one of tin; others no more than one part of the former to fix of the latter. Nor have they agreed upon the colour which this preparation ought to have; fome prefering that which is perfectly white; whilft others look upon a blueish tinge as a mark of the proportions having been duly observed, and the operation regularly performed: in the process above, it seems intended to be white: for without the obfervance of certain encheireses, not there mentioned, as particularly calcining the powder after the ablution, it will scarce have any thing of a blueish cast.

Practical physicians do not differ less in the accounts which they give of the virtues of this celebrated medicine. Some extol it as an excellent diaphoretic, &c. others are ready to vouch, that it has done most eminent service in hectical cafes; whilst many of no small note, . are not only confident that it has none of the virtues attributed to it. but utterly condemn it as unsafe, and capable of producing the very disorders said to be remedied by its This affair probably will not be satisfactorily determined, till the virtues of calx of tin and calx of antimony (which this medicine is a mixture of ) shall be better ascertained than they are at present. In the mean time, the use of the antibedic is in common practice laid alide; and is not likely to be ever introduced again.

BEZOARDICUM TOVIALE. Bezoar with tin.

Take of

of antimony, Regulus three ounces:

Pure tin, two ounces;

Corrolive fublimate mercury, five

Melt the regulus of antimony in a crucible, and put to it the tin. fo as to make a new regulus; to which after being levigated, add the corrofive sublimate, and distil the mixture in a retort. Let the butter which arises in this process be fixed by three repeated distillations with thrice its own quantity of spirit of nitre. The powder is then to be calcined; thrown, whilst ignited, into a proper quantity of spirit of wine; and afterwards dried for use.

This preparation is not greatly different from the foregoing. butter feems to contain more of the the tin, than of the antimonial regulus, united with the marine acid of the sublimate: the nitrous spirit expels the marine, and is itself afterwards expelled in the calcination, leaving the powder a mere calx, similar to one prepared from the same ingredients in a less troublesome manner, by deslagration with nitre.

### ÆTHIOPS ANTIMONIALIS.

Antimonial ethiops.

Let equal quantities of antimony and sea falt be melted together in a crucible for an hour; when grown cold, a regulus (improperly so called) will be found in the bottom; which is to be separated from the scorize that lie above it, and ground with an equal weight of purified quickfilver, until they are united.

This medicine is said to be of remarkable efficacy in venereal cases of long standing, in cancerous tumours, scorbutic and scrophulous disorders, obstinate glandular obstructions, and fundry other chronical distempers, which elude the force of the common medicines. A few grains may be given at first; and the dose gradually increased, according to its operation, to a scruple or more. It acts chiefly by promoting perspiration: in fome conflitutions, it proves purgative; and in others, if the dose is considerable, emetic.

Sundry other preparations of this kind have of late been held by fome people in confiderable effects, though not taken notice of by common practice. They have been generally composed of mercury united by triture either, with crude antimony, the medicinal regulus, or the golden or precipitated sulphur.

Mr. Malouin, of the faculty of Paris, made trial of different methods for uniting mercury and crude antimony into an ethiops. Those which succeeded I shall here extract from his chimie medicinale.

On grinding together two parts of antimony and one of mercury, the mercurial globules difappeared in three hours, and the compound proved fimilar in appearance to the ethiops made with the same proportions of mercury and common sulphur. Equal parts of the antimony and mercury were much more difficultly united, requiring the triture to be continued for two days; though it was found also, even with these proportions, that when the mercury was added, not all at once, but by little and little, the union might be effected in five hours. As common ethiops is made more perfect, in regard to the intimate union of the ingredients, by heat than by triture; the most persect antimonial ethiops also was obtained by means of fire, in the following manner:

A heated crucible is to be rubbed in the infide with tallow, immediately covered, and fet in the fire. When red hot, throw in the antimony beateninto coarse powder, and cover the vessel again. When the antimony is melted, take the crucible out of the fire. throw in a small bit of tallow, pour an equal weight of heated mercury on different parts of the furface, cover the crucible for a moment, and, while the mixture is Rill fluid, pour it out into a heated iron mortar. When grown cold, reduce it into a powder, which is to be levigated on a marble.

On this black powder the author directs fome spirit of wine to N n be be burned two or three times; an article which may very safely be omitted, as it can nowise affect the medicine. The only difficulty in the process relates to the degree of heat of the melted antimony: if it is not sufficiently sluid, the mercury cannot equably unite with it; and if over hot, great part of the mercury will be diffipated.

Mr. Malouin commends this ethiops, as a medicine of great efficacy in glandular obfiructions, obstinate cutaneous maladies of different kinds, inveterate rheumatisms, &c. It acts most commonly by urine and perspiration, rarely purges, or occasions only some slight nauseæ. The dose is from one grain to twenty, two or three times a day, that is, from one to fixty grains in a day. In some persons a drachm has no sensible operation: others are moved by fix grains.

### PART IV.

### Medicinal Compositions.

### CHAPTER I.

### Powders.

HIS form receives fuch materials only, as are capable of being sufficiently dried to become pulverable, without the loss of their virtue. There are many Substances, however, of this kind. which cannot be conveniently taken in powder: bitter, acrid, fetid drugs, are too disagreeable: emollient and mucilaginous herbs and roots are too bulky: pure gums cohere, and become tenacious in the mouth: fixed alkaline salts liquefy upon exposing the composition to the air: and volatile alkalies exhale.

The dose of powders, in extemporaneous prescription, is generally about half adrachm: itrarely exceeds a whole drachm; and is not often less than a scruple. Subflances which produce powerful esfects in smaller doses are not trusted to this form, unless their bulk is increased by additions of less efficacy: those which require to be given in larger ones, are better sitted for other forms.

The usual vehicle for taking the lighter powders in, is any agree-

(A

able thin liquid. The ponderous powders, particularly those prepared from metallic substances, require a more consistent vehicle, as syrups; for from thin ones, they soon subside. Resinous substances likewise are most commodiously taken in thick liquors: in thin ones, they are apt to run into lumps, which are not easily again dissoluble.

General rules for making powders.

I.

Particular care ought to be taken that nothing carious, decayed, or impure, be mixed in the compofition of powders: the stalks and corrupted parts of plants are to be separated [E.]

TT

The dry aromatics ought to be fprinkled, during their pulverization, with a few drops of any proper water [E.]

TTT

The moister aromatics may be dried with a very gentle heat, before they are committed to the mortar [E.]

N n 2 · IV. Gams,

IV.

Gums, and such other substances as are difficultly pulverable, should be pounded along with the drier ones, that they may pass the seve together [E.]

No part flould be separated for use, until the whole quantity put into the mortar has passed the sieve, and the several sistings been mixed together; for those parts of one and the same subject, which powder first, may prove different, at least in degree of efficacy, from the rest.

VI.

Powders of aromatics are to be prepared only in small quantities at a time, and kept in glass vesfels very closely stopped [E.]

If powders are long kept, and not carefully secured from the air, their virtue is in great measure destroyed, although the parts in which it confiss should not in other circumsances prove volatile. Thus, though the virtues of ipecacoanha are so fixed as to remain entire even in extracts made with proper menstrua, yet, as the college of Wirtemberg observes, if the powdered root be exposed for a length of time to the air, it loses its emetic-quality.

PULVIS ANTILYSSUS.

Powder against the bite of a mad dog.

L. E.

Take of
Afh-coloured ground liverwort,
two ounces;
Black pepper, one ounce.

Beat them together into a powder.

In our former pharmacopoeia, which more clearly distinguishes it the quantity of pepper was equal from the other. The caution of to that of the herb: which rendering the powder in a close vering the powder greatly too hot, is a very necessary one; for

the above diminution of it became necessary. The virtue which this medicine has been celebrated for, is expressed in its title; the dose is a drachm and a half, to be taken in the morning fasting, in half a pint of cows milk warm, for sour mornings together. See page 166.

PULVIS ARI COMPOSITUS.

Compound powder of arum.

Lond.

Take of

Arum root, fresh dried, two ounces;

Yellow water flag roots, Burnet faxifrage roots, each one ounce:

Crabs eyes prepared,
Cinnamon, each half an ounce;
Salt of wormwood, two drachms.
Beat them into a powder, which is
to be kept in a close vessel.

In former editions of the London pharmacopæia, one of the ingredients in this composition was called acorus vulgi or vulgaris; a name which has been applied, by different writers, both to calamus arematicus, and to the gladiolus luteus, or common yellow water-flag. In this uncertainty, the compounders generally took the former. But as the medicine was first contrived by a German phyfician, Birkmann, and as in some of the German pharmacopœias, the acorus vulgaris is explained to be the water-flag, the London college have now, rather in conformity to the original prescription, than from any opinion of the virtues of the water-flag (which appear, when the root is dried and powdered, to be very inconsiderable) made choice of this last, and expressed it by the name which more clearly diffinguishes it The caution of from the other. keeping the powder in a close vesif exposed to the air, the alkaline falt, imbibing moisture from it. would run into a liquid state. Two alkaline salts have been generally directed; but as they differ from one another only in name (see page 426.) one of them is here justly omitted, and supplied by a proportionable increase of the other. Posfibly the prepared crabs eyes might also have been dropped, unless they are intended to augment the volume of the medicine (an intention not very necessary in this compofition) for they do not appear to have any medicinal virtue which alkaline salts do not posses in a greater degree.

Agreeably to the above remark, in the former edition of this book, the college of Edinburgh, in the late revifal of their pharmacopoeia, have omitted the crabs-eyes, and continued the former practice of using calamus aromaticus for the alcorus vulgaris. They have likewise exchanged the cinnamon for canella alba; and the alkaline salt for a neutral one, better suited to the form of a powder. Their prefent formula is as follows:

Take of
Arum roots, newly dried, two
ounces;
Calamus aromaticus,
Burnet, faxifrage roots, each one
ounce;
Canella alba, fix drachms;
Vitriolated tartar, two drachms.

Mix and make them into a pow-

THE pulvis ari compositus was originally intended as a stomachic: and in weaknesses and relaxations of the stomach, accompanied with a surcharge of viscid humours, it is doubtless a very useful medicine. It frequently also has good effects in theumatic cases, of which I have

known some instances: the dose may be from a scruple to a drachm, two or three times a day, in any convenient liquor. It should be used as fresh as possible, for its virtue suffers greatly in keeping: the arum root in particular, its capital ingredient, soon loses the pungency in which its efficacy principally consists.

### PULVIS e BOLO COMPOSITUS fine OPIO.

Compound powder of bole without opium.

Lond.

Take of
Bole armenic, or French bole,
half a pound;
Cinnamon, four ounces;
Tormentil root,
Gum Arabic, each three ounces;
Long pepper, half an ounce.
Reduce these ingredients into powder.

### PULVIS e BOLO COMPOSITUS cum OPIO.

Compound powder of bole with opium.

Lond.

Take of opium strained, three drachms.

Dry it a little, so as to render it easily pulverable; and add to it the foregoing species, that they may all beat into a powder together.

This powder, with opium, is an elegant reform of the species of Fracastorius's confection, commonly called diascordium; confishing only of such of the ingredients of that composition, as are most conducive to the intention for which it is at present prescribed. Forty-five grains of the powder contain one of opium.

Nn 3

The

The powder is directed to be kept in the shops without opium, for cases, where the affistance of that drug is not wanted. It is a warm, glutinous astringent; and is given in sluxes, or other disorders, where medicines of this class are proper, in doses of a scruple or half a drachm.

PULVIS e CERUSSA
COMPOSITUS.
Compound powder of ceruffei
Lond.

Take of
Ceruffe, five ounces;
Sarcocolla, an ounce and a
half;

Gum tragacanth, half an ounce. Beat them together into a powder.

This composition is the trochiscial albi of Razi, brought back to its original simplicity with regard to the ingredients, and without the needless trouble of making it into troches. It is employed for external purposes, as in collyria, lotions, and injections, for repelling acrimonious humours, and in inflammations.

PULVIS e CHELIS
CANCRORUM COMPOSITUS.
Compound powder of crabs-claws.

Lond.

Take of
The tips of crabs-claws prepared, one pound;
Pearls prepared,
Red coral prepared, each three
ounces.
Mix them together.

Edinb.

Take of
Red coral prepared, one ounce;
Black tips of crabs-claws prepared, two ounces.
Mix them together.

THESE powders have lost several of their ingredients, without any injury to their virtues; and possibly they would still bear a farther reduction; for both the crabs-eyes and claws are by themselves at least as effectual as any composition of them with pearls and coral. In some of our hospitals, the following composition is substituted:

PULVIS TESTACEUS.

Compound testaceous powder.
Take of

Oystershells prepared, one pound; White chalk, half a pound. Mix them together.

THIS cheap absorbent powder is at least equally valuable, as a medicine, with the more costly and compounded crabs-claw and bezoardic powders of the shops. These kinds of preparations are given from half a scruple to halfadrachm. for absorbing or destroying acidities in the first passages; which seems to be the only good effect that can be reasonably expected from these fimple antacid earths. If they meet with no acid to diffolve them, they promise to be injurious rather than beneficial (see page 62.) They have often been given in fevers, under the notion of alexipharmacs and fudorifics, from a supposition that these disorders are occasioned by a latent acid; and, though this theory is now exploded, the practice built upon it is, in good mea-So far are fure, still continued. absorbents from being useful in these cases, that substances of a directly contrary quality, mild acidulous liquors, are in general the most successful remedies, wherever the vis vitæ is not too far depressed; and where it is, the infipid indolent earths can contribute nothing to support or raise it.

It may here be proper to take notice of a quality hitherto little expected from these kinds of substances; that of strongly promoting putrefaction. Flesh mixed with a fmall proportion of chalk, and exposed to a heat equal to that of the human body, not only corrupts fooner than without this addition, but likewise in a far greater degree, resolving in a few days into a perfect mucus. This quality of the absorbent powders (for the discovery of which, with many other curious experiments on the fame subject, the public are obliged to the ingenious Dr. Pringle) feems to forbid their use in all those kinds of fevers, where the animal juices are already too much disposed to a putrefactive state. We have above observed, that, in these cases, though very frequently employed, they are at best unserviceable; perhaps their ill effects would be oftner feen, if it was not for the quantity of acids usually given in acute diseases.

## PULVIS BEZOARDICUS. Bezoardic powder. Lond.

Take of

Compound powder of crabsclaws, one pound; Oriental bezoar prepared, one

Oriental Dezoar prepared, on

Mix them together.

BEZOAR has hitherto been an ingredient in the foregoing composition, which was then called Gascoigne's powder; though notwith-standing the addition which this article made to the price, it added nothing to the virtue of the medicine. The college of London have therefore very pradently disected an absorbent powder, without this costly article; and composed another, diffinguished by its name, for the use of those who

expect any particular virtues from it. The Edinburgh college have entirely expunged this unnecessary drug, and take no farther notice of it in their pharmacopæis, than barely giving it a place in the catalogue of simples.

### PULVIS CONTRAYERVÆ COMPOSITUS.

Compound powder of contrayerva.

Lond.

Take of

Compound powder of crabsclaws, a pound and a half; Contrayerva root, five ounces. Make them into a powder.

Edinb.

Take of

Contrayerva root, fix drachms; Virginian fnakeroot, two drachms;

English faffron, one drachm; Compound powder of crabsclaws, two ounces.

Make them into a powder.

THESE powders were formerly directed to be made up into balls with water, (and then called LAPIS CONTRAYERVÆ) a piece of trouble now laid afide as needless; for it was necessary to reduce the balls into powder again before they could be used. Nor did that form . contribute, as has been imagined, to their preservation; for it is scarce to be supposed, that the powder will lose more by being kept for a reasonable length of time in a close stopped glass, than the balls will, in the humectation with water, and exficcation in the air. before they are fit for being put by These medicines have a to keep. much better claim to the title of an alexipharmac and fudorific, than the two foregoing compositions. contrayerva, inakeroot, and faffron, by themselves are such, and prove very serviceable in low fevers,

where the vis vitæ is weak, and a diaphoresis to be promoted. It is possible, that the crabs-claw powders are of no farther service than as they divide these powerful ingredients, and render them supportable to the stomach.

PULVIS ad EPILEPTICOS de GUTTETA dictus.

Epileptic powder. Edinb. Take of

Wild valerian root,
Peony root, of each equal parts.
Make them into a powder.

This powder has undergone a great reduction of its ingredients, to its advantage as a medicine; the articles rejected being either infignificant, or at best far inferior to those retained, and consequently increasing the bulk of the compofition, without communicating a proportionable share of efficacy. Perhaps for the same reason, the peony roots are not altogether unexceptionable. The powder, however, as now reformed, may be looked on as a medicine of some importance for the purposes expressed in its title, far superior to those of fimilar intention in other pharmacopæias. The dose is from ten grains to half a drachm for children, and from half a drachm to two drachms for adults. The abforbent powders, generally directed in these kinds of compositions, are here more prudently omitted, as they may eafily be mixed extemporaneously, where particular cases may require them. For children, these additions are often necessary, as in most of their disorders, acidities in the first passages have a confiderable share: in adults, they are rarely of use.

> PULVIS e MYRRHA COMPOSITUS.

Compound posuder of myrrh.

Lond.

Take of
Rue leaves, dried,
Dittany of Crete,
Myrrh, each an ounce and a
half;
Ata'œrida,
Sagapenum,
Ruffia caftor,
Opopanax, each one ounce.
Beat them together into a powder.

This is a reformation of the trochifci e myrrha, a composition contrived by Kazi against uterine obstructions. It may be taken in any convenient vehicle, or made into boluses, from a scruple to a drachm or more, two or three times a day.

PULVIS ad PARTUM.

Powder to promote delivery.

Edinb.

Take of
Borax, half an ounce;
Castor,
Saffron, each a drachmand a half;
Oil of cinnamon, eight drops;
Oil of amber, fix drops.
Beat the species together into a powder, to which add the oils, and mix the whole well together.

This medicine has long been held in efteem for the purpose expressed in its title: nevertheless, its real efficacy, and what share thereof is owing to each of the ingredients, has not been sufficiently determined: the borax, though by somethought to be of little importance, does not perhaps contribute the least to its virtue. The dose is from a scruple to a drachm, or so much as can be conveniently taken up at once on the point of a knife. It should be kept in a very close vessel.

vessel, otherwise it will soon lose a considerable deal of its more valuable parts.

### PULVIS e SCAMMONIO COMPOSITUS.

Compound powder of fcammony.

Take of

Scammony, four ounces;
Calcined hartshorn prepared,
three ounces.
Grind them diligently together in-

to a powder.

HERE the scammony is divided by the earthy calx, and thus rendered somewhat more soluble, and less adhesive; hence its purgative quality is promoted, at the same time that it becomes less griping. The dose of the compound is from sisteen grains to half a drachm.

This powder has been usually prepared with diaphoretic antimony and crystals of tartar (instead of the calcined hartshorn above directed) and called from its first publisher, PULVIS CORNACHINI, which, in the Edinburgh pharmacopæia, is thus directed:

Take of

Diaphoretic antimony, Creme of tartar, Scammony, each equal parts. Make them into a powder.

This may be given to the quantity of a drachm or more. In other prescriptions, the tartar and antimonial calx bear nearly the same proportion to the scammony, as the calcined hartshorn in the preceding formula. It appears probable, that neither of these ingredients are of any farther use, than as they divide the texture of the scammony; though Cornachini proposes notable advantage from some deobstruent quality in the tartar, whereby the vessels shall be

opened, and the noxious humours prepared for expulsion; and from the preparation of antimony, though it have no sensible operation, he expects some share of the same saccess, which sometimes attends the rougher preparations of that mineral.

### PULVIS e SENA COMPOSITUS.

Compound powder of Sena. Lond.

Take of

Crystals of tartar, Sena, each two ounces; Scammony, half an ounce; Cloves, Cinnamon,

Ginger, each two drachms. Powder the scammony by itself; and all the other ingredients together; then mix them.

### PULVIS DIASENNÆ.

Take of
Creme of tartar,
Sena, each two ounces;
Scammony,
Ginger, each half an ounce;
Make them into a powder.

THESE powders are given as cathartics, in the dose of two scruples, or a drachm. The spices are added, not only to divide, but to warm the medicine, and make it sit easier on the stomach. The scammony is used as a stimulus to the sena; the quantity of the latter necessary for a dose, when not assisted by some more powerful material, being too bulky to be conveniently taken in this form.

## PULVIS STERNUTATORIUS. Sternutatory powder. Lond.

Take of Alarum,

Marjoram,

Marjoram,
Marum Syriscum leaves, dried,
Lavender flowers, dried, each
equal weights.

Rub them together into a powder.

PULVIS CEPHALICUS.

Cephalic powder.

Edinb.

Take of
Afarum,
Marjoram, of the leaves of each,
equal parts.

Beat them together into a pow-

The titles of these powders sufficiently express their intention. They are both agreeable and efficacious errhines, and superior to most of those usually sold under the name of herb soust.

## PULVIS STYPTICUS. Styptic powder. Edinb.

Take of
Alum, half an ounce;
Dragons blood, two drachms.
Make them into a powder.

This powder has long been in repute as an aftringent, under the title of Pulvis stypticus Helvetii. It is undoubtedly a very powerful medicine; though the dragons blood feems to have little thare in its effects. Some direct the ingredients to be melted together before they are powdered: but this circumftance does not appear to be necessary.

PULVIS e SUCCINO COMPOSITUS.

Compound powder of amber,

Land.
Take of
Amber prepared,
Gum Arabic, each ren drachms;

Juice of hypocifis,
Balaustines,
Japan earth, each five drachms;
Olibanum, half an ounce;
Strained opium, one drachm.
Beat them together into a powder.

This powder is composed of the . more unexceptionable ingredients of the Trochisci B CARABE of . our former pharmacopæia. The articles omitted, which are as many in number as those now retained, were manifestly absurd or superfluous; and the making it up into troches, a very unnecessary trouble. The medicine, as now reformed, may be looked upon as an useful, and tolerably elegant aftringent; though possibly the ingredient, which it receives name from, contributes little to its virtue. scruples of the composition contain one grain of opium.

#### PULVIS e TRAGACANTHA COMPOSITUS:

Compound powder of gum tragacantb.

Lond.

Take of
Gum tragacanth,
Gum Arabic,
Marihmallow root, each an
ounce and a half;
Starch,
Liquorice, each half an ounce;
Double-refined fugar, three
ounces;
Grind them into a powder.

### PULVIS DIATR AGACANTHI. Edinb.

Take of
Gum tragacanth, one ounce and
a half;
Marshmallow root,
Liquorice,
Starch, each half an ounce.
Beat them together into a powder.

Вотн

BOTH these powders are mild emollients; and hence become ferviceable in hectic cases, tickling coughs, strangury, some kinds of sivine fluxes, and other disorders proceeding from a thin acrimonious state of the humours, or an abrasion of the mucus of the intestines: they foften, and give a greater degree of confishency to the former, and defend the latter from being irritated or excoriated by them. All the ingredients coincide in thefe general intentions; the marshmal-low root, however, is somewhat too bulky for this form, and likewife subjects the composition to grow mouldy in keeping, an inconvenience which the cold feeds formerly employed in these powders were particularly liable to. The dose is from half a drachm to two or three drachms, which may be frequently repeated.

### HIERA PICRA. Lond.

Take of

The gum extracted from Socotorine aloes, one pound; Canella alba, three ounces. Beat them separately into powder, and then mix them together.

Pulvis HIERA PICRA dictus.

Edinb.

Take of

Socotorine aloes, four ounces; Virginian fnakeroot, Ginger, each half an ounce. Mix, and beat them into a powder.

THESE compositions were originally directed to be made into an electary: with us, they have been rarely used in that form, and not often in this of a powder, on account of their great nauseousness. They are chiefly employed as the basis of a tincture called tinctura secra. See page 297.

### SPECIES AROMATICÆ.

Aromatic Species.

Take of

Cinnamon, two ounces; Leffer cardamom feeds, husked, Ginger,

Long pepper, each one sonce. Beat them together into a powder.

### PULVIS DIAROMATON. Aromatic powder.

Edinb.
Take of

Nutmegs, Leffer cardamom feede, Ginger, each equal parts. Beat them together into a powder.

BOTH these compositions are agreeable, hot, spicy medicines; and as such may be usefully taken in cold phlegmatic habits and decayed constitutions, for warming the stomach, promoting digestion, and strengthening the tone of the viscera. The dose is from tea grains to a scruple and upwards. The first is considerably the warmest.

SPECIES e SCORDIO fine OPIO.

Species of scordium without opium.

Lond.

Take of

Bole armenic, or French bole, four ounces; Scordium, two ounces; Cinnamon, an ounce and a half; Storax strained, Tormentil root, Bistort root, Gentian, Dittany of Crete, Galbanum strained, Gum Arabic, Red roses each one ounce; Long pepper, Ginger, each half an ounce.

Reduce them into powder.

SPECIES

SPECIES e SCORDIO cum OPIO.

Species of scordium with opium.

Lond.

Take of strained opium, three drachms.

Dry it a little, that it may easily pulverize; and add it to the foregoing species in the beating, that they may be all reduced into a powder together.

This is the species of Fracastorius's confection or diascordium, which has been hitherto kept in the shops in the form of an electary only, but is now judiciously directed in that of a powder also, both with and without the opium: when made into an electary, the medicine, in keeping, loses of its astringency, in which consists great part of its virtue. As this composition has in common practice been looked upon as a medicine of great consequence, and its effects determined by long experience, the college have made no farther alteration in its ingredients, than substituting red roses themselves to the sugar of roses, omitting forrel feeds, which are certainly infignificant, and supplying the Lemnian earth, which with us is scarcee ver met with genuine, by a proper increase of the bole. They have nevertheless given an elegant reformation of it, in the pulvis e bolo, cum et fine opio: there, the scordium, storax, gentian, dittany, ginger, and galbanum, are rejected, as being either superfluous or contrary to the intention; whilst an increase of the tormentil root amply supplies the loss of the bistort and roses. The quantity of opium is the same in both, viz. one grain in forty-five of the composition:

> PULVIS TESTACEUS ÇERATUS.

Cerated testaceous powder. Edinb.

Melt fome yellow bees wax over a gentle fire, and carefully flir into it, by little and little, as much ofthecompound powder of crabsclaws as the wax will take up.

THIS preparation, made with oystershells, instead of the erabsclaw powder, has been in use for fome time in the Edinburgh infirmary, and is thence now received into the pharmacopæia of the college. It is given to the quantity of a drachm twice a day, in diarrhoeas and dysenteries, wherever the viscera are subject to be eroded by acrimonious humours, and in immoderate uterine discharges. Its virtue seems to depend wholly upon the wax, the earthly powder being of no farther use than to divide that concrete, and render it miscible with the animal fluids.

Pulvis Arthriticus Amarus.

Bitter gout powder.

Paris.

Take of
Gentian root,
Round birthworth root,
Rhapontic root,
Germander leaves,
Ground pine leaves,
Leffer centaury tops, of each
equal parts.
Make them into a powder.

COMPOSITIONS of this kind were in use among the ancient Greek physicians, and made a considerable part of their practice in gouty and arthritic complaints. But while they bestow great praises on them in cold and phlegmetic constitutions, they very properly condemn them as being extremely hurful in the hot and bilious. Afterwards, on account probably of the ill confequences arising from their indif-

criminate use, these medicines fell into neglect, till the introduction of the Greek volumes into the western parts of Europe, when they were transcribed by some of the earlier medical writers, and brought into some esteem in Italy, Germany, Switzerland, &c. A form differing from the above only in the omission of the rhapontic root, was some years ago brought from thence as a family receipt, by a person of high rank, who having experienced remarkable benefit from it in a hereditary gout, ordered it to be printed, and copies delivered to all who should ask for them. (See the Medical Observations and Inquiries published by a society of physicians in London, vol. i. p. 126.) The directions for using this medicine are to the following effect:

"Take one drachm of the powder every morning fasting, in a
cup of any agreeable liquor,
keeping fasting an hour and a
half after it. Continue this for
three months without interruption; then diminish the dose to
three quarters of a drachm, for
three months longer; then to
half a drachm, for fix months
more. After the first year, it
will be sufficient to take half
a drachm every other day. As

44 years before any great benefit
 45 is received. In rheumatifus
 46 that are only accidental, a few
 46 of the drachm dofes may do:

" this medicine operates insensi-

" bly, it will take perhaps two

but in habitual rheumatifms,
 and fuch as are of long fland ing, it must be taken as for

"the gout: the remedy requires patience as it operates but flow in both cases."

Dr. Clephane remarks (in the learned and judicious paper above referred to,) that this me-

dicine will probably do good in many cases; for in many cases there is reason to believe it extremely proper; but that an indiscriminate use of it will probably again do what a like abuse formerly did, bring a good medicine into disrepute.

Pulvis Catharticus salinus.

Saline cathartic powder.

Take of

Vitriolated tartar,
Crystals of tartar, each 1 drachm;
Sal prunel, or purified nitre, one scruple.
Make them into a powder.

This is an useful cathartic in inflammatory disorders, and a viscid impure state of the juices. The quantity above directed is intended for one dose, which should be accompanied with plentiful dilution.

Pulvis CARMINATIVUS.

Carminative powder.

Take of
Anifeed,
Sweet fennel feed, each two
fcruples;
Ginger, one fcruple;
Nutmegs, half a fcruple;
Fine fugar, half a drachm.
Reduce them into a powder, for
four dofes.

This powder is employed for expelling flatulencies arising from indigestion, particularly those to which hypochondriacal and hysterical persons are subject. It is likewise usefully given in the gripes of young children, either mixed with their food or otherwise.

Pulvis Diureticus.

Diuretic powder.

Take of

Sal prunel, ten grains;

Salt of amber, four grains;

Oil

Oil of turpentine, three drops; Fine fugar, one scruple. Drop the oil upon the fugar, then

add the falts, and grind the

whole together.

This powder is a very efficacious diuretic, and may be given to advantage in cases where the affiftance of fuch forcing medicines is required. The sales somewhat abate the heating quality of the oil, and at the same time cool and relax the passages.

PULVIS ROBORANS. Strengthening powder. Take of

Extractof Peruvian bank, twelve grains;

Salt of steel, two grains; Oil of cinnamon, one drop; Fine fugar, half a drachm.

Having mixed the oil with the fugar, add the other ingredients, and grind the whole well together, for two doles.

This medicine has a much better title to the appellation of a firengthener, than those usually met with under that name in dispensatories. In lax habits, debilities of the nervous system, the weaknesses peculiar to either sex, it has generally good effects.

Pulvis ad strumas. Powder against the king's evil. Take of Burnt sponge, one scruple; Nitre. Coralline. Fine fugar, each half a scruple.

Reduce them into powder.

This powder is recommended in scrophulous disorders and obstructions of the glands: it is supposed to open and deterge the mi-

nute vellels, and carry off the

offending matter by urine. Dr. Mead informs us, in his Monita medica, that he very frequently experienced its good effects: he used to give the quantity above prescribed twice a day, with three or four glasses of the less compounded lime-water along with each dose: if the patient was much emaciated, the lime-water was mixed with about an equal quantity of milk.

PULVIS TEMPERANS. Take of

Vitriolated tartar,

Purified nitre, each 3 drachms; Cinnabar, fincly levigated, two scruples.

Make them into a fubtile powder.

THESE kinds of powders are in frequent use among foreign phyficians in all disorders, accompanied with immoderate heat or exagitation of the humours. They are called also, especially when absorbents are joined, *pulveres* præcipitantes, and antispasmodici. They are given in doses of only a few grains at a time, but repeated at short intervals.

> PULVIS VERMIFUGUS. Vermifuge powder.

Take of Tanfy flowers, Worm feed, each 3 drachms: Salt of steel, one drachm. Make them into a powder.

Take of Tin reduced into fine powder, two drachms; Ethiops mineral, half a drachm; Fine fugar, one feruple. Mix them well together.

Take of Choice rhubarb, three drachms; ScamScammony,
Calomel, each one drachm.
Mix and make them into a powder.

ALL these compositions are well calculated for the purpose expressed in the title. The first is given in the hospitals, in doses of half a drachm twice a day; which quantity contains about four grains and a

half of the falt of feel. The fecond is divided into three or four doses, one of which is taken every morning, and a cathartic on the day following. The third, which is a brisk purgative, is used, in the quantity of half a drachm, after the others have been premised; or it is taken once or twice a week without their affishance.

CHAP.

#### CHAPTER II.

### Troches and lozenges.

NOCHES and lozenges are composed of powders made up with glutinous fubstances into little cakes, and afterwards dried. This form is principally made use of for the more commodious exhibition of certain medicines, by fitting them to dissolve slowly in the mouth, so as to pass by degrees into the stomach; and hence these preparations have generally a confiderable proportions of fugar or other materials grateful to the palate. Some powders have likewise been reduced into troches, with a view to their preservation; though possibly for no very good reasons: for the moistening, and afterwards drying them in the air, must in this light be of greater injury, than any advantage accruing from this form can counterbalance.

### General rules for making troches.

The three first rules laid down for making powders, are also to be observed in the powders for troches [E.]

If the mass proves so glutinous as to flick to the fingers in making up, the hands may be anointed with any convenient fweet or awith any convenient fweet or aTROCHISCI BECHICI ALBI. with powder of starch, or with that of liquorice [E.]

In order to thoroughly dry the troches, put them on an inverted fieve, in a shady, airy place, and frequently turn them [E.]

IV. Troches are to be kept in glass vessels, or in earthen ones well glazed [E.]

TROCHISCI ALBI RHASIS. feu SIEF ALBUM. The white troches, or dry collyrium of Razi. Edinb.

Take of Cerusse, three ounces: Sarcocolla, one ounce; Gum tragacanth, three drachms: Camphor, one drachm; Rose-water, as much as is suffi-

Make them into troches according to art.

THE making these ingredients into troches is an unnecessary trouble; since before they are used. they must be powdered again, for being mixed with rose water or other liquors, for the purposes of a cooling, antacrid, and moderately aftrictive collyrium, injection, &c. The London college have therefore directed them to be kept in the form of powder (under the title of pulvis e cerussa compositus) omitting the camphor, which is not found in the original of Razi.

White pelloral troches. Lond.

Take of Double-refined fugar, a pound and a half; Starch, an ounce and a half; Liquorice, fix drachms; Florence Florence orfis root, half an ounce.

Reduce these ingredients into powder, which is to be made up into troches with a proper quantity of mucilage of gum tragacanth.

#### Edinb.

Take of White fugar, in powder, one pound and a half; Compound powder of gum tragacanth three ounces; Florence orris root, one ounce;

Rose-water, & sufficient quan-

Make them into troches.

These compositions are very agreeable pectorals, and may be used at pleasure. They are calculated for foftening acrimonous humours, and allaying the tickling in the throat, which provokes coughing.

### TROCHISCI BECHICI NIGRI.

Black pettoral troches. Lond.

Take of

Extract of liquorice; Double refined fugar, each ten

Gum tragacanth, half a pound. Drop upon these ingredients so much water as will make the mass soft enough to be formed into troches.

#### Edinb.

Take of

Extract of liquorice, Gum Arabie, each four ounces; White fugar, eight ounces.

Boil the extract and gum in a sufficient quantity of water till they are dissolved: then having strained the liquor, add to it the fugar, and evaporate the mixture over a gentle fire, till it is of a proper confiftence for being formed into troches.

THESE compositions are designed for the same purposes as the white pectoral troches above described. In foreign pharmacopæias there are some other troches of this kind, under the titles of irochisci becchici slavi and rubri; the first are coloured with fassron, the latter with bole armenic. dissolving and straining the extract of liquorice and gum Arabic, as now ordered in the last of the above prescriptions, is a confiderable improvement; not only as they are by that means more uniformly mixed than they can well be by beating, but likewise as they are thereby purified from the heterogeneous matters, of which both those drugs have commonly no fmall admixture.

#### TROCHISCI de MINIO. Red lead troches. Edinb.

Take of

Red lead, half an ounce; Corrofive mercury sublimate, one ounce ;

Crumb of the finest bread, four ounces.

- Make them up with rose-water in÷ to oblong troches.

THESE troches are employed. only for external purposes as escharotics: they are powerfully fuch, and require a good deal of caution in their use,

#### TROCHISCI de MYRRHA. Troches of myrrb. Edinb.

Take of • Myrrh, one ounce and a half; Lovage seed, Pennyroyal leaves. Ruffia castor, Gal-00

1

Galbanum, each one ounce; Effential oil of favin, half a drachm;

Elixir proprietatis, as much as is sufficient.

Let the gum be fostened with the elixir into a mass of the consistence of honey; then add the oil and powders, and make the whole into troches according to art.

THESE troches are very well contrived in regard to efficacy, and superior to those in most other pharmacopœias, under the same title. Madder and cummin feed, two of their former ingredients, which were objected to in the preceding edition of this work, are now expunged; the one as being an unnecessary article; the other as being an offensive one, and not of similar intention with the rest. In the place of this last, lovage feed is introduced, which is doubt less more agreeable to the intention of the medicine. Asafoctida is supplied by an increase of the galbanum; and the essential oil of rue, by an increase of the oil of There seems to be no ocfavin. casion for making a medicine of this kind into troches, as it cannot be conveniently taken in that form: . the London college have therefore exchanged their TROCHISCI e Inyrrha for a Pulvis e myrrha compositus, which see.

## TROCHISCI e NITRO. Troches of nitre. Lond.

Take of

Nitre purified, four ounces; Double-refined sugar, one pound. Make them into troches with mucilage of gum tragacanth.

This is a very agreeable form for the exhibition of nitre; though,

when the falt is thus taken without any liquid (if the quantity is confiderable) it is apt to occasion uneasing about the stomach, which can only be prevented by large dilution with aqueous liquors.

## TROCHISCI e SCILLA. Troches of Squills. Lond.

Take of

Baked squills, half a pound;
Wheat flower, four ounces.
Beat them together, and form the
mass into troches, which are to
be dried with a gentle heat.

This preparation is used only as an ingredient in the theriaca. The design of baking the squill is, to abate its acrimony; and making it afterwards into troches seems the most convenient way of drying it: common wheat slower is as sit for this purpose as any, though that of the white vesch has been generally directed.

## TROCHISCI e SULPHURE. Troches of fulphur. Lond.

Take of

Flowers of fulphur, washed, two ounces;

Double - refined fugar, four ounces.

Beat them together, and adding fome mucilage of quince feeds, form them into troches.

### TROCHISCI DIASULPHU-RIS.

Troches of Sulpbur. Edinb.

Take of

Flowers of sulphur, 1 ounce; Flowers of benzoine, 1 drachm; White sugar, 3 ounces; Mucilage of gum tragacanth, as much as is sufficient.

Mix

three

Mix and make them into troches, according to art.

THESE compositions are to be confidered only as agreeable forms for the exhibition of fulphur, no alteration or addition being here made to its virtue; unless that by the flowers of benzoine in the fecond prescription, the medicine is supposed to be renderd more efficacious as a pectoral.

#### TROCHISCI e TERRA JAPONICA. Troches of Japan earth. Lond.

Take of Japan earth,

Gum Arabic, each two ounces; Sugar of roses, sixteen ounces. Beat them together, and dropping in some water, make them into troches.

Edinb.

Take of

Japan earth, two ounces; White sugar, one pound; Rose-water, a sufficient quantity.

Make them into troches.

A preparation of this kind, with the addition of ambergris and musk, which are here more prudently omitted, has long been in some esteem as a mild restringent, &c. under the title of CATECHU. Medicines of this class in general are excellently fitted for the form of troches; for when flowly and gradually received into the stomach, as this form occasions them to be, they produce much better effects, than if an equal quantity was taken down at once. The above troches are sufficiently palatable, and of confiderable fervice in some kinds of coughs, thin acrid defluxions, diarrhœas, &c. TABELLA CARDIALGICA. Cardialgic lozenges. Lond.

Take of

Chalk prepared, four ounces; Crabs-claws prepared, ounces: Bole armenic, or French bole, half an ounce: Nutmegs, one scruple; Double-refined fugar,

Reduce these ingredients into powder, and make them into troches with water.

#### TROCHISCI CARDIALGICI. Edinb.

Take of

Oystershells prepared, White chalk, powdered, each two ounces: Gum Arabic, half an ounce: Nutmegs, half a drachm; White fugar, fix ounces; Common water, a sufficient quantity.

Gum tragacanth, half an ounce; Make them into troches according to art.

> These compositions are calculated against that uneasy sensation at stomach improperly called the heartburn; in which they oftentimes give immediate relief, by abforbing and neutralizing the acidjuices that occasion this disorder. The absorbent powders here made use of, are of the most powerful kind, though there does not feem to be any occasion for using more than one of them. Some have prescribed the following formula.

### TABELLÆ ANTACIDÆ. Antacid lozenges.

Take of Prepared white chalk, four drachms; Candied ginger, three drachms; Cinnamon, one drachm; Fine 002

Fine fogar, diffolved in water, as much as is sufficient to reduce the whole into a due confishence for being formed into lozenges.

HERE it may be observed, that all these compositions, though very effectual for the intention, are accompanied with an inconvenience, which is frequently complained of in their use; their binding the belly. The use of the chalk, oystershells, and crabs-claws, is to absorb acidities; and both these and the other common absorbents, united with acids, compose therewith astringent concretes. The following composition is free from this inconvenience.

### TABELLÆ ANTACIDÆ LAXANTES. ·

Laxative antacid lozenges.

Take of

Magnefia alba, fix ounces;
Double-refined fugar, three
ounces;

Nutmegs, one fcruple.

Mix them well together, and form them into troches with mucilage of gum tragacanth.

## SACCHARUM ROSACEUM. Sugar of rojes. Lond.

Take of

Red rose buds, freed from the heels, and hastily dried, one ounce:

Double-refined sugar, one pound. Reduce them separately into powder, then mix, and moisten them with water, that they may be formed into troches, which are to be dried by a gentle heat.

In the Edinburgh pharmacopoeia, this preparation is directed as follows. TABELLÆ ROSACEÆ.

Rose tablets.

Edinb.

Take of

Conserve of red roses, four ounces;

White fugar, in powder, one pound.

If any moisture is required, take of syrup of dry roses a sufficient quantity for forming them into troches, which are to be dried, with a gentle heat.

THE fugar of roses was formerly made, by boiling a pound of fine fugar with four ounces of the juice of red roses, over a gentle fire, till the juice was almost all evaporated; then throwing in an ounce of dry red roses reduced to a very fine powder; after which the matter was poured out upon a marble, and formed into lezenges. The two methods above directed. are more simple and commodious: though, if any virtue be expected from the roses, the medicine is not at all improved by the alteration. As the conserve contains only one fourth its weight of roles, in a fresh state, it is obvious that the quantity of fresh roses in the fecond prescription is less than that of the dry ones in the first.

These preparations are chiefly valued for their agreeableness to the eye and palate. Some likewise esteem them, medicinally, as light restringents; and look upon them, not undeservedly, as an excellent addition to milk in phthiscal and hestic cases. Some have been accustomed to add a portion of acid in making these preparations: this improves the colour, but renders them unsit to be taken with milk.

T.ABELLE ANTHELMINTICE.

Anibelminic fugar-cakes
Take of
1.

Powdered

Powdered tin, half a drachm; Fine fugar, half an ounce; Rose-water, a sufficient quantity to make them into a mass for tablets.

Take of 2.
Scammony,
Mercurius dulcis, each four grains;
Fine sugar, half an ounce;
Rose water, a sufficient quantity to make them into tablets.

THESE composititions are calculated for children, who are not easily prevailed on to take anthelamintic medicines in less agreeable forms. If the first is made use of, it must be repeated three or four mornings successively, after which a purge is to be taken: the second, if it requires repetition, is to be given only every other morning. The proportions of the ingredients are to be varied, according to the age and strength of the patient.

TROCHISCI NERVINI. Nerve troches.

Take of
Compound spirit of lavender,
sixty drops;
Oil of cinnamon,
Oil of rosemary, each fourdrops;
Florence orris root, two drachms;
Fine sugar, one ounce;
Mucilage of gum tragacanth,
as much as will reduce them
into a mass, which is to be

formed into troches of about

ONE or two of these troches, taken occasionally, and suffered to dissolve in the mouth, prove serviceable to those who are subject to paralytic and other nervous disorders. Warm aromatic medicines, given in this form and manner, are supposed, from their

half a scruple each.

flow diffolution in the mouth, to affect the nervous system more immediately than if received at once into the stomach.

Morsuli purgantes.

Purging tablets.

Take of

Crystals of tartar, half an ounce; Scammony, three drachms; Oil of cinnamon, four drops; Double-refined fugar, eight ounces.

Make them up, with rose water, into troches, weighing each about a drachm.

This is a sufficiently elegant form for purgative troches. Each of the morfuli contains two grains and a half of scammony.

Morsuli de Rhabarbaro.

Rhubarb troches.

Take of
Creme of tartar,
Rhubarb, each two drachms;
Fresh lemon peel, half a drachm;
Fine sugar, sour ounces.
Make them into troches with rosewater.

Two drachms of these troches contain about seven grains of rhubarb, and as much creme of tartar. Both this and the preceding composition are among the officinals of the Brandenburgh pharmacoposia.

MORSULI RESTAURANTES

KUNCKELII.

Kunckel's antimonial tablets.

Take of

The best Hungarian antimony, levigated into an impalpable powder, three drachms and a half:

Sweet almonds peeled, Fresh pine nuts, each half an ounce;

O o 3

Cinna-

Cinnamon, one drachm; Leffer cardamom feeds, hufked, half a drachm;

Double-refined fugar, four ounces.

Diffolve the fugar in equal quantities of cinnamon water and rose-water; then mix therewith the other ingredients, and form the whole into tablets weighing one drachm each.

THESE tablets were brought into esteem by Kunckel, at a time when the internal use of crude antimony was almost universally reckoned poisonous. He had recourse to them as a desperate medicine, in violent pains and contractions of the arms, after all the common methods of cure had been used without any relief; and being happily, in a short time, perfectly freed from his complaints, he made trial of them in several other cases, with remarkable success. He seems to have begun with doses of four or five grains (that is, one of the tablets above prescribed) which were repeated thrice a day, and gradually increased to a drachm or more of the antimony every day.

TROCHISCI SIALAGOGI.
Sialagogue troches.

Take of
Pellitory of Spain, half an ounce;
Mastich, two drachms;
Oil of cloves and marjoram,
each one drachm;

Yellow wax, a fufficient quantity.

Make them into troches or pellets.

ONE of these troches is to be occasionally held in the mouth, and chewed, to promote a discharge of saliva; which they effect by warming and simulating the salival glands.

TROCHISCI STOMACHICI.

Stomachic troches.

Take of

Hard extract of Peruvian bark,

one drachm:

one drachm;
Oil of cinnamon,
Oil of mint, each ten drops;
Fine fugar, four ounces.
Make them into troches, with
mucilage of gum tragacanth.

THESE troches are of fervice for warming and strengthening the stomach, expelling statulencies, and promoting digestion: for these purposes they are as effectual as any thing that can be well contrived in this form.

TROCHISCI SUAVEOLENTES.

Sweet-fmelling troches.

Take of
Strained florax, one scruple;
Ambergris, fifteen grains;
Musk, seven grains;
Oil of cinnamon, six drops;
Fine sugar, one onnce.

Make them into small troches with
mucilage of gum Arabic.

### CHAPTER III.

### Pills.

adapted those drugs which operate in a small dose; and whose nauseous and offensive taste or smell require them to be concealed from

the palace.

Pills dissolve the most dissicultly in the stomach, and produce the most gradual and lasting effects of all the internal forms. This is in fome cases of great advantage; in others it is a quality not at all defirable, and sometimes may even be of dangerous confequence, particularly with regard to emetics, which if they pass the stomach undiffolved, and afterwards exert themselves in the intestines, operate there as violent cathartics. Hence emetics are, among us, fcarce ever given in pills. hence to the refinous and difficultly foluble substances, saponaceous ones ought to be added, in order to promote their folution.

Gummy refins and inspissated juices are sometimes soft enough to be made into pills, without addition: where any moisture is requisite, spirit of wine is more proper than syrups or conserves, as it unites more readily with them, and does not sensibly increase their bulk. Light, dry powders require syrup, or mucilages; and the more ponderous, as the mercurial and other metallic preparations, thick honey, con-

ferve or extracts.

Light powders require about half their weight of fyrup; of honey, about three-fourths their weight, to reduce them into a due confistence for forming pills. Half a drachm of the mass will make sive or six pills of a moderate size.

General rules for making pills, from the Edinburgh pharmacoparia.

The three first rules, formerly laid down for making powders are here also to be carefully observed.

Ħ

Gums and inspissated juices are to be first sestened with the liquid prescribed: then add the powders, and continue beating them all together till they are persectly mixed.

III.

The masses for pills are best kept in bladders, which should be moissened, now and then, with some of the same kind of liquid that the mass was made up with, or with some proper aromatic oil.

## PILULÆ AROMATICÆ Aromatic pills. Lond

Take of

Socotorine aloes, an ounce and a half

Gum guaiacum, one onnce; Aromatic species,

Balsam of Peru, each half an ounce.

Reduce the aloes and gum guaiacum feparately into powder, then mix them with the reft, and make the whole into a mass with a sufficient quantity of syrup of orange peel,

Ιτ

IT is somewhat difficult to unite these ingredients into a mass fit for making pills of. The best way is, to first rub the aromatic species with the balsam, then to add the powdered aloes, and afterwards the guaiacum; when thefe are well mixed together, drop in the fyrup by little and little at a These pills are contrived to supply the place of the PILULE DIAMBRÆ of our former pharmacopæia. They are far more elegant as well as simple, truly uniform in their ingredients, and excellently adapted to the purposes they seem designed for. Taken in small doses, as half a scruple, or little more, and occasionally repeated, they warm the stomach, and by degrees the whole habit, promote perspira-tion, and all the natural secretions. If the dose is considerable, they operate gently by stool: and if continued for some time in fmaller doses, they prove at length purgative, or introduce a falutary loofeness.

## PILULÆ ALOETICÆ. Albeite pills. Edinb.

Take of
Socotorine aloes in powder,
Caffile foap, each equal parts;
Thin honey, as much as is fufficient.
Make them into a mass.

This composition has been in use for some time in the Edinhurgh infirmary, as a deobstruent in cachectic indispositions; and from thence it is received into the pharmacopæia of the college. A scruple or half a dachm of the mass is directed to be made into pills of a moderate size for one cost.

### PILULÆ de JALAPPA. Jalap pills. Edinb.

Take of
Extract of jalap, two ounces;
Socotorine aloes, one ounce;
Vitriolated tartar, one drachm;
Syrup of ginger, a fufficient
quantity.

Beat them into a mais.

This composition also is now first received into the pharmacopocia. One of the same kind, with powdered jalap in substance instead of the extract, is used in some of our hospitals, as a cheap and effectual purge.

### PILULE E SCAMMONIO CHM

Pills of scammony with aloes.

Take of
Socotorine aloes, one drachm;
Aromatic species, half a drachm;
Scammony, one scruple;
Soft extract of liquorice, asmuch as is sufficient to reduce them into a mass of a due consistence for being formed into pills.

This warm purgative is recommended for removing the crudities, &c. after a furfeit or debauch, and for preventing arthritic and other complaints incident to those who live high. The quantity above described may be made into thirty pills, of which five or fix are to be taken for a dose.

### PILULÆ ex COLOCYNTHIDE SIMPLICIORES.

The more simple colocynub pills.

Lond.

Take of
Pith of colocynth,
Scammony, each two ounces;
Oil of cloves, two drachms.
Pulverize

Pulverize the coloquintida and fcammony by themselves, then mix in the oil, and make the whole into a mass with syrup of buckthorn.

THE operator should be careful, in pulverizing the colocynth, to avoid the finer particles that fly off from it; which, though they do not confiderably affect the mouth or fauces, have sometimes been observed to occasion violent purg-The drug should first be well dried, cut with a sheers into fmall pieces, and freed from the feeds; then rub it in an oiled mortar, adding a few drops of sweet oil occasionally during the trituration: afterwards mix this powder with the powdered scammony, add the effential oil prefcribed, and make the mixture into a mais, as above directed. The composition is apt to grow fliff and dry in keeping, and therefore ought to be made pretty foft at first: the pills should be formed as they are wanted; for when long kept, they become fo hard, as to have fometimes passed through the intestines undissolved.

These pills (formerly called PILULE DE DUOBUS, or pills of two ingredients) are very strong cathartics, and ought not to be ventured upon in cases where less violent medicines will take effect. They have been often made use of in large dofes, along with large doles also of mercurials, in venereal complaints, both in recent gonorrhoea, and in the swellings and inflammations which fometimes follow from the suppression of the discharge; but in both thefe cases they are apparently improper, as they generally injure the conftitution, and as the latter complaint is for the most part aggravated by them. The effential oil, which is added as a corrector to the purgative ingredients, does not contribute so much, as is commonly supposed, to abate the roughness of their operation. See page 359. The dose of these puls is from fifteen grains to half a drachm; some have imprudently gone as far as two scruples. Half a drachm contains ten grains of coloquintida, and as much scammony.

PILULÆ COCCIÆ.
The pills called cochiæ.
Edinb.

Take of
Coloquintida,
Scammony,
Socotorine aloes, each one
ounce;
Vitriolated tartar, two drachms;
Oil of cloves, one dracnm;
Syrup of buckthorn, a lufficient quantity.
Beat them into a mass.

This composition, like the foregoing, is strongly cathartic; not less effectual, though somewhat less irritating. Half a drachm contains above six grains and a half of coloquintida, the same quantity of scammony, and the same of aloes.

# PILULÆ ex COLOCYNTHIDE cum ALOE. Colocynth pills with aloes. Lond.

Take of
Socotorine aloes,
Scammony, each two ounces;
Pith of colocynth, one ounce;
Oil of cloves, two drachms.
Tet the dry feering he forested

Let the dry species be separately reduced into powder; then mix in the oil, and make the whole into a mass with syrup of buckthorn. By the diminution of coloquintida in this prescription, the ingredients are reduced to the proportions wherein they are set down in the original of Galeu; and what is of greater consequence, the medicine becomes less ungrateful to the stomach, and less virulent in its operation. Half a drachm of the mass contains nearly four grains of coloquintida, eight of aloes, and eight of scammony.

### PILULÆ ECPHR ACTICÆ, Deobstruent pills.

Lond.

Take of the
Aromatic pills, three ounces;
Rhubarb,
Extract of gentian,
Salt of steel, each one ounce;
Salt of wormwood, half an ounce.

Beat them together into a mass, with solutive syrup of roses.

IT is difficult to bring this mase into the due confistence, the two falts a ting upon one another, fo as to make it swell and crumble. Notwithstanding the alkaline salt employed, the pill does not prove at all alkaline; for the acid of the falt of steel for sakes its metal, and unites with the alkali, into a vitriolated tartar; whence some have proposed using, instead of the two falts here directed, an ounce of vitri lated tartar already made, and half an ounce of any of the calces of iron: this they observe prevents the inconveniency above m nt oned, without making any apparent alteration in the quality of the medicine.

PILUI.Æ ECPHRACTICÆ CHALYBEATE. Chalyheate ecpbractic pills. Edinb.

Take of
The mass of common pills,
called Rufus's pills, described hereafter, one ounce and
a half;

Gum ammoniacum, Refin of guaiacum, each half an ounce;

Salt of steel, five drachms; Syrup of orange peel, as much as is sufficient to reduce the whole into a mass.

THE falt of fteel, which is one of the most active preparations of that metal, remains here undecompounded. Both these and the foregoing pills are very well calculated for answering the intention expressed in the title. A drachm of the mass may be made into twelve pills, and two or three of these taken every night, or oftener, in chlorotic, or other cases, where warm, aperient, or deobstruent medicines are proper.

### PILULÆ ECPHRACTICÆ PURGANTES.

Purging deabstruent pills.
Edinb,

Take of
Socotorine aloes,
Extract of black hellebore,
Scammony, each one ounce;
Gum ammoniacum,
Refin of guaiacum, each half an
ounce;
Viriolated taxtes, two drachmon

Vitriolated tartar, two drachms pr Effential oil of juniper berries, one drachm.

Beat them into a mass, with a fufficient quantity of syrup of buckthorn.

This composition may be given from eight or ten grains to a scruple or half a drachm, according as it is intended to keep the belly open or to purge. Half a drachm

a drachm of the mais contains about fix grains of each of the capital purgative ingredients; aloes, fcammony, and extract of hellebore.

## PILULÆ FŒTIDÆ. Fætid pills. Edinb.

Take of
Asafoetida,
Russia castor, each one drachm
and a half;
Camphor, half a drachm;
Oil of hartshorn, twenty-four

Beat the camphor with the asafcetida, then add the castor and oil of hartshorn, and make the whole

into a mass.

### PILULÆ GUMMOSÆ. Gum pills.

Lond.

Take of
Galbanum,
Opoponax,
Myrrh,
Sagapenum, each one ounce;
Afafætida, half an ounce.
Make them into a mass with syrup
of saffron.

Edinb.

Take of
Gum ammoniscum, one ounce;
Ruffia caftor,
Myrrh,
Afafætida, each half an ounce;
Oil of amber, half a drachm;
Syrup of orange peel, as much
as is sufficient.

Beat them into a mass.

ALL these pills are designed for antihysterics and emmenagogues, and very well caculated for answering those intentions: half a struple, a scruple, or more, may be taken every night or oftener. The sected pills of our former pharmacopæia were considerably

purgative: the purgative ingredients are now omitted, as the physician may easily, in extemporaneous prescription, compound these pills with cathartic medicines, in such proportions as particular cases shall require.

The following compositions are calculated for the same intentions as the foregoing deobstruent, sould,

and gum pills.

Take of

Afafætida,

Wood-foot,

Myrrh, each two ounces;

Oil of amber, one drachm and

a half;

Syrup of fugar, a fufficient
quantity.

Mix, and make them into a mass,
according to art.

Take of 2.
Asasætida, one drachm;
Martial flowers, half a drachm;
Oil of amber, eight drops;
Balfam of Peru, a sufficient quantity to reduce them into a mass.

Take of

Asafontida,

Gum ammoniacum,

Myrrh,

Aloes,

Rust of steel prepared,

Extract of gentian, each oneforuple;

Syrup of ginger, as much as
will make the other ingredients
into a mass.

Take of
Galbanum, one drachm;
Salt of steel, half a drachm;
Asascetida,
Aromatic species, each one scruple;
Tincture of myrrh, as much as will make them into a mass.

In

hysterical disorders, after bleeding and purging, where a fanguine and plethoric habit indicates these evacuations, chalybeate : medicines are in general the most to be relied upon; especially when joined, as in these compositions, with bitters and deobstruent gums. At first taking, they are apt to increase the complaints, (as the experienced Sydenham observes) and occasion great disorders both of body and mind; which, however, foon go off, or may be relieved by a proper dose of opium given at bed-time. A drachm of either of the masses is to be made into twelve pills, one or two of which may be taken for a dose, twice or thrice a day.

### PILULÆ MERCURIALES.

Mercurial pills.

Edinb.

Take of
Purified quickfilver,
Refin guaiacum,
Cattile foap, each one ounce.
Grind them together in a glass
mortar, till the mercurial glohules cease to appear; then add
a sufficient quantity of common
syrup, and make the whole into
a mass according to art.

#### Lond.

Take of Quickfilver, five drachms; Strafburgh turpentine, two drachms; Cathartic extract, four scruples; Rhubarb powdered, one drachm. Grind the quickfilver with the turpentine, until they are perfectly incorporated; then let the other ingredients be beat up with this maxture into a mass. It the turpentine happens to be too thick, foften it with a little oil olive.

### PILULÆ MERCURIALES LAXANTES.

Laxative mercurial pills.

Take of
Pure quickfilver, one ounce;
Refin of guaiacum,
Extract of black hellebore,
Powdered Rhubarb, each half an
ounce;

Common syrup, a sufficient quantity.

Grind the quickfilver with the refin of guaiacum, until they are perfectly incorporated; then add the other ingredients, and beat the whole into a mass according to art.

THE three foregoing compositions are useful mercurial pills; the first as an alterative, the other two as purgative mercurials. They are all, however, liable to an inconvenience, uncertainty in regard to their strength; for the mercury is but loofely united with the other ingredients, and very apt to separate and run together in its original form, in which state it never exerts its proper virtue; though it appears perfectly extinguished by the matters it is ground with at first, part of it is apt to be spued out on beating up the mixture with the other ingredients into a mass.

#### PILULÆ de GAMBOGIA, Gamboye pills. Edinb.

Take of
Socotorine aloes,
Extract of black hellebore,
Gamboge,
Mercurius dulcis, each two
drachms;
Effential oil of juniper berries,
half a drachm;
Syrup of buckthorn, a fufficient
quantity.
Beat them into a mass.

THIS

This is a strong mercurial purgative, in which the mercurial preparation is not liable to the uncertainty which the crude quickfilver is accompanied with in the foregoing compositions. The dose is from ten or fifteen grains to half a drachm. This last quantity contains of aloes, extract of hellobore, gamboge, and mercurius dulcis, about five grains of each.

## PILULÆ ÆTHIOPICÆ. Ethiopic pills. Edinb.

Take of
Pure quickfilver, fix drachms;
Golden fulphur of antimony,
Refin of guaiacum,

Spanish soap, each half an ounce.

Grind the quickfilver, with the refin and foap, in a glass mortar, until the mercurial globules entirely disappear; then add the golden sulphur, with as much common syrup as is sufficient to make the mixture into a mass of the proper confistence for forming pills.

THESE pills are much more efficacious than those of the preced ing edition; the ethiops mineral, -there ordered, being exchanged for a more active composition. In their present form, they resemble Dr. Plummer's pills, described in the Edinburgh essays, (see page 534 of this work) to which they are preferable in one respect, that they are less apt to run off by stool. They are an useful alterative both in cutaneous and venereal disorders. Onefourth part of the quantity above · prescribed may be made into fixty pills; of which, from one to four may be taken every night and morning, the patient keeping moderately warm during the

whole time that this course is continued.

I shall here insert some other formulæ of mercurial pills, which may be occasionally had recourse to, and of which the greater part have been kept as secrets in particular hands.

Take of
Crude quickfilver,
Hard extract of guaiacum, each
one drachm and a half;
Effential oil of faffafras, twenty
drops;
Venice turpentine, a fufficient

quantity.

Grind the quickfilver with the turpentine, till they are perfectly incorporated: then add the other ingredients, and reduce the whole into an uniform mass; which is to be made into forty pills. Two, three, or more of these, may

be taken for a dose.

Take of 2.

Mercurius dulcis,
Prepared chalk, each one scruple;
Mucilage of gum Arabic, a
fufficient quantity.

Make them into twelve pills, of
which the dose is from one to
three.

Take of

Mercurius dulcis, half a scruple;

Softer extract of guaiacum, one drachm;

Essential oil of sassafras, ten drops.

Mix, and make them into a mass, for twenty pills; the dose of which is from one to six.

Take of 4.

Mercurius dulcis, half a scruple;
Camphor, half a drachm;
Soft extract of guaiacum, as
much as is sufficient to make

121

 $\mathbf{r}$ 

6 3

Ŋ,

5

:::

C,

٠:

٠.

them into a mass, which is to be formed into twenty pills: the dole is from one to fix.

Take of Mercurius dulcis, half a scruple; Venice turpentine, as much as will roduce it into a mass of a proper confistence; which is to be formed into five pills, for as many doses.

Take of Calcined mercury, commonly called præcipitate per se, Thebaic extract, each two grains: Balsam of Peru, as much as will make them into a mass: which is to be formed into two pills, for two doses,

Take of Turbith mineral, two scruples: Thebaic extract, one scruple; Mucilage of gum Arabic, as much as is sufficient to reduce them into a mass, which is to be formed into twenty pills, for as many doses.

The mercurius corallinus may be made into pills in the same manner, and taken in the same dose.

Take of 8. Mercurius dulcis, half a scruple; Crude antimony, finely levigated, one drachm;

Conferve of orange peel, as much as will reduce them into a mass.

This is to be formed into ten pills; of which the dole is from one to three.

Take of Mercurius dulcis. Precipitated fulphur of antimony, each five grains; Socotorine aloes, lifteen grains; Balfamic fyrup, a sufficient quantity to reduce them into a mass; which is to be made into five pills, for as many dofes.

THE method of managing the above mercurial medicines, as alteratives, is, to give small doses every morning and evening; and rather prolong the time of continuing their use, than increase the dose. The patient ought to keep warm, and drink of warm diaphoretic liquors, as infusion of fassafras, decoction of the woods, the fimple or compound lime water, &c.

#### PILULÆ PACIFICÆ.

The pacific pills, commonly called Mathews's pills.

Edinb.

Take of Gum ammoniacum. three Russia castor, two ounces; English saffron, Opium, each one ounce; Common syrup, as much as is sufficient. Mix, and make them into a mass, according to art.

THESE pills are contrived by a chemical empiric, Starkey, and communicated by him to Mathews, under whose name they were some time ago greatly celebrated. The form here given differs from the original in omitting a small portion of black hellebore, an ingredient of no great service; for though this article, as the London committee observes, might perhaps promote a stool the day after the medicine is taken, that advantage, in cases which require it, may with greater certainty be obtained by more obvious means.

Soap of tartar, as it is called, another ingredient in the original, on which the contriver of the composition laid no small stress. and which the Edinburgh college retained in their former edicions, is here, without any injury to the medicine, exchanged for an equal quantity of ammoniacum. Nor indeed are any of the ingredients of much consequence, except the opium; their quantity being too inconsiderable to answer any useful purpole. Eight grains of the composition contain nearly one of opium.

### PILULÆ SAPONACEÆ.

Saponaceous pills. Lond.

Take of

Almond foap, four ounces; Strained opium, half an ounce; Effence of lemons, one drachm. Soften the opium with a little wine, and then beat it with the rest, until they are perfectly mixed.

This pill is introduced in the room of Mathews's. The foap promotes the folution of the opium in the stomach, and thus occafions it to act the more quickly; which is the only intention that the more laborious foap of tartar can answer. The essence of lemons gives an agreeable flavour, makes the medicine fit eather on the ilomach, and prevents a naufea, which it would otherwise be apt to occasion. Ten grains of the pill contain nearly one grain of opium.

> PILULÆ e STYRACE. Storax pills. Lond.

Take of Strained storax, two ounces;

Saffron, one ounce: Strained opium, five drachms. Beat them together till perfectly united.

THESE are contrived for diffolving more flowly in the stomach than the faponaceous or Mathews's pills, and confequently producing more gradual and latting effects. One grain of opium is contained in five grains and four fitchs of a grain of the mass.

#### PILULÆ ex OLIBANO. Olibanum pills. Edinb.

Take of Olibanum, two ounces; Myrrh, one ounce; Opium, five drachms: Balfam of Peru, two drachms : Common syrup, a sufficient quantity.

Make them into a mass, which supplies the place of the storax pills.

### PILULÆ PECTORALES.

Pettoral pilis Edinb.

Take of

Gum ammoniacum, half an Balfam of Tolu, two drachms: Flowers of benzoine. English saffron, each 1 drachm :

Common fyrup, a fufficient quantity.

Make them into a mass according to art.

This composition is very well contrived for promoting expectoration, and may be ulcfully given in common colds, and in difficulty of breathing proceeding from viscid phlegm: the dose is from fix or eight grains to a teruple or more. It is here confiderably improved

::

Ç

t

: ;

7

3

11

proved from the last edition: the balfam of Tolu is introduced in the room of myrrh, the flowers of benzoine for benzoine in substance, and anisated balsam of sulphur, which encumbered the old form, is omitted. Here it may be observed, that though several compositions are denominated pectorals, they are nevertheless, in virtue, very distimilar. Thus, the pectoral decoction, the fyrup, and the troches, are calculated for foftening, lubricating, and incrassating thin tickling humours; whilst the pectoral pills, the clixir and the oxymel, tend to stimulate and deterge the vessels, and attenuate or dissolve thick, tenacious juices.

PILULÆ RUFI.
Rufus's pills.
Lond.

Take of
Socotorine aloes, two ounces;
Myrrh,
Saffron, each one ounce.
Make them into a mass with syrup of saffron.

### PILULÆ COMMUNES, vulgò RUFI.

The common pills, vulgarly called Rufus's pills. Edinb.

Take of
Socotorine aloes, two ounces;
Myrrh, one ounce;
Saffron, half an ounce.
Beat them into a mass with a proper quantity of syrup of orange peel.

THESE pills have long continued in practice, without any other alteration than in the fyrup which the mass is made up with, and in the proportion of saffron. In our last pharmacopæia, the syrup of wormwood was ordered, which is here judiciously exchanged for that

of faffron, this preferring and improving the brightness of colour in the medicine, which is usually looked upon as the characteristic of its goodness. The saffron, in the composition which is attributed to Rufus, is equal in quantity to the myrrh; and in these proportions the pill was received in our first pharmacopæia. As the diminution afterwards made in the faffron was grounded on very absurd reasons, (viz. " lest the " former quantity should occasion "a spasmus cynicus,") the London college have now again increased it, and restored the pill to its original form. The virtues of this medicine may be easily understood from its ingredients See Elixir proprietatis, pages 292 and 323. The pills, given to the quantity of half a drachm or two scruples, prove considerably cathartic; but they answer much beiter purposes, in smaller doses, as laxatives or alteratives.

## PILULÆ STOMACHICÆ, Stomachic pills, Edinb.

Take of
Rhubarb, one ounce;
Socotorine aloes, fix drachms;
Myrth, half an ounce;
Vitriolated tartar, one drachm;
Effential oil of mint, half adrachm;
Syrup of orange peel, a fufficient quantity.
Make them into a mass.

This pill is intended for moderately warming and trengthening the flomach, and evacuating crude viscid humours. A scruple of the mass may be taken twice a day.

### PILULÆ SCILLIT!CÆ. Squill pills.

Take of Spanish foap, one ounce;

Gum

Gum ammoniacum, Millepedes prepared, Fresh squills, each half an Balfam of Copaiba, as much as is sufficient. Reduce them into a mass according to art.

This is an elegant and commodious form for the exhibition of squills, whether for promoting expectoration, or in the other intentions to which that medicine As the virtue of the is applied. compound is chiefly from the squills, the other ingredients are often varied in extemporaneous prescription: the soap is commonly omitted, as being of no great use in the quantity that goes to a dose of the composition; and other powders, as the leifer cardamom seeds, are substituted to the millepedes, whose virtues, in such small doses, are very infignificant. In any of these forms. if the squills are fresh and juicy, there is no need of balfam; but as the mass soon dries and hardens, it must be formed immediately into pills.

Agreeably to the above remarks in the former edition of this work, the college of Edinburgh has now given the following improvement of this composition.

Take of

Gum ammoniacum, Cardamom feeds, in powder, Fresh squills, each half an ounce.

Beat the squills and ammoniacum together in a marble mortar, then add the cardamom feeds, and make the whole into a mass. If the compound should prove too fliff, fosten it with a little balfam of Copaiba.

PILULE AD DYSENTERIAM. Pills against the dysentery. Take of Yellow wax, half an ounce; Spermaceti. Japan earth, each one drachm; Oil of cinnamon, twelve drops.

Make them into a mass.

This medicine has often been of great benefit for the purpose expressed in its title, at the same time firengthening the intestines, and covering them with a foft mucus, which defends them from being irritated by the acrimony of the humours. Each half drachm of the mass may be formed into five or fix pills for one or two doses.

> PILULE PICER. Tar pills.

Take any quantity of tar, and mix with it as much powdered elecampane root as will reduce it to a proper thickness for being formed into pills.

THE powder here mixed with the tar, though of no great virtue, is nevertheless a very useful addition, not only for procuring it a due consistence for taking, but likewise as it divides the resinous texture of the tar, and thus contributes to promote its folution by the animal juices. In the Edinburgh infirmary, half a drachm of the mass, made into middle-sized pills, is given every morning and evening, in disorders of the breast, scurvies. &c.

> PILULÆ ROBORANTES. Strengthening pills.

Take of Hard extract of Peruvian bark, one drachm; Salt of steel, ten grains; Oil of cinnamon, five drops; Balfam Balfam of Peru, as much as will reduce them into a mass.

Z.
Take of

Olibanum, one drachm;
Styptic powder, two scruples;
Salt of steel, one scruple;
Syrup of sugar, a sufficient
quantity to make them into a
mass.

In a lax state of the sibres, debilities of the nervous system, and some decays of constitution, the sirst of these compositions is an effectual strengthener and referrited is made into twenty pills, one or two of these may be taken for a dose, and repeated twice a day. The other is a stronger styptic, and is used for restraining immoderate alvine evacuations, and sanguineous or serous discharges from remoter parts.

3.

Take of
Aromatic species,
Extract of gentian, each one

drachm; Extract of Peruvian bark, half

a drachm; Elixir of aloes, as much as will reduce them into a mass. THESE pills are ferviceable for warming and strengthening a weak cold stomach, expelling statulencies, and promoting digestion. If ten pills are made out of a drachm of the mass, two may be taken thrice a day, about an hour before meals.

Pilulæ e spermate ceti.

Spermaceti pills.

Take of

Spermaceti, one drachm; White fugar-candy in powder, two drachms;

Balfamic fyrup, as much as is fufficient.

Grind the spermaceti with the sugar, till they are persectly mixed; then adding the syrup, rub them with a warm pestle into an uniform mass.

WHERE spermaceti cannot be commodiously exhibited in any other form, three or four moderate-fized pills, made from this mass, may be taken two or three times a day, in erosions of the viscera by acrimonious humours, tickling coughs, and other like disorders.

### [ 579 ]

### CHAPTER IV.

### Boluses.

DOLUSES differ little in confiftence from electaries, being only fomewhat fiffer, so as to retain their figure without spreading

or falling flat.

This form is very convenient for the exhibition of the more powerful medicines, which require their dose to be exactly adjusted, as the stronger alexipharmacs, cathartics, and opiates. As boluses are chiefly intended for immediate use, volatile salts, and other materials, which, if the mass was to be kept, would exhale or swell it, are frequently admitted into them.

The quantity of a bolus very feldom exceeds a drachm: if the ingredients are of the lighter kind, even this will be too bulky to be commodiously swallowed down.

The lighter powders are made up with fyrup; a scruple or twentyfix grains of the powder, with as much syrup as will bring it to a due consistence, makes a bolus suf-

ficiently large.

The more ponderous powders, as the mercurial ones, are commonly made up with conferve: fyrups fearce holding them together. For the testaceous powders also an addition of conserve is used; though if made up with this alone, they would be too bulky.

Both the light and ponderous powders may be conveniently made up with mucilage, which increases the bulk less than the other additions, and occasions them to pass

down more freely.

The officinal pharmacopæias have no formula of this kind: most of the following compositions are taken from our hospitals. Bolus Alexipearmacus.

Alexipharmac bolus.

Take of

Compound powder of contrayerva, half a scruple; Syrup of wild poppies, a sufficient quantity to make it into a bolus.

. .

Take of Contrayerva root, half a fcruple; Syrup of faffron, as much as is fufficient.

Make them into a bolus.

3.

Take of
Virginian snakeroot, half a scruple;
Consection of kermes, as much
as is sufficient.
Mix and make them into a bolus.

Take of

Virginian snakeroot, Contrayerva root, each eight grains; Saffron, three grains;

Syrup of meconium, a sufficient quantity to reduce them into

a bolus.

5.

Take of Virginian fnakeroot, fifteen grains;

Caftor, ten grains; Syrup of fugar, as much as is fufficient.

Mix and make them into a bolus.

Take of

Camphor, two grains;
Saffron, five grains;
Cordial confection, one fcruple.
Mix and make them into a bolus.
P.p. 2
7. Take

7.

Take of
Camphor, two grains;
Nitre,

Contrayerva root, each ten

Syrup of clove-july-flowers, as much as will make them into a bolus.

8.

Take of
Musk, ten grains;
Cordial confection, one scruple.
Make them into a bolus.

9.

Take of
Musk, ten grains;
Salt of harthorn, or of sal ammoniac, five grains;
Thebaic extract, half a grain;
Syrup of saffron, a sufficient
quantity.

Make them into a bolus.

THESE boluses are designed for low depressed fevers, in which medicines of this kind are generally prescribed, for keeping up the vis vitæ, raiting the pulse, and promoting a diaphoresis. The compositions differ in strength, nearly according to the order in which they stand. In etwo last are of great power, and are designed chiefly for cases accompanied with convulsive symptoms, which are often abated by them.

Bolus ex alumine.

Alum bolus.

Take of
Alum,
Extract of Peruvian bark,
Nutmeg, each ten grains;
Simple tyrup, as much as will
reduce them into a proper confidence for a bolus,

This composition is a very strong astringent, and as such is used with success in violent uterine hamor-

rhages, and other immoderate secretions which require to be speedily restrained. It may be taken twice a day; or if the flux is very violent, every sour or fix hours, till it abates.

Bolus E CAMPHORA.

Campbor bolus.

Take of
Camphor, half a scruple;
Gum Arabic, half a drachm;
Syrup of marshmallows, a sufficient quantity to make them
into a bolus.

This is a very convenient form for the exhibition of camphor: this drug, however, when thus given by itself in large doses, is apt to nauseate the stomach; and rarely has so good effects as when mixed in small quantities with nitre or other like substances, and frequently repeated.

> Bolus e Castoreo, Caftor bolus.

Take of
Caftor, one scruple;
Salt of hartshorn, five grains;
or oil of hartshorn, five drops;
Simple syrup, a sufficient quantity.

Make them into a bolus.

This medicine is given in hysterical and hypochondriacal disorders, and likewise as an alexipharmac in severs. Its virtues, which are great and unquestionable, seem to depend more upon the set animal oil, or volatile salt, than on the drug from which it takes its name.

Bolus Catharticus.

Purgative bolus.

Take of

Rhubarb, half a drachm;

Solutive syrup of roses, a sufficient

cient quantity to make a bo-

Take of
Jalap, one scruple;
Jamaica pepper,
Crystals of tartar, each five
grains;
Syrup of buckthorn, as much as
will reduce them into a mass
of a due consistence.

Take of
Scammony, ten grains;
Soluble tartar, one scruple;
Soft extract of liquorice, a sufficient quantity.
Let the scammony be well ground

with the foluble tartar, then add the extract, and make them into a bolus.

Take of
Gamboge,
Crystals of tartar, each eight
grains;
Syrup of ginger, a sufficient

Syrup of ginger, a sufficient quantity to reduce them into a bolus.

Take of

Elaterium, two grains;

Extract of jalap, half a scruple;

Crystals of tartar, one scruple;

Syrup of orange peel, a sufficient
quantity to make them into a
bolus.

THE virtues of these compositions are sufficiently obvious; the first is a mild purgative; the two last too strong to be in general ventured on; and the other two of intermediate degrees of strength.

Bolus CATHARTICUS CUM

MERCURIO.

Purgative bolus with mercury.

Take of

Jalap, one scruple;

Mercurius dulcis, five grains;

Solutive fyrup of roles, as much as is sufficient to make them into a bolus.

Take of
Gamboge, seven grains;
Mercurius dulcis,
Aromatic species, each half a
scruple;
Syrup of buckthorn, a sufficient
quantity to make a bolus.

THE first of these compositions is a safe and mild mercurial cathartic: the second is too strong for general use.

Bolus Diaphoreticus.

Diaphoretic belus.

Take of
Compound powder of contrayerva,
Crude fal ammoniac, each one
fcruple;
Simple fyrup, a fufficient quantity to form them into a bolus.

This bolus is given in fevers, and other cases where a diaphoresis is to be promoted. Sal ammoniac is for this purpose one of the most efficacious of the neutral salts. It requires, however, when thus given in a solid form, to be assisted by warm diluents, frequently repeated; which not only promote its action, but likewise prevent its sitting uneasy on the stomach.

Bolus Diureticus.

Diuretic bolus.

Take of
Fresh squills, fix grains;
Compound powder of arum, ten
grains;
Ginger, five grains;
Syrup of orange peel, a sufficient
quantity.
Make them into a bolus.
Pp 3
This

This composition is recommended by Dr. Mead, to be taken every morning in hydropic cases, for promoting urine. He observes, that in these disorders, diuretic medicines vary greatly in their essess, those which answer sufficiently in one person, failing in another; and that the squill and its preparations are, of all others, those which most generally succeed.

Bolus AD DYSENTERIAM.

Bolus against the dysentery.

Take of

The cordial confection,
French bole, each one fcruple;
Thebaic extract, one grain.
Make them into a bolus.

This composition is excellently well calculated for the purpose expressed in its title. Dr. Mead assures us, that he has never found any one medicine more effectual, either for restraining the slux, or healing the exulcerated membranes. Previous to the use of this or other like medicines, the first passages must be cleansed by mild emetics and cathartics, as ipecacoanha and rhubarb. See page 586.

Bolus emmenagogus.

Emmenagogue bolus.

Take of

Socotorine aloes, eight grains;
Saffron, four grains;
Guinea pepper, two grains;
Oil of favin, two drops;
Conferve of rue, as much as is
fufficient to reduce them into

a due confistence.

Take of
Salt of steel, one grain;
Myrrh, half a scruple;
Cordial confection, fifteen grains.
Make them into a bolus.

Take of Black hellebore root, 8 grains; Fresh squills, four grains; Essential oil of pepper-mint, two drops;

Conferve of orange peel, as much as is sufficient to make them into a bolus.

ALL these are medicines of great power of promoting or exciting the menstrual flux. The two sirst are calculated for lax phlegmatic habits; the third, for persons of a sanguine temperament, where chalybeate medicines cannot be borne.

Bolus Febrifuge bolus.

Take of
Peruvian bark, one scruple;
Cascarilla, half a scruple;
Mucilage of quinceseeds, a sufficient quantity to make them into a bolus.

This elegant composition is excellently well adapted to the cure of intermittent fevers; and may be given in cases where the Peruvian bark by itself would be less proper. Where aromatics, chaly beates, bitters, &c. are also requisite, they are either to be premised, or occasionally interposed. See page 196.

Bolus Hystericus.

Hysteric bolus.

Take of
Musk,
Asafætida, each six grains;
Castor, half a scruple;
Syrup of saffron, as much as is
sufficient to make them into a bolus.

This medicine is a very well contrived one for the purposes expressed in its title. It is of great service both in hysterical and hypochondriacal disorders; and often gives relief in the depressions, faintings, statulent choics, head achs, and other symptoms, attending them.

It may be taken twice a day, along with any suitable liquor.

Bolus Iliacus.

Iliac bolus.

Take of

Cathartic extract, one scruple; Thebaic extract, one grain. Make them into a bolus.

This bolus is prescribed by Dr. Mead, for easing the pain, and procuring stools, in the illiac passion, and dry belly-ach; where the irritating cathartics, exhibited by themselves, are thrown up by vomit. The use of this medicine is to be preceded by plentiful bleeding, and accompanied with purgative glysters of the more acrickind; and its operation promoted by infusion of sena mixed with a little of the elixir salutis, or tincture of sena.

BOLUS MERCURIALIS.

Mercurial bolus.

Take of

Calomel, from five to fifteen grains:

Conserve of roses, half adrachm.

Mix and make them into a bolus.

This bolus is given every night, or oftener, for raising a salivation, in venereal, and other disorders, which require that herculean operation. It is likewise taken at night as an alterative, to be carried off next morning by a cathartic: mercurials exhibited in this manner, have generally better effects than when joined with purgatives directly.

BOLUS MERCURIALIS EMETICUS.

Emetic mercurial bolus.

Take of

Yellow emetic mercury, fix grains;

Conserve of roses, a sufficient quantity.

Make them into a bolus.

This strong emetic is given in venereal and leprous diseases; particularly in the case of soul ulcers of long standing, the cleansing and cure of which are frequently promoted by it. The violence of its operation limits its use to robust constitutions.

BOLUS PECTORALIS.

Pettoral bolus.

Take of

Spermaceti, fifteen grains;
Gum ammoniacum, ten grains;
Salt of hartshorn, five grains;
Simple syrup, as much as is sufficient.

Mix and make them into a bolus.

In colds of long standing, old coughs, ashmas, and beginning consumptions, this bolus generally gives relief; especially if bleeding is premised, and repeated, if necessary, at proper intervals.

Bolus RHEI CUM MERCURIO.

Bolus of rhubarh with mercury.

Take of

Choice rhubarb, twenty-five grains;

Calomel, five grains; Simple fyrup, as much as will form them into a bolus.

This is a very mild mercurial purgative. It is given to deftroy worms, and in cachectic, chlorotic, and other like disorders.

Bolus RHEUMATICUS.

Rheumatic bolus.

Take of

Extract of guaiacum, half a drachm;

Salt of hartshorn, seven grains; Simple syrup, a sufficient quantity.

Make them into a bolus.

In chronical rheumatisms, whether the remains of a rheumatic se-

ACL!

ver, or a continuation of pains that proceeded at first from neglected colds, this bolus has been given with good effects, once a week or oftener: the patient keeping warm, and drinking warm liquers, to promote its operation as a cathartic and diaphoretic. use ought to be accompanied by venæsection, which is to be repeated every eight or ten days as long as the blood is fizy. This medicine is likewise exhibited in sciatic, arthritic, and other pains not accompanied with a fiziness of blood: in these it much more frequently fails than in the true rheumatism.

Bolus scilliticus.

Scillitic bolus.

Take of

Fresh squills, twelve grains;
Aromatic species, half a scruple;
Oil of pepper-mint, one drop.
Beat them well together into an
uniform mass, of a due consistence for a bolus.

This is a warm, stimulating, and attenuating medicine, and may be given to great advantage in cases where the natural secretions are obstructed or suppressed from a viscidity or suggishness of the juices. The efficacy of the squills is promoted by the additional ingredients, which at the same time warm and strengthen the stomach and intestines, and prevent the composition from being thrown up by vomit, which this quantity of squills, given by itself, would in many constitutions be.

Bolus THERIACALIS.

Treacle bolus.

Take of Theriaca, two scruples; Salt of hartshorn, seven grains ;
Camphor, three grains.
Mix and form them into a bolus.

CAMPHOR and falt of hartshorn, when thus joined with opiates, have in many cases better effects than if exhibited by themselves, their diaphoretic virtue being greatly promoted by the relaxation which the opium occasions. The quantity of theriaca in this bolus contains somewhat more than a quarter of a grain of opium.

Bolus sudorificus.
Sudorific bolus.

Take of

Camphor, five grains;
Thebaic extract, one grain;
Syrup of orange peel, a fufficient quantity to reduce them into a bolus.

This medicine is one of the most effectual sudorifics, generally exciting a copious sweat. In many cases, where this intention is to be answered, whether acute or chronical, it may be given to great advantage.

Bolus terebinthinatus.

Turpentine bolus.

Take of

Chio turpentine, one scruple; Powdered liquorice, a sufficient quantity.

Make them into a bolus.

This is a convenient form for the exhibition of turpentine, the liquorice powder answering the same intention here as the elecampane root in the pilulæ piceæ, page 577.

### CHAPTER V.

#### Elettaries.

LECTARIES are composed chiefly of powders mixed up with fyrups, &c. into such a confistence, that the powders may not separate in keeping, that a dose may be easily taken up on the point of a knife, and not prove too stiff to swallow.

Electaries receive chiefly the milder alterative medicines, and fuch as are not ungrateful to the palate. The more powerful drugs, as cathartics, emetics, opiates, and the like, (except in officinal electaries to be dispensed weight) are seldom trusted in this form, on account of the uncertain-. ty of the dose; disgustful ones, acrids, bitters, fetids, cannot be conveniently taken in it; nor is the form of an electary well fitted for the more ponderous substances, as mercurials, their being apt to subfide in keeping, unless the composition is made too stiff.

The lighter powders require thrice their weight of honey, or fyrup boiled to the thickness of honey, to make them into the confiftence of an electary; of fyrups of the common confiftence, twice the weight of the powders is suffitient.

Where the common fyrups are employed, it is necessary to add likewise a little conserve, to prevent the compound from drying too soon. Electaries of Peruvian bark, for instance, made up with syrup alone, will often in a day or two grow too dry for taking.

Some powders, especially those of the less grateful kind, are more conveniently made up with muci-

lages than with fyrups, honey, or conserve. The three latter stick about the mouth and fauces, and thus occasion the taste of the medicine to remain for a considerable time; whilst mucilages pass freely, without leaving any taste in the mouth. A little soft extract of liquorice, joined to the mucilage, renders the composition sufficiently grateful, without the inconveniences of the more adhesive sweets.

The quantity of an electary directed at a time, in extemporaneous prescription, is rarely less than an ounce, or more than three ounces.

General rules for making electaries.

I.

The rules already laid down for decoctions and powders in general, are likewise to be observed in making decoctions and powders for electaries [E.]

Gums, inspissated juices, and such other substances as are not pulverable, should be dissolved in the liquor prescribed: then add the powders by little and little, and keep the whole briskly stirring, so as to make an equable and uniform mixture [E.]

Aftringent electaries, and such as have pulps of fruits in their composition, should be prepared only in small quantities at a time [E.]

For aftringent medicines lose greatly of their virtue, on being kept kept in this form, and the pulps of fruits are apt to become four. IV.

The superfluous moissure of the pulps should be exhaled over a gentle fire, before the other ingredients are added to them [E.]

Electaries, if they grow dry in keeping, are to be reduced to the due confidence, with the addition of a little Canary wine; [L.E.] and not with fyrup or honey; by this means, the dose will be the least uncertain; a circumstance deserving particular regard, in those especially which are made up with syrup, and contain a large quantity of opium, as the consession Paulina, and philonium [L.]

#### ELECTARIUM ad DYSENTERICOS. Antidysenteric electary. Edinb.

Take of

Japonic confection, two ounces; Locatelli's balfam (beaten up with a fufficient quantity of yolks of eggs) one ounce; Powdered rhubarb, half an ounce; Syrup of marshmallows, a suffi-

cient quantity.

Mix and make them into an electary.

This composition is extremely well contrived for the purpose expressed in its title. Astringents or opiates by themselves rarely have place in dysenteries, even after the first passages have been evacuated by an emetic or a full dose of rhubarb; they ease the pain and moderate the flux for a time; but the short relief is apt to be followed by dangerous or even satal confequences from the retention of the acrid and corrupted humours. The

shubarb, which the college of Edinburgh has now added from the practice of the infirmary, in good measure prevents this accumulation, without much counteracting the falutary effects of the other materials; in many cases. however, it may be still necessary to interpose that laxative drug by The dose of the electary itself. is the bulk of a large nutmeg, once or twice a day, according to the urgency of the symptoms. One drachm contains about onefixth part of a grain of opium.

### ELECTARIUM e BACCIS. LAURI.

Electary of bay-berries. Lond.

Take of
Rue leaves dried,
Caraway feeds,
Parfley feeds,
Bay-berries, each one ounce;
Sagapenum, half an ounce;
Black pepper,
Ruffia cattor, each two drachms;
Clarified honey, thrice the weight of the powdered species.

Mix the species with the honey, and make them into an electary.

This composition is sometimes taken, in statulent cholics and hysterical disorders, from a scruple to two drachms: but its principal use is in carminative glysters, nor is it often employed in these. The college of Edinburgh have entirely dropped it.

## ELECTARIUM e CASIA. Electary of cafia. Lond.

Take of
Solutive fyrup of roles,
Pulp of casia, fresh extracted,
each half a pound;
Manna, two ounces;

Pulp

Pulp of tamarinds, one ounce. Grind the manna in a mortar, and, with a gentle heat, dissolve it in the syrap; then add the pulps, and continue the heat until the whole is reduced to a due consistence.

### · DIACASIA. Edinb.

Pulp of casia, twelve ounces;
Pulp of casia, twelve ounces;
Pulp of tamarinds, six ounces;
Calabrian manna, eight ounces;
Syrup of pale roses, one pound.
Dissolve the manna in warm water, strain the solution, and evaporate it, along with the syrup,
over a gentle fire, to the consistence of honey: then mix in the
pulps, so as to make the whole
into an uniform electary, according to art.

THESE compositions are very Convenient officinals, to serve as a basis for purgative electaries and other like purposes; as the pulping a small quantity of the fruits, for extemporaneous prescription, is sufficiently troublesome; the tamarinds give them a pretty tafte, and do not subject them, as might be expected, to turn four: after flanding for four months, the compolition was found no fourer than when first made up. They are likewise usefully taken by them. selves, in the quantity of two or three drachms occasionally, for gently loosening the belly in costive habits.

## ELECTARIUM LENITIVUM. Lenitive electory. Lond.

Take of
Figs, one pound;
Sena, eight ounces;
Pulp of tamarinds,
Pulp of casia,

Pulp of French prunes, each half a pound; Coriander feeds, four ounces; Liquorice, three ounces; Double-refined fugar, two pounds and a half.

Pulverize the sena along with the coriander seeds, and sift out ten ounces of the powder; the remainder is to be boiled with the sigs and siquorice, in four pints of water, to one half; then strain and press out the siquor, and evaporate it to the weight of a pound and a half, or somewhat less: in this dissolve the sugar, so as to make it into a syrup, and add this syrup, by little and little, to the pulps: lastly, mix in the powder before separated by the sieve.

This electary may be occasionally taken to the quantity of a nutmeg or more, for loosening the belly in costive habits. It is frequently employed in glysters, though for that use the following is rather more convenient.

#### Edinb.

Take of
Sena, four ounces;
Liquorice, two ounces;
Coriander feeds, one ounce;
Pulp of prunes, one pound;
tamarinds,

casia, each half a pound;
White sugar, fix pounds;
Boiling water, four pints.
Insufe the sena, liquorice, and coriander seeds in the water, in a close vessel, for a night: press out and strain the liquor, and boil it with the sugar to the consistence of a thick syrup: then add the pulps, and mix the whole well together, so as to make them into an elec-

tary.

THIS

This electary is much preferable, for glysters, to those which have powders in their composition. It is now freed from some superfluous ingredients which were left in it at former revisals; viz. polypody roots, French mercury leaves, fenugreek feeds, and linfeed. Honey is exchanged for fugar, which is less apt to grow four in keeping; perhaps melasses would be preferable or either, as it coincides with the intention, and is not only of itself inapt to ferment, but likewise prevents such substances as are this way disposed, from running into fermentation.

#### ELECTARIUM PECTORALE. Pedoral eledary. Edinb.

Take of

Rob of elder berries, two ounces;

Spermaceti dissolved in a sussicient quantity of yolks of eggs, half an ounce;

Flowers of benzoine, one drachm; Balfamic fyrup, as much as is fufficient to make the other ingredients into an electary.

This is a very useful medicine, in tickling coughs and common colds, calculated both to obtund acrimony and promote expectoration. It may be used two or three times a day, in doses of about the quantity of a small nutmeg. Taken to the bulk of a large nutmeg, at bed-time, it generally, not only relieves the breast, but tends to procure a falutary diaphoresis or sweat in the night. It is here improved from the former editions, by fubilituting rob of elder berries to conserve of roses, and spermaceti to compound powder of gum tragacanth.

ELECTARIUM e SCAMMONIO. Electary of scammony. Lond.

Take of

Scammony, an ounce and a half; Cloves, Ginger, each fix drachms;

Essential oil of caraway seeds, half a drachm;

Honey, half a pound. Let the spices be ground together, and mixed with the honey; then add the powdered scammony, and afterwards the oil.

This electary is a warm, brifk purgative. It is a reform of the electarium caryocoftinum of our preceding dispensatories, a composition which was greatly complained of, as being inconvenient to take, on account of the largeness of its dose. A drachm and a half of this, which contains fifteen grains of scammony, is equivalent to half an ounce of the other.

ELECTARIUM e SCORDIO.
Electory of scordium, commonly called
Diascordium.
Lond.

Take of

The species of scordium with opium, any quantity;
Syrup of meconium, boiled to the consistence of honey, thrice as much by weight.

Mix the species with the syrup, so as to make an electary.

In our former dispensatories, the species were ordered to be made up with honey: this is now exchanged for a syrup, more agreeable to the intention of the medicine, which is that of an opiate astringent, whilst honey is manifestly aperient and detergent. It is not perhaps necessary, for the purposes of the shops, to make the

the species into an electary at all: by keeping in this form, the ingredients lose greatly of their aromatic flavour and attringency, becoming foft and fmooth upon the palate; and the red colour, imparted by the bole, decays. The London college have therefore very justly ordered them to be kept in powder as well as in an electary; and directed the powder, both with and without opium, for different occasions. See Species e fcordia, and Pulvis e bolo, cum and fine opia. Either of these powders may be made up extemporaneoully into an electary, with any fyrup that shall be judged proper.

Diascordium was intended by its author Fracastorius for an antipestilential; but we have been so happy as to have little occasion for medicines in that intention; nor could this be anywise depended on. It is a moderately warm astringent and opiate; and in this light only is considered by the present practice. One grain of opium is contained in nine scruples of the elec-

tary.

The species e scordie, which make the basis of this electary, contain, as we have already feen (page 5 (6) several superfluous ingredients; for though the London coilege has given a judicious reformation of the powder, under the title of Pulous e bolo, the electary is made with the powder unreformed; partly, that no material alteration might be made in a medicine, which is so much depended on, and whose effects have been so long experienced; and partly because the physician, if he prefers the pulvis e bolo, may direct an electary to be made with it in extemporaneous prescription. In the Edinburgh pharmacopæia, this medicine is not ordered to be kept in powder, but the electary is reformed to a great degree of elegance and timplicity. And as the ingredient from which it received its name, being a very unimportant one, is now omitted, the composition is distinguished by another title, viz.

## CONFECTIO JAPONICA. Japonic confedion. Edinb.

Take of Japan earth, three ounces; Tormentil root,

Nutmeg,

Olibanum, each two ounces; Opium diffolved in a fufficient quantity of Canary wine, one drachm and a half;

Syrup of dry roses, hoiled to the thickness of honey, thrice the weight of the powders. Mix and make them into an electary, which supplies the place

of diascordium.

THE ingredients in this electary feem extremely well chosen, and are so proportioned to one another, that the quantity of opium is the same as in the diascordium of the former pharmacopoeias of Edinburgh, viz. one grain in ten scruples.

## BALSAMUM LOCATELLI. Locatelli's balfam. Lond.

Take of

Oil olive, one pint; Strasburgh turpentine, Yellow wax, each half a pound; Red saunders, six drachms.

Melt the wax over a gentle fire, with some part of the oil; then add the rest of the oil, and the turpentine; afterwards mix in the faunders, and keep them stirring together, until the mixture is grown cold.

Edinb.

Edinb.

Take of

Yellow wax, one pound;
Oil olive, a pint and a half;
Chio or Strasburgh turpentine, a
pound and a half;
Police of Peru, two ources:

Balíam of Peru, two ounces; Dragons blood, in powder, one ounce.

Melt the wax in the oil over a gentle fire, then add the turpentine; and having taken them from the fire, mix in the balfam of Peru and dragons blood, keeping them continually stirring till grown cold.

DRAGONS blood gives a more elegant colour to this composition than red saunders, though on another account it is fomewhat less proper, having been found, when dissolved in oil, to communicate fome degree of heat and pungency, qualities quite foreign to the intention of the medicine. balsam is used in internal bruises and hæmorrhages, erofions of the intestines, dysenteries, and in some kinds of coughs and afthmas: the dose is from two scruples to two drachms: it may be commodiously taken with about double its weight of conserve of roses, as directed hereafter in page 598. Some have likewise applied it externally for deterging and incarnating recent wounds and ulcers.

BALSAMUM CEPHALICUM.
Cepalic balsam.
Edinb.

Take of

Expressed oil of nutmegs, one ounce;

Dissilled oil of cloves, of lavender, of rosemary, each half a drachm; of amber, half a scruple;

Balsam of Peru, one drachm.
Lquesy the oil of nutmegs in a

filver vessel; and when taken from the fire, mix into it the distilled oils and the balsam, according to art.

This medicine is recommended to be rubbed on the temples, and on paralytic limbs, for warming the part and comforting the nerves; and to be fmelled to, for refreshing and enlivening the spi-Some have also given it inwardly as a warm cordial, in languid cases, and in debilities of the nervous system. There are abundance of preparations of this kind in foreign pharmacopæias, composed each of only one essential oil, incorporated with the expressed oil of nutmegs; which last is to be previously freed from its flavour (by distillation with water) that the smell of the former may not be injured thereby: in the room of this prepared sebaceous matter, a mixture of white wax and oil olive might be used. In the pradical chemistry, a general process is given for the making of these kinds of preparations, under the title of

BALSAMUM ODORIFERUM.

An odoriferous balfam.

Take of Oil olive,

White bees wax, each two ounces. Put the oil into a China bason. placed in a pan of boiling water, and flice the wax into it. Stir them together with a clean knife, or small spatula, till the wax is melted: then remove the vessel out of the hot water, and when the matter begins to thicken, drop in four drachms of any odoriferous effential oil, as that of cinnamon, nutmegs, mace, lemon peel, rhodium, lavender, rosemary, &c. or of a mixture of two or three of these oils: to which

which may be added one drachm of effence of ambergris, which will beighten the smell of the oils, without communicating any of its own. Keep the whole constantly stirring, that they may be perfectly mixed; and as soon as this is done, plunge the veffel into cold water, to prevent the distipation of the essential oils.

THESE kinds of balfams may be made of any colour, so as to resemble in this respect also, as well as in smell, the vegetable, from which the essential oil, you make use of, was drawn. A little of the pigment, called by the painters sap-green, being previously ground with the oil olive, will give a sine green; a little cinnabar, a scarlet; turmeric, a lemon colour; Prussian blue, a violet; and cochineal, a sine purplish hue.

#### CONFECTIO CARDIACA.

Cordial confection.

Edinb.

Take of

Conserve of rosemary slowers, three ounces:

Candied nutmegs, one ounce and a half;

Candied ginger, fix drachms; Compound powder of contrayerva, half an ounce;

Oil of cinnamon, fifteen drops; Syrup of orange peel, as much as is sufficient.

Mix them into an electary, according to art.

This electary, though very different in composition, is nearly similar in virtue to that formerly described under the same name, page 414. Particular care ought to be had in the choice of the effential oil; for on its goodness,

that of the medicine in great meafure depends.

CONFECTIO PAULINA.

The confection called Paulina.

Lond.

Take of

Costus, or in its stead zedoary, Cinnamon,

Long pepper,
Black pepper,
Storax,
Galbanum,
Opium.

ftrained,

Russia castor, each two ounces; Simple syrup, boiled to the consistence of honey, thrice the weight of the other ingredients.

Warm the fyrup, and carefully mix with it the opium first diffolved in wine: gradually add this mixture, whilst it continues warm, to the storax and galbanum previously melted together; and afterwards sprinkle in the other species reduced into powder.

This is the confectio ARchicenis of our former dispenfatory, brought back to its first form and author. It is a warm opiate medicine, and as such is sometimes made use of in practice: thirty-two grains contain one grain of opium.

MITHRIDATIUM, five CONFECTIO DAMOCRATIS.

Mithridate, or the confedion of Damocrates.

Lond.

Take of
Cinnamon, fourteen drachms;
Myrrh, eleven drachms;
Agaric,
Indian nard,
Ginger,
Saffron,
Seeds of mithridate mustard,
Frank-

Take of

Frankincense, Chio turpentine, each ten drachms: Camels hay, Costus, or in its stead zedoary, Indian leaf, or in its stead mace. Stechas. Long pepper, Hartwort seeds. Hypocistis, Storax thrained, Opoponax, Galbanum strained. Opobaliam, or in its stead expressed oil of nutmegs, Russia castor, each one ounce; Poley mountain, Scordium, Carpobalsam, or in its stead cubebs. White pepper, Candy carrot feed, Bdellium strained, each seven drachms: Celtic nard. Gentian root, Dittany of Crete, Red roses. Macedonian parsley seed, Leffer cardamom feeds, husked, Sweet fennel seed. Gam Arabic, Opium ftrained, each 5 drachms; Calamus aromaticus, Wild valerian root, Anisced, Sagapenum, strained, each three drachms; Meum athamanticum, St. John's wort, Acacia, or in its stead terra Japonica, Bellies of skinks, each two drachms and a balf; Clarified honey, thrice the weight of all the other ingredients. Warm the honey, and mix with it the opium dissolved in wine; melt the storax, galbanum, turpentine and opobalfam (or expressed oil of nutmegs) together

in another vessel, continually stirring them about, to prevent their burning; with these, so melted, mix the hot honey, at first by spoonfuls, and afterwards in larger quantities at a time; when the whole is grown almost cold, add by degrees the other species reduced into powder.

### THERIACA ANDROMACHI. Venice treacle. Land

Troches of fquills, half a pound; Long pepper, Opium strained, Vipers dried, each three ounces; Cinnamon, Opobalsam, or in its stead expressed oil of nutmegs, each two ounces; Agaric, Florence orris root, Scordium. Red roses. Navew feeds. Extract of liquorice, each an ounce and a half; Indian nard, Saffron, Amomum, Myrrh, Costus, or in its stead zedoary, Camels hay, each one ounce; Cinquefoil root, Rhubarb. Ginger. Indian leaf, or in its stead mace, Dittany of Crete, Horehound leaves, Calamiat leaves, Stechas, Black pepper, Macedonian parfley feed, Olibanum, Chio turpentine, Wild valerian root, each fix drachms: Gentian root, Celtic

Celtic nard. Spienel Poley Mountain ) St. John's wort > leaves, Groundnine Germander tops, with the feed, Carpobaliam, or in its flead cubebs. Annisced. Sweet fenael feed. Leffer cardamom feeds, hufked, Bishops weed Hartwort ·fceds. Treacle mufterd.) Hypocifiis, Acacia, or in its Read Japan earth. Gum Arabic. Storax Arained. Sagapenum firsined, Terra Lemuia, or in its fload bole armenic or French bole, Green vitriol calcined, each half an ounce; Small (or in its flead, the long) birthwort root. Leffer centaury tops, Candy carrot feed, Opopanax, Galbanum frained. Ruffia caftor, Jews pitch, or in its stead, white amber prepared, Calamus aromaticus, each two drachms: Clarified honey, thrice the weight of all the other ingredients. Let these ingredients be mixed together, after the same manner as directed in making the mi-

THESE celebrated electaries are almost the only remains, which the late reformation has left in the shops, of the wild exuberance of composition, which the superstition of former ages brought into vogue. The theriaca is a refor-

thridate.

mation of mithridate, made by Andromachus, physician to Nero: the mithridate itself is faid to have been found in the cabinet of Mithridatus king of Pontus. The first publishers of this pompous areanum were very extravagant in their commendations of its virtues; the principal of which was made to confift in its being a most powerful prefervative against all kinds of venom: whoever took a proper quantity in a morning, was enfured from being personed during that whole day: this was confirmed by the example of its supposed inventor, who, as Celfus informs us, was by its conftant use fo fortified against the commonly reputed poisons, that none of them would have any effect upon him when he wanted their affiftance. But the notions of poisons, which prevailed in those ruder ages, were manifeftly erroneous. Before experience had furnished mankind with a competent knowledge of the powers of fimples, they were under perpetual alarms from an apprehension of poisons, and hufied themselves in contriving compositions which should counteract their effects, accumulating together all those substances which they imagined to be possessed of any degree of alexipharmac power. Hence proceed the voluminous antidotes which we meet with in the writings of the antient phyficians: yet it does not appear, that they were acquainted with any real poifon, except the cicuta, aconitum, and bites of venomous beafts; and to these they knew of no antidote whatever. Even admitting the reality of the poisons, and the efficacy of the feveral antidotes feparately, the compositions could no more answer the purposes expected from them, than the accumulating of all the medicinal fimples into Qq

one form could make a remedy dice to the medicine.

against all diseuses.

Yet notwithstanding the absurdity in the original intention of these medicines, and their enormity in point of composition; as they contain several powerful materials, whose virtues, though greatly prejudiced, yet are not destroyed, by their multiplicity and contrariety; the compounds have been found, from repeated experience, to produce very considerable effects, as warm opiate diaboretics.

These compositions might without doubt be lopt of numerous fuperfluities, without any diminu tion of their virtues; yet as the effects of them, in their present form, are so well known, so much regard has been paid to ancient au hority, as not to attempt a reformation of that kind. London college have however thought proper to retrench, from forms originally complex, all subfequent additions that have crept into them. Neither the description in verse of the elder Andromachus, not the prose explanation of the younger, make any mention of the white pepper afterwards added to the theriaca; and the orris root, in the mithridate of our former pharmacopœias, is also a supernumerary ingredient, not warranted by the original: these therefore are rejected. Nor is the afarum in mithridate grounded on any good authority: the verse it is taken from, is mutilated and corrupt; and the word which fome, upon conjecture only, suppose to have been asarum, others, also upon conjecture, chuse to read differently: till fome emendation shall be better founded ed than merely upon critical guesfes, this fingle ipocies may be tafely passed over, without any prejudice to the medicine. None of the antient descriptions afford any other light in this particular; for they either omit this ingredient, and others also, or abound with additions.

One innovation in both these medicines, the college have allowed themselves. In each of these compositions are found both cinnamon and cafia lignea; and it is very evident, from feveral parts of Galen's works, that the latter was used by the ancients only upon account of the great difficulty of procuring the other; so that to retain the casia, now that cionamon is so common, is a blind following of these writers, without any attention to their meaning: the cana therefore is now rejected. and half the quantity of cinnamon put in its room, which is the proportion that Galen directs to be observed in substituting the one for It is probable, that the the other. case is the same with regard to the Celtic and Indian nard; that the first had a place in these compositions, on account of the difficulty of procuring the Indian; for Galen expressly prefers the latter.

There is a material error in regard to the theriaca, which has passed through all the editions of our pharmacopæia, except the prefent: this is, the substituting Roman vitriol to the ancient chalcitis, now not certainly known, and in the catalogue of simples, describing the Roman to be a blue vitriol: whereas the Italian writers are unanimous it is a green vitriol; and were it not, it would not answer to the effects of the chalcitis, which was certainly a chalybeate, and gives the medicine its black colour. What has chiefly occasioned chalcitis to be supposed a cupreous vitriol, seems to be its name, derived from xalus, copper: but it is to

be

be observed, that all vitriols were formerly imagined to proceed from copper, and were named accordingly: the green or martial vitriols are fill called by the Germans hupferwaffer, and by us copperas; It is probable, that the ancient chalcitis was no other than a native martial vitriol, calcined by the heat of those warm climates, to a degree of yellowish red or coppery colour: and therefore the common green vitriol, thus calcined by arty very properly supplies its place.

The London college have likewife fomewhat facilitated the preparation of these medicines, by omitting the trochifei cyphen used in the mithridate, and the bedychroi and viperini for the theriaca; and inserting their ingredients, after Z-welffer's manner, in the compositions they are intended for. This is done in the theriaca very commodiously, the ingredients in these troches uniting with those in the theriaca itself, into unbroken numbers. But to render the numbers equally simple in the mithridate. it was necessary to retrench a few odd grains from some of the articles, and make a small addition to fome others: they adjusted the proportions of the ingredients in the trochisci cyphos from the original description in Galen; the numbers in our former pharmacopœia being yery erroncous.

The college of Edinburgh, paying very little deference to antiquity or common prejudice, have ventured at length to discard these venerable reliques; and have substituted in their room an elegant and simple form, equivalent to them both in essicacy, under the title of

#### THERIACA EDINENSIS.

Edinburgh theriaca. Edinb. Take of
Virginian snakeroot, ten ounces;
Contrayerva root, six ounces;
Refin of guaiacum, sour ounces;
Lesser cardamom seeds, two ounces;
Myrrh,
English saffron,
Opium, each one ounce;
Rob of elderberries, thrice the weight of the powders;
Canary wine, as much as is sufficient to dissolve the opium.
Make them according to art into

an electary.

THIS composition confished very powerful ingredients, and is doubtless capable of answering every thing that can be reasonably expected from the more voluminous theriaca of Andromachus. London college also had formerly their theriaca composed of the less exceptionable ingredients of Andromachus. But as these medicines have for a long time been chiefly employed for external purposes, by the way of cataplasm, the theriaca Londinensis is now omitted, and its place supplied by a cataplaim composed of a few well-chosen articles, under the name of cataplasma e cymino, of which hereafter. For internal use, none of the theriacas are at prefent fo much regarded as they have been heretofore; practitioners having introduced in their room, extemporaneous bolutes of Virginian inakeroot, camphor, contrayerva. and the like; which answer all their intentions, with this advantage, that they may be given either with or without opium, an ingredient which renders the others prejudicial in cases where they might otherwise be proper.

With regard to the quantity of opium in the foregoing compositions, one grain thereof is con-

Qq 2 tained

tained in four drachms of the mithsidates: in threefcruples; fifteen grains of the Venice treacles and in five scruples of the theriace Edinonsis. The proportion of apium will vary a little, according to the time that they have been kept; their moisture by degrees exhaling, so as to leave the remainder stronger of the opium, than an equal weight was at A change of this kind is taken notice of by many writers, but falfely attributed to an imagimany fermentative quality of the ingredients; by which they were supposed, from their multiplicity and contrariety, to be continually exalting and improving the virtues of one another.

A good deal of care is requifite in making these compositions, to prevent the waste which is apt to happen in the pounding, and which would render the proportion of opium to the other ingredients precarious. The intention of disloving the opium in wine, for these and other electaries, is, that it may be more uniformly mixed with the rest.

PHILONIUM LONDINENSE.

Lendon philonium. Lond.

Take of

White pepper, Gioger,

Caraway feeds, each twoodness; Strained opium, fix drachms; Syrup of meconium, boiled to the confiftence of honey, thrice the weight of the other ingredients.

Heat the fyrup, and carefully mix with it the opium, previously diffolved in wine; then add the other ingredients, reduced into powder.

Tute is a reformation of the philonium described by Galen, which was secrimed in our preceding pharmacopecies with the addition of fome superfluous ingredients, and distinguished, but not very properly, by the opithet Romanum. The additional articles, and fome unnecefary ones that were in the original, are here omitted, and the quantities of the other varied, so as to preserve the same proportion of opium to the whole, as in our last pharmacopteia. Thirty-fix grains of the composition contain one grain of opium.

Ten mithridate, theriace, dinfcordium, confectio Paulina, and philonium, are the only compositions now remaining, of what have been called the officinal capitals. They are all medicines of great power; and as, on the one hand. they are applicable, by the judicious physician, to excellent purpoles, lo on the other, their imprudent use has often been productive of mischievous consequences. It has been customary among nurses, and others, to give diascordium to children, to cafe their complaints, and to procure sleep: intentions which it effectually anfwers, but at the same time never fails to bring on a coflive habit, the foundation of many diseases: this medicine has likewife been too unwarily given for restraining fluxes; whose suppression was afterwards followed by more dangerous lymptoms. The celebrated elexipharmacs, mithridate, and theriaca, have oftentimes aggravated the diforders they were intended to remedy, have converted a commen cold into a high fever, have raised flight febrile complaints into a malignant fever. However firongly therefore these kinds of medicines are recommended for eating pain, warming, promoting fweat, expelling malignity, &c.

the atmost cention is requisite in the use of them; the cases which demand their affiftance, are much less frequent than is generally suppoled.

> BLECTARIUM ACIDUM. Acid elettary.

Take of

Conferre of woodforrel, one ounce :

Pulp of tamarinds, half an ounce; Weak spirit of vitriol, as much as is sufficient to give a grateful acidity;

Syrap of lemon juice, as much as will reduce the whole into the confiftence of a fost electary.

Terrs grateful acid composition is an useful refrigerant and antiseptic in different kinds of inflammetory and putrid disorders.

ELECTARIUM ALEXETERIUM. Alexeterial declary.

Take of

Confection of kermes, one drachm:

Candied ginger, fix drachms; Contrayerva root,

Virginian snakeroot, each one drachm;

Syrup of orange peel, as much as is fufficient to make the other ingredients into the confiftence of an electary.

This is a moderately warm electary, contrived by Boerhaave for railing and recruiting the firength in low fevers, where the pulse is funk, and the patient languid and dejected. It may be taken to the quantity of a small nutmeg every four or five hours with any proper julep.

RLECTARIUM ALTERANS. Alterative electory.

Take of

Crude antimony, finely levigeted, three drachme: Refin of guaiacum, two drachmes Oil of faffafras, fix drops;

Conferve of sed roles, one ounce and a half :

Ballamic fgrup, as much as is fufficient.

Grind the ratio and the levigated antimony well together; and having mixed these with the oil (dropped on a little fugar) and the conferre, let the whole be fostened with the syrup into a due confiftence.

Turs medicine is used against cutaneous foulnesses, obstructions of the glands, and impurities of the blood and juices. Dispensatory writers in general lay the principal stress, in compositions of this kind, upon the calx, ceruffe, orcinnabar of antimony, preparations which are far inferior to the crude mineral, and very ill deserve the great character which has been usually given of them. The bulk of a small nutmeg of this electary may be taken every morning and evening with a little of the fimple or compound lime water.

> ELECTARIUM ANTIEPILEPTICUM. Antiepileptic electory.

Take of

Peruvian bark, one ounce; Wildvalerian root, two drachms; Syrup of orange peel, a fufficient quantity to reduce the others into an electary.

This medicine has been frequently prescribed by Dr. Mead, in epileptic cases, with success: he directs one drachm to be taken every morning and evening, for three months together: after which, to confirm the cure and prevent a relapic,

**Q**93

lapse, the same dose is to be repeated for three or four days before every new and full moon for a confiderable time.

> ELECTARIUM ANTIDYSENTERICUM. Antidysenteric electary.

Take of

Yellow wax, three drachms; Spermaceti, two drachms; Conferve of red roses, an ounce and a half;

Oil of almonds, half an ounce; Balfamic fyrup, afufficient quantity.

Let the wax and spermaceti be melted in the oil, over a gentle fire, and then mixed with the conserve and syrup.

WHERE sharp irritating humours have eroded the intestines, and laid open the mouths of the blood-vessels, this soft healing electery is often of great use. It is said that sluxes of long standing, contracted in the Indies, which had yielded nothing to medicines of the restringent kind, have been removed by this, which supplies the matural mucus of the bowels that the slux has carried off, heals the excoriations, and obtunds the acrimonious humours.

ELECTARIUM AROMATICUM.

Aromatic electary.

Take of

The aromatic species, one drachm and a half:

Conferveoflavender, two ounces; Syrup of orange peel, a fufficient quantity.

Make them into an electary.

This warm cordial medicine is of use in nervous complaints and decays of constitution. The bulk of a small nutneg may be taken two or three simes a day with a

glass of wine, or any other properliquor, after it.

ELECTARIUM BALSAMICUM.

Balfamic electary.

Take of

Conserve of roses, two ounces; Locatelli's balsam, one ounce. Dissolve the balsam in the yolk of an egg, and then mix therewith the conserve.

This electary is used in some coughs and disorders of the breast; as also in the vomica, or suppuration in the stomach, which sometimes happens after dysenteries; and where there is an erosion or rupture of the blood-vessels, as in hamoptoes. In these cases, the bulk of a nutmeg may be taken for a dose twice or thrice a day.

ELECTARIUM CHALYBEATUM.

Chalybeate electary.

Take of 1.

Salt of steel, one drachm; Candied nutmegs,

Candied ginger, each half an ounce:

Oil of cinnamon, five drops; Conferve of orange peel, one ounce:

Balfamic fyrup, as much as is fufficient to make them into an electary.

Take of

Rust of steel, or steel prepared with sulphur, fix drachms; Candied ginger, one ounce; Conserve of orange peel, three ounces;

Syrup of orange peel, as much as will reduce them into a proper confiftence.

These elegantehalybeate medicines are given not only in cachectic and chlorotic cases, and menstrual obstructions; but like-

Wile

wife in low hysteric, and melancholic disorders; and for warming and invigorating the habit in great debilities and decays of constitution. In either of these intentions, the bulk of a small nutmeg is to be taken twice a day, and its effects promoted by moderate exercise.

ELECTARIUM DEOBSTRUENS.

Deobstruent electary.

Take of

Gum ammoniacum, Hard foap, each a drachm; Powdered fquills, one scruple; Conserve of orange peel, half an ounce;

Syrup of ginger, as much as is fufficient to reduce the other ingredients into the confiftence of an electary.

WHERE the breast is oppressed by thick phlegm, or the viscera obstructed, this electary may be taken twice or thrice a day to the bulk of a small nutmeg at a time. The quantity above prescribed is sufficient for fix or eight doses.

ELECTARIUM AD GONORRHORAM.

Electory for a genorrhora.

Take of
1.

Lenitive electary, three ounces;
Jalap, three drachms;
Nitre, one drachm and a half;
Simple fyrup, a fufficient quantity to make them into an electary.

Take of
Lenitive electary, three ounces;
Balfam of Copaiva, one ounce
and a half;
Rhubarb,
Gum guaiacum,
Nitre, each one ounce;
Syrup of orange peel, as much
as will reduce them into a proper confiftence for an electary.

These compositions are said to be used in some of the military hospitals; the first as a cooling laxative, for the instammation and tension of the urinary passages, which always accompany a virulent gonorrheae: in this intention, adrachm and a half is directed to be taken every morning and evening. The second is designed for strengthening the parts after the virulence is expelled, and the heat and instammation have ceased: the bulk of a nutmeg may be taken twice or thrice a day.

ELECTARIUM E GUMMI GUAIACO.

Eledary of gum guaiacum.

Take of
Gum guaiacum,
Compound powder of arum,
Canella alba, each fix drachms;
Conferve of fcurvy-grass, two

ounces;
Syrup of orange peel, as much
as will bring them into a proper confidence,

In chronical rheumatisms, pains and aches in general, that are not accompanied with inflammation, and some kinds of paralytic numbnesses, this warm simulating electary may be taken to the quantity of a nutmeg twice a day.

ELECTARIUM EX HELLEBORO NIGRO.

Electory of black bellebore.

Take of

Black hellebore root,

Extract of favin,

Compound powder of myrrh,

each half an ounce;

Canella alba, two drachms;

Syrup of orange peel, as much
as is fufficient.

Mix and make them into an electary.

THIS

This electricy is employed in one of our holpitals for promoting the natural evacuations from the uterus: for which purpose, it is undoubtedly a medicine of great power. It may be taken to the quantity of half a drachm twice a day.

BTECTARIUM INCRASSANS.

Incraffasing slettary.

Take of

Gum tragacanth,

Pulp of fresh comfry root, each one onnce:

Conferve of mallows, half an ounce:

Syrup of marshmallows, as much as is sufficient to reduce the whole into the consistence of an electary.

This electary is taken to the quantity of a cheffinut, three or four times a day, along with a milk diet, for incraffating and obtunding thin ferous humours, in hettic diforders, in coughs proceeding from thin tickling rheums, in fluxes and heat of urine, where the natural mucus of the parts is abraded.

### ELECTARIUM AD NEPHRITICOS. Neporitie electary.

Take of

Lenitive electary, an ounce and a half;

Venice turpentine, one ounce; Egg shells prepared [or prepared oyster shells] balf an ounce; Choice rhubarb, one drachm; Syrup of marshmallows, as much as is sufficient.

Diffolve the turpentine in the yolk of an egg, and then mix the whole together, according to art, fo as to make thereof an electary.

This composition, taken from the Edinburgh infirmary, is contrived for cleaning the urinary passages

in nephritic disorders. Turpentine, properly divided by earthy powders, is a safe, and, at the same time, one of the most powerful divided to the same time, one of the most powerful divided to the same time one; the rhubarb and lax-ative electary are very useful additions; for the belly ought here to be always kept open, though the stronger purgatives are very improper. A drachm of the electary may be taken once or twice a day, along with an infusion of marshmallow roots, sweetened with a spoonful of honey.

BLECTARIUM PARALTTICUM.

Paralytic electory.

Take of

Mustard seed,

Conferve of rolemary tops, each one onnce;

Compound spirit of lavender, two drachms.

Beat the mustard feed with a little water, that the pulp may be pressed through a hair sieve; then mix with it the conserve and the spirit.

This is a very efficacious medicine for paralytic difforders, fremors and numbres of the kimbs, the decays accompanying old age, and in all cases where the folids require to be stimulated, or sluggish stagnant juices to be put in motion. It ought to be taken every morning and evening, or oftener, to the bulk of a large nutmeg; with a glass of rich wine, or any proper julep, after it.

ELECTARIUM E CORTICE.
PERUVIANO.

Electory of Peruvian bark. Take of 1.

Peruvian bark, three ounces; Cafcarilla, half an ounce; Syrup of orange peel, a fufficient quantity.

2. Take

Take of

Peruvien bank, three ounces; Virginian inakeroot, one ounce; Symp of orange peel, a fufficient quartity.

3.

Take of

Peravias bark, three ounces; Grade falammoniae, 3drachme; Syrup of lemon juice, a fufficient quantity.

Take of

Peruvian bark, three ounces; Coloothar of vitriol, 6 drachme; Simple fyrup, a fufficient quantity.

5.

Take of

Peruvian bark, three outless;
Alum, one ounce;
Syrap of lemon juice, as much
as is sufficient.

6.

Take of

Extract of Peruvian bark, one ounce;

Extract of logwood,

Extract of liquorice, each half an ounce;

Mucilage of quince feels, as much as is sufficient to rectore the other ingredients into the consistence of an electary.

ALL these compositions are very elegant and efficacious in the intentions for which they are designed. The first is calculated for common intermittent severs, in the cure of which the virtues of the bark are greatly affisted by the cascarilla. The second and third are given in those intermittents, which happen in cachecia habits, and persons subject to obstructions of the viscera, where the bark by itself, on account of its great astringency, would be prejudicial. The sourch is a good strengthener in laxities of

the folids and decays of conflication; and the Afth, a powerful flyptic in fluxes and hamorrhages, particularly in the diabetes and fluor albus. The bulk of a nuture of each may be taken as a time, and repeated according to the exigency of the case. The fixth is a very agreaable form for the exhibition of Peruvius bank to those who are more than ordinarily offended with its taffe; the fubitunces here joined effectually covering its take, at the fame time that they coincide with it in virtue. The composition is a very elegant and pleafant one, and well deferves a place in the shope: it may either begiven in the form of a bolus or electary, in the dole of a drachm or more; or diffolved in any faitable liquor into a draught.

### RESCHARIUM PURGANS

An acid purgative electory.

Take of

Pulp of meaninds, swo dences; Crystals of tarter, two drachms. Make them into an electary.

This is an useful cooling laxative in hot billious dispositions, or inflammatory diseases, The bulk of a nutmeg may be taken every hour, or oftener, till it begins to operate, or the same quantity may be taken once a day occasionally in dry costive habits.

### ELECTARIUM SAPONACEUM. Saponaceous electary.

Take of

Hard Spanish soap, two ounces; Pareira brava, one ounce; Rhubarb.

Gum of aloes, each 3 drachms; Syrup of orange peel, a fufficient quantity.

Mix and make them into an electary.

THIS

This electary is calculated for jaundices arifing from an obstruction of the biliary ducts, or a vifcidity of the bile itself; such are those which most commonly occur, in which the stools are of a whitish or ash-colour, and voided with difficulty. The dose is from half a drachm to a drachm, twice a day. How far the pareira brava in this composition contributes to its virtues. I shall not take upon me to determine. Some have recommended this root, as a most powerful attenuant, in a great variety of diforders (see page 193.) whilst others look upon it as not superior, if equal, to the common aperient roots (page 249.) The sensible qualities of the pareira discover little foundation for the great character given of it; and a competency of fair trials of its virtue, is as yet The London college wanting. have not received it into their pharmacopæia.

ELECLARIUM SISTEMS.

Binding electory.

Take of

The japonic confection, two ounces;
Extract of logwood, one ounce;

Syrup of dry rofes, as much as will reduce them into a proper confifence for an electary.

This electary is calculated for the relief of dysenteries, and other intestinal sluxes, after the acrid humours have been duly evacuated by mild cathartics, &c. The quantity of a nutmeg may be taken every four or sive hours.

Electarium e sulphure.

Electary of Sulphur.

Take of

Flowers of fulphur, half an ounce;

Lenitive electary, two ounces; Syrup of marshmallows, a sufficient quantity to make them into an electary.

This electary is defigned against the piles, and generally distinguished in the hospitals by the title of electarium hamorrhoidale: where the disorder is accompanied with sebrile or inflammatory symptoms, some nitre is occasionally added, in the proportion of two drachms, to the quantity here directed. It may be given from a drachm to half an ounce at a time.

### CHAPTER VI.

### Lobochs.

Lohoch, eclegma, lindus, or lambative, is a fost compound defigned to be licked or flowly swallowed down, of a middle confistence between a syrup and electary, at least never so thin as the former, nor so thick as the latter.

I hese preparations are generally composed of expressed oils, mixed with fyrups, and other like fubstances. In making them, the syrup is first to be mixed with a little sugar, and then briskly beat up in a mortar, with the oil; which will thus readily incorporate, especially if the lyrup is of the acid kind Two ounces of syrup, a drachm of fugar, and an ounce of expressed oil, form a linctus of a due confiftence; which may be made thicker at pleasure by adding more oil. or thinner by an increase of the fyrup.

Any oily substance, as Locatelli's balsam, spermaceti, &c. may likewise be reduced into this form; and instead of sugar, powders more agreeable to the intention of emollients or pectorals, may be used; as the compound powder of gum tragacanth, or the white or black bechic troches of the shops. But the form at best is very unsightly and disagreeable, and substances of this kind render it more so.

#### LOHOCH COMMUNE.

Common lohoch.

Edinb.

Take of
Fresh-drawn oil of almonds,
Syrup of marshmallows, or balfamic syrup, each one ounce;
White sugar, two drachms.

Mix and make them into a lohoch.

## LOHOCH EX AMYLO. Starch lohoch. Edinb.

Take of
Starch, two drachms;
Japan earth, one drachm;
Balfamic fyrup,
Whites of eggs, beaten up into
a thin fluid, each one ounce.
Mix and make them into a lohoch.

## LOHOCH DE LINO. Lohoch of linfeed. Edinb.

Take of
Fresh-drawn linseed oil,
Balsamic syrup, each one ounce;
Flowers of sulphur, washed,
White sugar, each two drachms.
Mix and make them into a lohoch.

## LOHOCH DE MANNA. Lohoch of manna. Edinb.

Take of
Calabrian manna,
Fresh-drawn oil of almonds,
Syrup of violets, each equal
parts.
Mix and make them into a lohoch.

### LOHOCH SAPONACEUM.

Saponaceous lohoch.

Edinb.

Take of
Castile soap, one drachm;
Oil of almonds, one ounce;
Syrup of lemon juice, one ounce
and a half.
Mix and make them into a lohoch
according to art.

LOHOCH

LOHOCH DE SPERMACETI.

Lohoch of Spermacesi.

Edinb.

Take of
Spermaceti, two drachms;
Freth-drawn oil of almonds,
half an ounce;
Balfamic fyrup, one ounce.
Mix the foermaceti with a fuffi-

Balfamic fyrup, one ounce.

Mix the fpermaceti with a fufficient quantity of yolks of eggs, then add the oil and fyrup, and make them into a lohoch.

LOHOCH BALSAMICUM.

Balfamic lehoch.

Take of
Spermaceti, two drachms;
Balfam of Peru, one drachm;
Syrup of marshmallows, two
ounces.

Let the spermaceti and balsam be well worked up with a sufficient quantity of yolks of eggs; and then mix with them the syrup.

LINCTUS SOLUTIVUS.

Solutive lobech.

Take of
Conferve of hips, one ounce;
Solutive fyrup of rofes,
Oil olive, each four ounces.
Mix and make them into a lohoch.

THE principal efe of lohochs is in diforders of the internal parts of the mouth, fauces, and cefophagus; as in aphthæ, and tickling coughs from defluxions in the

first passages; for however they may have been celebrated, under the vague appellation of pectorals, in affections of the breast and lungs, it is not to be expected. that their emollient lubricating quality can reach those parts, or that they can give any relief in the true pulmonary cough. The flow manner in which they are fwallowed down renders them well adapted to correct acrimony and irritation in the throat and about the mouth of the stomach; though the free afe of fach uncluous compositions is soon liable to pall the appetite. Indeed the form is an inelegant one, and in the present practice is little regarded.

LINCTUS ACIDULUS.

Acidulous linctus.

Take of

Conserve of red roses, two

Weak spirit of vitriol, four feruples, or as much as is sufficient to give a grateful acidity.

Mix them together.

This lineaus is of a different nature from the foregoing preparations, and is used as a light restringent and detergent. It rather strengthens than relaxes the stomach, is sufficiently agreeable in taste, and of a fine red colour.

### CHAPTER VIL

### Emulsions.

IN the foregoing chapter, oils were united with watery liquors, by the mediation of fugar and fyrops, into thick unchaous compounds. The present chapter contains mixture of oily, resinous and other like bodies, with water, in a liquid form, of a white colour resembling milk, and hence called emulsions.

Emulsions have been generally prepared by grinding the oily seeds of plants, or kernels of fruits, along with common water, or any agreeable simple distilled water. In this process, the oil of the subject is, by the mediation of the other matter, united with the aqueous shuid; and hence they possess forme share of the emollient vitrue of the pure oil; with this advantage, that they are agreeable to the palate, and not apt to turn rancid or acrimonious by the heat of the body, which the pure oils in some instammatory cases may do.

Emulsions, besides their use as medicines themselves, are excellent vehicles for certain substances which cannot otherwise be so conveniently taken in a liquid form. Thus camphor, triturated with almonds, readily unites with water into an emulsion, and in this form is conveyed into the remotest parts of the body, with sufficient efficacy to answer intentions of moment, at the same time that its heat and pungency are softened by the unctuosity of the almonds.

Pure oils, balfams, refins, and cother fimilar fubfiances, are likewife rendered mifeible with water, into emultions or milky liquors, by the intervention of mucilages. The

white or yolk of an egg unites these bodies also with water, but less elegantly.

Several of the gummy refins, as ammoniacum, galbanum, myrrh, and others, are reducible into emultions by trituration with water alone; their refinous part being rendered dissoluble by the mediation of the gummy.

### EMULSIO COMMUNIS.

Common emulsion. Lond.

Take of

Sweet almonds blanched, one ounce:

Gum Arabic, half an ounce; Double-refined fugar,6 drachmes Barley water, two pints.

Diffolve the gum in the barley water warmed; as foon as the water is grown thoroughly cold, pour it by little at a time upon the almonds and fugar, first beat together, continuing to grind the whole, that the liquor may grow milky; after which, it is to be passed through a firginer,

#### Edinb.

Take of

Sweet almonds, one ounce; White fugar, two drachms; Simple cinnamon water, one ounce;

Common water, two pints.

Beat the almonds with the feeds in a marble mortar, and gradually pour on them the common water, working the whole well together. Then firain off the liquor, and add to it the cinnamon water and the fugar,

If half an ounce of gum Arabic be previously dissolved in the water, the preparation is called EMULSIO ARABICA, the Arabic e-malfum.

GREAT care should be taken. that the almonds are not become rancid by keeping; which will not only render the emulsion extremely unpleasant, a circumstance of great consequence in a medicine that requires to be taken in large quantities, but likewise give it injurious qualities little expected from preparations of this class. These liquors are principally made use of for diluting and obtunding acrimonious humours; particularly in heat of urine and stranguries arising either from a natural sharpness of the juices, or the operation of canthárides, or other irritating medicines: in these cases, they are to be drank frequently, in the quantity of half a pint or more at a time.

Some have ordered emulsions to be boiled, with a view to deprive them of fome imaginary crudity; but by this process they quickly cease to be emulsions, the oil separating from the water, and floating diffinct upon the furface. Acids, and vinous spirits, produce a like decomposition. On standing also for some days, without addition, the oily matter separates, and rises to the top, not in its pure form, but in that of a thick creme. These experiments prove the composition of the emultions made from the oily feeds of kernels, and at the same time point out some cautions to be attended to in their preparation and use.

## EMULSIO CAMPHORATA. Camphorated enulsion.

Edinb.

Take of Camphor, half a drachm; Sweet almonds, fix in number; White sugar, half an ounce; Simple pennyroyal water, half a pint,

Grind the camphor and almonds well together in a stone mortar, and add by degrees the pennyroyal water; then strain the liquor, and dissolve in it the sugar.

This is a very commodious form for the exhibition of camphor; the unctuous quality of the almonds in great measure covering its pungency. In fevers that require the assistance of this powerful diaphoretic drug, a spoonful of the emulsion may be taken every three or four hours.

# LAC AMMONIACI. Milk of ammoniacum.

Lond.

Take of

Gum ammoniacum, 2 drachms; Simple pennyroyal water, half a pint,

Grind the ammoniacum with the water, in a mortar, until the gum is dissolved.

This liquor is employed for attenuating tough phlegm, and promoting expectoration, in humoural althmas, coughs, and obstructions of the viscera. It may be given to the quantity of two spoonfuls twice a day.

## EMULSIO PURGANS. A purping emulsion.

A purging emulfion.

Take of
Sweet almonds blanched, two
drachms;
Fine fugar, one drachm;
Gum Arabic, half a drachm;
Scammony ten grains;
Simple cinnamon water, one
ounce.

Diffolve

Diffolve the gam in the cinnamon water, and having ground the scammony with almonds and sugar, pour on the liquor by little at a time, continuing to grind them together, so as to make them into an emulsion.

This emulsion is an agreeable and an effectual purgative. It may be prepared with different proportions of the scammony, at pleasure: other purgative refins, as that of jalap, may be fubilituted to the fcammony; a proper quantity of any fyrup to the fugar; and to the cinnamon water, any other simple water that may be more acceptable: but spirituous waters, for reafons already mentioned, have no place. Some have employed an infusion of liquorice, which appears to be a very proper addition in these kinds of preparations, as it coincides with the almonds in correcting the irritating power of the purgative material.

EMULSIO OLEOSA.

Oily emulsion.

Take of

Oil olive, a quarter of a pint; Spirit of hartshorn, 2 drachms; Simple pennyroyal water, twelve ounces;

Pectoral fyrup, an ounce and a half;

Mix them together.

This composition is often used against recent colds, for alleviating the cough, and promoting expectoration. Wherethecomplaints are of long standing, these kinds of medicines have no place; nor is their use in any case to be long continued, as they relax the stomach, pall the appetite, and increase the disorder.

A much more elegant oily emulfion, for all the intentions in which the simple lubricating quality of expressed oils is wanted, may be prepared in the following manner:

Take an ounce of powdered gum Arabic, and the same quantity of common water: diffolve the gum in the water, that it may form a thick mucilage; to which add by degrees four ounces of fresh-drawn oil of almonds, rubbing them well together in a mortar till they incorporate into a smooth white mass. Then pour in by little and little, continuing the agitation, four ounces of common water; to which may be added nutmeg water, rose water, and fimple fyrup, of each two ounces.

This appears to be the pleasantest form that oils can be given in. The union is also more perfect, and the oil less disposed to separate on standing, than in the emulsions obtained by other means. Even strong acids added to the emulsion, produce no decomposition in it. But alkalies can have no place in this form; for these, as we have observed upon another occasion, precipitate pure gums themselves from water.

Emulsio spermatis ceti.

Emulsion of Spermaceti.

Take equal parts of spermaceti and of mucilage of gum Arabic. Rub them together in a mortar till they are incorporated into a thick mass, which may be diluted at pleasure with water, as in the foregoing process.

EMULSIONS of spermaceti, or spermacetidraughts, are commonly prepared by means of yolks of eggs; and the emulsions, so prepared, are sufficiently uniform. Those made with mucilage, as here directed, have this advantage, that

they are less differentials in take, and not liable to grow rancid. The mixture of the spermateri and maxilage may be kept, for many days, in a state at forboing diluted, by gradual additions of water, into a smooth emultion.

Emplesso Cum are.

Empless count area root.

Take of

Frosh arem soot,
Gum Arabic, each 2 drachms;
Spermacesti, two scruples;
Common water, five onnces;
Numeg water,
Symp of orange peel, each half
an ounce.

Dissolve the gum Arabic, with part
of the water, into a mucilage,
which is to be beaten wish the

spermaceti into a smooth paste.

To this add the argin root, previously beaten by itself into a pulp; and rub them well together, that they may be thoroughly mixed. Then gradually pour in the waters and the fyrap.

FRESH arum root may be taken in this form without the least inconvenience from the pangency, with which the root itself to violently affects the mouth. I have given a spoonful of the emultion every fix boars, or oftener, in cases of the rheumatic kind, and generally with great benefit. The more immediate effect experienced from it was that of warming the stomach, and promoting sweet, which in some instances it did profusely.

### CHAPTER VIII

### Juleps, Mixtures, and Draughts.

PY julep is commonly understood, an agreeable liquor, defigned as a vehicle for medicines of greater efficacy, or to be drank after them, or to be taken occasionally as an auxiliary. In this light their basis is generally common water, or a simple distilled water, with one-fourth or one-third its quantity of a diffilled spirituous water: this mixture is sweetened with fugar, or any proper fyrup, or acidulated with vegetable or mineral acids, or impregnated with other medicines suitable to the intention; care being taken that these additions be such, as will not render the compound unfightly or unpalatable. The quantity usually directed at a time, in common prescription, is fix or eight ounces, to be taken by spoonfuls.

A mixture, more strictly fo called, receives more efficacious materials, whether foluble in water, as extracts or falts, or indisfoluble, as powders; more regard being here had to the medicinal intention. than to the fightliness or palatable-There is ness of the compound. indeed no precise distinction between the two; the same compofition being often called by one a julep, and by another a mixture; though in general, few would give the name of julep to a very difagreeable liquor, or that of mixture to a very pleafant one.

A draught differs from a julep or mixture only in being prescribed in less quantity, the whole being intended for one dose.

JULEPUM e CAMPHORA.

Julep of camphor.

Lond.

Take of
Camphor, one drachm;
Double refined fugar, half an
ounce;

Boiling water, one pint.

Grind the camphor first with a literestified spirit of wine, until it grows soft; and afterwards with the sugar, till they are perfectly mixed: then add the water by little and little, let the mixture cool in a close vessel, and lastly pass it through a strainer.

This is a more easy and effectual way of mingling camphor with aqueous liquors, than grinding it with water alone, or fetting it on fire, and then quenching it in water, as directed in our former dispenfatory, and in other books of pharmacy: though even this method is liable to some inconveniencies; part of the camphor exhaling, unless an extraordinary deal of care is taken, upon the affusion of the boiling water; and part remaining upon the strainer. The julep taftes strong of the camphor, and may be given, in cases where this drug is proper, in the dole of a spoonful or two. In extemporaneous prescription, vinegar is sometimes employed instead of water; this acid not only rendering the julep more grateful to the palate and flomach, but likewise promoting and extending the efficacy of the camphor, rendering it serviceable in some severs where that hot pungent medicine by itself would be less proper. In this view the following form is a very elegant one.

Rг

. Jule-

JULEPUM E CAMPHORA ACE-TOSUM.

Camphor julep with vinegar.

Camphor, one drachm; Gum Arabic, two drachms; Double refined fugar, half an ounce;

Vinegar, one pint.

Grind the camphor with a few drops of rectified spirit of wine, till it grows soft; then add the gum, previously reduced to a mucilage, with equalits quantity of water, and rub them together till they are persectly mixed. To this mixture add by degrees the vinegar with the sugar dissolved in it.

By this management, the whole fubfiance of the camphor is united with, and kept suspended in, the liquor; and consequently every spoonful of the mixture is equivalent to one grain and seven-eighths of a grain of camphor in substance. The same treatment succeeds equally when water is used for the menfiruum; and if the affistance of nitre is required, this also may be added in either form.

### JULEPUM e CRETA. Chalk julep. Lond.

Take of

The whitest chalk, prepared, one ounce;
Double, refined from: 6 drachms:

Double-refined fugar, 6 drachms; Gum Arabic, two drachms; Water, two pints.

Mix them together

This julep is defigned for heartburns and other like diforders arifing from acid juices in the first passages. The chief use of the gum is to give a greater degree of consistence to the water, and enable it to keep the powdered chalk suspended. JULEPUM e MOSCHO.

Musk julep.

Lond.

Take of

Damask-rose water, six ounces by measure; Musk, twelve grains; Double-resined sugar, 1 drachm;

Grind the fugar and the musk together, and gradually add to them the rose-water.

This is an improvement upon the Hysteric julep with Musk of Bates. Orange flower water is directed by that author; and indeed this more perfectly coincides with the musk than rose-water; but as the former is difficultly procurable in perfection, the latter is here pre-The julep appears turbid ferred. at first; on standing a little time, it deposits a brown powder, and becomes clear, but at the same time loses great part of its virtue. This inconvenience may be prevented. by thoroughly grinding the muse with two or three drachms of mucilage of gum Arabic, before the addition of the water, as directed in the preceding chapter for making emultions; by means of the gume the whole substance of the musk is made to remain suspended in the water. Volatile spirits are in many cases an useful addition to musk. and likewise enable water to keep fomewhat more of the musk diffolved, than it would otherwise retain. The following composition of this kind is used in some of our hospitals.

# Julepum moschatum. Mufk julep.

Take of

Rofe-water, fix ounces;
Volatile oily spirit, one drachm
and a half;
Muss. fisteen grains;
White sugar, halt an ounce.
Grind

## Chap. 8. Juleps, Mintures, and Draughts.

Grind the male with the fugar, and then mix therewith the other ingredicats.

JULEPUM ALEXIPHARMACUM. Mexiphermat julep.

Take of Simple alexeterial water, fix OUNCES :

Spirituous alexeterial water, two Syrup of cleve-july-flowers, two

drachms.

Mix them together.

Take of Simple alexeterial water, fix ounces; Spirituous alexeterial water with vinegar, two ounces; Syrup of lemon juice, two drachms. Mix them together.

> TULEPUM CARDIACUM. Cordiat julep.

· 1. Take of Simple cinnamon water, Simple orange peel water, each three ounces; Nutmeg water, two ounces; Syrup of orange pecl, half an Mix them together.

Take of Dill feed water, 6 ounces; Cardamom-seed water, 2 ounces; Compound spirit of lavender, Syrup of faffron, each 2 drachms. Mix them together.

TULEPUM CARMINATIVUM. Carminative julep.

Take of Fennel-feed water, fix ounces; Compound juniper water, two ounces; Syrup of clove-july-flowers, half an ounce.

Take of water, Jamaica-pepper ounces; Compound annifeed water, two ounces; Syrup of orange pect, half an ounce.

Take of Diff-feed water, fix ounces; Compound caraway water, two ounces; Syrup of ginger, half an ounce.

> TULEPUM HYSTERICUM. Hyfteric julep.

Take of Simple pennroyal water, Castor water, each three ounces; Spirituous pennyroyal water, two ounces; Simple fyrup, two drachms.

Take of Simple alexeterial water, ounces: Cardamom-feed water, ounces: Compound spirit of lavender, Volatile aromatic spirit, each one drachm: Syrup of clove-july-flowers, half an ounce.

3. Take of Dill-seed water, four ounces; Simple peppermint water, two Ounces; Ticture of cardamoms, Syrup of ginger, each 2 drachms.

JULEPUM REFRIGERANS. . A cooling julep.

Take of Rhenish wine, five ounces; Damaik rose water, two ounces; Seville orange juice, Syrup of violets, each 6 drachms;

R 1 2 TULEPUM TULEPUM STOMACHICUM. Stomachic julep.

Simple mint water, fix ounces;

Spirituous mint water, two ounces:

Syrup of faffron, two drachers: 2.

Take of

Tincture of mint, fix ounces; Cardamom water, two ounces; Simple fyrup, half an ounce.

Take of Cinnamon water, fix ounces; Nutmeg water, Stomachic tincture, each one Syrup of orange peel, half an

The titles of these mixtures express the intentions for which they are calculated: four or five spoonfuls of either may be taken occasionally, or used as vehicles and diluters

of medicines of greater efficacy. The following julapia are used in the Edinburgh infirmary.

JULAPIUM AMMONIACUM. Ammoniacum julep. Take of

Milk of ammoniacum, four ounces;

Syrup of squills, three ounces; Mix them together.

Two spoonfuls of this mixture may be given twice a day, in coughs, asthmas, and oppressions at the breast. It is a medicine of considetable efficacy, but not a little unpleasant, though called a julep in the hospitals where it is used.

JULAPIUM ANTIHYSTERICUM. Antibysteric julep. Take of

Pennyroyal water, four ounces; Compound valerian water, two ounces ;

Tincture of castor, 2 drachmez Salt of hartshorn, 10 grains; White sugar, 6 drachers. Mix them together.

THE virtues of this composition are fufficiently obvious from its title: the dose is two spoonfuls, to be taken twice or thrice a day.

> TULAPIUM CARDIACUM. Cordial julep.

Take of Alexeterial water, four ounces; Aromatic water, two ounces:

Volatile oily spirit, Tineture of fastron, each two drachms:

White fugar, half an ounce: Mix, and make them into a julep.

THIS mixture is an ufeful cordial in all depressions of the spirits, in the finkings of low fevers, and the languors, to which hysterical and hypochondriacal persons are sub-An ounce, or two sponfuls, may be taken for a dose, two or three times a day.

JULAPIUM DIAPHORETICUM. Diapboretic julep.

Alexeterial water, 4 ounces; Spirit of Mindererus, 2 ounces; Salt of hartshorn, 10 grains; White sugar, 6 drachms; Mix them for a julep.

This excellent composition is a very powerful sudorific, and anfwers its intention more effectually. and with greater certainty, than many others calculated for the same purpose. Where a copious sweat is to be excited, as in rheumatic diseases, two spoonfuls are to be taken warm in bed every hour, or two hours, till the sweat breaks out; if warm diluting liquors are not afterwards sufficient to keep it up, the same medicine is to be occasionally repeated,

JULAPIUM DIAPHORETICUM ACIDUM.

Acid diaphoretic julep.

Take of

Alexeterial water, four ounces; Treacle vinegar, two ounces; Tincure of faffron, half an ounce;

Spirit of amber, one drachm; White sugar, one ounce. Mix them together.

THE acid quality of this diaphoretic julep adapts it more particularly to those disorders in which any of the internal parts are inflamed, as in pleurisies and peripneumonies. It is given in the same dose as the preceding.

JULAPIUM DIURETICUM.

Diuretic julep.

Take of

Spirit of Mindererus, four ounces;
Compound horseradish water, two ounces;

Syrup of marfimallows, three ounces.

Mix them together,

THE spirit of Mindererus is an excellent aperient faline liquor, capable of promoting evacuation either by the cutaneous pores, or the prinary passages, according to the manner of exhibiting it. have seen above, that when taken warm in bed, it proves a powerful fudorific; especially if affilted by volatile falts, small doses of opiates, or other substances which tend to determine its action to the ikin. If the patient walks about, in a cool air, it operates gently, but for the most part effectually, by urine: the additions here joined to it, correspond with this intention, and promote its operation. As this medicine excites the urinary discharge, without heating or irritating the parts, it takes place not only in dropsies, but likewise in inflammatory disorders, wherever this salutary secretion is to be promoted. It is given to the quantity of two spoonfols, thrice a day.

A drachm of spirit of amber is sometimes mixed with this julep, which nevertheless does not seem to receive from that ingredient any additional virtue: whatever virtues the falt of amber may possess (which probably are not so great as is generally supposed) the spirit is impregnated therewith in an extremely low degree; see page 482.

JULAPIUM FORTIDUM.
Fatid julep.

Take of

Afafortida, one drachm and a

Rue water, fix ounces; Compound valerian water, two ounces;

Oil of hartshorn, twenty drops; White sugar, ten drachms.

Rub the afarcetida in the rue water till it disfolves, and having dropped the oil upon the fugar, mix the whole together.

THIS composition is not a little feetid and unfightly; it is nevertheless a medicine of great efficacy, in hypochondriacal and hysteric disorders, asthmas, and other nervous complaints; the dose is one spoonful, to be taken thrice a day. It is sometimes prepared without the oil of hartshorn.

JULAPIUM HYDRAGOGUM.

Hydragogue julep.

Take of

Chamomel-flower water, fix ounces;

Emetic tartar, ten grains;
R r 3 Syrup

Syrup of buckthorn, two ounces. Mix them together.

Two spoonfuls of this julep are given, in hydropic cases, every two hours, till it takes sufficient effect as a purgative; which it generally does before the quantity here prescribed has been made use of. Emetic tartar, thus exhibited in small doses, and frequently repeated, proves as certain and powerful a cathartic, as it does an emetic when given in a larger quantity at once. It operates nevertheless, for the most part, with sufficient ease.

JULAPIUM SISTENS.

Binding julep.

Take of

Alexeterial water, four ounces; Aromatic water, two ounces; Japonic confection, two drachms; Japan earth, in powder, one drachm;

Liquid laudanna, forty drops; White fugar, half an ounce. Mix them well together.

This julep is calculated against dyfenteries and diarrhoss; in which, after proper evacuations, it generally eases the gripes, and restrains the flux. It is to be given three or four times a day, in the quantity of a spoonful at a time.

## MISTURA ALEXETERIA. Alexeterial mixture.

Take of

Common water, four ounces; Spirituous alexeterial water with vinegar,

Julep of camphor, each one ounce and a half;

Compound powder of contrayerva, four scruples; Nitre, two scruples; Syrup of orange peel, 6 drachms; Mix them together.

In hospitals and places ill aired. common inflammatory fevers fometimes change into putrid and malignant ones. To guard against any accident of this kind, as foon as the inflammation begins to abate, or the pulse to loften, three or four spoonfuls of this alexipharmac mixture may be given every Camphor seems to fix hours. answer best when thus given in a liquid form; and to be most efficacious in such small doses, for abating inflammation and nervous fymptoms, and likewise for promoting a gentle diaphorens.

MISTURA ANTIDYSENTERICA.

Antidysenteric mixture.

Take of 1.

Simple cinnamon water, seven ounces;

Spirituous cinnamon water, one ounce;

Electary of fcordium with opium, half an ounce. Mix them together.

Take of 2.

Extract of logwood, three drachms;

Tincture of Japan earth, two drachms;

Spirituous cinnamon water, one ounce;

Common water, seven ounces.

Dissolve the extract in the cinnamon water, and then add the common water and the tincture.

In recent dysentaries, after the necessary evacuations, a spoonful or two of either of these mixtures may be given after every motion, or once in four or five hours: if the first, which is a mild opiate, fails of procuring rest, it is a sign that some of the corrupted humours fiill

fill remain in the bowels, and that it is more proper to go on with the evacuation, than to suppress the These medicines will sometimes likewise take place in the last flage of the disease, when through neglect or mismanagement it has continued till the strength is much impaired, the intestines greatly relaxed, and their villous coat abraded; provided there are neither ichorous or involuntary stools. apthz, petechiz, hiccup, or great anxiety at the breaft. Rhubarb, and these astringents, are to be so interposed, that at the same time that the putrid humours are diflodged, the strength may be supported, and the intestines braced. See Dr. Pringle's excellent observations on the diseases of the army, page 254, & seq. where the reader will find a full and fatisfactory history of the symptoms and cure of this distemper, so. frequent and fatal in the camp.

# MISTURA ANTIBMETICA

Saline antiemetic mixture.

Take of

Saltofwormwood, halfadrachm; Lemon juice, fix drachms; Simple cinnamon water, one ounce:

Fine fugar, one scruple. Mix them together.

This mixture is frequently prefcribed, not only for the purpole expressed in its title, but likewise as a saline aperient in icterical, inflammatory, and other disorders, where medicines of that class are proper.

MISTURA CARDIACA.

Cordial mixture.

Take of Simple cinnamon water, four ounces; Spirituous cinnamon water, 2
ounces;

Extract of faffron, 1 scruple; Confection of kermes, 6 drachms; Mix them together.

In great languors and depreffions, a spoonful of this rich cordial mixture may be taken every half hour.

MISTURA AD PHTHISIM.

Mixture against the phthis.

Take of

Balfam of Copaiba, a drachm;

Common water, 4 ounces;

Spirituous cinnamon water, I

ounce;

Syrup of orange peel, half an ounce.

Let the balfam be diffolved in a proper quantity of yolk of egg, and then mixed with the other ingredients.

Take of
Thebaic extract, one grain;
Conserve of roses, half a drachm;
Mix them together for a bolus.

Take of 3.
Oxymel of fquills, a drachm and a half;

Thebaic tincture, fifteen drops; Spirituous cinnamon water, two drachms;

Common water, two ounces. Mix them together.

In the advanced flate of a confumption, we may distinguish two forts of coughs, one occasioned by the ulcers, and the other by a thin rheum falling upon the fauces and trachea, which parts being then deprived of their mucus, become extremely sensible to irritation. It is this last kind, perhaps, which is most painful and teazing to the patient. The first fort requires balsamics, if the ulcer is open, and the matter can be expectorated.

nated. For this purpose, the first of the above mixtures is a very elegant and effectual formula; two spoonfuls are to be taken at a time, twice a day: if the balfam purges, two drachms of the paregoric elixir, added to the quantity of the mixture here prescribed, will prevent that effect, other kind of cough can only be palliated by incrassants; and for that purpose, the second of the above compositions is one of the most successful medicines; the conserve is altogether safe, and otherwise well adapted to the nature of the disease, but of weak virtues: the opiate extract is the most efficacious ingredient, but is to be given with great caution, as opiates in general are apt to heat, to bind the body, and to obstruct expectoration. As these bad qualities are in good measure corrected by squills; as soon as the patient begins to complain of restless nights from coughing, the third mixture may be given at bed-time. See Pringle's Observations on the diseases of the army.

MISTURA B VALERIANA.

Valerian mixture.

Take of

Simple pepper-mint water, twelve ounces; Wild valerian root, in powder.

one ounce; Compound spirit of lavender.

Compound spirit of lavender, half an ounce;

Syrup of orange peel, one ounce. Mix them together.

Wild valerian root, one of the principal medicines in epilepsies and vertigos, seems to answer better when thus exhibited in substance, than if given in form of tincture or insusance. The liquors here joined to it excellently coincide with, and by their warmth

and pungency greatly improve, its virtues. Two spoonfuls of the mixture may be taken twice or thrice a day.

HAUSTUS CATHARTICUS.

Cathartic draught.

Take of
Scammony, 10 grains;
Spirit of rosemary, 2 drachms;
Syrup of buckthorn, 6 drachms;
Grind the scammony with the spirit in a glass mortar, and when persectly incorporated, mix in the syrup.

Take of

Jalap, in powder, one fcruple;
Ipecacoanha, three grains;
Compound juniper water, one
ounce;
Infusion of linfeed, an ounce
and a half;
Simple syrup, one drachm.
Mix them together.

Both these compositions are strong cathartics, yet for the most part easy and safe in operation. They are calculated chiesly for hydropic cases, in which they procure copious evacuations, without weakening or fatiguing the patient so much as many other medicines of this kind.

HAUSTUS CATHARTICUS
SALINUS.

Saline cathartic draught.

Take of

Glauber's cathartic falt, Manna, each fix drachms; Boiling water, three ounces; Tincture of cardamoms, one drachm.

Diffolve the falt and manna in the water, and having strained off the liquor, add to it the tincture of cardamoms.

THIS.

This is a very elegant and agreeable faline purgative. Tincture of cardamoms is one of the best additions to liquors of this kind, or to the purging mineral waters, for rendering them acceptable to the stomach.

HAUSTUS DIAPHORETICUS.

Diaphoretic draught.

Take of
Spirit of Mindererus,
Syrup of meconium, each half
an ounce;
Salt of harthorn, five grains.

Mix them together.

This draught is a very powerful faline diaphoretic. It is given with fafety, and often with great benefit, in the beginning of inflammatory fevers, after bleeding; where theriaca, and other warm substances usually employed, if they fail in bringing out a sweat, increase the fever.

HAUSTUS DIURETICUS.

Diuretic draught.

ke of ...

Take of

Oxymel of fquills, one drachm
and a half;
Simple cinnamon water, one
ounce;
Compound spirit of lavender,
Syrup of orange peel, each one
drachm.
Mix them together.

Take of
Vinegar of fquills, one drachm
(or one drachm and a half;)
Saltofwormwood, halfadrachm;
Lemon juice, fix drachms;
Simple cinnamon water, an
ounce and a half;
Spirituous pepper-mint water,
half an ounce;
Syrupoforange peel, one drachm.
Let the falt of wormwood and
lemon juice be first mixed toge-

ther, and then add to them the other ingredients.

Take of
Diuretic falt, two scruples;
Oxymel of squills, one drachm
by measure;
Water, an ounce and a half.
Mix them together.

Take of
Tincture of cantharides, fifteen
drops;
Saltofwormwood, halfadrachm;
Lemon juice, fix drachms;
Simple pennyroyal water, an
ounce and a half;
Simple fyrup two drachms.
Mix them together-

THE two first of these elegant and efficacious compositions are commended by Dr. Mead, for promoting urine in hydropic cafes. He directs them to be taken every night or oftener, according to the urgency of the symptoms. fquill, one of the most powerful diuretics, is, by the additions here joined to it, rendered not only more grateful to the palate and stomach, but likewise enabled more effectually to answer the purposes intended by it. The other two are taken from our hospitals; in which the former, composed on the same plan with the two preceding, is justly distinguished by the title of mitior or milder; and the latter, containing besides the faline matter, a moderate dose of cantharides, by that of furtion or stronger.

HAUSTUS ANODYNO-DIURE-TICUS.

An anodyne diuretic draught. Take of

Ley of tartar, half a drachm; Thebaic tincure, forty drops; Pepper-mint water, one ounce; Simple Simple cinnamon water, half an ounce;
Spirituous cinnamon water, two drachms;
Syrup of marshmallows, one drachm.
Mix them together.

THOUGH practitioners have rarely ventured to exhibit opium in dropfies, yet, in those which are accompanied with great pain, this anodyne drug, by easing the pain, and removing the firsture of the passages, which painful sensitions always occasion, proves a medicine of great service, and notably promotes the urinary discharge. Dr. Mead has given a remarkable instance of the good

effects of the mixture above prescribed, in a person labouring under an ascites and tympany at the same time, where the pain was intolerable, the thirst intense, and the urine in very small quantity: the stronger purgatives increased the distemper; soap, alkaline salts, nitre, and other diuretics, were tried in vain: this draught (when the patient seemed to be beyond any affistance from medicine) procured unexpected relief, not only a gentle sleep, and truce from the pain, but likewise a copious discharge of urine. By repeating the medicine for a little time, every eight hours, and afterwards using corroborants, the cure was perfectly completed.

### CHAPTER

### Lotions, Gargarisms, Injections, &c.

AQUA ALUMINOSA BATEANA. Bates's alum water. Lond.

AKE of Alum. White vitriol, each half an ounce: Water, two pints. Boil the falts in the water till they

are dissolved, let the solution fettle, and afterwards filter it through paper.

BATES directs the falts to be full powdered and melted over the fire; but this is needless trouble, fince the melting only evaporates the ageous parts, which are seflored again on the addition of the This liquor is used for cleanfing and healing ulcers and wounds; and for removing cutaneous eruptions, the part being bathed with it hot, three or four times a day. It is fometimes likewife employed as a collyrium; and as an injection in the gonorrhoea and fluor albus, when not accom-Panied with virulence.

#### AQUA ALUMINOSA. Ahım water. Edinb.

Take of Corrofive mercury fublimate, Alum, each two drachms; Water, two pints. Let the sublimate and alum be ground into powder, and boiled

with the water, in a glass vessel,

to the confumption of half the

water; then suffer the liquor to fettle, and pour it off clear from the fediment.

This is taken from Fallopius, with the exchange of role and plantane waters for common water, which is equally fit for the purpose. The composition is defigned chiefly for cutaneous puffules and ulcerations.

#### AQUA SAPPHIRINA. Sapphine coloured water. Lond.

Take of Lime-water, one pint; Sal ammoniac, one drachm. Let them stand together, in a copper vessel, or along with some plates of copper, until the liquor has acquired a sapphire colour.

#### Edinb.

Take of Lime-water, newly made, one Sal ammoniac, two drachms.

Dissolve the falt in the lime water. and let the folution stand in a brass vessel, until it has acquired a blue colour.

This water is at present pretty much in use, as a detergent of foul and obstinate ulcers, and for taking away specks or films in the The copper contributes eyes. more to its colour, than to its medicinal efficacy: for the quantity of the metal dissolved is extremely minute.

AQUA

### AQUA VITRIOLICA CÆRULEA.

Blue vitriolic water. Land.

Take of Blue vitriol, three ounces;

Strong spirit (or oil) of vitriol, each two ounces; Water, a pint and a half.

Boil the falts in the water, until they are diffolved, then add the acid spirit, and filter the mixture through paper.

#### AQUA STYPTICA. Styptic water. Edinb.

Take of Blue vitriol, Alum, each three ounces; Water, two pints. Boil them until the falts are diffolved, then filter the liquor, and add two drachms of oil of vitriol.

THESE compositions are formed upon the flyptic, recommended by Sydenham, for stopping bleeding at the nose, and other external hæmorrhages: for this purpose, cloths or doffils are to be dipped in the liquor, and applied to the part.

### AOUA VITRIOLICA CAMPHORATA.

Campborated vitriolic water. Lond.

Take of

White vitriol, half an ounce; Camphor, two drachms; Boiling water, two pints. Mix them, that the vitriol may be dissolved; and after the feces have subsided, filter the liquor through paper.

Tris is an unfrugal method of managing camphor, the greatest part

of which separates with the seces of the vitriol, very little of it remaining suspended in the water. The Edinburgh college, in the preceding edition of their pharmacopœia, had a preparation under the title of AQUA OPHTHALMI-CA, differing little otherwise from the above than in the quantity of water being greater, and in an addition of tutty and bole, ingredients which could be of no use, as not being soluble in water, and subfiding from it in standing. They have therefore, at the late revifal of their pharmacopæia, reduced this preparation to the following more simple form.

### AQUA VITRIOLICA.

Vitriolic water. Edinb.

Take of White vitriol, two drachms; Water, two pints. Boil till the vitriol is dissolved, and then filter the liquor.

WHERE the eyes are watery or inflamed, these solutions of white vitriol are very useful applications: the flighter inflammations will frequently yield to this medicine, without any other affiftance; in the more violent ones, venæsection and cathartics are to be premised to its use.

### AQUA PHAGEDÆNICA. Phagedenic water. Edinb.

Take of Lime-water, one pint; Corrofive mercury sublimate, half a drachm; Let a folution be made.

THIS is defigned for washing and cleanfing old foul ulcers, and preventing the growth of fungous stesh. It is for most purposes ra-

Chap. 9. Lotions, Gargarisms, Injections, &c. 621 ther too acrid to be used without dilution.

GARGARISMA ASTRINGENS. Astringent gargarism.

Take of Oak bark, one ounce; Alum, one drachm; Honey of roses, one ounce; Water, a pint and a half. Boil the water with the oak bark, till such time as the liquor, when strained, will amount only to one pint; to which add the alum and the honey. GARGARISMA COMMUNE.

Common gargarism. Take of Tincure of roses, one pint; Honey of roses, two ounces. Mix them together.

Take of Water, fix ounces; Nitre, one drachm; Honey of roles, one ounce. Mix them together. Where acids are requifite, forty drops of the weak spirit of vitriol are added

to this composition. GARGARISMA DETERGENS. Detergent gargarism. Take of

Emollient decoction, one pint; Tincture of myrrh, one ounce; Honey, an ounce and a half. Mix them together.

GARGARISMA EMOLLIENS. Emollient gargarism. Take of

Marshmallow root, two ounces; Figs, four in number; Water, three pints. Boil them till one pint is wasted, and then strain the liquor.

THESE liquors are used for washing the mouth and fances; the first, where the parts are extremely relaxed; the fecond and

third, where ulcerations require to be deterged, or the excretions of thick viscid saliva promoted; and the fourth, where the mouth is dry, parched and rigid, to moissen and fosten it. In some cases, volatile spirits may be advantageously joined to these kinds of preparations. Dr. Pringle informs us, that in the inflammatory quinfey, or strangulation of the fauces. he has observed little benefit arising from the common gargles; that fuch as were of an acid nature feemed to do more harm than good, by contracting the emunctories of the faliva and mucus, and thickening those humours: that the decoction of figs in milk and water seemed to have a contrary effect, especially if some spirit of sal ammoniac was added, by which the faliva was made thinner, and the glands brought to fecrete more freely; a circumflance always conducive to the cure.

> ENEMA DE AMYLO. Starch glyster.

Take of Gelly of starch, four ounces; Linfeed oil, half an ounce.

Liquefy the gelly over a gentle fire, and then mix in the oil. Forty drops of liquid laudanum are sometimes added.

ENEMA ANODYNUM, five OPIATUM.

Anodyne, or opiate glyfter. Take of

Infusion of linseed, six ounces; Liquid laudanum, forty drops.

Mutton broth, five ounces; Thebaic extract, three grains.

ENEMA ANTICOLICUM. Glyster against the colic. Take of

Common decoction, half a pint; Tinctura Tinctura facra, one ounce; Common falt, one drachm; Linseed oil, two ounces. Mix them together.

ENEMA ASTRINGRUS.

Aftringent glyfter.

Take of
Lime-water, ten ounces;
Japonic confection, half an
ounce.

Mix them together for a glyfter, of which one half is to be injected at a time-

ENEMA ASTRINGENS
BALSAMICUM.

Aftringent balfamic glyfter.

This is made by adding to the foregoing half an ounce of Locatelli's balfam, dissolved in the yolk of an egg.

Enema commune.

Common glyster.

Take of
Common decoction, twelve
ounces;
Lenitive electary, one ounce;
Common falt, half an ounce;
Oil olive, two ounces.
Mix them together.

Enema domesticum.

Domestic glyster.

Take of
Cows milk, half a pint;
Brown fugar,
Oil olive, each one ounce.
Mix them together.

Enema emolliens.

Emollient glyfter.

Take of
Palm oil, an ounce and a half;
Cows milk, half a pound.

Let the oil be beat up with the yolk of one egg, and then add the milk. ENSMA POETIDUM.
Fætid glyfter.

Take of
Afafætida, two drachms;
Rue,
Savin, each half an ounce;
Oil olive, one ounce;
Oil of amber, half a drachm;
Water, one pint and a half.

Boil the water with the rue and favin, till half a pint is wasted, then strain off the remaining decoction, and mix with it the associate and the oils. Half the quantity of the composition here directed, is to be injected at a time.

Busha Purgams.

Purging glyster.

Take of
Common decoction, half a pint;
White foap, one ounce;
Syrup of buckshorn, an ounce
and a half.
Mix them together.

EMEMA TERBBINTHINATUM.

Turpentine glyfter.

Take of

Common decoction, ten ounces; Venice turpentine (diffolved in the yolk of an egg) half an ounce;

Linfeed oil, one ounce. Mix them together.

THE uses of these compositions are sufficiently obvious from their titles. The starch, anodyne, emollient, and astringent glysters, are used in dysenteries, and other alvine sluxes, to strengthen the tone of the intestines, defend them from being corroded by the acrimonious humours, to heal their exulcerations, and ease the pains which accompany these disorders. The turpentine glyster is injected in perhritic cases; the section hysteric ones. The others are calculated

### Chap. 9. Lotions, Gargarisms, Injections, &c. 623

calculated for unloading the intestines of their contents, where the exhibition of purgatives in other forms is improper, or unsafe. Glyffers have been looked upon by fome as mere topical applications, whose operation was confined to the intestine, into which they are received. But experience has fhewn, that in many cases their action is extended much farther: thus the turpentine glyster, above described, promotes the discharge by the kidneys, and communicates to the prine a violet smell; and the anodyne glyster proves narcotic, as if a moderate dose of opium had been swallowed: persons have been inebriated by spirituous glysters; and some affirm, that life has been supported, for several days, by those of a nutritious kind.

INJECTIO BALSAMICA.

Balfamic injection.

Take of

Balfam of Copaiba, half an ounce;
Lime-water, fix ounces;
Honey of rofes, two ounces.
Let the balfam be well beaten up with the yolk of one egg; and then gradually add the limewater and honey.

INJECTIO MERCURIALIS.

Mercurial injection.

Take of
Quickfilver,
Balsam of Copaiba, each half an
ounce;
Rose-water, half a pint.

Rub the quickfilver with the balsam, till they are perfectly incorporated; then mix with them
the yolk of an egg, and afterwards add the rose-water.

This and the foregoing preparation are defigned to be injected into the urethra in virulent gonor-rheas, for cleaning and deterging the parts.

### CHAPTER X.

### Plasters.

PLASTERS are composed chiefly of oily and unctuous subflances, united with powders, into such a consistence, that the compound may remain firm in the cold, without slicking to the singers: that it may be soft and pliable in a small heat; and that by the warmth of the human body it be so tenacious, as readily to adhere both to the part on which it is applied, and to the substance on which it is spread.

There is however a difference in the confittence of plasters, according to the purposes they are to be applied to: thus, such as are intended for the breast and stomach, should be very foft and yielding; whilst those designed for the limbs are made firmer and more adhefive. An ounce of expressed oil, an ounce of yellow wax, and half an ounce of any proper powder, will make a plaster of the first confistence; for a hard one, an ounce more of wax, and half an ounce more of powder may be added. Plasters may likewise be made of refins, gummy-refins, &c. without wax, especially in extemporaneous prescription: for officinals, these compositions are less proper, as they soon grow too foft in keeping, and fall flat in a warm air.

It has been supposed, that plasters might be impregnated with the specific virtues of different vegetables, by boiling the recent vegetable with the oil employed for the composition of the plaster. The coction was continued till the herb was almost crisp, with care to

prevent the matter from contracting a black colour: after which the liquid was ftrained off, and fet on the fire again till all the aqueous moisture had exhaled. We have already observed, that this treatment does not communicate to the oils any very valuable qualities even relative to their use in a shuid state: much less can plasters, made with such oils, receive any considerable efficacy from the herbs.

Calces of lead, boiled with oils, unite with them into a plaster of an excellent confishence, and which makes a proper basis for several

other plasters.

In the boiling of these compofitions, a quantity of water must be added, to prevent the plaster from burning and growing black. Such water, as it may be necessary to add during the boiling, must be previously made hot: for cold liquor would not only prolong the process, but likewise occasion the matter to explode, and be thrown about with violence, to the great danger of the operator; this accident will equally happen upon the addition of hot water, if the plaster is extremely hot.

### EMPLASTERUM ANODY-NUM.

Anodyne plaster. Edinb.

Take of
White refin, eight onnces;
Tacamahaca, in powder,
Galbanum, each four ounces;
Cummin feeds, three ounces;
Black

Black foap, four ounces.

Melt the refin and the gums together; then add the powdered feeds and the foap, and make the whole into a plaster.

This platter fometimes gives ease in slight rheumatic pains, which it is supposed to effect by preventing the afflux of humours to the part, and putting in motion and repelling such as already stagnate there.

### EMPLASTRUM ANTIHYSTERICUM. Antihysteric plaster:

Edinb.

Take of

Galbanum, twelve ounces;
Tacamahaca, in powder,
Yellow wax, each fix ounces;
Venice turpentine,
Cummin feeds in powder,
Afafœtida, each four ounces.
Mix and make them into a plaster.

This plaster is applied to the umbilical region, or over the whole abdomen, in hysteric cases, and sometimes with good effect.

### EMPLASTRUM ATTRAHENS. Drawing plaster.

Lond.

Take of
Yellow refin,
Yellow wax, each three pounds;
Tried mutton fuet, one pound.

Melt them together, and, whilft
the mass remains fluid, pass it
through a strainer.

This is a very well contrived plaster for the purpose expressed in its title. It is calculated to supply the place of melilot plaster, whose great irritation, when employed for the dressing of blisters, has been continually complained

This was owing to the large quantity of refin contained in it, which is here, for that reason, retrenched. It should seem that, when defigned only for dreffing blifters, the resin ought to be intirely omitted, unless where a continuance of the pain and irritation, excited by the vesicatory, is required. Indeed plasters of any kind are not very proper for this purpose: their consistence makes them fit uneasy, and their adhesiveness renders the taking them off painful. Cerates, which are foster and less adhesive, appear much more eligible: the ceratum album will ferve for general use; and for some particular purposes the ceratum citrinum may be applied.

# EMPLASTRUM CEREUM. Wax plaster. Edinb.

Take of
Yellow wax, four pounds;
White refin,
Mutton suet, each two pounds.

Mutton fuet, each two pounds.

Melt them together into a plaster,
which supplies the place of melilot plaster.

This plaster is similar to the foregoing; but the further reduction of the resin renders it for some purposes more eligible.

### EMPLASTRUM CEPHALICUM. Cephalic plaster. Lond.

Take of

Burgundy pitch, two pounds; Soft labdanum, one pound; Yellow resin,

Yellow wax, each four ounces; The expressed oil, called oil of mace, one ounce.

Melt the pitch, refin, and wax together; then add, first, the S f labdanum,

labdanum, and afterwards the oil of mace.

Edinb.

Take of

Tacamahaca in powder,

Yellow wax,

Venice turpentine, each four ounces;

Oil of lavender, two drachms; Oil of amber, one drachm.

Melt the tacamahaca with the wax, and then add the turpentine, that a plaster may be formed: when this compound is taken from the fire, and grown almost cold, mix in the oils.

THESE plasters are applied, in weakness or pains of the head, to the temples, forehead, &c. and fometimes likewise to the feet. Schulze relates, that an inveterate rheumatism in the temples, which at times extended to the teeth, and occasioned intolerable pain, was completely cured in two days by a plaster of this kind, (with the addition of a little opium) applied to the part, after many other remedies had been tried in vain. He adds, that a large quantity of liquid matter exuded. under the plaster, in drops, which were so acrid as to corrode the cuticle.

EMPLASTRUM de CICUTA cum AMMONIACO. Plaster of bemlock, with am-

> moniacum. Edinb

Take of

Juice of hemlock leaves, four

ounces;

Gum ammoniacum, 8 ounces; Vinegar of squills, as much as is sufficient to dissolve the gum.

fufficient to dissolve the gum.

Add the juice to this solution, and having strained the mixture, boil it to the consistence of a plaster.

This plaster was formerly supposed to be a powerful cooler and discutient, and to be particularly ferviceable against swellings of the spleen, and distensions of the hypochondres For fome time past, it has been, among us, entirely neglected; and hence the London college, at the late revisal of their pharmacopæia, omitted it. But the high resolvent power which Dr. Storck has discovered in hemlock, and which he found it to exert in this as well as in other forms. entitle it to further trials. The plaster appears very well contrived, and the additional ingredients well chosen for affishing the efficacy of . the hemlock.

EMPLASTRUM COMMUNE.

Common plaster, usually called

Diachylon.

Lond.

Take of

Oil olive, one gallon:

Litharge, ground into a most Tubtile powder, sive pounds. Boil them over a gentle sire, with about two pints of water, keeping them continually stirring, till the oil and litharge unite, and acquire the consistence of a plaster. If all the water should be consumed before this happens, add some more water previously made hot.

Edinb.

Take of
Oil olive, fix pints;
Litharge, three pounds.
Boil them into a plaster.

THE heat in these processes should be gentle, and the matter kept continually stirring, otherwise it swells up, and is apt to run over the vessel. If the composition proves discoloured; the addition

addition of a little white lead and oil will improve the colour.

These plasters are the common applications in excoriations of the Kin, flight fiesh wounds, and the like. They keep the part foft, and fomewhat warm, and defend it from the air, which is all that can be expected, in these cases, from any plaster. Some of our industrious medicine-makers have thought these purposes might be answered by a cheaper composition, and accordingly have added a large quantity of common whiting and hogs lard: this, however, is by no means allowable, not only as it does not flick to well. but likewise as the lard is apt to grow rancid and acrimonious. The counterfeit is diffinguishable by the

## EMPLASTRUM COMMUNE ADHÆSIVUM.

Common flicking plaster.

Lond.

Take of

eye.

Common plaster, three pounds; Yellow refin, half a pound.

Melt the common platter over a very gentle fire; then add the refin, first reduced into powder, that it may melt the fooner; and mix them all together.

This plaster may otherwise be made, by taking, instead of the common plaster, its ingredients, oil and litharge, and adding the refin a little before they have come to the due consistence; then continue the boiling, till the plaster is finished.

It turns out the most elegant when made by this last method.

EMPLASTRUM ADHÆSIVUM. Sticking plafter. Edinb.

Take of

Common platter, two pounds;
Burgundy pitch, one pound.
Melt them together, so as to make
a platter.

THESE plasters are used chiefly as adhesives, for keeping on other dressings, &c.

# EMPLASTRUM COMMUNE com GUMMI.

Common plaster with gums.

Lond.

Take of

Common plaster, 3 pounds;
Galbanum strained, 8 ounces;
Common turpentine,

Frankincense, each 3 ounces.

Melt the galbanum with the turpentine over a gentle fire, and sprinkle in the frankincense, reduced to powder: then gradually mix with these the common plaster, previously liquested by a very gentle heat.

Or, inflead of the common plaffer already made, you may take the oil and litharge boiled together: as foon as these unite, before they have acquired the consistence of a plaster, the other ingredients are to be added.

### EMPLASTRUM GUMMOSUM. Edinb.

Take of

Palm oil, four pints; Litharge, 1 pound and a half; Gum ammoniacum, Galbanum, Venice turpentine,

Yellow wax, each half a pound. Boil the oil with a litharge to the confidence of a plaster; then add the other ingredients, and make the whole into a plaster, according to art.

Sf2 Both

BOTH these platters are used as digestives and suppuratives; particularly in absection, after a part of the matter has been maturated and discharged, for suppurating or discussing the remaining hard part.

EMPLASTRUM CROCEUM,
vulgo OXYCROCEUM.
Saffron plaster, commonly called
Oxycroceum.
Rdinb.

Take of
Burgundy pitch,
Yellow wax, each one pound;
Galbanum,
Tar, each half a pound;
Saffron rubbed into powder, two ounces.

Let the Burgundy pitch, wax, and galbanum, be melted together over a gentle fire; then add the tax and faffron, and make the whole into a plaster.

This infrugal and injudicious composition is said to strengthen the parts to which it is applied, especially the tendinous ones; to warm in a great degree; and to resolve and discuss cold tumours. Tar is now introduced as an ingredient, in the room of Venice turpentine, myrrh, and olibanum.

# EMPLASTRUM e CYMINO. Cummin plaster. Lond.

Take of
Burgundy pitch, three pounds;
Yellow wax,
Cummin feeds,
Caraway feeds,
Bay berries, each three ounces.
Melt the pitch with the wax;
then sprinkle in the other ingredients, first reduced into a powder, and mix the whole well together.

This plaster stands recommended as a moderately warm discutient; and is directed by some to be applied to the hypogastric region, for strengthening the viscera, and expelling statulencies.

#### EMPLASTRUM DEFENSIVUM. Defensive plaster. Edinb.

Take of
Litharge, two pounds;
Oil olive, four pints;
Yellow wax,
Olibanum, in powder,
Venice turpentine, each four
ounces;

Colcothar of vitriol, fix ounces. Boil the oil with the litharge, till they have acquired nearly the confishence of a plaster: in this liquefy the wax, and then add the other ingredients, so as to form the whole into a plaster according to art.

This plaffer is laid round the lips of wounds and ulcers, over the other dreffings, for defending them from inflammation, and a fluxion of humours; which, however, as Mr. Sharp very juftly observes, plaffers, on account of their confidence, tend rather to bring on than to prevent.

# Emplastrum e meliloto. Melilot plaster. Take of

Melilot leaves, fresh, fix pounds:

Beef fuet, three pounds;
White refin, eight pounds;
Yellow wax, four pounds.
Boil the herb in the melted fuet till
it is almost crisp; then strongly
press out the suet, and adding
the refin and wax, boil the
whole a little, so as to make a

plaster thereof.

THIS plafter has been frequently made use of for dreffing blifters. (See Emplastrum attrahens) The London college have diminished the quantity of refin, to render the composition less irritating; and likewise omitted the herb, as being of no fignificancy towards the use of the plaster, and of a very disagreeable scent; a circumstance of primary consequence to be avoided in disorders, where freedom from disturbance, and every means that can contribute to quiet rest, ought, by all possible endeavours, to be procured: not to mention the mifchievous adulterations sometimes practised in this platter with irritating materials, for procuring the green colour, which is made its marketable characteristic, more compendiously than by the decoction of the herb. The most certain method of discovering abuses of this kind, is to put a little of the plaster into some spirit of sal ammoniac: if it tinges the spirit blue, we may be certain it is adulterated. The London college has substituted to this plaster the emplastrum attrahens, and the Edinborgh the emplastrum cereum.

EMPLASTRUM ex AMMO-NIACO cum MERCURIO. Plaster of ammoniacum with mercury. Lond.

Take of

Gum ammoniacum, strained, one pound;

Quickfilver, three ounces; Simple baliam of sulphur, one drachm.

Grind the quickfilver with the balfam of fulphur, till it ceases to appear; then, having melted the ammoniacum, add it gradually, a little before it cools, to this mixture; and let the whole be perfectly mingled together.

This is a very well contrived mercurial platter: if in fome cases it should not prove adhesive enough, the addition of a small quantity of turpentine will readily make it so.

EMPLASTRUM COMMUNE cum MERCURIO.

Common plaster with mercury.

Lond.

Take of
Common plaster, one pound;
Quicksilver, three ounces;
Simple balfam of sulphur, one
drachm;

Make them into a platter, after the same manner as the foregoing.

EMPLASTRUM
MERCURIALE.
Mercurial plaster.
Edinb.

Take of Gum platter, a pound and a half;

Quickfilver, eight ounces; Venice turpentine, one ounce; Liquid florax, an ounce and a half.

Grind the anickfilver in a mortar, with the turpentine and storax, until they are perfectly incorporated; and then, having melted the gum plaster, and taken it from the fire, add to it this mixture.

THESE mercurial plasters are looked on as powerful resolvents and discutients, asting with much greater certainty in these intentions, than any composition of vegetable substances alone; the mercury exerting itself in a confiderable degree, and being sometimes introduced into the habit, in such quantity as to affect the Sf3 mouth.

limbs from a venereal cause, nodes, tophs, and beginning indurations of the glands, are faid fometimes to yield to them.

### EMPLASTRUM e MINIO. Red lead plaster. Lond.

Take of

Oil olive, four pints; Red lead, reduced to a most subtile powder, two pounds and a half.

Make them into a plaster, after the manner directed for preparing the common plaster: but more water is here required, and greater care is necessary to prevent the composition from burning and growing black.

This is used for the same purposes as the common or diachylon plaster, from which it differs little Otherwise than in colour. It has an inconvenience of not sticking so well; and therefore the Edinburgh college have now omitted this compolition.

### EMPLASTRUM DE MINIO CUM SAPONE.

Red lead plaster with foap.
This is made by adding to the foregoing plaster taken from the fire as foon as the moisture is evaporated, and whilft hot, half a pound of Spanish soap cut into thin lices: ftir the whole Arongly together, until the foap is liquefied, and a plaster formed according to art.

THIS is much esteemed by some, for discussing gouty tumours, and the juices flagnating after fprains. Whatever virtues it may have distinct from the general ones of the applications of this class, they depend entirely upon the foap; and

Pains in the joints and foap, in the form of plasters, does not appear to exert much of the efficacy which it does in forms of a fofter confiftence.

### EMPLASTRUM e MUCILAGINIBUS. Plaster of mucilages. Lond.

Take of

Yellow wax, forty ounces: Oil of mucilages, half a pint; Gum ammoniacum, strained, half a pound;

Common turpentine, 2 ounces. Melt the ammoniacum with the turpentine; and having, in another vessel, liquefied the wax . with the oil, add this latter mixture to the other.

Some have been accustomed to use, instead of the oil of mucilages, common oil olive, flavoured with fenugreek feeds; and possibly this substitution may be admitted as a venial one; for the oil of mucilages, genuinely made, contains fcarce any thing of any of the ingredients, except that part of the fenugreek feeds wherein their flayour resides, the mucilaginous materials serving only to provide it with a name. See p. 330.

### EMPLASTRUM ROBORANS. Strengthening plafter.

Take of

Common plaster, two pounds: Frankincense, half a pound; Dragons blood, three ounces.

Melt the common plaster, and add to it the other ingredients reduced into a powder.

The dragons blood should be reduced to a very fine powder, otherwise the mixture will not be of an uniform colour.

Tris is a reformation of the laborious and injudicious composition described

described in our preceding pharmacopæias, under the title of EM-PLASTRUM AD HERNIAM; and though far the most elegant and Simple, is as effectual, for that purpose, as any of the medicines of this kind. If constantly worn, with a proper bandage, it will, in children, frequently do fervice; though perhaps not fo much from any Arengthening quality of the ingredients, as from its being a fost, close, and adhefive covering. has been supposed, that plasters composed of fly pric medicines, conftringe and strengthen the part to which they are applied; but on no very just foundation: for plasters in general relax, rather than aftringe, the uncluous ingredients necessary in their composition, counteracting and deftroying the effect of the others.

### EMPLASTRUM e SAPONE.

Soap plaster. Lond.

Take of

Common plaster, three pounds; Hard foap, half a pound,

Having melted the common plaster, mix with it the foap, and boil them to the confidence of a plaster. Take care not to let it grow too cold before you form it into rolls; for then it will prove too brittle.

THIS platter differs only in colour from the red lead platter with soap, above-mentioned.

> **EMPLASTRUM** SAPONACEUM. Saponaceous plaster. Edinb.

Take of

Gum plaster, three pounds; Castile soap, sliced, half a pound. Melt the gum platter, and mix into it the foap.

HERE the addition of the gums is supposed to promote the resolvent virtue of the foap.

> **EMPLASTRUM** STOMACHICUM. Stomach plaster. Lond.

Take of

Soft labdanum, three ounces: Frankincense, one ounce; Cinnamon.

The expressed oil, called oil of mace, each half an ounce:

Estential oil of mint, 1 drachm. Having melted the frankincense, add to it, first the labdanum foftened by heat, and then the oil of mace; afterwards mix these with the cinnamon and oil of mint; and beat them together, in a warm mortar, into a mass, which is to be kept in a close vessel.

This is a very elegant stomach plaster. It is contrived so as to be easily made occasionally (for these kinds of compositions, on account of their volatile ingredients, are not fit for keeping) and to be but moderately adhefive, so as not to offend the skin, and that it may without difficulty be frequently taken off and renewed, which these forts of applications, in order to their producing any confiderable effect, require to be.

Edinb.

Take of

Yellow wax, eight ounces; Tacamahaca in powder, four ounces;

Cloves, powdered, two ounces; Palm oil, fix ounces; Expressed oil of mace, an ounce

and a half;

Essential oil of mint, 2 drachms. Melt the wax and tacamahaca with the palm oil; then removing the mixture from the fire, add

the

the other ingredients, and make them into a plaster, according to art.

These plasters are applied to the pit of the stomach, in weakness of that viscus, in vomitings, the disorder improperly called the heartburn, &c. and sometimes with good The pit of the stomach, fuccess however, as Hoffman has observed, is not always the most proper place for applications of this kind to be made to: if applied to the five lower ribs of the left fide, towards the back, the stomach will in general receive more benefit from them: for it appears from anatomical inspection, that greatest part of it is fituated there.

EMPLASTRUM
VESICATORIUM.
Blistering plaster, or epispastic
plaster.
Lond.

Take of
Drawing pla

Drawing plaffer, two pounds; Cantharides, one pound; Vinegar, half a pint.

Melt the drawing plaster, and a little before it grows sliff, mix in the cantharides, reduced into a most subtile powder; then add the vinegar, and work them well together.

Edinb.

Take of

Burgundy pitch, twenty ounces; Venice turpentine, Cantharides, each fix ounces.

Reduce the cantharides into a most fubtile powder, and add them to the other ingredients, previously melted together, so as to make the whole into a plasser, according to art.

Compound epispastic plaster. Edinb.

Take of

Burgundy pitch, twelve ounces; Yellow wax, four ounces; Venice turpentine, eighteen ounces;

Mustard feed, Black pepper, each one ounce; Verdigrise, two ounces;

Cantharides, twelve ounces.

Melt the wax and pitch together;
then add the turpentine; and
when this is liquefied, fprinkle
in the other ingredients, first
powdered and mixed together;
keeping them continually stirring, so as to make a plaster
thereof according to art.

The bliftering plafters are to be kept in oiled bladders.

THIS last composition has long been used in some particular shops, as the most infallible blister: tho? either of the other two answer the purpose very successfully. ther the vinegar in the first is of any advantage, is greatly to be doubted : in some cases, indeed, it has been observed, that the plaster without this addition feemed at first to fail of its effect, and that on taking it off, and rubbing the part with vinegar, the same plaster, applied again, has bliftered freely: but this does not appear to be fo much owing to any peculiar quality of the vinegar, as to its foftening the skin when applied in this manner, and fitting it for the action of the cantharides: when mixed with the other ingredients of the plaster, it has not this effect: it likewise exhales in keeping, infomuch that the composition, though fufficiently foft at first, becomes. in no long time, too dry. have been accustomed to spare the trouble of making any plaster on purpose for blittering, by occasionally spreading some of the cantharides in powder upon a common plaster. EM-

EMPLASTRUM ANODYNO DISCUTIENS.

An anodyne and discutient plaster.
Take of

Cummin plaster, two ounces; Camphor, three drachms; Thebaic extract, one drachm and a half.

Grind the camphor, with some drops of oil olive, into a very subtile powder, and then mix it with the other ingredients, according to art, into a plasser.

EMPLASTRUM CALIDUM.

Warm plaster.

Take of
Gum plaster, one ounce;
Blistering plaster, two drachms.

Melt them together over a gentle fire.

EMPLASTRUM SUPPURANS.

Suppurating plaster.

Take of

Gum plaster, an ounce and a half;

Burgundy pitch, half an ounce. Melt them together.

The uses of the three foregoing compositions, which are taken from our hospitals, are sufficiently obvious from their titles. The warm plaster is a very stimulating application, of great use in fixed pains, as in the rheumatism, sciatica, beginning chilblains, &c.

### CHAPTER XI.

### Ointments, Liniments, and Cerates.

OINTMENTS and liniments differ from plasters little otherwise than in consistence. Any of the officinal plasters, diluted with so much oil as will reduce it to the thickness of stiff honey, forms an ointment: by farther increasing the oil, it becomes a liniment.

#### UNGUENTUM ÆGYPTIACUM. Edinb.

Take of Verdigrife, finely powdered, five ounces:

Honey, fourteen ounces;
Vinegar, feven ounces.
Boil them over a gentle fire, to the confidence of an ointment.

## MEL ÆGYPTIACUM.

Take of
Verdigrife, reduced into a very
fubtile powder, five ounces;
Honey, fourteen ounces by
weight;

Vinegar, seven ounces by meafure.

Boil these ingredients together, over a gentle sire, till they have acquired a due consistence, and a reddish colour. On keeping this mixture for some time, the thicker part falls to the bottom; the thinner, which floats on the top, is called mel Egytiacum.

THESE preparations are defigned only for external use, for cleaning and deterging ulcers, and keeping down fungous flesh: they are serviceable also in venereal ulcerations of the mouth and tonsils. If for particular purposes they should be wanted more acrid, they may be occasionally rendered so by shaking the vessel, so as to mix up the thick matter at the bottom (which contains greatest part of the verdigrise) with the upper thin one. J

# UNGUENTUM ALBUM. White ointment. Lond.

Take of
Oil olive, one pint;
White wax, four ounces;
Spermaceti, three ounces.
Liquefy them by a gentle fire, and keep them constantly and briskly stirring, till grown thoroughly cold.

Edinb.

Take of
Oil olive, three pints;
Ceruffe, one pound;
White wax, nine ounces.

Melt the wax in the oil, then gradually add the ceruffe, and ftir them well together, that they may be thoroughly mixed into an ointment.

THESE are useful, cooling, emollient ointments, of good service in excoriations, and other likestrettings of the skin. The cerusse is omitted in the sirst prescription, on a suspicion that it might produce some ill essents frequently are, to the tender bodies of children. Though there does not seem to be much danger in this external use of cerusse, the addition of it is the less necessary here, as there is another oint-

### Chap. 11. Ointments, Liniments, and Cerates.

ciatment containing a more active preparation of the same metal, the unguentum saturninum; which may be occasionally mixed with this, or employed by itself, in cases where faturnine applications are wanted.

### UNGUENTUM ALBUM CAMPHORATUM.

Camphorated white ointment. Lond.

This is made by adding to the white ointment a drachm and a half of camphor, previously ground with fome drops of oil of almonds.

Edinb.

Take of The white ointment, one pound; Camphor, rubbed with a little oil, one drachm and a half. Mix them together.

THESE ointments are supposed to be more discutient than the foregoing, and serviceable against cutaneous heats, itching, and fer-piginous eruptions. They should be kept in close vessels, otherwise the camphor will foon exhale: their smelling strong of this in-

> UNGUENTUM ex ALTHÆA.

goodness.

Ointment of marshmallows. Lond.

gredient is the best mark of their

Take of Oil of mucilages, three pints; Yellow wax, one pound; Yellow resin, half a pound; Common turpentine, two ounces. Melt the refin and wax with the oil; then, having taken them from the fire, add the turpentine, and, while the mixture remains hot, strain it.

This ointment receives no virtue from the ingredient which it takes its name from (see p. 330.) and therefore the Edinburgh college has omitted it.

Unguentum antipsoricum. Ointment against the itch.

Take of

Elecampane root, fresh, Sharp-pointed dock root, fresh, each three ounces:

Water cresses, fresh and bruised. ten ounces;

Hogs lard, four pounds; Yellow wax,

Oil of bays, each four ounces; Vinegar, one pint;

Water, three pints. Bruise the roots, and boil them in the water and vinegar, till half the liquor is confumed: frain and strongly press out the remainder; add to it the watercresses and the lard, and boil them till the moisture is exhaled; then press out the ointment. and liquely in it the wax and the oil of bays.

Sulphur is added to this ointment occasionally.

Unguentum antipsoricum CUM MERCURIO.

Ointment against the itch with mercury.

This is made by adding to the foregoing ointment four ounces of quickfilver, killed with a fufficient quantity of Venice turpentine, and mixing them together. according to art, into an unguent.

THESE ointments are very inclegant ones, and rarely made use of. The first is likewise precarious in its effects; and though those with fulphur and mercury are of undoubted efficacy, yet they are by no means superior to the more fimple simple ointments of those drugs deferibed hereaster. The Edinburgh college has, therefore, at the late reformation, very properly omitted both the antipsoric ointments.

### UNGUENTUM BASILICUM FLAVUM.

Yellow basilicum ointment.

Lond.

Take of

Oil olive, one pint; Yellow wax,

Yellow refin,
Burgundy pitch, each I pound.
Common turpentine, three

ounces.

Melt the wax, refin, and pitch, along with the oil over a gentle fire; then take them from the fire, add the turpentine, and, whilft the mixture remains hot, firain it.

Edinb.

Take of

Yellow wax, one pound;
White refin, one pound and a
half:

Venice turpentine, halfa pound;

Oil olive, one pint.

Melt the wax and refin in the oil;
then add the turpentine, and

. firain out the ointment.

THESE are commonly employed in dreffings, for digesting, cleaning and incarnating wounds and ulcers. They differ very little, if at all, in their effects, from the linimentum Arcei.

### UNGUENTUM BASILICUM NIGRUM vel TETRAPHARMACUM.

Bluck basilicum ointment; or ointment of sour ingredients.

Lond.

Take of
Oil olive, one pint;
Yellow wax,

Yellow refin,
Dry pitch, each nine ounces.
Melt them all together, and, whilft
the mixture is hot, firain it off.

This ointment was formerly of confiderable efteem for healing and incarnating wounds, &c. but is faid to have an inconvenience of being apt to render them foul, and produce fungous flesh; at present is is rarely made use of; the yellow bafilicum, and the liniment of Arcaus, being in general preserved.

In the Edinburgh pharmacopoeis the black basilicum is direct-

ed as follows:

Take of
Yellow wax,
White refin,
Mutton fuet,
Tar, each half a pound;
Oil olive, a pint and a half.
Melt them over a gentle fire, ftirring them well together; and
then frain the ointment.

How far the alterations here made may contribute to prevent the inconveniences above complained of, or indeed whether the objections to the old ointment were well founded, I cannot take upon me to determine. Those who are the most conversant in the use of these forts of applications, are apt to ascribe more to the composition than it has any share in producing.

### UNGUENTUM BASILICUM. VIRIDE.

Green bafilicum ointment.

Lond.

Take of

Yellow bafilicum, eight ounces; Oil olive, three ounces by meafure;

Verdigrife prepared, one ounce. Mix and make them into an ointment.

This

THIS ointment is an efficacious detergent. Our hospitals have been accustomed to prepare an ointment greatly resembling this, under the title of Unguentum viride detergens.

#### UNGUENTUM CITRINUM. Yellow ointment.

Edinb.

Take of

Quickfilver, one ounce; Spirit of nitre, two ounces; Hogs lard, tried, one pound. Diffolve the quickfilver in the spirit of nitre, by digestion in a fand-heat; and, whilst the solution is very hot, mix with it the lard, previously melted by it-

felf, and just beginning to grow Stir them briskly together, in a marble mortar, fo as to form the whole into an ointment.

### UNGUENTUM CÆRULEUM FORTIUS.

The stronger blue ointment.

Take of

Hogs lard tried, two pounds; Quickfilver, one pound; Simple balsam of sulphur, half an ounce.

Grind the quickfilver with the balsam of sulphur till they are perfectly incorporated; then gradually add the lard heated, and mix them carefully together.

### UNGUENTUM CÆRULEUM MITIUS.

The milder blue ointment. Lond.

Take of

Hogs lard, tried, four pounds; Quickfilver, one pound; Common turpentine, one ounce; Grind the quickfilver with the turpentine in a mortar, till it ceases to appear; then gradually add

the lard warmed, and carefully mix them together.

This last unguent turns out of a much better blue colour than the foregoing, which is of a very dingy hue. Mercurial unguents have in many cases the same effects with the preparations of this mineral taken internally; and are at prefent frequently employed, not only against cutaneous disorders, as alterants, but likewife in venereal and other obstinate cases, for raising a falivation. The ptyalism excited by unction is faid to be attended with the fewest inconveniencies, and to perform the most complete cure. In some constitutions, mercurials, taken inwardly, run off by the intestines, without affecting the mouth; and in others, they affect the falival glands so quickly, as to occasion a copious ptyalism, without extending their action to the remoter parts, and consequently without removing the cause of the disease.

Unguentum desiccativum RUBRUM.

Red deficcative ointment.

Take of

Oil olive, a pint and a half; White wax, half a pound; Calamine prepared, fix ounces; Litharge prepared,

Bole armenic, each four ounces; Camphor, three drachms.

Melt the wax in the oil, and having taken them from the fire, gradually sprinkle in the other ingredients, fiftring them briskly together into an ointment. The camphor must be previously ground with a little oil of almonds.

This is faid to be an excellent drier and healer; but is at present in no great siteem, and rarely kept

in the shops. the last Edinburgh pharmacopæia, fluid, ftrain it off. but is now dropped.

UNGUENTUM DIAPOMPHOLYGOS. Ointment of pompholyx. Take of

Oil olive, twenty ounces; Juice of the berries of common, or deadly nightshade, eight ounces;

White wax, five ounces; Cerusse, four ounces; Burned lead,

Pompholyx, each two ounces; Pure frankincense, one ounce.

Roil the oil and the juice over a gentle fire, till the juice is exhaled; and towards the end of the coction, melt the wax in the oil; then take the mixture from the fire, and add to it, whilft hot, the other ingredients reduced to Mix and make them powder. into an ointment.

This is taken, as the preceding, from the last edition of the Edin-It stands burgh pharmacopæia. recommended against hot inflammatory ulcers and sharp defluxions on the eyes; but is very rarely made use of, having for some time given place to compositions more fimple, though at least equal in efficacy: for which reason it is now omitted by the college.

### UNGUENTUM e GUMMI ELEMI.

Ointment of gum elemi. Lond.

Take of

Mutton suct, fresh and tried, two pounds;

Gum elemi, one pound:

Common turpentine, 10 ounces. Melt the gum with the fuet, and having taken them from the fire, immediately mix in the turpen-

It was retained in 'tine; then, whilft the mass remains

UNGUENTUM, vulgo LINIMENTUM, ARCÆI.

The ointment, commonly called liniment, of Arceus. Edinb.

Take of

Hogs lard, one pound; Goats fuet, or mutton fuet, two pounds;

Venice turpentine, Gum elemi, each a pound and a

Melt and strain them, so as to make an ointment, according to art.

This unguent has long been in use for digesting, cleansing, and incarnating; and for these purposes is preferred by some to all the other compositions of this kind.

#### UNGUENTUM EMOLLIENS. Emollient ointment. Edinb.

Take of

Palm oil, four pints; Fresh-drawn linseed oil, three pints;

Yellow wax, one pound; Venice turpentine, half a pound. Melt the wax in the oils, over a gentle fire, then mix in turpentine, and firain the ointment, which supplies the place of the ointment of marshmallows.

IT is at least equal to that ointment for the purpose expressed in its title, nothing of the mucilage or emollient matter of the marshmallows being there retained. And indeed, if mucilages were blended with ointments, they would possibly diminish, rather than increase their emollient virtue; as they render oils fenfibly less unctuous, forming with them a new compound differ-

### Chap. 11. Ointments, Liniments, and Cerates.

639

ent from the ingredients, and missible with water into a milky liquor, as we have seen in chap. vii.

UNGUENTUM
MERCURIALE.
Mercurial ointment.

Take of

Hogs lard, two ounces; Quickfilver, one ounce.

Beat them diligently together, till the quickfilver disappears. It may likewise be made with two, three, or more times the quantity of quickfilver.

THIS is the most simple of the mercurial ointments, though poffibly as efficacious as any. It requires, indeed, a great deal more labour to extinguish the mercury in the lard alone, than when turpentine, or other like substances, are joined: but in recompence, the composition with lard is free from an inconvenience which the others are accompanied with, viz. being apt, by frequent rubbing, to fret tender skins. Some chuse to stiffen this ointment with a fourth part of fuet (proportionably diminishing the lard) which gives it a better confiltence for use.

The above prescription and remark are from the foregoing edition of this Dispensatory. The college has now received the addition of suct, in such proportion as to make one-fourth the quantity of the whole ointment; the proportion of quicksilver being continued the same. The composition is now as follows:

Edinb.

Take of

Hogs lard, three ounces; Mutton fuet, one ounce; Quickfilver, one ounce.

Beat them diligently together in a mortar, till the mercurial globules disappear. This ointment is made also with twice and with thrice the quantity of mercury.

UNGUENTUM e MERCURIO PRÆCIPITATO.

Ointment of mercury precipitate. ...

Take of

Simple ointment, an ounce and a half;

Precipitated fulphur, 2 drachms; White mercury precipitate, two feruples.

Mix them well together, and moisten them with ley of tartar, that they may be made into an ointment.

This is a very elegant mercurial ointment, and frequently made use of against cutaneous disorders. The preparations of mercury and sulphur, here directed, are chosen on account of their colour.

UNGUENTUM NERVINUM.

Nerve ointment.

Take of

Southernwood, Marjoram (or origanum)

Mint, Pennyroyal,

Rue,

Rolemary, each, fresh gathered, fix ounces;

Nears-foot oil, five pints; Beef fort three pounds; Oil of bays, half a pint.

Boil the herbs, with the neats foot oil and fuet, till the aqueous moisture is exhaled; then press and strain out the liquid, and, adding to it the oil of bays, make the whole into an ointment.

This ointment is defigned, as its title expresses, for warming and strengthening the nerves. The above form is from the preceding edition of the Edinburgh pharmacopæia. It is an ill contrived one:

for

for besides the ingredients being more numerous than there is any occasion for, the method of treating them is very exceptionable. The warm stimulating, nervine virtues of the herbs confift in their volatile parts, which are lost in the boiling The most of them with the oil. effectual method of impregnating ointments with these virtues of vegetables, is that which we have formerly proposed (page 332.) and which the college has now received, adding a fuitable quantity of the essential oil of the subject. The present reformed composition is as follows:

Edinb.

Take of

Mutton fuet, two pounds;
Oil of chamomile (by decoction)
one pint;

Oil of bays, a pint and a half; Essential oil of origanum, or of rosemary, two ounces.

Melt the sue, over a gentle fire, in the oil of chamomile, so as to make an ointment thereof; which being removed from the fire, stir into it the oil of bays and essential oil.

Some, instead of mixing any effential oil with the composition, are accustomed to rub a few drops of it upon the surface of the plaster when spread.

# UNGUENTUM NUTRITUM. The ointment called nutritum. Edinh.

Take of Litharge, Vinegar, each two ounces;

Oil olive, fix ounces.

Rub them in a mortar, adding the oil and vinegar, alternately, by little and little at a time, till the vinegar ceases to appear, and the ointment becomes uniform and white.

This ointment is troublesome to make, and does not keep well, the vinegar exhaling, so as to leave the compound too stiff: for which reason, it is now directed to be made in less quantity than in former editions. It is supposed to be a good cooler and deficcative; and is occasionally used in excertations, slight serpiginous eruptions, and for anointing the lips of wounds or ulcers that itch much, or tend to instantion.

Unguentum Ophthalmicum.

Eye ointment.

Take of

Ointment of tutty, an ounce and a half;

Saturnine ointment, half an ounce;

Camphor, half a drachm. Mix and make them into an ointment, according to art.

This ointment may likewise be made with two, three, or more times the quantity of camphor.

This unguent is very well contrived for the purpose expressed in its title; scarce any of those commonly met with being of equal efficacy in inflammations, and hot acrid defluxions on the eyes. But as a good deal of caution is requifite in the use of saturnine applications, for fo tender an organ as the eye; and as compositions of this kind may be easily formed extemporaneously, with such proportions of the ingredients as the prescriber shall think fit, the Edinburgh pharmacopæia (from the preceding edition of which the above form is taken) has now omitted it.

# UNGUENTUM e PICE. Ointment of tar. Lond.

Take of
Mutton suet, tried,

Tar,

#### Chap. 11. Ointments, Liniments, and Cerates. 641

Tar, each equal weights.

Melt them together, and firain the
mixture whilf hot.

This composition, with the addition of half its weight of resin, has long been used in the shops as a cheap substitute to the black basilicum.

#### UNGUENTUM SAMBUCINUM. Ointment of elder. Lond.

Take of

Elder flowers, full blown, four pounds;

Mutton suet, tried, three pounds;

Oil olive, one pint.

Melt the suet with the oil, and in this mixture boil the flowers till they are almost crisp: then strain and press out the ointment.

#### Edinb.

Take of

The inner bark of the elder tree, The leaves of elder, fresh, each four ounces;

Linfeed oil, two pints; White wax, fix ounces.

Let the bark and leaves be well bruised, and boiled in the oil till the humidity is consumed; then press out the oil through astrainer, and melt in it the wax, so as to make an ointment.

THESE ointments do not feem fuperior to fome others, which are much neater, and parable at less expence. They can fearce be supposed to receive any considerable virtue from the ingredients which they take their name from.

# UNGUENT. SATURNINUM. Saturnine ointment. Lond.

Take of Oil olive, half a pint; White wax, an ounce and a half; Sugar of lead, two drachms. Let the fugar of lead, reduced into a very jubil powder, be ground with fome part of the oil, and the wax melted with the reft of the oil; mix both together, and keep them firring till the oint-

#### Edinb.

Take of

Sugar of lead, one ounce; White wax, threeounces; Oil olive, one pint.

ment is grown cold.

Liquefy the oil and wax together, and gradually add the fugar of lead; continually firring them, till, growing cold, they unite into an ointment.

BOTH these ointments are useful coolers and desiccatives: much superior both in elegancy and esfecacy to the nutrium or tripbarmacum.

## UNGUENTUM SIMPLEX. The fimple simment. Lond.

Take of

Hogs lard, tried, two pounds; Rose water, three ounces by measure.

Beat the lard with the rose water, till they are well mixed; then melt them over a very gentle fire, and set them by for some time, that the water may subfide: pour the lard off from the water, and keep incessantly stirring and beating it about till it grows cold, so as to reduce it into a light incoherent mass: lastly, add so much essence of lemons as will be sufficient to give a grateful odour.

UNGUENTUM ROSACEUM.
valgo POMATUM.

The rose ointment, commonly called pomatum. Edinb.

On any quantity of hogs lard, cut into small pieces, and placed in a glazed earthen vessel, pour as much water as will rise above it some inches; and digest them together for ten days, renewing the water every day. Then liquesty the lard with a very gentle heat, and pour it into a proper quantity of rose water; work them well together; and afterwards, pouring off the water, add to the lard some drops of oil of rhodium.

THESE ointments are in common use for softening and smoothing the skin, and healing chaps.

# UNGUENTUM e SULPHURE. Ointment of fulphur. Lond.

Take of

The simple ointment, half a

Flowers of fulphur, unwashed, two ounces:

Effence of lemons, one scruple. Mix them together.

This is defigned for cutaneous disorders; it is much neater than the unguentum antipsoricum cum sulphure, though, at least, equally efficatious.

Unguentum ad Psoram.

Ointment against the itch.

Take of

Sulphur, one ounce;
White heliebore root, in powder, or crude sal ammoniac,
two drachms;

Hogs lard, two ounces.

Mix, and make them into an ointment.

Sulphur is a certain remedy for the itch, more fafe and efficacious than mercury: for, as Dr. Pringle observes, unless a mercurial unction was to touch every part of the skin, there can be no certainty of success; whereas, by a fulphureous one, a cure may be obtained by only partial unction, the animalcula, which occasion this disorder, being, like other in-fects, killed by the sulphureous steams, which extrale by the heat of the body. As to the internal use of mercury, which some have accounted a specific, there are several instances of men undergoing a complete falivation for the cure of the lues venerea, without being freed from the itch.

he quantity of ointment, above directed, serves for four unctions: the patient is to be rubbed every night; but to prevent any disorder that might arise from stopping too many pores at once, a fourth part of the body is to be rubbed at one Though the itch may thus be cured by one pot of ointment, it will be proper to renew the application, and to touch the parts most affected, for a few nights longer, till a fecond quantity also is exhausted, and in the worst cases to subjoin the internal use of sulphar, not with a view to purify the blood, but to diffuse the steams more certainly through the ikin; there being reason to believe, that the animalcula may fometimes lie too deep to be thoroughly destroyed by external applications.

#### UNGUENTUM TRIPHARMACUM.

Ointment of three ingredients.

Lond.

Take of

Common plaster, four ounces; Oil olive, two ounces by meaface;

Vinegar,

Vinegar, one ounce by measure. Boil them together over a gentle fire, keeping them continually firring till they are reduced to the confishence of an ointment.

This is a new method of preparing the unguentum nutritum, much less trouble some than the one already described under that title. The composition proves likewise more smooth and uniform, and not so liable to grow dry in keeping. This ointment is nevertheless interior, both in respect of elegancy and efficacy, to the unguentum saturninum.

#### UNGUENTUM TUTIÆ.

Ointment of tutty.

Lond

Let any quantity of prepared tutty be mixed with as much purified vipers fat, as is sufficient to reduce it into the consistence of a soft ointment.

THIS ointment is defigued for an ophthalmic. What particular virtues it receives from the vipers fat, we shall not presume to determine.

In the preceding edition of the Edinburgh pharmacopæia, this ointment was directed as follows:

Take of

White wax, three ounces; Best oil olive, ten ounces; Tutty prepared, two ounces;

Calamine prepared, one ounce. Liquefy the wax with the oil over a gentle fire: the gradually sprinkle in the tutty and calamine, continually stirring them, till the ointment grows cold.

This ointment may tike ife be made extemporaneously by mixing the calamine and tutty with four times their quantity of fresh butter.

In the present edition, the ointment is ordered to be made only with butter, stiffened with a little wax: and the calamine is omitted, for certainly calamine and tutty were not both necessary in one composition. The formula is now as follows:

Take of

Fresh butter, four ounces; White wax, two drachms; Tutty prepared, one ounce,

Melt the wax with the butter over a gentle fire, and then fprinkle in the tutty, continually ftirring them together, till the ointment is grown cold.

THERE is one inconvenience in the use of butter, that it soon grows rancid in keeping, infomuch that the ointment made with it, should seem unfit for an officinal. That either butter, or vipers fat. is more effectual, or more inoffenfive to the eyes, than the belt fort of Florence oil; that the tutty is in any respect superior to calamine: or that either have, in this form, all the effect which they are commonly supposed to have, I will not affirm. Both the calamine and tatty act only by virtue of the zinc they contain; and calamine appears to contain the most of the two, and likewise to be the least variable in its contents. But the pure flowers prepared from zier itself (see page 542.) are doubtless preferable to either.

THE Edinburgh college have now added

UNGUENTUM TUTIÆ CAMPHORATUM

Camphorated ointment of tutty.

Edinb.

This is made by adding to the foregoing ointment two drachms of T t 2 camphor.

camphor. It is prepared also with a double quantity of camphor.

Unquentum vermitugum.
Ointment against worms.

Ointment against avorms.

Take of
Lavender cotton,
Wormwood,
Rue,
Savin,
Tansy leaves, fresh gathered,
each two ounces;
Oil olive, a pint and a half;
Hogs lard, one pound;
Yellow wax, three ounces;
Ox gall,
Socotorine aloes, each an ounce
and a half;
Coloquintida,
Wormseed, each one ounce.

Bruise the herbs, and boil them with the oil and lard, till the aqueous moisture is evaporated; then press the liquor through a firainer, melt in it the wax, and afterwards add the other ingredients, boiling and firring them together, so as to make an ointment. The aloes, coloquintida, and worm-seed, must be previously reduced into a very subtile powder.

This ointment is rubbed on the bellies of children for defiroying worms, and fometimes, as is faid, with good fuccess. It is taken from the preceding edition of the Edinburgh pharmacopæia: in the prefent it is omitted.

UNGUENTUM ad VESICATORIA [L.] UNGUENTUM EPISPASTICUM [E.] Ointment for blifters. Lond.

Take of
Hogs lard, tried,
Bliftering plafter, each equal
weight.

Melt them together over a very gentle fire, and keep them conflantly firring till grown cold.

Edinb.

Take of
Hogs lard,
Vehice turpeatine, each three
ounces;
Yellow wax, one ounce;
Cantharides, three drachms.
To the lard and wax melted tope-

To the lard and wax melted together, add first the cantharides reduced into powder, and then the turpentine: lastly, mix the whole into an ointment.

THESE ointments are added in the dreffings for blifters, intended to be made perpetual, as they are called, or to be kept running for a confiderable time, which in many chronic, and fome acute cases, they are required to be. Particular care should be taken, that the cantharides employed in these compositions be reduced into very subtile powder, and that the mixtures be made as equal and uniform as possible.

UNGUENTUM
EPISPASTICUM MITIUS.
The milder opifpaftic ointment.
Edinb.

Take of
Cantharides,
White refin,
Yellow wax, each one ounce;
Hogs lard,
Venice turpentine, each two
ounces;
Boiling water, a fufficient quan-

Infuse the cantharides in the water, in a close vessel, for a night; then firongly press out, and strain the liquor, and boil it with the lard till the watery moisture is confumed; then add the resin, wax, and turpentine, and make the whose into an ointment.

THIS

This cintment, containing the foluble parts of the cantharides uniformly blended with the other ingredients, is more commodious, and occasions less pain, though not less effectual in its intention, than the two foregoing compositions with the fly in substance.

#### UNGUENTUM VIRIDE.

Green ointment.

Lond.

Take of
The green oil, three pints:
Yellow wax, ten ounces.

Melt them together over a gentle fire, and keep the mixture continually fiirring until it is grown cold.

This ointment does not feem to receive any particular virtue from the ingredients to which its colour is owing.

### LINIMENTUM ALBUM, White liniment.

Lond.

Take of
Oil olive, three ounces by mea-

fure; Spermageti, fix drachms; White wax, two drachms;

Melt them together over a gentle fire, and keep them constantly and briskly stirring, till grown cold.

THIS differs only in confidence. from the unguentum album.

### BALSAMUM VIRIDE. Green balfam.

Green balfam. Edinb.

Take of
Linfeed oil,
Oil of turpentine, each one
pound;
Verdigrife, in powder, three
drachms.

Boil and flir them well together till the verdigrife is diffolved.

A balfam, fimilar to this, is faid to have been greatly valued by our furgeons as a detergent.

#### LINIMENTUM TRIPHARMACUM.

Liniment of three ingredients.

Lond.

Take of

Common platter, four ounces;
Oil olive, a quarter of a pint;
Vinegar, one ounce by measure.
Boil them over a gentle fire, continually flirring them until they acquire the consistence of a liniment.

This is the same with the unguentum tripharmacum, except that the quantity of oil is here increased, to give the compound the softer consistence of a liniment.

### LINIMENTUM VOLATILE. Volatile liniment.

Take of
Oil of harthorn,
Spirit of harthorn, each equal
parts.
Mix them together.

DR. Pringle observes, that in the inflammatory quinfey, or firangelation of the fauces, a piece of flannel, moistened with this mixture, and applied to the throat, to be renewed every four or five hours, is one of the most efficaci-By means of this ous remedies. warm stimulating application, the neck, and fometimes the whole body, is put into a fiveat, which, after bleeding, either carries off, or lessens the inflammation. Where the ain cannot bear the acrimony of this mixture, the volatile liniment of the shops (page 440) may be made trial of.

T:3 CERA-

#### CERATUM ALBUM.

White cerate.

Lond.

Take of

Oil olive, a quarter of a pint; White wax, four ounces; Spermaceti, half an ounce.

Liquefy them all together, and keep them stirring till the cerate is grown quite cold.

This differs from the whiteointment and liniment, only in being of a thicker confidence.

### CERATUM CITRINUM. Yellow cerate.

Lond.

Take of

Yellow basilicum ointment, half a pound;

Yellow wax, one ounce. Melt them together.

THIS is no otherwise different from the yellow basilicum, than being of a stiffer considence, which renders it for some purposes more commodious.

#### CERATUM EPULOTICUM.

Epulotic cerate.

Lond.

Take of Oil olive, one pint;

Yellow wax, Calamine prepared, each half a

pound.

Liquefy the wax with the oil, and, as foon as the mixture begins to grow sliff, sprinkle in the calamine, keeping them constantly stirring together, till the cerate is grown quite cold.

#### UNGUENTUM e LAPIDE CALAMINARI.

Ointment of calamine. Eainb.

Take of Yellow wax, eighteen ounces; Oil olive, two pints; Calamine prepared, ten ounces and a half.

Melt the wax with the oil, and gradually sprinkle in the calamine, mixing and stirring them well together till grown cold.

THESE compositions are formed upon the cerate which TURNER strongly recommends in cutaneous ulcerations and excoriations, and which has been usually distinguished by his name. They appear from experience to be excellent epulotics, and as such are frequently made use of in practice.

#### CERATUM MERCURIALE.

Mercurial cerate.

Lond.

Take of

Yellow wax,

Hogs lard, tried, each half a pound;

Quickfilver, three ounces; Simple baliam of fulphur, one drachm.

Melt the wax with the lard, then gradually add this mixture to the quickfilver and balfam of fulphur previoully ground together.

UNGUENTUM PARALYTICUM.

Palfy ointment.

Take of

Hogs lard,

Oil of bays, each four ounces; Strong spirit of vitriol, one ounce.

Mix, and make them into an unguent.

This irritating composition is applied to numbed or paralytic limbs: it soon reddens and inflames the skin, and when this effect is produced, must be taken off; after which, the part is to be anointed with any emollient unguent; as that or elder.

UNGUEN-

# I N D E X.

▲ Bforbent earths	page 62	Aloes elixir	page 323
Acacia, Ægyptian	73	pills	ib.
German 2		wine	292
Acids	61	wood	75
animal	143	Alterative mercurials, &	c <b>.</b>
mineral	24	see Mercurials, &	c.
marine	25, 448	Alum	82.
nitrous	25, 447	burned or dried	464
vitrioli <b>c</b>	25, 443	curd	650
vegetable	8, 14	water or folution	464
Acrids	69	whey	288
Adder	245	Amber	234
Adderstongue	188	prepared	256
Æthiops, See Ethiops	ib.	balfam	48€
Affinities of bodies	27	compound powder	554
Agaric	76	tindure	3 (8
of the oak	ib.	troches	554
Agrimony	77	oil, falt and fpirit	481
hemp-agrimony		Ambergris	83
water agrimony		cffence	328
Alder	79	Ammoniac falt, fee Salt	ib.
black alder	79	Ammoniacum gum	84
Alcohol	38í	purified	261
Ales medicated	· 301	julep	ib.
aperient	302	milk, or milky fo	
bitter	ib.	plaster with merc	ury 629
Buder's	ib.	Amomum	84
cephalic	ib.	Anacardium	86
diuretic	ib.	Angelica	87
<b>scorbatic</b>	303	candied	337
Alexanders	153	distilled water	
Alkaline infusion	276	Anhalt water,	<b>3</b> 5 <b>3</b> 68
falts, fixed		Animals	18
volatile	429	Anime	88
Alkanet	87	Aniseed	ib.
Allspice	201	essential oil	360
Almonds	`85	compound water	392
expressed oil	266	Antihectic of Poterius	543
Almond foap	436		343 89
Aloes	79	butter	538
purified '	262	calx	5 <sup>2</sup> 9
gum	497	caustic	538
relin	~ '#,	ceruffe	530
			Antimon

### 652 I N D E X.

Antimony		page	539	В.	
	Crocus		527	Dalamaina	
	diaphoretic		529		page 99
	emetic mild		528	Balm	175
	ethiops glass		545	fimple water	378
	glass cerated		536	compound water Balfam of amber	383
	panacea		537		48 E
	prepared	200	535 256	of Berne	21, 440
	purging	-22,		of Canada	322
	regulus	•	537 531	cephalic	239
	fulphurs	£ 22.	534	of Copaiba	590
	tinctures	319,		Friar's	99 322
	wine	3.30	293	of Gilead	189
Ants			143	green	645
	fee Decoction	S.	- 73	of guaiacum	321
Apple, cra	ib		172	Guido's	421
	rd <b>en</b>		171	of life	330
Aquafortis			449	Locatelli's	589
	compound		451	odoriferous	590
	purified		450	Perfian	322
Aqua regia			451	of Peru	100
Arabic gui			i49	faponaceous	439
	llion		605	of fulphur	485
Arcæus's li	niment		638	of Tolu	ioi
Archangel			163	of turpentine	419
. Aromatics			69	traumatic	322
Arquebula	de water		398	Wade's	ib.
Arrach, ga	ırden		97	Barbadoes tar	198
ftin	king .		97	diffilled oil	419
Arfenic		- 1	93	Barberries	103
Arimart, b	oiting		195	gelly of	338
	ootted		194	Barilla	427
Arum			94	Barks, season of gathering	
electar			334	Barley	153
emulfi			607	Barley water	282
	and powder		548	Bafil	186
Afafætida			95	Batilicum ointment	636
	urified		261	Baftard-faffron	118
	ncture		309	Bastard-stone-parsiey	85
Afarabacca			95	Baume de commandeur	322
Afta tree		0	144	Bay-berries	164
Ashes of ve		0,	22 126	electary	586
pearl-s			160	effential oil	362
pota <b>li</b> Ruffia			126	Bay leaves Bdellium	164
·	Smes			<b>=</b> '	101
Asparagus Ashmatic e	livia		<b>9</b> 6	Beans buck-beans	140
Astringents	TLAIF		324 67	Malaca bean	<b>342</b> 86
Athanor fu	rn <i>ace</i>		.32	Bearsbreech or brankursine	
Avens	HELV		119	Beddraw, ladies	, T
76113			٠.٦	Manual was the same	Beca

			20
Bees		Boluses, emmenagogue	page 582
Beets	103	febrifuge	ib.
Benit herb	119	hysteric	ib.
Benzoine	102	iliac	583
flowers	479	mercurial	ib.
magistery	327	emetic	ib.
tinclure	ib.	pectoral	ib.
Betony	103	of rhubarb with n	nercury ib.
Bezoar stone	104	rheumatic	ib.
prepared	255	<b>f</b> cilliti <b>c</b>	. 584
Bezoar mineral	540	<b>fudorific</b>	ib.
with tin	544	treacle	ib.
Bezoardic powder	551	turpentine	ib.
Birch		Bones	63
Birthworts	92		107
Bihopfweed	83	Borax	ib.
Bifmath	105	its fedative falt	480 .
magistery	541	Box wood	110
Biftort	106	empyreumatic oil	418
Bitters	70		142
Bitter ale	202	Bramble	21[
infusion	_	_	
purging	273 ib.		242
tindure			74 `218
wine	305	Bread	
	293	fow-bread	242
Bitter-apple	128		94
Bitter-fweet	136	Briar, wild	134
Bitter-vetch	190		234
Bitumens	21		102
Black-berries	211		146
Bleffed-thiftle	117	butchersbroom	211
Blistering plasters	632		283
epithem	647	Bryony	108
ointments	644	compound water	393
Bloodstone	150	Buckbeans	242
prepared	255	Buckthorn	229
Bluebottle	133	fyrup	<b>34</b> 7
Bolar earths	23, 65	Bugle	109
Bole-armenic	106		i.
French, &c.	ib.		ı Çı
compound powders	549	decoction	285
Boluses	579	Burgundy pitch	202
alexipharmac	ib.	Barnet	199
of alum	580	Burnet-faxifrage	ib.
antidyfenteric	582	black	209
of camphor	580	Butchersbroom	211
of castor	ib.	Butler's ale	302
<b>c</b> athartic	ib.	Butter of antimony	538
with mercu	ry 581	Butterbur	198
diaphoretic	ib.	-	-,-
digretic	ib.		
, <u> </u>			Cabbages

2)4		25 22,	
C.		Carrot, deadly	Page 239
Cabbana	4	Cafcarilla Caffia	137
	page 108	electary	120
Calamine	110		58 <b>6</b>
prepared	255 6.6	pulp Coffic liance	<b>260</b>
cerate	646 ib.	Cassidony	125
Onlawin	111		232
Calamint Calamina aromations	īb.	Castumunar Castor	12 E ib.
Calamus aromaticus Calcination		bolus	580-
Calomei	56		
Calx of antimony	519 529	fimple water	307, 308
	512	Cataplaims	375
mercur <b>y</b> tin	503	aromatic	647 ib.
Camels hay	159	comphorated	
Cammock	187	of cummin	649
Lamomile, fee Chamomile		discutient	647 ib.
Campeachy wood	167	emodlient	
Camphor Camphor	•	ischiadic	649 <i>ib</i> .
bolus	12, 13		
	,	ripening . ftomachic	648
cataplaf <b>m</b> - emulfi <b>en</b>	649 606		649
	609	fuppurating Catechu	648
julep ointment	635	Catmint	563 182
folution in oil	332	Cauliflower	102
folution in spirit	332 327	Caustic, antimonial	538
Candies, see Preserves	32/ 336	common milder	
Canella, white	114	ftronger	4 <sup>2</sup> 9-
Cantharides	115	lunar	
tinctures	306	Cautery, potential	49 <b>0</b> 42 <b>8</b>
plaiters	632	Cayan pepper	201
ointments	644	Celandine	124
_	116	Centaury	122
Capers Caper-bark	ib.	extract	404
Capillary herbs	250	Cerate of calamine	646
Capiliary neros	2Q1	epulotic	ib.
Caranna	116	mercurial	ib.
Caraway feeds	119	Turner's	ib.
essential oil	360	white	ib.
fpirituous water	394	yellow	ib.
Cordamom seeds:	116		23, 501
spirituous water		compound powder	-3, 5°1
tincture	397	Cerusse of antimony	530
Carduus benedictus	117	Ceterach	
extract	375	Chalk '	123
infusion	269	prepared	133
fimple water	375	decoction	255 279
Carline thiftle	118	julep	610
Carpobaliam	ib.	Chalybeate preparations,	
Carrot	135	Steel tinctures	312
of Crete	ib.		J·~
		. Ch	aly beate

II	N D	E X. 6	55
	ze 294	Collection of simples page	
Chamomile	124	73-11	620
essential oil	361	dry of Rhazi	<u>560</u>
oil by decoction	330		420
· fimple water	37 <b>7</b>		128
extract	404	extract 408,	414
wild	130		569
Chafte tree	77	Coltsfoot	243
Cheeserennet Cherries	145	Columbines	90
	122	Comtry	129
black cherry water winter cherries	•	Comminution	5 <b>3</b>
Chervil	77	Confection of Archigenes	591
Chick peas	123	cordial 414, of Damocrates	59,1
Chickweed	125 81	- C I-	ib.
China root	125		349
Chio turpentine	. ,	5	589
Chocolate nut	238 110	A Davidstate '	59 L
Cicely fweet	180	Conformer	414
Cichory	125	0 6: 61:	333
Cinnabar of antimony	539	1	336
factitious	509	mallows	335 ib.
native	126		334
Cinnamon	127		335 335
essential oil	361	rolemary	337 <b>16.</b>
fimple water	377	rofes	ib.
spirituous water	394	rue	334
infulion	277	fcurvy-grafs	ib.
tincture '	308		336
Cinquefoil	194		334
Circulatory vessels 44	, 304	wormwood	335
Citrons	127	Confound, middle	109
Citruls	ib.	Contrayerva	120
Civet	249	compound powder	5 5 E
Clary	153	Copaida dallam	99
Clays	23	compound oil	420
Clivers	90		129
Clove-july-flowers	119	Copper	1 32
fyrup	342	Caix	4
fpice effential öil	119	crystals	F.
Coating glasses	<b>3</b> 60		<b>19</b> 9
Cobalt	34	Copperas	247
Cobweb	93	nranarad	129
Cochineal	90	C	25 <b>5</b>
Coffee	127 128	4.2- 112	129
· Colbatch's styptic powder	543		513
Colcothar of vitriol	543 443	Cork	29
Cold feeds	250	Cornachini's nomice	233
Coleworts, garden	108	C	553
lea ·	ib.	Cornerate	130
		Costm	193 arv
			/

•			•	
656	I	I	DEX.	
Colimary	pag	e 99	Decantation pag	. 49
Cottus		130	Decoctions 268,	
Cowilips		193	antifebrile	284
fyrup		346	aptihectic	283
Crab apples		172	astringent	285
Crabs claws		114	of burdock	ib.
prepared		255	of chalk	279
compound p	<b>Mder</b>	_	common	287
Crabs-eyes		114	diuretic	285
prepared		255	febrifage	284
Cranesbill		140	for fomentations	287
Cream of tartar		473	for glysters	ib.
Creffes		181	of Japan earth	286
sciatica cresses		156	Japonic	279
Crocus of antimony		527	icteric	280
washed		528	of logwood	28.5
medicinal		526	nephritic	280
of iron, aperier		495	nitrous	281
altringe	ene	ib.	pectoral	ib.
Crocus of metals		527	Peruvian Constru	286 ib.
Crowfoot-cranefbill		140	feneka Godennost	28 I
Crystal Crystallination	46	64	fnakeroot of tamarinds with fena	
Crystallization Crystals of copper	40,	459	vulnerary	284
tartar		498	white	278
Cubebs		472 131	compound	
Cucamber		ib.	of the woods	279 280
wild cucumber		132	Deflagration .	
Cucarbit		ib.	Deliquiation	57
Cummin		134		43
essential oil			Despumation	45
cataplaim			Detonation	57
plaiter		628		178
Currants, red		209		626
. gelly		338		588
Currant railins		248		De-
Cyperus root		134	coctions.	
Cypress tree		132	Digettion	· <b>4</b> 4
Ly prus turpentine		238	Dill	87
		•	essential oil	360
D.			fimple water	374
Daffy's clixir		317	Distillation	49
Daines		102		370
Damask zose		209	of vinous spirits	186
Damfons		205	of acid spirits	443
<b>Dandelion</b>		135		430
Danewort		136		333
Dates		134		418
Dead nettle		163		165
Deadly carrot		239	Dittany of Crete	135
Deadly nightshade		227	white or baftard	143
			L	ock

_			- 31
Dock	page 163	Electary, alexeterial	age 597
water dock	ib.	alterative	ib.
Dodder of thyme	133	antiepileptic	ib.
Dog-role	134	antidysenteric 5	86, 598
Dog's mercury	176	aromatic	598
Double tongue	453	of arum root	334
Dragons	136	balfamic	598
Dragant gum, fee Tragac	anth.	of bayberries	<b>586</b>
Draughts	609	binding	602
cathartic	616	caryocostine	588
faline	ib.	of cafia	586
diaphoretic	617	· chalybeate	598
diuretic	ib.	deobstruent	599
anodyne	ib.	for a gonorrhœa	ib.
oily	607	of guaiacum	ib.
Drops, palfy	386	hæmorrhoidal	602
of life	328	incraffating	60g
Dropwort	141	lenitive	587
Drying of simples	252	nephritic	600
Dungs	232	paralytic	ib.
Dwarf-elder	136	pectoral	¢88
	- 30	Peruvian	600
E.		purgative acid	601
		laponaceous	ib.
Earths, general division	of 21.62	of scammony	588
Earth, fullers	126	of scordium	ib.
Japan	155	of fulphur	602
Lemnian	107	Elemi	137
fealed	167	ointment	638
Silefian	107	Eleutheria	137
Earthworms	170	Elixir of aloes	323
Eaton's flyptic	318	afthmatic	324
Eau d'arquebusade	398	Daffy's	317
de carmes	383	of health	316
de luce	442	of guaiacum	321
de Rabel	455	of myrrh compour	id 326
Edulcoration	48	paregoric	344
Rei	88	pectoral	₄\$b.
Eggshells prepared	256	of propriety	23
Elaterium	402	Helmont's	292
Elder	216	Paracelfus's	323
berries, rob of	399	vitriolic	ib.
flowers, fimple wa		facred	326
oil by dec	oction 331	flomachic	305
ointment	641	· uterine	325 326
Elder, dwarf	136	of vitriol	. 325
Elecampane	138	acid	324
candied	337	dulcified	3 <b>26</b>
extract	405	Mynficht's	32 <u>5</u>
Electaries	, 5°5	Vigani's	326
Electaries  Electary, acid	597		_
	377	U u E	77 lutriation

- 35			
Elutriation	page 55	Extract of Rudius	page 408
Emerald	64	of rue	407
Emetics	70	of (affron	411
Emulions	605	of favin	407
Emulsion, Arabic	606	of fteel	294
with arum	608	Thebaic	261
camphorated	606	of wormwood	
common			404
	605	Eyebright	140
oily	607	F.	
purging	606	Essa	.0 .0
of (permaceti	607.	Fats	18, 98
Endive	138	prepared or purified	
Ens Veneris	449	Featherfew	174
Epithems	648	Fennel	142
blistering	ib.	essential oil	361
volatile	440	fimple water	<b>3</b> 7 <b>8</b>
Epsom salt	214	Fennel flower	18 <b>3</b>
Eryngo	139	hogs-fennel	199
<b>ca</b> ndied	336	Fenugreek	143
Essence, alexipharmac	329	Fermentation	5
of ambergris	328	Fern	142
of lemons	363	Feverfew	174
of Neroli	97	Figs	118
roval	328	Figwort	223
of the woods	329	Filtration	45
Essential oils		Fir	72, 238
falts	353	Fish-glue	156
	475	Flag, sweet	111
Ethereal spirit	455	yellow water-flag	147
Ethiopic pills	573		169
Ethiops antimonial	545	Flax, common	ib.
mineral	508	purging	
martial	494	Fleawort	206
Evaporation	48	Flies, Spanish	115
Exficcation	5,2	Flint	64
Axpression .	ib.	Flixweed	228
Extraction	43	Flower de-luce	158
Extracts	403	yellow water flower-de	-luce 147
Einact, cathartic	413	Flowers of benzoine	479
of centaury	404	fal ammoniac	
of chamomile	ib.	fulphur	484
of elecampane	405	wash	ed ib.
of gentian	ib.	fteel	496
uaia. um	407	zinc	542
of black helleb		Fluellin	137
- of hemlock	400	honey of	350
of jalap	412	Fluxes for metals	57, 532
of liquorice	405	Fomentation, anodyne	287
	406, 413	aromatic	ib.
of opium by dige	eftion 415	common	ib.
of Peruvian bark	406.4.2	firengthen	
of plantane	400,4 3	Poxglove	135
o. prantene	,400	2 38.3.2	Frank-
		•	

I	N	D	<b>E</b> :	X.	659
Frankincense	page 24	8 G	lyster,	fætid	page 622
Friar's balfam	32			piate	621
Frogs spawn	20	7	F	urgative	622
Follers earth	12	6	f	larch	621
Fulminating gold	48	9		urpentine	622
Fumitory			ozts rue		145
Furnaces	3	o Go	old 🛴		98
Fusion	5	5		minating	489
•	•		mo		506
G.		C	pot olden-re	abie	489
Galangal	• •	_	oldilock		246
Galbanum			oolegra		232
purified			ourd fe		90
Galenic pharmacy				paradise	132 146
Gails				nidian	220
Gamboge	14	<i>5</i> .		ily, parging	
pills			rass	ily, Puiging	146
Gargarism, astringent			romwel	1	169
common			roundiv		150
emollient			roundp		124
detergent			roundle		139
Garlick				n, gum and	wood 148
oxymel	35			balfam	321
fyrap.	34			elixir	ib.
Garnet Rone		4	•		natical oil 418
Gascoign's powder .	55	-		extract	407
Gelly of barberries	33			refin	410
currants		ь.		tincture,	volatile 310
hartshorn •	28	3	•	wines	301
quinc <b>es</b>	33	š G	uido's		421
Gentian .	14	6 <b>G</b>	ninea p	еррег	201
extract	40	ος G	υm		13, 6 <b>6</b>
Germander	12	13 G	um of	aloes	407
water germai	nder 22	3		moniacum	84
Gilla of vitriol	46	54		ime	· 88
Ginger	24			abic	149
fyrup	i	<i>b</i> .		erry and plu	m tree ib.
Ginleng	14	ļ6		mi	£137
Glass vessels, coating		14		aia <b>cum</b>	148
Glass of antimony		36		deræ	150
cerate				ip <b>er</b>	159
Glass-wort		50	lac	_	- 162
Glauber's falt	48			eka	149
Glutinous substances	. 6	_		gacanth	· ib.
Glytter, anodyne			um-rei		14
astringent		22		aining of	261
for the colic			ypfum	LJ 4	23, 65
common		22 6. H	artihor	H. 4	
domestic emollient		10. N 21	er mol	n calcin <b>ed</b>	130
EIMMILIEUF ,	0	4 L	II n		Want 259

Hartshorn gelly pag	e 283	Hystop fage 154.
oil, falt, spirit	430	effential oil 362
Hartstongue	168	fimple water 378
Hartwort	226	3/5
Hawthorn	229	J. I.
Hedgehystop	148	J. 4.
Hedgemustard	139	Jack-by-the-hedge 78
Hellebore, black	151	Jalap 155
extract	406	extracts 412
tincture	313	refin 410
white	151	tinctures 310, 311
honey	350	Jamaica pepper, se Pimento 201
tincture	321	Japan earth
Hematites	150	decoction 286
Hemlock	125	tincture 311
extract	400	troches 563
Hemp	114	Jalmine 156
water-hemp	139	Isteric decostion 280
hemp-agrimony	ib.	Jerusalem-oak 108
Henbane, black and white	154	Jerusalem-sage 206
Herb-benet	119	Jesuits bark, see Peruvian 195
Herb-mastich	173	Jews-ear 98
Herb-paris	152	Jews-pitch 106
Herb-truelove	ib.	Incineration 56
Herb-twopence	184	Indian leaf 171
Hermodactyls	153	Infernal flone 490
Hiera picra	555	Infusions 268
Hipps	134	Infusion, alkaline 276
conferve	336	antiscorbutic ib.
Hogs-fennel	199	bitter 273
Honey	174	purging ib.
clarified	258	of carduus 260
Egyptian	634	cephalic 276
of fluellin	350	of cinnamon 277
hellebore	ib.	diuretic ib.
of herb mercury	ib.	of linseed 276
of roles	ib.	pectoral ib.
folutive	ib.	paralytic 277
Howey water	386	of Peruvian bark 270
Honey fuckle	116	of rhubarb 275
Hops	170	of fena 274
Horehound .	173	with lemon ib.
Horns .	63	Injection, balfamic 623
Horseradish	207	mercurial ib.
compound water		Inspissation . 53
Horfetail	121	Ipecacoanha 156
Hot feeds	250	tincture 295
Houndstonge	134	wine ib.
Houseleek	224	Irith flate 153
Hungary water	384	Iron 140
Hypociftis	154	preparations of 492
		Iron,

•		_		-
Iron, fa Steel	page	140	Lavender	page 164
Minglass		156	conferve	335
T aices		263	effential oil	362
fcorbutic		264	spirit, simple	385
Jajabes		159	compo	and ib.
Julep, alexipharmac		611	French	232
of ammoniacum		612	Lavender-cotton	72
<b>a</b> ntihysteric	611,	612	Laurel, spurge	164
binding		614	Lazuli stone	165
of camphor		609	Lead	203
. acid		610	burned	500
carminative		611	red	ib.
of chalk		610	white	501
cooling		611	fugar	ib.
<b>c</b> ordia <b>l</b>	611,	612	vinegar	291
diaphoretic		612	tincture	316
divretic		613	ointment	641
fætid		ib.	Leeks	204
hydragogue	_	ib.	Lemnian earth	107
hysteric	611,		Lemon juice	158
of mulk		610	fyrup	343
flomachic		612	Lemon peel	168
July-flowers		119	candied	337
Juniper-berries		159	effential oil	363
effential oil		36 <b>2</b>	Lemon thyme	240
rob		408	Lentils	165 <i>ib</i> .
compound water		394	Lentilk tree	
Juniper gum		159	Leopards-bane	136
wood		ib.	Lettuce, garden wild	162
Ivy		ib.	_	163
			Levigation	255
K.			Ley, foap of iron	428
			of tartar	496
Kermes grains		160	Lily of the valley	427 167
confection	on	349	water lily	186
fyrup		343	white lily	167
mineral		535	Lime	142
Knee-holly		211		2, 62, 112
Knotgrafs		122	Lime-water, fimple	271
- ·			more compounde	
L;			less compounded	ib.
_,			Lime-tree	241
Labdanum		160		603
Lac		162	Liniment anodyne	647
tincture		312	Arcæus's	638
Ladies bed-straw		145	hæmorrhoida	
Ladies mantle		77	<b>mer</b> curial	646
Lard, tried or purified		257	faponaceous	439
Laudanum liquid		314	of three ingre	edients 645
of Sydenhau	M.	298	volatile	440, 645
<i>p</i> .	-		Uu3	Liniment,

Tinimana milia			
Liniment, white Linfeed	page 645	Mallow pag	e 17 t
infusion	169	conferve	335
	276	vervain-mallow	77
expressed oil	266	Mandrake	172
Liquid amber	169	Manna	ib.
Liquorice	147	lohoch	603
extract	405	Marble 2	2, 62
Litharge	169	Marigold	111
vinegar of	291	Marjoram :	171
Liver of antimony	527	estential oil	363
of fulphur	486	Marjoram, wild	190
Liverwort	166	essential oil	364
ashcoloured,	ground ib.	Marmalade of quinces	338
noble	152	Marshmallow	8 z
Locatelli's balfam	589	ointment	635
Lockyer's pill	555	fyrup	340
Logwood	167	Marshtrefoil	342
decoction	285	Masterwort ·	156
extract	496, 413	Mastich	173
Lohochs	603	wood	165
Lohoch, balfamic	604	herb	173
acidulous	ib.	Materia medica	59
common	603	Maudlin	76
of linfeed	ib.	Mayweed	130
of manna	ib.	Meadowsweet	248
faponaceous	ib.	Measures	36
of spermaceti		Mechoacan	•
of flarch	603	Medicinal stone	174
Lotion, saponaceous	439	Medlar	543
Lovage	165	Melilot	177
Lozenges, see Troches	560	plaster	175 628
Lunar caustic		Melons	
pills	490	Mentrua	175
Lugwort, spotted	492 206	Mercury	39
Lupine		alkalized	91
Lutes	170	calcined	508
	. 34	calx	511
M.		coralline	512
441.		corrosive precipitate	513
Mace		fublimate	
essential oil	170	cinnabar	2-2
Madder	363	dulcified fublimate	509
Maghtery of benzoine	210	dulcined nutringle	518
of bismuth	327	dulcified precipitat	-
of binners	541	emetic, yellow	52 <b>3</b>
Magnelia alba	506	ethiops	508
Maidenhair, Canada	22, 469	panacea, red	514
	345	\$ white	520
English	24 L	precipitate, brown	522
true	75	green	ib.
Malaca bean	345.	red	512
	. 86	fweet	522
		<b>3</b>	Mer-

	_		_
Nutmegs, essential oil page	363	Oil of mace, exp. fage	185
expressed oil	185	of marjoram, eff.	<b>363</b>
spirituous water	396	of mint, eff.	ib.
•		of mucilages, decoct.	331
О.		of mustard seed, exp.	266
		of Neroli, off.	97
Oak	207	of nutmegs, eff.	185
Oak of Jerusalem	108	of nutmegs, exp.	363
Oats	97	odoriferous, infus.	332
			186
Odoriferous balfam	186	olive, exp.	
· · · · · · · · · · · · · · · · · · ·	590	of orange-flowers, eff.	97
effence	328	of orange-peel, eff.	266
oil	332	of origanum, eff.	364
troches	566	palm	191
water	386	of pennyroyal, eff.	364
Oils, by infusion and coction	330	of peppermint, eff.	363
expressed 10,	265	of rhodium, esc.	364
empyreumatic	418	rock.	198
rechified	421	of rolemary, eff.	364
essential	353	of rue, off.	ib,
concrete	12	of faint John's wort, infus.	43 I
Oil of almonds, expressed	266	of fassafras, eff.	365
of amber, empyreumatic	48 t	of favin, eff.	364
animal, empyreum. red.	421	of foot, emp.	434
of annifeed, essential	360		
	•		
of Barbadoes tar, emp.	419	of wax, emp.	420
of bayberries, eff.	362	of wormwood, eff.	359
of bays, exp.	164	of sulphur, so called	444
of box, emp.	418	of tartar per deliquium	427
of bricks, emp.	419	of vitriol	443
camphorated	332	Oily draught	607
of caraway feeds, eff.	360	emulfion .	ib.
of chamomile, eff.	361	purging grain	226
of chamomile, decott.	330	Ointment of Arczus	638
of cinnamon, eff.	361	basilicum, black	636
of cloves, eff.	360	green	ib.
of Copaiba compound, emp	-	yellow	ib.
of cummin feeds, eff.	361	for blifters	644
of dill feed, eff.	360	blue, milder	627
of elder, decott.	33 I	ftronger	637. ib.
f fennel seeds, eff.	361	of calamine	646
	-	digestive	
preen, decod.	331		647
of guaiacum, emp.	418	drying	637
of hartshorn, emp.	430	Egyptian	034
of hytrop, eff.	362	of elder	041
of Jamaion pepper, ef.	364	emollient	638
of juniper berries, eff.	362	epilpattic	044
of lavender, eff.	ib.	for the eyes	640
of lemon peel, eff. 266	, 303	of four ingredients	636
of linfeed, exp.	26 <b>6</b> .	green	645
of mace, eff.	363	of gum elemi	638
· <del>-</del>			ment
		· •	

		_		003
Ointment for t	he itch <i>page</i> 635,	64z	Oxymel of garlick	page 35 t
	arfhmallows	ib.	pectoral	遨.
merc	curial	639	fimple	352
of m	ercury precipitat	e ib.	of fquills	351
nerv	-	ib.	Oystershells prepared	255
nutr	itu <b>m</b>	640		
para		646	<b>P.</b> .	
	atum	642		
of p	ompholi <b>x</b>	638	Palm oil	191
fatu	rnine	641	Palmer's powder	4 <b>6</b> 3
fimp		ib.	Palfy drops	386
		642	Panacea of antimony	< 35
of t		640	of mercury, red	514
of th	hree ingredients	642	wh	ite 520
of ti	atty	643	nitrous	463
	with campho	r ib.	Panic	192
<b>Vern</b>	nifuge	644	Paradise grain	148
whit		635	Pareira brava	193
	with camphor		Pariley, common	199
yelle	OW .	637	fimple	water 379
Olibanum		187	Macedonian	199
Olives		186	bastard stone	. 85
Oneberry		152	Parineps	194
Onions		122	Peach	195
Opium			Pearls	172
extract	by long digestion		prepared	255
Araine		<b>261</b>	Pearl afhes	126
tinctur	'es	314	Pearl-barley	15
pills		574	Peas	. 202
Opobaliamun	3.	189	Pellitory of Spain	207
Opodeldoc		439	of the wall	194
Opopanax		190	bastard	206
Orach		97	Pennyroyal	ib.
Oranges, Sev	ille	ib.	essential oil	364
	rasio <b>a</b>	98	fimple water	
Orange peel		97	fpirituous wa	iter 397
	candied	337	Peony	191
	conferve	335	compound water	39 <b>6</b>
	essential oil	266	Pepper	200, 201
	fyrup	34 I	Jamaica, see Pim	
1	imple water	375	Pepperwort	• 165
	spirituous water		Peruvian bark	195
Orchis		221	decoction	286
Origanum		190	electaries	600
	ential oil	364	extracts .	406, 413
Orpiment		93	refin	411
Orpine		1 30	tincture	308, 309
Orris root	•	158	' wine	300
Ofmund-roya	II	142	infusion infusion	270
Oftencolla		191	Petroleum Philonium	198
Oxeye daily		102	Philonium	596
			•	Pile-

	ge 125	Plaster, anodyne discutient	. 633
Pills	567	antihysteric	625
aloeti <b>c</b>	568	bliftering	632
<b>a</b> romati <b>c</b>	567	cephalic	625
chalybeate	570	common	626
cochiæ	569	common sticking	627
of coloquintida, simple		common with gums	ib.
with alo	es 569	common with mercur	y 629
common	. 576	of cummin	628
deobstruent	570	defenfive	ib.
purgative	ib.	diachylon	626
dyfenteric	577	drawing	625
ethiopic .	57.3	epispastic	632
fœtid	571	with gums	627
gamboge	572	of hemlock	626
gummole	571	melilot	628
of jalap	568	mercurial	629
Lockyer's	535	of mucilages	630
lunar	333 492	Oxycroceum	628
Mathews's		red lead	630
	574	faffron	628
mercurial 572, 57 of olibanum		flicking	627
pacific '	575	foap	- •
pectoral	574	firengthening	631
Plummer's	575	flomach	630
	576		631
Rudius's	408	lappurating volatile	633
Rufus's	576	warm	440
of scammony with alor			633
faponaceous	575	Wax	625
fcillitic	576	Plans Poles mountain	205
of spermaceti	578	Poley-mountain	203
Romachic	576	Polypody	ib.
ftorax	<b>575</b>	Pomatum	642
<b>Arengthening</b>	57,7	Pomegranates	148
of tar	ib.	Pompholyx	204
of two ingredients	569	Poplar buds	ib.
Pimento	201	Poppies, garden	192
essential oil	364	fyrap	344
fimple water	379	wild	193
fpirituous water	396	fyrup	345
mpernel	86	Potash	126
Pine-tree	200	Poterius's antihectic	543
Pinachio nuts	185	Poultice, see Cataplasm	
Pitch	202	Powders	547
Jews pitch	106	Powder of Algeroth	539
Plantane	202	of amber compound	554
extract	, 400	antiepileptic	552
Plasters	624	aromatic	555
Platter adhefive	627	of arum compound	548
of amm, with mercury		bezoardic	551
anodyne	624		

Powder

Powder for the bite of a mad dog  page 541  of bole with opium 549  Queen of the meadows page 2	48
page 541	48
	48
	7-
	28
	12
cathartic faline ib. Quickfilver	91
	-
cephalic 554 Quinces of ceruffe compound 550 gelly	33
	38
of contrayerva comp. 551 Quince feed	43
	34
	82
of crabs-claws comp. 550	
to promote delivery 552	
diaretic 57	
fever 531	
Gascoign's 551 Ragwort	55
	48
	81
	10
of myrrh compound 502 fyrup	44
prince's 514 Rattlesnake root	25
of scammony 553 Realgar	93
ofscordium with opium 556 Reduction	56
without opium 555 Regulus of antimony	31
of fena compound 553 martial	32
sternutatory ib. medicinal	31
	33
flyptic of Colbatch 543 Refins 13,	
funcia of Halvetine and Dofin of slope	107
and blade	20
and the same of th	110
testaceous cerated 556 of jalap	ib.
ac dia	111
of course	110
• • • • • • • • • • • • • • • • • • • •	365
	187
Preserves 336 Revivification	56
33	209
	167
	264
	364 208
Brimer C. in C. C. in	259
Pennal fals (6) from with form	275
70	347
Duff k-11	315
	296
	190
Pumpion 194 Rifingal	93
Purging-flax 160 Rob of elder berries	399
Purging oily grain 226 of juniper berries	408
Purstain 204 Rocket	139
Putrefaction 7, 18 Rock-oil	198
ه مر	lod,

200			
Rod, golden	page 246	Salts, alkaline, fixed	page 423
Rose, damask	209	volatile	7-7
fimple water	380	essential	14. 475
lyrap	346	neutral	458
whey	288	Sal ammoniac	213
Rose, red	209	purified	463
conferve	<b>335</b>	flowers of	
hoey	350	fpirit	434 ified 441
fugar	564	volatile s	775
fyrap tincture	346	Sal gem	
vinegar	275	Salt of amber	215 481
Rolemary	289	purified	482
conferve	210	bitter purging	214
essential oil	335	common or fea	215
fpirit	364 384	fpirit	448
Rolewood	167		ulated 470
Rue	211	dulci	fied 457
conferve	334	diaretic	471
extract	407	of hartshorn	430
effential oil	364	polychreft	467
fimple water	380	prunel	468
Rue, goats	145	purging of Glauber	ib.
Rupturewort	153	of faline wa	aters 214
Rush, sweet	159	of Rochel	474
Ruffia afhes	126	fedarive	480
Ruft of fleel prepared	493	of Seignette	474
Rye	224	of foot	434
	_	of forrel	475
S.	•	of ficel	497
0.00		of tartar	422
Saffron	131	of tin	503
extract	411	of vitriol	464
fyrop tinctu <b>re</b>	343	of wormwood of zinc	425
wine	309		542
Saffron, hastard	<b>2</b> 94	Saltpetre, see Nitre Samphire	183
Sagapenum	118	Sand bath	131
<b>Lage</b>	213 213	Sanicle	33 ·
of Jerusalem	206	Sapphire	64
wood fage	216	Sapphire-coloured water	619
Sago	213	Sarcocol	219
Saint John's wort	154	Sarfaparilla	220
oil by infusion	n 331	Saffafras	ib.
Salep	221	essential oil	365
Saline mixture, antiemeti		Saturnine tincture	316
cathartic	616	ointment	641
. diuretic	617	Satyrion	221
Salts, table of the quanti	ties	Sauce alone	78
. foluble in water	39	Savin_	212
alkaline	8	. extract	407
		· 🛰	Şavir

• / •	: -		
Spikenard	page 181	Squills, pills tage	576
Spirits, acid	443	fyrup	346
alkaline	430	troches	562
vinous	381	vinegar	289
table of their w	eight 37	Starch	
Spirit of amber	48 i	glyster	242 621
aromatic faline	•	lohoch	_
ethereal	441	Stavefacre	603
of hartshorn	455	Steel	231
	430		140
of lavender, fimp		alkalized	495
comp	ound ib.	foluble	ib.
of Mindererus	473	fugared	337
of nitre	447	fulphurated	494
dolcified	456	tartarized	495
of rosemary	384	Crocus	ib.
of faffron	411	flowers	496
faline aromatic	441	ley	ib.
of fal ammoniac	434	rust prepared	493
with quick	-lime 435	falt	496
dulcified	441	tinctures	312
of scurvy-grass	387	vitriol	497
purg		wine	294
of fea falt	448	Sternutatory powder	553
coagul	ated 470	Stochas	232
dulcifie		Stone-parsley, bastard	85
of foot	434	Stone, infernal	
of fulphur	444	medicinal	490
of turpentine	365	feptic	543 428
of vinegar		Stones, precious	64
of vitriol	- 451	Storax	
dulcifie	443	ftrained	233 260
volatile aromatic	. •,,,	liquid .	
fætid	441	Strawberry bush	232
	ib.	Styptic powder	143
fuccinate	• •		554
of wine, proof	230	tincture	317
	229, 381	water Cublimation	620
camphoi		Sublimation	51
<b>tartarize</b>	· J-J	Sublimate mercury, corrolive	515
Sweenwort	123	dulcified	
Springe	230	Succory	125
burned	259	Sugar	212
Spunk	76	of milk	478
Spurge	241	of rofes	564
Spurge flax	ib.	of lead	501
Spurge-laurel	164	Sugar-cakes, anthelmintic	564
Spurge-olive	177	Sugared mercury	508
Squills	222	Reel	337
a baked	258	Sulphur	234
dried	ib.	balfams .	485
bolus	584	flowers	484
exymel .	351	* <b>R</b> c	487
•	<i>-</i>	Su!	phur
		<b>*</b>	• ′

<b></b>			Tala balka	
Tinctur	e of cantharides page			agr 101
•	of cardamom feeds	307	Tormentil	241
•	of caftor	ib.	Touchwood	76
	of castor compound	308	Tragacanth gum	149
•	cephalic	296 ib.	compound pow	
•	cephalic purging		Treacle, Andromachus's	592
	of cinnamon	308	Edinburgh ,	595
	of copper, volatile		Venice .	592
	fostid	309	Treacle bolus	584
	of guaiacum, volat. of black hellebore	310	vinegar water	291
		3!3	Treacle mustard	391
	of white hellebore	321		240 .
	of jalap	310	Trefoil, sweet marsh	169
	of jalap, compound,	ib.	Troches	242
	of Japan earth		of amber	560
	of ipecacoanha of iron	295	antacid	554
	of lac	312 ib.	antacid laxative	503 564
٠.	of martial flowers	ib.	anthelmintic	ib.
	of melampodium		antimonial	. 565
	of mint	313 269		505 56I
	of myrrh .		cardialgic	
	of myrrh and aloes	313	of Japan earth	563 <i>ib.</i>
	of opium	314 ib.	of myrrh	561
	of Peruvian bark	308	nerve	565
	of rhubarb, spirituou		of nitre	562
• .	Vinons	296	of red lead	561
•	of roles	275	pectoral, black, wi	ite.
	royal	328	&c.	560
	facred	297	purging	565
	of faffron	309	of rhubarb	ib.
•	of falt of tartar	320	of roles	564
	faturnine	316	fialogogue	566
	of fena	ib.	of fquills	562
	of inakeroot	317	ftomachic :	566
	of foot	310	of fulphur	562
	of ficel	312	fweet smelling	566
	flomachic 298,		· white of Rhafi	560
7	flyptic 290,	317	Truelove, herb	152
2	fudorific	319	Turmeric	133
J.	of falphur	ib.	Turneps	208
٠.	of sulphur, volatile	487	Turner's cerate	646
	thebaic	298	Turpentines	238
	h balsam of Tolu	320	balfam	419
	for the tooth ache	329	bolus	584
	of valerian	321	glyster	622
	of veratrum	ib.	oil and spirit	
Toads		100	Turpeth mineral	523
Toadfla	<b>x</b>	168	root	243
_			7	
Tobacc	0	182	Tatty	ib,
		182 126	Latty Brebared	- 1
	o o-pipe clay		Tatty prepared	256 Tutty

Хx

Water,

Water, nutmeg	ge 396	Wine, antimonial pa	ge 293
odoriferous	386	aromatic	299
orange peel, ample	375	bitter	293
orange peel, spirituo		chaly beate	294
parsley	37,9	emetic	293
pennyroyal, fimple	ib.	febrifuge	300
pennyroyal, spirituo		guaiacum	301
peony, compound	396	ipecacoanha	295
peppermint, simple	397	millepedes	ib.
peppermint, spirituo	us 395	<b>faffron</b>	294
Jamaica pepper, samp		fcorbuti <b>c</b>	299
Jamaica pepper, spis		fteel	294
phagedænic	620	viper	295
plague	392	Winter cherries	77
horfe-radifh	397	Winter's bark	248
rofe	380	Wolf's bane -	136
Inc	ib.	wholefom <b>e</b>	90
fapphise-coloured	619	Woods, decoction of	280
favin	380	essence -	329
flyptic	620	Woodbine	169
fulphurated	446	Woodroof	96
tar	270	Woodforrel	170
treacle	391	conferve	334
vitriolic blue	620	Worms, earth-	870
vitriolic camphorat.	ib.	Wormseed	218
valerian, compound	393	Wormwood, common	72
vulnerary	398	Roman and fea	
wormwood	390	conferve	335
Waterflag	147	extract	404
Water-germander, fee Score	lium.	effential oil	359
Wax, white and yellow	122	fixed fak	425
oil	420	compound water	13 3QQ
plaster,	625	•	. 47
Weights	. 36	Υ.	
table of the weights of liqu	ids 37		
Wheat	242	Yagrow	
Whey	162	Tallon	177
alum	288	<b>.</b> Z.	
laxative	ib.	L.	
mustard	ib.		
fcorbutie ,	289	Zarnich	83
Whin, petty	187	Zedoary	248
Wines	245	Zinc	249
aloctic alkaline	292	preparations of	542

### Index Medicamentorum.

A BIES	page 71	Althæa	page 82
Abies canadenfis	238	Alumen	ib.
A brotanum	7,2	uftum	464
Abfinthium	ib.	Amaradulcis	136
Acacia	73	Ambragrisea	83
Acacia germanica	205, 400	Ammi	ib.
Acanthus	7,4	Ammoniacum gummi	. 84
Acetofa	ib.	parificatum	264 84
Acetocella, vide Lujula	170	Amomum verum	85
Acetum	. 74	vulgare	187
distillatum	451	Amurca	85
lithargyrites	291	Amygdalæ Amylum	242
prophylacticum	290 280	Anacardium	86
rofaceum feilliticum	ib.	Anagallis	ib.
therizeale		Anchusa	87
Aconitum falutiferum	291 90	Anethum	· ib.
Acorus verus	111	Angelica	ib.
vulgaris	147	Anguilla	88
Adianthum	7.5	Anguria	127-
Ærugo	ib.	Anime	88
præparata	255	Anifum	ib.
Æs ustum	498	Anonis	187
Æthiops antimonialis	545	Anserina	91
martialis	494	Anthora	90
mineralis	509	Anuhecticum Poterii	-543
Agallochum	75	Antimonium	89
Agaricus	ib.	catharticúm	537
Ageratum	¢. 76	diaphoreticu	
Agnus castus	77	lotu	
Agresta	247		ratum ib.
Agrimonia	77 ib.	præparatum	255
Alcea	_	Aparine	. ib.
Alchimilla	ib.	Apes	
Alcis ungula	ib.	Apium hortense	199
Alcohol	381	palustre	· 70 284
Alkekengi	77	Apozema aperiens	•
Alliaria	78 <i>ib</i> .	Aqua abfinthii alexeteria fimplex	390
Allium		fpirituosa	374 391
Alnus	. 79	fpirituofa cum	aceto ib.
Aloc	80 262	aluminosa	619
purificata		anethi fem.	374
Alces lignum Alfine	.75 81	angelicz	375
Winge	<b>3.</b>	X x 2	Aqua.
		<b>*-</b>	

676	. I 1	N	D	E	X.		
Aqua	Anhaltina pa	ge	38 <b>8</b>	Aqua	piperis Jam	aic. spirituos	a 396
•	anifi composita	_	392	-	pulegii sim		379
	aromatica		· <del>3</del> 95			ોદાલાં	397
	artemifæ		375	÷	Rabelliana		455
	aurantiorum corticum	7		•	reginæ Hu		384
	fimplex	3	375			makenarum	380
	fpirit <b>ao</b> f		393 <sup>°</sup>		rutæ		ib.
	bryonæ composita		ib.		<b>fabinæ</b>		ib.
	calcis fimplex		271		fambuci flo	rum	ib.
	composita		272		<b>fapphirina</b>		619
	magis composit	a	ib.		Ayptica		620
	minus composit		ib.		fulphurata		446
	cardamomi		393		theriacalis		396
•	cardui		375		valerianæ (	compofita	393
	caroi feminum				vitriolica c	erulea	620
	castorei		39 <b>4</b>			amphorata	ib.
	catapultarum		375 398		vulneraria	am provide	_
	cerasorum nigrorum		376	Aani	ls alba		398
	chamæmeli			Aqui			529
	cinnamoni fimplex		377 ib.		icum gumn	.;	90
					earum telæ	14	149
	fpirituola		394		a bovis		90
	epidemia fœniculi		392		num coralli	n	187
			378	Arci		uum	514
	fortis		449			•	101
	composita:		451		fia hybernic		153
	duplex		449		ntina		9,1
	fimplex		ib.		ntum		ib.
	purificata		450	Arge	ntum vivun		ib.
	hordeata		282			catum	507
	hyffopi		378		lla alba		1 26
	juniperi composita		394		olochiæ		92
	liberans		272	Arni	_		136
	meliffæ		.378		nicuma		93
	composita		383	Arte	_		94
	menthæ piperitidis	Ł	270	Arth	anit <b>a</b>	•	ib.
	fimplex		379	Arun	a		ib.
	ſpirituoſa		395	Afafe	œtid <b>a</b>		95
	menthæ vulgaris ?		270		purifi	cata	261
	simplex \$		<i>379</i>	Asar	um		95
	fpirituo <b>fa</b>		395	Afcle	epias		244
	mirabilis -		ib.	Aſpa	lathus		167
•	naphæ		97		ragus		96
	nephritica		396	Afpe	rula		ib.
	nucis moschatæ		ib.		altum		106
•	ophthalmica		620	Afpl	enium.		97
	odorifera		386	Aftac		•	114
	pæoniæ composita		396	After			138
	petroselini		379	Aftra	• -		156
	phagædenica		620	Atri	_		97
•	picea		270	Aver			ib.
	piperis Jamaicensis		379		ntia curafla	ventia	98
			J17				antia
						4241	

I,	N	D	E X.	677
Aurantia hispalentia	page	97	Betonica	page 103
Auricula judz	- •	98	Betonica Pauli	244
muris			Betula	104
A uripigmentum	93,	98	Bezoar lapis	ib.
Aurom		ib.	præparatus	255
fulminans	4	.89	Bezoardicum joviale	544
. mulivum		06	minerale	540
potabile		.89	Bismuthum	105
Axungia		98	Bistorta	·ib.
- curata	.3	57	Bitumen	ZI
· 10			Judaicum Polya armay a fa	106
В,			Bolus armena, &c.	ib.
Balaustia		~~	alexipharmacus ex alumine	579
Balfamita		99 ib.	e camphora	580 ib.
Balfamum anodynum		21	e caftoreo	ib.
Bateanum		40	catharticus	mib.
cephalicum		90		curio 581
Copaiba	_	99	diaphoreticus	ib.
commendatoris		22	diureticus	: <i>ib</i> .
Canadense		39	dyfentericus	582
Gileadense		89	emmenagogus	, ib.
guaiacinom		21	febrifagus a	in ib.
Guidonis		21	hystericus	
Judaicum	1	86	iliacus	583
Loçatelli	5	89	<b>m</b> ercurialis	10
e Mecha	· ' i	89	emetic	ив 🔐 ја.
odorifer <b>em</b>	9	90	pectoralis	jb.
Peruvianum		QO	rhei cum mercurio	
Perficum	· · 3	22	rheumaticus	. 12 'ib.
faponaceum		139	fcilliticus .	:584
fuccini		181	fudorificus	ib.
fulphuris		185	terebinthinatus	: <b>ib.</b>
Syriacum	•	89	~ 1	ib.
terebinthinæ		119	Bombycis folliculi	., 226
Tolutanum		101		107
traumaticum viride			Borago Borax	山坡
universale, V.			Botrys	10 14.
guent, fatur			Brafficæ	)…108 <i>ib.</i>
vitæ		120	Britannica	v. <b>163</b>
Barba cappe		48		5.2005
Bardana		•	Bruscus	.:211
Basilicum unguentum		126		801
Bdellium			Bufo	LÓO
Becabunga		02	Ruglossum	rib.
Belladonna n	: :::::::::::::::::::::::::::::::::::::	127	Bugula	44
Bellis		102	Bursa pafforis	110
Benzoinum :.	٠٠ تار ټ		Butua, V. Pareira.	193
Berberis	1,0		Butyrum antimonii.	538
Beta	• : •	-14	Buxus	. 110
			Xx3	

.

C.		Cataplasma ischiadicum page 650 maturana 640
C.		- 77
0		7
Cacao pag	il 10 ib.	fuppurans 649
Cadmia, V. Calaminaris		Catechu 563
Calambac	167	Cauda equina 121
Calaminaris lapis	110	Causticum antimoniale 538
præparatus	255	commune fortius 429
Calaminthæ	1 1 1	mitius ib.
Calamus aromaticus	ib.	lunare 490
Calendula seu caltha	ib.	potentiale 428
Calomelas	519	Centinodium 122
Calx antimonii	529	Centaurium ib.
jovis	503	Cepa ib.
mercurii	512	Cera ib.
· viva	112	Cerasa ib.
Camphora	113	Ceratum album
Cancrorum chelæ	114	citrinom ib.
. præp <b>aratæ</b>	255	epuloticum ib.
Cancrorum oculi	114	mercuriale ib.
præparati	255	Cerevisia amara / 302
Cannabis	114	aperièns ib.
Canella alba	ib.	Butleri ib.
Cantharides	115	cephalica ib.
Capillus veneris	75	diuretica ib.
Capparis	116	scorbutica 303
Caprifolium	ib.	Coruffa 123, 501
Capficum	201	antimonii 530
Carabe, V. Soccinum	234	Cervus 130
Caranna	116	Ceterach 123
Cardamomum	ib.	Chacarilla 137
Cardiaca	117	Chærefolium 123
Carduus benedictus	ib.	Chalcitis 594
Caricæ	1 18	Chalybs 140
Carlina	ib.	cam falphure 494
Carpobalfamum	ib.	tartarizatus 495
Carthamos	ib.	rubigo præparata 402
Carui	119	Chamæcypariffus 123
Caryocostinum	588	Chamadrys ib.
Caryophylla	119	Chamæleon 118
Caryophillata	B.	Chamæmelum 124
Cascarilla	137	fætidum: 130
Cafia fistularis	120	Chamæpitys 124
lignea	121	Cheiri ib.
Castoreum	· 18.	Chelæ cancrorum
Cafumunar	ib.	præparatæ 255
Cataplasma aromaticum	648	Chelidonium 124
c t camphoratum	650	minus 128
e cymino .	648	Chenopodium 57, 105
difcorlene	ib.	Linermes 160
• • • • • • • • • • • • • • • • • • •	650	China radix 125
o` .	•	China

	_	<del>-</del> •	~/7
China chinz, V. Pera-	100		125
vianus cortex } page	195	germanorum	249
Cicer	124	Copal-	129
Cichoreum	ib.	Corallina	ib.
Cicla	103	Corallium	ib.
Cicuta	125	, præparatum	255
Cimolia alba	126	Coriandrum	129
purparafcens	ib.	Cornu cervi	130
Cineres clavellati	160	calcinatum	-
Ruffici	126	Cornus	259
			130
Cionabaris antimonii	539	Cortex Peruvianus	195
factitia	509	Coffus	430
Dativa	126	Collus hortorum	99
Çinnamomum	127	Cotula	130
Citrea mala	ib.	Courbaril	88
Citrullus .	<i>ib</i> .	Craffula	130
Cnigus, V. Carthamus	118	Cremor tartari	474
Cnidia grana	24 I	Creta	130
Coagulum aluminofum	650	præparata	255
Cobaltum	93	Crithmam	131
Coccinella	127	Crocus	ib.
Cochlece, free limaces	170	antimonii	527
Cochlearia	128	lotus	528
Coffee	ib.	nition	527
Colcothar	443	medicinalis	
Collyriam	620	martis aperiens	<b>J</b>
Colocynthis	128	aftringens	495 ib.
Colophonia	420	metallorum	-
Condita :		Crystalli tartari	527
Confectio alkermes	<b>5</b> 36	veneris	473
	349		498
Archigenis cardiaca 414.	591	Grystallus lapis Cubebæ	64
	594-		131
Dameeratis	591	Cucumis agrestis	132
Japonica	589	hortenfis	131
Paulina	291	Cucurbita	132
Raleighans	414	Cuminum	134
Conferva abfinthii	335	Cunila babula	190
cochleariz	334	. fativa	221
corticum aurantiorum	<b>33</b> 5	Cupressus	132
cynolbati	336	Cupram	ib.
lavendulæ	335	· Curcuma	133
. lajalæ	334	Cuscuta .	ib.
malvæ	335	Cyanus	ib.
menthæ	334	Cyclamen	94
pruņorum filvestrium	336	Cydonia	133
rorifmarini	335	Cyminum	134
rofarum .	ib.	Cynosbatos	ib.
rutæ .	334	Cyperus	ib.
Confolida major	129	Cypressus	138
media	100	D.	- 3 -
minima	102	Dadyli	774
	- 44		134 aucus
•		, L	-u-u

### I N DE EIX

Daucus	page	135	Electarium	alterans page	59#
Decoctum		278		antiepilepticum	ib.
· . t	album compositum		: •	aromaticum	598
	antifebrile	284		e baccis lauri .	586
	antihecticum	283	:	balfamicum	598
•	aftringens	285	•	caryocoltinum	5.88
	bardanæ	ib.		e cafia	586
	campechense	ib.		chalybeatum:	598
	commune,	287	•	deobstruens.	599
4	pro clyftere .	ib.	•	ad dysentericos	586
	cretaceum	279		ad gonorrhæam	599
	diareticum	285	.:	e guaiaco	ib.
•	febrifugum	284		hæmorrhoidale	602
	Japonicum	279	÷.	ex helleboro ?	
•	ad ictericos	280		nigro 🖇	59 <del>9</del>
	lignorum	16.	4	incrassans	600
	ad nephriticos	. ibs	€14 .	lenitivum	587
	nitroſum	28 I.	11.	ad nephriticos	600
	pectorale .	ib.		paralyticum	ib.
	Peruvianum	286	; .	pectorale	588
	fenekæ .	دفاه		Peruvianum	600
• •	ferpenteria com- ?	-	c <del>.</del> •	purgans acidum	· 601
	politum	281	74.	faponaceum	ib.
	tamarindorum ?			e scammonio	588
	cum fena	282	· ·	e scordio	ib.
•	terræ Japonica	286		fiftens	602
7 1.	vulnerarium	284		¢ fulphure	ib
Dens leon		135	Èlemi	, tarpaare	
Diacassia		587			137
Diacodion		•	Eleutheria		90
Diambra		354 5 <b>68</b>	Elichryfum		137
Diapensia			Elixir aloes	• •	232
Diaromato		217		aeicum	323
Diascordiu		555		cinum	324
		588		se combolitum	321
Dictamnus	intui baisis 1	554		soci <b>cum</b>	326
Dictamuu	creticus	143			324
Digitalis	cicticus	135	•		
Doronicun		ib.	Prop	rietatis	323
Draconis (	i anamia	136		Helmontii	- 7
	•	217	٠,	Paracelfi .	323
Dracuncul		136	· faa	_vitriolicum	
Dulcamara	,	ib.	facru		326
•	19		. faluti	-	316.
	<b>E.</b>			ichicum :	305
Ebulus	•		Vitrio	di	325
	W Yahash	136.		acidum	324
Elaterium	V. Lohoch.		•	dulce.	326
Elatine	. 132,	402	1	Mynsichti	225
		137,		Vig <b>ani</b>	326
Electarium		597	uteri		ib.
	alexeterium	ib	Emplastrum	adhæ <b>sivom</b> .	627
				Emplaft	rum
				-	

7	TA	D.	E	Δ.	160
Emplaftrum ex ammo- ?		_	Enema	terebinthinatum	Mare 622
miaco cum mercurio	page	629	Ens ve		449
anodynum		624		campana	138
anodyno difcuti	ione	692		ma volatile	440.
antihystericum		625	Epithe	vesicatorium	648
attrahens		ib.	Epithy		ib.
calidum			Equife		121
cephalicum		93 <b>3</b> 625	Erigen		_
cereum		ib.	Eruca		139
e cicuta .		626		6 orohue	ib.
• • • • • • • • • • • • • • • • • • • •		622	Elanti	five orobus	190
e cymino		626	Eryngi		139
commune	_	_	Eryfim		ib.
adhæfivu		627 ib.	Enenti	a ambræ	328
cum gum				alexipharmaca	329
cum merc	eano			lignorum	ib.
croteum		628		limonum ·	· 363
defenfivum	•	ib.		martis	496
diachylon		626		Neroli	97
epispasticum 2		63,2		regia	328
compofi	tom	ib.	Efula		241
<b>gam</b> molum		627	Eupato	orium cannabinum	1 139
ad herniam		631		veterum	77
e meliloto		628	Eupho		130
mercuriale		629	Euphra		ib.
e minio		630	Extrac	tum abfinthii	404
e mucilaginibu	15	ib.		catharticum	413
oxycroceum ·		628		centaurii	404
roborans		630		cicutæ	400
e lapone	•	6 <u>3</u> 1	•	chamæmeli	404
flomachicum		ib.		coritcis Peru-	
fuppurans		633		. viani	406, 413
veficatorium		632		croci	411
volatile	• •	440		enulæ campai	næ · 405
Emuliio arabica	•	706		gentianæ	ib.
cum aro '		607		glycyrrhiza	ib.
' camphorata		606		guaiaci	407
'communis		605		· jalapii	412
oleofa		607.		hellebori nigi	i 406
purgans	•	606	•	ligni Campec	henfis ib.
spermatis ceti		608		martis ·	294
Endivia -		138		opi <b>i</b>	415
Enema de amylo		621	•	plantaginis	400
anodynum	•	ib.	•	Ruďii	408
anticolicum.	•	ib.		rutæ	407
astringens		622		fabinæ .	ib.
commune		ib.	•	thebaicum .	261
domesticum		ib.		:	·
emolliens		· ib.		F.	٠.
fætidum	•	ib.	Faba		140
opiatum		621	Farfara	<b>.</b>	243
purgans		622	Ferrun	a	140
A :=					Ficus

Ficus	page 118	Glycyrrhiza	page 147
Filipendula	141	Gramen	148
Filix	142	Grana cnidia	241
Flores benzoini	479	paradifi	148
martjales	496	Granata	ib.
falis ammoniaci	463	Gratiola	ib.
fulphuris	484	Guaiacum	ib.
loti	ib.	Gummi aloes	407
zinci	542	ammoniacum	84
Fæniculum	142	anime .	88
Fœnum-græcum	143	arabicam.	149
Formicæ	ib.	elemi	37
Fotus anodynus	<b>2</b> 87	hederæ	150
aromaticu\$	ib.	juniperi	159
communis	ib.	<b>fenica</b>	149
roborans	288	tragacantha	ib.
Fragaria	143	Guttæ vitæ	328
Frangula	79	Gyplum	23
Fraxinella	143		•
Fraxinus	144	н.	
Fuga dæmonum	154	n.	
Fuligo	144		
Fumaria	ib.	Hæmatites	150
Furfur	242	præparatus .	255
£ 43.42		Halicacabus	.77
<b>G.</b> ,		Haustus diaphoreticus diureticus	617 <i>ib</i> .
Galanga .	144	anod	ynus <i>ib.</i>
Galbanum	ib.	catharticus	616
purificatum	261	falig	
Galega	145	, oleofus	607
Gallæ	ib.	Hedera	150
Gallium	ib.	Helenium	138
Gambogia	ib.	Helleborus albus	151
Gargarisma astringens	621	Helleborus niger	ib.
commune	ib.	Helxine	194
emolliens	ib.	Hepar antimonii	527
detergens	ib.	fulphuris	486
Gas splphuris	446	Hepatica	152
Gelatina berberum	338	Herbo Britannica	163
cornu cervi	283	Paris	152
cydoniorum	338	pedicularia	232
ribefiorum	ib.	Hermodactylus	153
Genista	146	Herniaria	įЬ.
Gentiana	ib.	Hesperis	78
Geranium	ib.	Hiera picra	55 <b>5</b>
Gilla vitrioli	464	Hippoglossum	15.3
Ginfeng	i46	Hippolelinum	ib.
Gith fee nigella	. 188	Hirundinaria	744
Gladiolus	147	Hordeum	153
Glaftum	ib.	•	
		1	Horminum

Lappa

Yanga malan		Lohoch ex amylo	page 603
Lappa major pag Laudanum liquidum	314	balfamicum	604
Sydenhami	298	commune	603
Lavendula	164	de lino	ib.
Laureola mas	ib.	de manna	ib.
femina	177	pectorale	ib.
Lauras	164	faponaceum	603
Lazuli lapis	165	de spermate ceti	604
Lens	ib.	Lotio saponacea	439
Lentiscus	ib.	Lotus urbana	169
Lepidium	ib.	Lujula	170
Legcanthemum	124	Lumbrici	ib.
Leucoium	ib.	Lupinus .	ib.
Levisticum	165	Lupulus	ib.
Lichen	166	Lycoperdon	ib.
cinereus terrestris	ib.	_,-,-	
Lignum aloes	75	M.	
aspalathus	167		
Campechense	ib.	Macis	170
guaiacum	148	Magisterium benzoini	3 <b>2</b> 7
juniperus	150	bismuthi	541
nephriticum	182	ftanni	506
rhodium	167	Magistrantia	156
fantalum	217	Magnefia alba	469
Ligusticum	165	Majorana	171
Lilium album	167	Mala aurantia	97
convallium	ib.	curaflavenfia	98
Limaces	170	hortenfia	17,1
Limones	168	fylvestria	ib.
Linaria	ib.	Malabathrum	ib.
Linctus acidulus	604	Malicorium	148
folutivus	ib.	Malvæ	171
Lingua cervina	1 <b>6</b> 8	Malva verbanaça	77
Linimentum album	645	Mandragora	17,2
. anodynum	647	Manna	ib.
Arcæi	638	Manna thuris	187
hæmorrhoid <b>ale</b>	647	Margaritæ	172
. mercuriale	646	præparatæ	255
faponaceum	439	Marrubium	173
tripharmacum	645	Mars faccharatus	. 337
volatile 440	, 645	alcalizatus	495
Linum catharticum:.:	168	folubilis	ib.
vulgare	169	folpheratus	494
Liquamen salis tartari	427	Martis limatura præparat	
Liquidambra	169	Marum Mastiche	173 <i>ib</i> .
Liquor anodynus mineralis	455.		
Lithargyrus	169	Matricaria Mechanganna	17 <b>4</b> <i>ib</i> .
Lithospermum .	ib.	Mechoacanna	
Lixivium martis	496	Meconium Mal	344
foponariu <b>m</b>	428	Mel Ægyptiacum	624
' tartari	427	2DEJ Priacum	634
			Mel

torrefacta

Nymphæa

259

186 O. Ochra

614

ib.

Mistura alexiteria

antidysenterica

## INDEX.

		N	D	E	X.	
686	. 1	14	-	-	mucilaginibus	page 33 %
330	0.		OI			496
				11	ucis moschatæ	185
		page 18	36	1	doriferum	332
Ochra	the same of the sa	1	<i>b</i> .		olivarum	186
Ocimum		1	14		1	191
Oculi can	Crorum		55	-	petrolei Barbad	ensis 41%
Pra	parati	_	.66		fambucinum	33
Oleum ar	nygdalinum		21		finapios	48
31	Illiaio	4	118		fuccini	
bi	amphoratum		332		C 2 1	44
C	hamæmeli		330		tartari per dell	quium 42
C	ornu cervi		ib.		terebinthinæ	
		2 12 1	420		viride	33
	ceræ Copaivæ compo	litum	ib.		vitrioli	42 x3
			421	Oliba		1
	essentiale absint	hii	350	Oliva		2
	anethi		360 ib.	Omn	hacium	1
	anifi		266	Onn	n15	1
	aurantiorun	n	360	Onh	iogloffum	
			ib.	Opi	in m	2
	carvophylli	orum	361	1	colatum	
	chamæmei	7	ib.	One	balfamum	3
	cinnamom	i	ib.	Opt		
	cymini		ib		purificat	2
	fœniculi		36		chis	
	hyflopi		ib	. Or	iganum ,	100
	innipert		ib	Or	obus	-
	lavendul3	2		. 01	yza	
	Jauri bac	carum	36	. 0	Rencolla.	
	limonum			6 0	Greatum tella	
100	macis			b. O	vorum teltæ	
	2	næ .			170	-2
	menthæ	bibeir	[1019	ib. C	constha Car	eni
	vulgaris			07	10.5	2713
	- acrals			62 (	Dxycroceum	ALC: U
	nucis n	olchat	35	:1 4	Devilanathum	
					Owner ex allie	
	piperis	Jamas	Centra	ib.	DECLOS	
	pulegii			ib.	feillitic	ons
	rhodil			ib.	fimplex	4
	rorifm	arini		ib.	Oxys	1
	rutæ			ib.	200	, d
	fabina	e		365		P. (
	fastafr	35	-	ib.		
	terret	ointhin	Chin	434		C
	fuliginis			418	n-onia	. 1
	qualact			331	Pæonia P. limpista	(
	hyperic	1		419	Palimpiffa	1
100	lateriii!	1719		266	Palma	monii 🦂
11	lateritic lini fen macis	1719	411		Danacea antil	ilaom

000	. 1	, т	, r, v.	
Pulvis antilyssus	page	548	Realgar	page 93
antispasmodicu	s	558	Regina prati	248
ari compolitus		548	Regulus antimonii	531
arthriticus ama	rus	556	martialis	532
bezoardicus		551	medicinalis	531
e bolo composi	tus 🕽 🗀	541	ftellatus	533
cum opi			Refina aloes	407
fine opio	)	549	flava	365
carminativus		557	guaiaci	410
Carthusianus		535	jalapii	ib.
catharticus falis	านร	557	nigra	420
cephalicus	C	554	Peruviani corticis	411
e cerussa compo		550	fcammonii	410
e chelis cancro		ib.	Rhabarbarum	208
contrayervæ con	происия		torrefactum	٠, -
cornachini		553	Rhamnus catharticus	229
diaromaton diafennæ	•	555	Rhaponticum	209
		553	vulgare	122 208
diatragacanthi diureticus		555	Rheum, see'rhabararum Rhodium	167
ad epilepticos		557	Rhus .	
de gutteta		552 ib.	Ribefia	236 209
hieræ picræ		554	Risagon, V. Casumunar	121
e myrrha comp	ofitus	55 <b>2</b>	Rifigal	93
ad partum		ib.	Rob baccarum?	
præcipitans		514	juniperi }	408
principis		ib.	fambuci	399
roborans		558	Rofa folis	414
e scammonio co	mpolitus	553	rubra	209
e sena composit	us	ib.	damascena	ib.
<b>Rernutatorius</b>		ib.	Rofmarinus	210
succini composi	tus	554	Rubia	ib.
stypti <b>cu</b> s		ib.	Rubigo chalybis	493
temperans		558	Rubus idæus	210
testaceous		550.	vulgaris	211
cerat		556	Rufcus	ib.
e tragacantha co	mpontu		Ruta	ib.
vermifugus	,	558	<b>S.</b>	
Pyrethrum		207		
_			Sabina ·	212
Q.			Saccharum	ib.
0			lactis	478
Quercus Quinquefolium		207	rofaceum	564
Quinquenervia		194	faturni	501
Quinquenervia		202	Sagapenum	213
, <b>n</b>			Sago	ib.
` R.			Sal abfinthii	425
Ranarum fperma		40=	acetofæ	475
Raphanus rusticanus		207 ib.	alcalinus falis marini ammoniacus	427
Rapum		208	purificatus	213
	•	<b>700</b> .	Larinceras	463
			••	- Sal

0.1		. 1	
Sal ammon, volatilis	page 434		k 12 <b>2</b>
- catharticus ?	214	Scolopendrium	168
amarus S	•	Scordium	225
Glauberi	468	Scorodotis	216
Chalybis	497	Scorzonera	223
communis	215	Scrophularia	2
cornu cervi	430	aquatica major	<b>}</b> 224
diuretícus	471	vulgaris	223
gemmæ	215	Sebesten	224
jovis	503	Secale	ib.
· marinus	215	Sedum	ib.
martis	496	Selenites	23
mirabilis	468	Sempervivum	224
polychreftus	467	Sena	ib.
prunellæ	468	Seneka	225
rupellenfis	475	Senecio	
fedativus	480	Senica gum	139
fuccini	481	Septinervia	149
	•	Sericum	202
tartari	423		226
vitrioli	464	Serpentaria virginiana	ib.
zinci	542	Serpyllum	ib.
Salep	221	citratum	240
Salvia	216	Serum aluminosum	288
Sambucus	ib.	fcorbuticum.	289
Sanguis draconis	217	finapinum	288
Sandaracha .	93	folutivum	ib.
Sanicula	217	Sefamum	226
Santalum album	ib.	Sefeli	ib.
citrinum	218	Sevum ovillum curatum	257
rubrum	ib.	Sicula	103
Santonicum	ib.	Sief album	560
Sapo albus	ib.	Sigillum Salmonis	227
amygdalinus	436		ib.
cum oleo olivarum		Sinapi	ib.
mollis	218	Sinapifmus	649
· niger	ib.	Sifon	85
purificatus	438	Sium aromaticum	ib.
tartarens	ib.	Smyrnium	
Saponaria	219	Soda	153
Sarcocolla	ib.	Solanum lethale	427
	220		227
Sarfaparilla Saffafaa		lignofum	136
Saffafras Samuela	ib.	vulgare	227
Satureia Carrier	221	veficarium	77
Satyrium	ib.	Soldanella	801
Saxifraga	ib.		512
Scabiosa	ib.	Sophia	228
Scammonium	ib.	Sorbus	ib.
Scilla	222	Species aromaticæ	555
cocta	258	e scordio cum opio	55 <b>6</b>
exficcata	ib.	. fine opiq	555
Scincus	223	Sperma ceti	228
Schœnanthus	222	Y y	Spica

600	NT T	DE X.	•
		Succi scorbutici	page 264
Indica	<i>ib.</i> 228	Saccinom	234
Spina alba cervina		præparatum Succus prunorum filv.	256
Spiritus aceti	229 451	Succisa pronorum mv.	400 178
æthereus	455	Sulphur	234
camphoræ	327	præcipitatum	487
cochleariæ	. 387	vivom	484
aureu		antimonii aurat	
cornu cervi	430	præcipitatum	533
croci	411	Sumach	236
fuliginis	434	Symphytum	129
lavendulæ	385	Syrupus ex allio	340
· compo	fitus <i>ib</i> .	ex althæ	ib.
Mindereri	473	aurantiorum cortice	ım 341
nitri dulcis	456	balfamicus	342
Glaub		caryophillorum rub	
rorifmarini	384	croci	34,3
falinus aromaticu		cydoniorum	ib.
· fanis ammoniaci	434	diacodion	344
cum calce dulcis	435	kermefinus limonum	343 <i>ib.</i>
falis marini	441	e meconio	
coagulat	448 us 470	mororum	34 <b>4</b> <i>ib</i> .
dulcis	457	papaveris albi	ib.
Glauber		erratici	345
fuccini	481	paralyfis	346 346
fulpharis	444	pectoralis	345
\$erebinthinæ	365	rofarum folutivus	346
vini camphoratus	327	e rolis ficcis	ib.
rectificatus 2	29, 381	rubi idzi fractus	344
tartarizatus	383	<b>facchari</b>	347
tenuior	230	de iena & rheo	ib.
vitrioli dulcis	453	fcilliticus	346
fortis	443	fimplex	347
teņuis	ib.	e spina cervina	ib.
volatilis aromatic	• • •	violarum	348
causticu	177	zingiberis	349
fœtidus	441	<b>m</b> .	
fuccinat	• • •	<b>T.</b> •	
Spongia ufta	230	Tabella antocida	46.0
•	359	Tabellæ antacidæ laxantes	563
Stannum pulveratum	` 231	anthelminticæ	564 <i>ib</i> .
Staphisagria Staphisagria	503	cardialgicæ	563
Stercora	231 232	rofaceæ	564
Stibium	. 89	Tacamahaca	236
Stoechas	232	Tamarindus	ib.
Styrax calamita	ib.	Tamarifcus	237
colatus	260	'I anacetum	ib.
liquida	233	Tapius barbatus	ib.
Suber	ib.	Taraxacum	135
			Tartarus

Tartarus

## INDEX.

•			
IN	D	EX.	pģi.
Tartarus page	237	Tinctura ipecacoanhæ page	295
emeticus	540	laccæ	312
regeneratus	470	• martis	ib.
folubilis	474	melampodi <b>i</b>	313
vittiolatus	465	menthæ	269
Tegula hybernica	153	myrrhæ	313
Telæ aranearum	90	& aloes	314
Telephium	130	odontalgica	329
Terebinthinæ	238	• <b>o</b> pii	314
Terra japonica	155	regia .	328
foliata tartari	472	shabarbari ?	315
lemnia	107	fpirit <b>uofa</b> ∫	_
figillata	ib.	vinosa	296
Silefiaca	ib.	rofarum	275
Thapfia	239	facra	297
Thea	240	falis tartari	320
antiphthifica	277	falutifera	305
Theriaca	592	faturnina	316
Edinensis	595	ſen <b>z</b>	ib.
Thlaspi	240	ferpentariæ	317
Thus	ib.	flomachica 298,	317
Thymelæa	241	Ryptica	317
Thymus	240	<i>f</i> occini	318
citratus	ib.	fudorific <b>a</b>	319
Tilia	24 t	falphuris	ib.
Tinctura amara	305	volatilis	487
ambræ	328	thebaic <b>a</b>	298
antimonii	319	Tolutana.	320
diaphoretici	320	Valerianæ 🤰	
antiphthifica	316	fimplex }	321 .
aromatica	305	volatilis volatilis	ib.
balfamica	306	veneris volatilis	499
benzoini	327	veratri	321
cantharidum		Tithymalos	241
cardamomi	207 <sup>-</sup>	Tormentilla	ik.
cafforei ·	ib.	Tragacantha	140
composits	308	Trichomanes	241
cephalica	296	Trifolium palustro	242
purgans	ib.	Trissago -	123
cinnamomi	3¢8	Triticum	242
corticis Peruviani	ib.	Trochifci albi	-
composita	309	Rhazis 5	560
volatilis	308	bechici albi	ib.
- annai	309	nigr <b>i</b>	561
Tœtida	ib.	e carabe	554
fuliginis	310	<b>ca</b> rdialgici	563
guaiacina volatilis	ib.	diafulphuris .	562
hellebori nigri	313	de minio	561
jalapii	310	de myrrh <b>a</b>	ib.
comboura .	311	· nervini	565
japonica	ib.	e nitro	562
J#h		Tree	hisci



## 692 I N D E 🛪

<b>-9-</b>			• •	
Trochifci e fo	illa <i>page</i>	562	Unguentum vermifugum page	644
	agogi	ς66	ad vesicatoria	ib.
	nachici '	jb.	viride	637
	veolentes	ib.	Urtica	248
	lphure	562	Uvæ paffæ .	ib.
	rra japonica	563	Valeriana.	243
Turpethum	ira japonica		Veratrum	
	in and la	243	Verbafculum.	321
T(1)	ineral <del>e</del>	523		237
Tuffilago		243	Verbena	244
Tutia		, ib.	Veronica	ib.
bræbara	a	256	aquatica.	102
•	77		Vincetoxicum	244
	υ.		Vinum	243
Ulmaria		0	aloeticum alcalinum	2 <b>92</b>
	TC	248	amarum	293
Unguentum	regyptiacum	634	aromaticum	299
alb		ib.	antimoniale	293
	camphoratum		antiscorbuticum	299
	althæa ·	ib.	chalybeatum	294
Arc		63 <b>8</b>	croceum	ib.
., bai	ilicum flavem	636	emeticum .	293
•	nigrum	ib.	febrifugum	300
	viride	ib.	guaiacinum	301
cita	inom	637	cum helleboro	ib.
cær	uleum fortius	ib.	_	
	mitius	ib.	ipeccaoanæ	295
defi	cćatívum	ib.	mellepedum	ib.
dia	pompholygos	638	fcorbuticum	299
dio	estivum	647	viperinum	295
	olliens	638	Viola	245
	fpafticum		Vipera	ib.
epi	ummi elemi	644	Virga aurea	246
, eg	mida salaminasi	638	Vifcus	ib.
, € 18	pide calaminari		Vitis	ib.
	rcuriale	639	Vitriolum	247
	ercurio præcipits		calcinatum	465
	vinum	jb.	purificatum	464
	ritum	640	martis	497
	thalmicum	ib.	zinci	542
' - par	alyticum	646	Vitrum antimonia	536
ể p	ic <b>e</b>	640	. ceratum	
por	natu <b>m</b>	642	·	5 <b>37</b>
ad	pioram	635	<b>w.</b>	
	aceum	642	Winteranus cortex	248
· fau	buccinum	641	•	-40
	rninum .	ib.	<b>Z.</b>	
	plex	ib.	Zarnich	٥
	lphure	642	Zedoaria	93
ta:	apharmacum	636	Zibethum	248
	parmacum		7inoum	249
	harmacum	642		542
tuu	<i>-</i>	643	Zingiber	249

